




A Review of Discrete Time Linear Systems

- Description of Discrete Time Signals
- A Description of Discrete Time Systems
- Frequency Domain Analysis


Discrete Time Signals



Discrete Time Signals




Discrete Time System



A discrete time system produces an output discrete time signal from an input discrete time signal.

Linear Discrete Time System




$$y[n] + a_1y[n - 1] = b_0x[n] + b_1x[n - 1]$$

Linear Constant Coefficient Difference Equation
LCCDE




Recursions in the form of an LCCDE are the most popular for linear discrete time systems because only multiplications by fixed constants and additions are required.




The linear systems of interest to us have an input/output relationship defined by LCCDE

$$\text{e.g., } y[n] - ay[n - 1] = x[n]$$




The linear systems of interest to us have an input/output relationship defined by LCCDE

$$\text{e.g., } y[n] - ay[n - 1] = x[n]$$




Systems defined by an LCCDE are also time-invariant




The linear systems of interest to us have an input/output relationship defined by LCCDE


$$\text{e.g., } y[n] - ay[n - 1] = x[n]$$



Systems defined by an LCCDE are also time-invariant




The input/output relationship of a linear time-invariant system may also be described using the impulse response and convolution.




The linear systems of interest to us have an input/output relationship defined by LCCDE

$$\text{e.g., } y[n] - ay[n - 1] = x[n]$$





Systems defined by an LCCDE are also time-invariant



The input/output relationship of a linear time-invariant system may also be described using the impulse response and convolution.


$$\text{Kronecker "delta": } \delta[n] = \begin{cases} 1 & n = 0 \\ 0 & n \neq 0 \end{cases}$$






The linear systems of interest to us have an input/output relationship defined by LCCDE


$$\text{e.g., } y[n] - ay[n - 1] = x[n]$$



Systems defined by an LCCDE are also time-invariant




The input/output relationship of a linear time-invariant system may also be described using the impulse response and convolution.



$$y[n] = \sum_{m=-\infty}^{\infty} x[m]h[n-m] = \sum_{m=-\infty}^{\infty} h[m]x[n-m]$$


Convolution Example



$$x[n] = U[n] \quad h[n] = a^n U[n]$$


$$y[n] = \sum_{m=-\infty}^{\infty} x[m]h[n-m] = \sum_{m=-\infty}^{\infty} h[m]x[n-m]$$

Convolution Example




$$y[n] = \sum_{m=-\infty}^{\infty} x[m]h[n-m] = \sum_{m=-\infty}^{\infty} h[m]x[n-m]$$

$$x[n] = U[n] \quad h[n] = a^n U[n]$$




Convolution Example




$$y[n] = \sum_{m=-\infty}^{\infty} x[m]h[n-m] = \sum_{m=-\infty}^{\infty} h[m]x[n-m]$$

$$x[n] = U[n] \quad h[n] = a^n U[n]$$




Convolution Example




$$y[n] = \sum_{m=-\infty}^{\infty} x[m]h[n-m] = \sum_{m=-\infty}^{\infty} h[m]x[n-m]$$

$$x[n] = U[n] \quad h[n] = a^n U[n]$$




Convolution Example




$$y[n] = \sum_{m=-\infty}^{\infty} x[m]h[n-m] = \sum_{m=-\infty}^{\infty} h[m]x[n-m]$$

$$x[n] = U[n] \quad h[n] = a^n U[n]$$




Convolution Example




$$y[n] = \sum_{m=-\infty}^{\infty} x[m]h[n-m] = \sum_{m=-\infty}^{\infty} h[m]x[n-m]$$

$$x[n] = U[n] \quad h[n] = a^n U[n]$$




Convolution Example




$$y[n] = \sum_{m=-\infty}^{\infty} x[m]h[n-m] = \sum_{m=-\infty}^{\infty} h[m]x[n-m]$$

$$x[n] = U[n] \quad h[n] = a^n U[n]$$




Convolution Example




$$y[n] = \sum_{m=-\infty}^{\infty} x[m]h[n-m] = \sum_{m=-\infty}^{\infty} h[m]x[n-m]$$

$$x[n] = U[n] \quad h[n] = a^n U[n]$$




Convolution Example



$$y[n] = \sum_{m=-\infty}^{\infty} x[m]h[n-m] = \sum_{m=-\infty}^{\infty} h[m]x[n-m]$$


$$x[n] = U[n] \quad h[n] = a^n U[n]$$



$$n < 0, \quad y[n] = 0$$


$$n \geq 0, \quad y[n] = \sum_{m=0}^n a^m = \frac{1-a^{n+1}}{1-a}$$

Convolution Example




$$y[n] = \sum_{m=-\infty}^{\infty} x[m]h[n-m] = \sum_{m=-\infty}^{\infty} h[m]x[n-m]$$

$$x[n] = U[n] \quad h[n] = a^n U[n]$$




$$n < 0, \quad y[n] = 0$$

$$n \geq 0, \quad y[n] = \sum_{m=0}^n a^m = \frac{1-a^{n+1}}{1-a}$$



Convolution Example



$$x[n] = U[n] \quad h[n] = a^n U[n]$$

$$y[n] = \sum_{m=-\infty}^{\infty} x[m]h[n-m] = \sum_{m=-\infty}^{\infty} h[m]x[n-m]$$

$$\sum_{n=0}^{\infty} r^n = \frac{1}{1-r}$$

$$\sum_{n=0}^{N-1} r^n = \frac{1-r^N}{1-r}$$

$$\sum_{n=1}^{\infty} r^n = \frac{r}{1-r}$$

$$\sum_{n=0}^{\infty} r^{n-1} = \frac{1}{r(1-r)}$$

$$\sum_{n=0}^{\infty} n r^{n-1} = \frac{1}{(1-r)^2}$$


$$\sum_{n=0}^{\infty} n r^n = \frac{r}{(1-r)^2}$$

geometric sums and their variations are helpful in this discrete time business

$$|r| < 1$$

$$n < 0, \quad y[n] = 0$$

$$n \geq 0, \quad y[n] = \sum_{m=0}^n a^m = \frac{1-a^{n+1}}{1-a}$$



Frequency Domain Analysis

z Transform

$$\mathbf{X}(z) = \sum_{n=-\infty}^{\infty} x[n]z^{-n}$$

$$x[n] = \frac{1}{j2\pi} \oint \mathbf{X}(z)z^{n-1}dz$$

Frequency Domain Analysis


z Transform

$$\mathbf{X}(z) = \sum_{n=-\infty}^{\infty} x[n]z^{-n}$$

$$x[n] = \frac{1}{j2\pi} \oint \mathbf{X}(z)z^{n-1}dz$$

Comments:

- z is a complex variable and $\mathbf{X}(z)$ is a complex-valued function of the complex variable z .



Frequency Domain Analysis

z Transform


$$\mathbf{X}(z) = \sum_{n=-\infty}^{\infty} x[n]z^{-n}$$

$$x[n] = \frac{1}{j2\pi} \oint \mathbf{X}(z) z^{n-1} dz$$

Comments:

- z is a complex variable and $\mathbf{X}(z)$ is a complex-valued function of the complex variable z .
- LCCDEs always produce a ratio of polynomials in z^{-1} :

$$\mathbf{X}(z) = \frac{B(z)}{A(z)} = \frac{b_q z^{-q} + \dots + b_1 z^{-1} + b_0}{a_p z^{-p} + \dots + a_1 z^{-1} + a_0}$$



Frequency Domain Analysis

z Transform


$$\mathbf{X}(z) = \sum_{n=-\infty}^{\infty} x[n]z^{-n}$$

$$x[n] = \frac{1}{j2\pi} \oint \mathbf{X}(z) z^{n-1} dz$$

Comments:

- z is a complex variable and $\mathbf{X}(z)$ is a complex-valued function of the complex variable z .
- LCCDEs always produce a ratio of polynomials in z^{-1} :

$$\mathbf{X}(z) = \frac{B(z)}{A(z)} = \frac{b_q z^{-q} + \dots + b_1 z^{-1} + b_0}{a_p z^{-p} + \dots + a_1 z^{-1} + a_0}$$



Frequency Domain Analysis

z Transform


$$\mathbf{X}(z) = \sum_{n=-\infty}^{\infty} x[n]z^{-n}$$

$$x[n] = \frac{1}{j2\pi} \oint \mathbf{X}(z) z^{n-1} dz$$

Comments:

- z is a complex variable and $\mathbf{X}(z)$ is a complex-valued function of the complex variable z .
- LCCDEs always produce a ratio of polynomials in z^{-1} :

$$\mathbf{X}(z) = \frac{B(z)}{A(z)} = \frac{b_q z^{-q} + \dots + b_1 z^{-1} + b_0}{a_p z^{-p} + \dots + a_1 z^{-1} + a_0}$$



Frequency Domain Analysis

z Transform


$$\mathbf{X}(z) = \sum_{n=-\infty}^{\infty} x[n]z^{-n}$$

$$x[n] = \frac{1}{j2\pi} \oint \mathbf{X}(z)z^{n-1}dz$$

Comments:

- z is a complex variable and $\mathbf{X}(z)$ is a complex-valued function of the complex variable z .
- LCCDEs always produce a ratio of polynomials in z^{-1} :

$$\mathbf{X}(z) = \frac{B(z)}{A(z)} = \frac{b_q z^{-q} + \dots + b_1 z^{-1} + b_0}{a_p z^{-p} + \dots + a_1 z^{-1} + a_0}$$



The z transform sum

$$\mathbf{X}(z) = \sum_{n=-\infty}^{\infty} x[n]z^{-n}$$

only converges for certain values of z . These values are called *the region of convergence* (ROC).

- The ROC cannot contain any poles.
- The ROC for the z transform of a stable (bounded) time-domain signal contains the contour $e^{j\omega}$. This contour is called the *unit circle*.

Frequency Domain Analysis

z Transform

$$\mathbf{X}(z) = \sum_{n=-\infty}^{\infty} x[n]z^{-n}$$



$$x[n] = \frac{1}{j2\pi} \oint \mathbf{X}(z) z^{n-1} dz$$

Comments:

- z is a complex variable and $\mathbf{X}(z)$ is a complex-valued function of the complex variable z .
 - LCCDEs always produce a ratio of polynomials in z^{-1} :

$$\mathbf{X}(z) = \frac{B(z)}{A(z)} = \frac{b_q z^{-q} + \dots + b_1 z^{-1} + b_0}{a_p z^{-p} + \dots + a_1 z^{-1} + a_0}$$

ROC for left-sided signals



Frequency Domain Analysis

z Transform

$$\mathbf{X}(z) = \sum_{n=-\infty}^{\infty} x[n]z^{-n}$$



$$x[n] = \frac{1}{j2\pi} \oint \mathbf{X}(z) z^{n-1} dz$$

Comments:

- z is a complex variable and $\mathbf{X}(z)$ is a complex-valued function of the complex variable z .
- LCCDEs always produce a ratio of polynomials in z^{-1} :

$$\mathbf{X}(z) = \frac{B(z)}{A(z)} = \frac{b_q z^{-q} + \dots + b_1 z^{-1} + b_0}{a_p z^{-p} + \dots + a_1 z^{-1} + a_0}$$

ROC for right-sided signals



Frequency Domain Analysis

z Transform

$$\mathbf{X}(z) = \sum_{n=-\infty}^{\infty} x[n]z^{-n}$$



$$x[n] = \frac{1}{j2\pi} \oint \mathbf{X}(z) z^{n-1} dz$$

Comments:

- z is a complex variable and $\mathbf{X}(z)$ is a complex-valued function of the complex variable z .
- LCCDEs always produce a ratio of polynomials in z^{-1} :

$$\mathbf{X}(z) = \frac{B(z)}{A(z)} = \frac{b_q z^{-q} + \dots + b_1 z^{-1} + b_0}{a_p z^{-p} + \dots + a_1 z^{-1} + a_0}$$

ROC for two-sided signals



Frequency Domain Analysis

z Transform

$$\mathbf{X}(z) = \sum_{n=-\infty}^{\infty} x[n]z^{-n}$$

$$x[n] = \frac{1}{j2\pi} \oint \mathbf{X}(z)z^{n-1}dz$$

Comments:

- z is a complex variable and $\mathbf{X}(z)$ is a complex-valued function of the complex variable z .
- LCCDEs always produce a ratio of polynomials in z^{-1} :

$$\mathbf{X}(z) = \frac{B(z)}{A(z)} = \frac{b_q z^{-q} + \dots + b_1 z^{-1} + b_0}{a_p z^{-p} + \dots + a_1 z^{-1} + a_0}$$

- Because z is a complex variable, z as a *variable of integration* defines a *contour integral* in the complex plane.

Frequency Domain Analysis

z Transform

$$\mathbf{X}(z) = \sum_{n=-\infty}^{\infty} x[n]z^{-n}$$

$$x[n] = \frac{1}{j2\pi} \oint \mathbf{X}(z)z^{n-1}dz$$

Comments:

- z is a complex variable and $\mathbf{X}(z)$ is a complex-valued function of the complex variable z .
- LCCDEs always produce a ratio of polynomials in z^{-1} :

$$\mathbf{X}(z) = \frac{B(z)}{A(z)} = \frac{b_q z^{-q} + \dots + b_1 z^{-1} + b_0}{a_p z^{-p} + \dots + a_1 z^{-1} + a_0}$$

- Because z is a complex variable, z as a *variable of integration* defines a *contour integral* in the complex plane.
 - Contour integral wizards compute the inverse z transform from the definition.
 - Everyone else uses partial fraction expansion to construct tables, then uses the tables.

Frequency Domain Analysis

z Transform

$$\mathbf{X}(z) = \sum_{n=-\infty}^{\infty} x[n]z^{-n}$$

$$x[n] = \frac{1}{j2\pi} \oint \mathbf{X}(z)z^{n-1}dz$$

Comments:

- z is a complex variable and $\mathbf{X}(z)$ is a complex-valued function of the complex variable z .
- LCCDEs always produce a ratio of polynomials in z^{-1} :

$$\mathbf{X}(z) = \frac{B(z)}{A(z)} = \frac{b_q z^{-q} + \dots + b_1 z^{-1} + b_0}{a_p z^{-p} + \dots + a_1 z^{-1} + a_0}$$

- Because z is a complex variable, z as a *variable of integration* defines a *contour integral* in the complex plane.
 - Contour integral wizards compute the inverse z transform from the definition.
 - Everyone else uses partial fraction expansion to construct tables, then uses the tables.

Signal	Transform	ROC
$\delta[n]$	1	all z
$U[n]$	$\frac{1}{1-z^{-1}}$	$ z > 1$
$a^n U[n]$	$\frac{1}{1-az^{-1}}$	$ z > a $
$na^n U[n]$	$\frac{az^{-1}}{(1-az^{-1})^2}$	$ z > a $
$\delta[n-m]$	z^{-m}	all z^a
$\cos(\omega_0 n) U[n]$	$\frac{1 - \cos(\omega_0)z^{-1}}{1 - [2\cos(\omega_0)]z^{-1} + z^{-2}}$	$ z > 1$
$\sin(\omega_0 n) U[n]$	$\frac{\sin(\omega_0)z^{-1}}{1 - [2\cos(\omega_0)]z^{-1} + z^{-2}}$	$ z > 1$
$r^n \cos(\omega_0 n) U[n]$	$\frac{1 - r \cos(\omega_0)z^{-1}}{1 - [2r\cos(\omega_0)]z^{-1} + z^{-2}}$	$ z > r$
$r^n \sin(\omega_0 n) U[n]$	$\frac{r \sin(\omega_0)z^{-1}}{1 - [2r\cos(\omega_0)]z^{-1} + z^{-2}}$	$ z > r$

^aexcept 0 if $m > 0$ or ∞ if $m > 0$

Property	Signal	z -transform	ROC
	$x[n]$	$\mathbf{X}(z)$	R_x
	$y[n]$	$\mathbf{Y}(z)$	R_y
Linearity	$ax[n] + by[n]$	$a\mathbf{X}(z) + b\mathbf{Y}(z)$	at least $R_x \cap R_y$
Time Shifting	$x[n - n_0]$	$z^{-n_0}\mathbf{X}(z)$	R_x
Time Scaling (Upsampling)	$\begin{cases} x(n/K) & n \text{ is a multiple of } K \\ 0 & \text{otherwise} \end{cases}$	$\mathbf{X}(z^K)$	$R_x^{1/K}$
Conjugation	$x^*(n)$	$\mathbf{X}^*(z^*)$	R_x
Convolution	$x[n] * y[n]$	$\mathbf{X}(z)\mathbf{Y}(z)$	at least $R_x \cap R_y$
First Difference	$x[n] - x[n - 1]$	$(1 - z^{-1})\mathbf{X}(z)$	at least $R_x \cap \{ z > 0\}$
Accumulation	$\sum_{k=-\infty}^n x[k]$	$\frac{1}{1 - z^{-1}}\mathbf{X}(z)$	at least $R_x \cap \{ z > 1\}$
Initial Value Theorem	$x[0]$	$\lim_{z \rightarrow \infty} \mathbf{X}(z)$	
Final Value Theorem	$\lim_{N \rightarrow \infty} x[n]$	$\lim_{z \rightarrow 1} (1 - z^{-1})\mathbf{X}(z)$	poles of $(1 - z^{-1})\mathbf{X}(z)$ inside unit circle.

Frequency Domain Analysis

z Transform

$$\mathbf{X}(z) = \sum_{n=-\infty}^{\infty} x[n]z^{-n}$$

$$x[n] = \frac{1}{j2\pi} \oint \mathbf{X}(z)z^{n-1}dz$$

Discrete Time Fourier Transform (DTFT)

$$\mathbf{X}(e^{j\omega}) = \sum_{n=-\infty}^{\infty} x[n]e^{-j\omega n}$$

$$x[n] = \frac{1}{2\pi} \int_{-\pi}^{\pi} \mathbf{X}(e^{j\omega})e^{j\omega} d\omega$$

Comments:

- z is a complex variable and $\mathbf{X}(z)$ is a complex-valued function of the complex variable z .
- LCCDEs always produce a ratio of polynomials in z^{-1} :

$$\mathbf{X}(z) = \frac{B(z)}{A(z)} = \frac{b_q z^{-q} + \dots + b_1 z^{-1} + b_0}{a_p z^{-p} + \dots + a_1 z^{-1} + a_0}$$

- Because z is a complex variable, z as a *variable of integration* defines a *contour integral* in the complex plane.
 - Contour integral wizards compute the inverse z transform from the definition.
 - Everyone else uses partial fraction expansion to construct tables, then uses the tables.

Frequency Domain Analysis

z Transform

$$\mathbf{X}(z) = \sum_{n=-\infty}^{\infty} x[n]z^{-n}$$

$$x[n] = \frac{1}{j2\pi} \oint \mathbf{X}(z) z^{n-1} dz$$

Comments:

- z is a complex variable and $\mathbf{X}(z)$ is a complex-valued function of the complex variable z .
- LCCDEs always produce a ratio of polynomials in z^{-1} :

$$\mathbf{X}(z) = \frac{B(z)}{A(z)} = \frac{b_q z^{-q} + \dots + b_1 z^{-1} + b_0}{a_p z^{-p} + \dots + a_1 z^{-1} + a_0}$$

- Because z is a complex variable, z as a *variable of integration* defines a *contour integral* in the complex plane.
 - Contour integral wizards compute the inverse z transform from the definition.
 - Everyone else uses partial fraction expansion to construct tables, then uses the tables.


Discrete Time Fourier Transform (DTFT)

$$\mathbf{X}(e^{j\omega}) = \sum_{n=-\infty}^{\infty} x[n]e^{-j\omega n}$$

$$x[n] = \frac{1}{2\pi} \int_{-\pi}^{\pi} \mathbf{X}(e^{j\omega}) e^{j\omega} d\omega$$

Comments:

- w is a real variable and $\mathbf{X}(e^{j\omega})$ is a complex-valued function of the real variable ω .



Frequency Domain Analysis

z Transform

$$\mathbf{X}(z) = \sum_{n=-\infty}^{\infty} x[n]z^{-n}$$

$$x[n] = \frac{1}{j2\pi} \oint \mathbf{X}(z) z^{n-1} dz$$

Comments:

- z is a complex variable and $\mathbf{X}(z)$ is a complex-valued function of the complex variable z .
- LCCDEs always produce a ratio of polynomials in z^{-1} :

$$\mathbf{X}(z) = \frac{B(z)}{A(z)} = \frac{b_q z^{-q} + \dots + b_1 z^{-1} + b_0}{a_p z^{-p} + \dots + a_1 z^{-1} + a_0}$$

- Because z is a complex variable, z as a *variable of integration* defines a *contour integral* in the complex plane.
 - Contour integral wizards compute the inverse z transform from the definition.
 - Everyone else uses partial fraction expansion to construct tables, then uses the tables.


Discrete Time Fourier Transform (DTFT)

$$\mathbf{X}(e^{j\omega}) = \sum_{n=-\infty}^{\infty} x[n]e^{-j\omega n}$$

$$x[n] = \frac{1}{2\pi} \int_{-\pi}^{\pi} \mathbf{X}(e^{j\omega}) e^{j\omega} d\omega$$

Comments:

- ω is a real variable and $\mathbf{X}(e^{j\omega})$ is a complex-valued function of the real variable ω .
- $\mathbf{X}(e^{j\omega})$ is *periodic* in ω with period 2π .



Frequency Domain Analysis

z Transform

$$\mathbf{X}(z) = \sum_{n=-\infty}^{\infty} x[n]z^{-n}$$

$$x[n] = \frac{1}{j2\pi} \oint \mathbf{X}(z)z^{n-1}dz$$

Comments:

- z is a complex variable and $\mathbf{X}(z)$ is a complex-valued function of the complex variable z .
- LCCDEs always produce a ratio of polynomials in z^{-1} :

$$\mathbf{X}(z) = \frac{B(z)}{A(z)} = \frac{b_q z^{-q} + \dots + b_1 z^{-1} + b_0}{a_p z^{-p} + \dots + a_1 z^{-1} + a_0}$$

- Because z is a complex variable, z as a *variable of integration* defines a *contour integral* in the complex plane.
 - Contour integral wizards compute the inverse z transform from the definition.
 - Everyone else uses partial fraction expansion to construct tables, then uses the tables.

Discrete Time Fourier Transform (DTFT)

$$\mathbf{X}(e^{j\omega}) = \sum_{n=-\infty}^{\infty} x[n]e^{-j\omega n}$$

$$x[n] = \frac{1}{2\pi} \int_{-\pi}^{\pi} \mathbf{X}(e^{j\omega})e^{j\omega} d\omega$$


Comments:

- w is a real variable and $\mathbf{X}(e^{j\omega})$ is a complex-valued function of the real variable ω .
- $\mathbf{X}(e^{j\omega})$ is *periodic* in ω with period 2π .
- Because ω is a real variable, the integral that defines the inverse transform is the familiar integral introduced in your first and second calculus courses.
 - Integral wizards compute the inverse DTFT transform from the definition.
 - Everyone else uses tables.

Signal	DTFT
$x[n]$	$\mathbf{X}(e^{j\omega})$
$\delta[n]$	1
$\delta[n - n_0]$	$e^{-j\omega n_0}$
$U[n]$	$\frac{1}{1 - e^{-j\omega}} + \pi \sum_{\ell=-\infty}^{\infty} \delta(\omega - 2\pi\ell)$
$a^n U[n], a < 1$	$\frac{1}{1 - ae^{-j\omega}}$
$\sum_{k=k_0}^{k_0+N-1} c_k e^{jk(2\pi/N)n}$	$2\pi \sum_{k=-\infty}^{\infty} c_k \delta\left(\omega - \frac{2\pi k}{N}\right)$
$\sum_{k=-\infty}^{\infty} \delta[n - kN]$	$\frac{2\pi}{N} \sum_{\ell=-\infty}^{\infty} \delta\left(\omega - \frac{2\pi\ell}{N}\right)$
1	$2\pi \sum_{\ell=-\infty}^{\infty} \delta(\omega - 2\pi\ell)$
$\begin{cases} 1 & n \leq N \\ 0 & n > N \end{cases}$	$\frac{\sin\left(\omega\left[N + \frac{1}{2}\right]\right)}{\sin\left(\frac{\omega}{2}\right)}$
$e^{j\omega_0 n}$	$2\pi \sum_{\ell=-\infty}^{\infty} \delta(\omega - \omega_0 - 2\pi\ell)$
$\cos(\omega_0 n)$	$\pi \sum_{\ell=-\infty}^{\infty} \left\{ \delta(\omega - \omega_0 - 2\pi\ell) + \delta(\omega + \omega_0 - 2\pi\ell) \right\}$
$\sin(\omega_0 n)$	$\frac{\pi}{j} \sum_{\ell=-\infty}^{\infty} \left\{ \delta(\omega - \omega_0 - 2\pi\ell) - \delta(\omega + \omega_0 - 2\pi\ell) \right\}$

Property	Signal	DTFT
	$x[n]$	$\mathbf{X}(e^{j\omega})$
	$y[n]$	$\mathbf{Y}(e^{j\omega})$
Linearity	$ax[n] + by[n]$	$a\mathbf{X}(e^{j\omega}) + b\mathbf{Y}(e^{j\omega})$
Time Shifting	$x[n - n_0]$	$e^{-j\omega n_0} \mathbf{X}(e^{j\omega})$
Time Scaling (Upsampling)	$\begin{cases} x[n/K] & n \text{ is a multiple of } K \\ 0 & \text{otherwise} \end{cases}$	$\mathbf{X}(e^{jK\omega})$
Conjugation	$x^*[n]$	$\mathbf{X}^*(e^{-j\omega})$
Convolution	$x[n] * y[n]$	$\mathbf{X}(e^{j\omega}) \mathbf{Y}(e^{j\omega})$
Multiplication	$x[n]y[n]$	$\frac{1}{2\pi} \mathbf{X}(e^{j\omega}) * \mathbf{Y}(e^{j\omega})$
First Difference	$x[n] - x[n - 1]$	$(1 - e^{-j\omega}) \mathbf{X}(e^{j\omega})$
Accumulation	$\sum_{k=-\infty}^n x[k]$	$\frac{1}{1 - e^{-j\omega}} \mathbf{X}(e^{j\omega})$
Parseval's Relation	$\sum_{n=-\infty}^{\infty} x[n] ^2 = \frac{1}{2\pi} \int_{-\pi}^{\pi} \mathbf{X}(e^{j\omega}) ^2 d\omega$	

Relationships: LCCDE, Impulse Response, Convolution, Frequency Domain




The linear systems of interest to us have an input/output relationship defined by LCCDE

$$\text{e.g., } y[n] - ay[n - 1] = x[n]$$

$$y[n] - ay[n - 1] = x[n]$$

Relationships: LCCDE, Impulse Response, Convolution, Frequency Domain




The linear systems of interest to us have an input/output relationship defined by LCCDE

$$\text{e.g., } y[n] - ay[n - 1] = x[n]$$

$$y[n] - ay[n - 1] = x[n]$$

$$\mathbf{Y}(z) - a(z^{-1}\mathbf{Y}(z) + y[-1]) = \mathbf{X}(z)$$

Relationships: LCCDE, Impulse Response, Convolution, Frequency Domain



The linear systems of interest to us have an input/output relationship defined by LCCDE


$$\text{e.g., } y[n] - ay[n - 1] = x[n]$$

$$y[n] - ay[n - 1] = x[n]$$

$$\mathbf{Y}(z) - a(z^{-1}\mathbf{Y}(z) + y[-1]) = \mathbf{X}(z)$$

0

Relationships: LCCDE, Impulse Response, Convolution, Frequency Domain



The linear systems of interest to us have an input/output relationship defined by LCCDE


$$\text{e.g., } y[n] - ay[n - 1] = x[n]$$

$$y[n] - ay[n - 1] = x[n]$$

$$\mathbf{Y}(z) - a(z^{-1}\mathbf{Y}(z) + y[-1]) = \mathbf{X}(z)$$

$$\mathbf{Y}(z) - az^{-1}\mathbf{Y}(z) = \mathbf{X}(z)$$

Relationships: LCCDE, Impulse Response, Convolution, Frequency Domain



The linear systems of interest to us have an input/output relationship defined by LCCDE

$$\text{e.g., } y[n] - ay[n - 1] = x[n]$$


$$y[n] - ay[n - 1] = x[n]$$

$$\mathbf{Y}(z) - a(z^{-1}\mathbf{Y}(z) + y[-1]) = \mathbf{X}(z)$$

$$\mathbf{Y}(z) - az^{-1}\mathbf{Y}(z) = \mathbf{X}(z)$$

$$(1 - az^{-1})\mathbf{Y}(z) = \mathbf{X}(z)$$

Relationships: LCCDE, Impulse Response, Convolution, Frequency Domain



The linear systems of interest to us have an input/output relationship defined by LCCDE

$$\text{e.g., } y[n] - ay[n - 1] = x[n]$$

$$y[n] - ay[n - 1] = x[n]$$

$$\mathbf{Y}(z) - a(z^{-1}\mathbf{Y}(z) + y[-1]) = \mathbf{X}(z)$$


0

$$\mathbf{Y}(z) - az^{-1}\mathbf{Y}(z) = \mathbf{X}(z)$$

$$(1 - az^{-1})\mathbf{Y}(z) = \mathbf{X}(z)$$

$$\mathbf{Y}(z) = \frac{1}{1 - az^{-1}} \mathbf{X}(z)$$

Relationships: LCCDE, Impulse Response, Convolution, Frequency Domain



The linear systems of interest to us have an input/output relationship defined by LCCDE

$$\text{e.g., } y[n] - ay[n - 1] = x[n]$$

$$y[n] - ay[n - 1] = x[n]$$

$$\mathbf{Y}(z) - a(z^{-1}\mathbf{Y}(z) + y[-1]) = \mathbf{X}(z)$$

0

$$\mathbf{Y}(z) - az^{-1}\mathbf{Y}(z) = \mathbf{X}(z)$$


$$(1 - az^{-1})\mathbf{Y}(z) = \mathbf{X}(z)$$

$$\mathbf{Y}(z) = \frac{1}{1 - az^{-1}} \mathbf{X}(z)$$



Transfer function $\mathbf{H}(z)$

Relationships: LCCDE, Impulse Response, Convolution, Frequency Domain



The linear systems of interest to us have an input/output relationship defined by LCCDE

$$\text{e.g., } y[n] - ay[n - 1] = x[n]$$

$$y[n] - ay[n - 1] = x[n]$$

$$\mathbf{Y}(z) - a(z^{-1}\mathbf{Y}(z) + y[-1]) = \mathbf{X}(z)$$

0

$$\mathbf{Y}(z) - az^{-1}\mathbf{Y}(z) = \mathbf{X}(z)$$

$$(1 - az^{-1})\mathbf{Y}(z) = \mathbf{X}(z)$$


$$\mathbf{Y}(z) = \frac{1}{1 - az^{-1}} \mathbf{X}(z)$$



Transfer function $\mathbf{H}(z)$

$$\mathbf{Y}(z) = \mathbf{H}(z)\mathbf{X}(z) \Rightarrow y[n] = \sum_{m=\infty}^{\infty} h[m]x[n-m]$$

Relationships: LCCDE, Impulse Response, Convolution, Frequency Domain



The linear systems of interest to us have an input/output relationship defined by LCCDE

$$\text{e.g., } y[n] - ay[n - 1] = x[n]$$

$$y[n] - ay[n - 1] = x[n]$$

$$\mathbf{Y}(z) - a(z^{-1}\mathbf{Y}(z) + y[-1]) = \mathbf{X}(z)$$

0

$$\mathbf{Y}(z) - az^{-1}\mathbf{Y}(z) = \mathbf{X}(z)$$

$$(1 - az^{-1})\mathbf{Y}(z) = \mathbf{X}(z)$$

$$\mathbf{Y}(z) = \frac{1}{1 - az^{-1}} \mathbf{X}(z)$$


↑

Transfer function $\mathbf{H}(z)$

$$\mathbf{Y}(z) = \frac{1}{1 - az^{-1}} \mathbf{X}(z) \Rightarrow y[n] = \sum_{m=0}^{\infty} a^m x[n - m]$$

$$\mathbf{Y}(z) = \mathbf{H}(z)\mathbf{X}(z) \Rightarrow y[n] = \sum_{m=\infty}^{\infty} h[m]x[n - m]$$

Relationships: LCCDE, Impulse Response, Convolution, Frequency Domain



The linear systems of interest to us have an input/output relationship defined by LCCDE

$$\text{e.g., } y[n] - ay[n - 1] = x[n]$$

$$y[n] - ay[n - 1] = x[n]$$

$$\mathbf{Y}(z) - a(z^{-1}\mathbf{Y}(z) + y[-1]) = \mathbf{X}(z)$$

0

$$\mathbf{Y}(z) - az^{-1}\mathbf{Y}(z) = \mathbf{X}(z)$$

$$(1 - az^{-1})\mathbf{Y}(z) = \mathbf{X}(z)$$

$$\mathbf{Y}(z) = \frac{1}{1 - az^{-1}} \mathbf{X}(z)$$


↑
Transfer function $\mathbf{H}(z)$

$$\mathbf{Y}(z) = \frac{1}{1 - az^{-1}} \mathbf{X}(z) \Rightarrow y[n] = \sum_{m=0}^{\infty} a^m x[n - m]$$


$$\mathbf{Y}(z) = \mathbf{H}(z)\mathbf{X}(z) \Rightarrow y[n] = \sum_{m=\infty}^{\infty} h[m]x[n - m]$$



Is this really the impulse response?



Relationships: LCCDE, Impulse Response, Convolution, Frequency Domain



The linear systems of interest to us have an input/output relationship defined by LCCDE

$$\text{e.g., } y[n] - ay[n - 1] = x[n]$$

$$y[n] - ay[n - 1] = x[n]$$

$$\mathbf{Y}(z) - a(z^{-1}\mathbf{Y}(z) + y[-1]) = \mathbf{X}(z)$$

0


$$\mathbf{Y}(z) - az^{-1}\mathbf{Y}(z) = \mathbf{X}(z)$$

$$(1 - az^{-1})\mathbf{Y}(z) = \mathbf{X}(z)$$

$$\mathbf{Y}(z) = \frac{1}{1 - az^{-1}} \mathbf{X}(z)$$

$$x[n] = \delta[n] \Rightarrow \mathbf{X}(z) = 1$$

Relationships: LCCDE, Impulse Response, Convolution, Frequency Domain



The linear systems of interest to us have an input/output relationship defined by LCCDE

$$\text{e.g., } y[n] - ay[n - 1] = x[n]$$

$$y[n] - ay[n - 1] = x[n]$$

$$\mathbf{Y}(z) - a(z^{-1}\mathbf{Y}(z) + y[-1]) = \mathbf{X}(z)$$

0

$$\mathbf{Y}(z) - az^{-1}\mathbf{Y}(z) = \mathbf{X}(z)$$


$$(1 - az^{-1})\mathbf{Y}(z) = \mathbf{X}(z)$$

$$\mathbf{Y}(z) = \frac{1}{1 - az^{-1}} \mathbf{X}(z)$$

$$x[n] = \delta[n] \Rightarrow \mathbf{X}(z) = 1$$

$$\mathbf{Y}(z) = \mathbf{H}(z)\mathbf{X}(z) \Rightarrow \mathbf{Y}(z) = \mathbf{H}(z)$$

Relationships: LCCDE, Impulse Response, Convolution, Frequency Domain



The linear systems of interest to us have an input/output relationship defined by LCCDE

$$\text{e.g., } y[n] - ay[n - 1] = x[n]$$

$$y[n] - ay[n - 1] = x[n]$$

$$\mathbf{Y}(z) - a(z^{-1}\mathbf{Y}(z) + y[-1]) = \mathbf{X}(z)$$

0

$$\mathbf{Y}(z) - az^{-1}\mathbf{Y}(z) = \mathbf{X}(z)$$

$$(1 - az^{-1})\mathbf{Y}(z) = \mathbf{X}(z)$$

$$\mathbf{Y}(z) = \frac{1}{1 - az^{-1}} \mathbf{X}(z)$$


$$x[n] = \delta[n] \Rightarrow \mathbf{X}(z) = 1$$

$$\mathbf{Y}(z) = \mathbf{H}(z)\mathbf{X}(z) \Rightarrow \mathbf{Y}(z) = \mathbf{H}(z)$$

$$\mathbf{Y}(z) = \mathbf{H}(z) \Rightarrow y[n] = h[n]$$

Relationships: LCCDE, Impulse Response, Convolution, Frequency Domain

Relationships: LCCDE, Impulse Response, Convolution, Frequency Domain




The linear systems of interest to us have an input/output relationship defined by LCCDE

$$\text{e.g., } y[n] - ay[n - 1] = x[n]$$

$$y[n] - ay[n - 1] = x[n]$$


Relationships: LCCDE, Impulse Response, Convolution, Frequency Domain



The linear systems of interest to us have an input/output relationship defined by LCCDE

$$\text{e.g., } y[n] - ay[n - 1] = x[n]$$


$$y[n] - ay[n - 1] = x[n]$$



z domain transfer function

$$\mathbf{Y}(z) = \mathbf{H}(z)\mathbf{X}(z)$$


Relationships: LCCDE, Impulse Response, Convolution, Frequency Domain



The linear systems of interest to us have an input/output relationship defined by LCCDE


$$\text{e.g., } y[n] - ay[n - 1] = x[n]$$

$$y[n] - ay[n - 1] = x[n]$$



z domain transfer function


$$\mathbf{Y}(z) = \mathbf{H}(z)\mathbf{X}(z)$$



Time domain convolution

$$y[n] = \sum_{m=-\infty}^{\infty} x[m]h[n-m] = \sum_{m=-\infty}^{\infty} h[m]x[n-m]$$


Relationships: LCCDE, Impulse Response, Convolution, Frequency Domain



The linear systems of interest to us have an input/output relationship defined by LCCDE


$$\text{e.g., } y[n] - ay[n - 1] = x[n]$$

$$y[n] - ay[n - 1] = x[n]$$



z domain transfer function

$$\mathbf{Y}(z) = \mathbf{H}(z)\mathbf{X}(z)$$



Time domain convolution

$$y[n] = \sum_{m=-\infty}^{\infty} x[m]h[n-m] = \sum_{m=-\infty}^{\infty} h[m]x[n-m]$$




To find the impulse response of an LTI system described by an LCCDE

1. Solve the LCCDE using the z transform with all-zeros initial conditions.
2. Write the z domain solution as $\mathbf{Y}(z) = \mathbf{H}(z)\mathbf{X}(z)$.
3. Identify $\mathbf{H}(z)$ in the solution.
4. The impulse response $h[n]$ is the inverse Laplace transform of $\mathbf{H}(z)$.

Relationships: LCCDE, Impulse Response, Convolution, Frequency Domain + DTFT


Relationships: LCCDE, Impulse Response, Convolution, Frequency Domain + DTFT



The linear systems of interest to us have an input/output relationship defined by LCCDE

$$\text{e.g., } y[n] - ay[n - 1] = x[n]$$


Relationships: LCCDE, Impulse Response, Convolution, Frequency Domain + DTFT



The linear systems of interest to us have an input/output relationship defined by LCCDE

$$\text{e.g., } y[n] - ay[n - 1] = x[n]$$


$$y[n] - ay[n - 1] = x[n]$$



DTFT domain frequency response

$$\mathbf{Y}(e^{j\omega}) = \mathbf{H}(e^{j\omega})\mathbf{X}(e^{j\omega})$$


Relationships: LCCDE, Impulse Response, Convolution, Frequency Domain + DTFT



The linear systems of interest to us have an input/output relationship defined by LCCDE


$$\text{e.g., } y[n] - ay[n - 1] = x[n]$$

$$y[n] - ay[n - 1] = x[n]$$



DTFT domain frequency response


$$\mathbf{Y}(e^{j\omega}) = \mathbf{H}(e^{j\omega})\mathbf{X}(e^{j\omega})$$



Time domain convolution


$$y[n] = \sum_{m=-\infty}^{\infty} x[m]h[n-m] = \sum_{m=-\infty}^{\infty} h[m]x[n-m]$$

Relationships: LCCDE, Impulse Response, Convolution, Frequency Domain + DTFT




The linear systems of interest to us have an input/output relationship defined by LCCDE

$$\text{e.g., } y[n] - ay[n - 1] = x[n]$$



DTFT domain frequency response

$$\mathbf{Y}(e^{j\omega}) = \mathbf{H}(e^{j\omega})\mathbf{X}(e^{j\omega})$$




Time domain convolution

$$y[n] = \sum_{m=-\infty}^{\infty} x[m]h[n-m] = \sum_{m=-\infty}^{\infty} h[m]x[n-m]$$




- One can solve an LCCDE using the DTFT with all-zeros initial conditions. But it is more common to use the z transform.

Relationships: LCCDE, Impulse Response, Convolution, Frequency Domain + DTFT




The linear systems of interest to us have an input/output relationship defined by LCCDE

$$\text{e.g., } y[n] - ay[n - 1] = x[n]$$



DTFT domain frequency response

$$\mathbf{Y}(e^{j\omega}) = \mathbf{H}(e^{j\omega})\mathbf{X}(e^{j\omega})$$



Time domain convolution

$$y[n] = \sum_{m=-\infty}^{\infty} x[m]h[n-m] = \sum_{m=-\infty}^{\infty} h[m]x[n-m]$$




- One can solve an LCCDE using the DTFT with all-zeros initial conditions. But it is more common to use the z transform.
- After identifying the z domain transfer function $\mathbf{H}(z)$, the DTFT domain frequency response is

$$\mathbf{H}(e^{j\omega}) = \mathbf{H}(z) \Big|_{z=e^{j\omega}}$$


This works only as long as the ROC of $\mathbf{H}(z)$ contains the unit circle ($z = e^{j\omega}$).

Relationships: LCCDE, Impulse Response, Convolution, Frequency Domain + DTFT




The linear systems of interest to us have an input/output relationship defined by LCCDE

$$\text{e.g., } y[n] - ay[n - 1] = x[n]$$



DTFT domain frequency response

$$\mathbf{Y}(e^{j\omega}) = \mathbf{H}(e^{j\omega})\mathbf{X}(e^{j\omega})$$



Time domain convolution

$$y[n] = \sum_{m=-\infty}^{\infty} x[m]h[n-m] = \sum_{m=-\infty}^{\infty} h[m]x[n-m]$$



- One can solve an LCCDE using the DTFT with all-zeros initial conditions. But it is more common to use the z transform.
- After identifying the z domain transfer function $\mathbf{H}(z)$, the DTFT domain frequency response is

$$\mathbf{H}(e^{j\omega}) = \mathbf{H}(z) \Big|_{z=e^{j\omega}}$$

This works only as long as the ROC of $\mathbf{H}(z)$ contains the unit circle ($z = e^{j\omega}$).

- For stable systems, this is always true.