

ODE notes by Deepak Bhojia

- Gauss's Hypergeometric Equation
- Special functions:
 - Legendre Polynomials
 - Bessel functions
 - Gamma functions

Solving a Differential Equation using Hypergeometric Functions

Gauss's Hypergeometric Equation (GHE)

The differential equation

$$x(1-x)y'' + [c - (a+b+1)x]y' - aby = 0 \quad (1)$$

where a , b , and c are constants, is called **Gauss's Hypergeometric equation (GHE)**.

Learnings:?

- We will find the **general solution of GHE** at its regular points.
- As an application, in particular, we find the general solutions of the equation

$$(x - A)(x - B) y'' + (C + Dx) y' + Ey = 0$$

where $A \neq B$ by using the general solution of GHE.

Singularities

If we simplify GHE in the form of a standard second-order linear equation, we obtain

$$P(x) = \frac{c - [a + b + 1]x}{x(1 - x)} \quad \text{and} \quad Q(x) = \frac{-abx}{x(1 - x)}.$$

What are the singular points: $x = 0$ and $x = 1$ are only singular points.

Are these regular points? Yes, $x = 0$ and $x = 1$ are regular points, since

$$\begin{aligned} xP(x) &= \frac{c - [a + b + 1]x}{(1 - x)} = [c - (a + b + 1)x] (1 + x + x^2 + \cdots) \\ &= c + [c - (a + b + 1)]x + \cdots \end{aligned}$$

and

$$x^2Q(x) = \frac{-abx}{1 - x} = -abx(1 + x + x^2 + \cdots) = -abx - abx^2 - \cdots.$$

Similarly, we can show that $x = 1$ is a regular point. Aim to find the general solutions of GHE at $x = 0$ and $x = 1$.

General solution at the regular point $x = 0$

- From the above expansions of $xP(x)$ and $x^2Q(x)$, $p_0 = c$ and $q_0 = 0$ so the indicial equation is

$$m(m-1) + mp_0 + q_0 = 0 \quad \Rightarrow \quad m[m - (1-c)] = 0$$

and the exponents are $m = 0$ and $m = 1 - c$.

- Solution corresponding to the exponent $m = 0$:** The exponent $m = 0$ corresponds to a Frobenius series solution by Theorem A if $1 - c < 0$ or the difference $(1 - c) - 0 = 1 - c$ is **not a positive integer**.

Note that the second condition implies the first one, so if $1 - c$ is not a positive integer, equivalently c is neither zero nor a negative integer, then $m = 0$ corresponds to a Frobenius series solution of the form

$$y = x^0 \sum_{n=0}^{\infty} a_n x^n = a_0 + a_1 x + a_2 x^2 + \cdots \quad (2)$$

where a_0 is nonzero.

On substituting the above into (1) and equating the coefficients of x^n to zero, we obtain the following relations:

$$a_1 = \frac{ab}{c} a_0; \quad a_2 = \frac{(a+1)(b+1)}{2(c+1)} a_1 = \frac{a(a+1)b(b+1)}{2c(c+1)} a_0$$

$$a_{n+1} = \frac{(a+n)(b+n)}{(n+1)(c+n)} a_n. \quad (3)$$

With these coefficients and by letting $a_0 = 1$, the solution (2) becomes

$$y = 1 + \frac{ab}{c}x + \frac{a(a+1)b(b+1)}{2c(c+1)}x^2 + \cdots$$

$$= 1 + \sum_{n=1}^{\infty} \frac{a(a+1) \cdots (a+n-1) b(b+1) \cdots (b+n-1)}{n! c(c+1) \cdots (c+n-1)} x^n. \quad (4)$$

This is known as the *hypergeometric series*, and is denoted by the symbol $F(a, b, c, x)$. It is called by this name because it generalizes the familiar geometric series as follows:

- When $a = 1$ and $c = b$: $F(1, b, b, x) = 1 + x + x^2 + \cdots = \frac{1}{1-x}$.

What is the Hypergeometric Series?

- The **Gauss hypergeometric series** is

$$F(a, b; c; x) = \sum_{n=0}^{\infty} \alpha_n \frac{x^n}{n!},$$

where the coefficients α_n are given by $\alpha_n = \frac{(a)_n(b)_n}{(c)_n}$, $n \geq 0$.

- Here $(q)_n$ is the **Pochhammer symbol** (rising factorial):

$$(q)_n = q(q+1)(q+2) \cdots (q+n-1), \quad (q)_0 := 1.$$

- Familiar Functions as Special Cases of $F(a, b; c; x)$:**

Function	Hypergeometric Form	Comment
e^x	$F(1, 1; 1; x)$	Standard series expansion
$(1-x)^{-a}$	$F(a, 1; 1; x)$	Binomial theorem
$\ln(1+x)$	$x F(1, 1; 2; -x)$	Integral representation
$\arcsin x$	$x F(1/2, 1/2; 3/2; x^2)$	Trigonometric inverse
$\arctan x$	$x F(1/2, 1; 3/2; -x^2)$	Another inverse trig example
$\frac{1}{1-x}$	$F(1, 1; 1; x)$	Geometric series

Solution corresponding to the second exponent $m = 1 - c$

- If $1 - c$ is neither zero nor a negative integer, i.e., c is not a positive integer, then **Theorem A** also tells us that there is a second independent solution of **GHE** (1) near $x = 0$ with exponent $m = 1$.
- This solution can be found directly by substituting

$$y = x^{1-c}(a_0 + a_1x + a_2x^2 + \cdots)$$

into **GHE** (1) and calculating the coefficients.

- The other way of finding the solution is to change the dependent variable in (1) from y to z by writing $y = x^{1-c}z$. When the necessary computations are performed, equation (1) becomes

$$x(1-x)z'' + \left[(2-c) - ([a-c+1] + [b-c+1] + 1)x\right]z' - (a-c+1)(b-c+1)z = 0 \quad (5)$$

which is the **hypergeometric equation** with the constants a , b and c are replaced by $a - c + 1$, $b - c + 1$ and $2 - c$.

- We already know that (5) (from the previous Subsection) has the power series solution

$$z = F(a - c + 1, b - c + 1, 2 - c, x)$$

near the origin, so our desired second solution is

$$y = x^{1-c} z = x^{1-c} F(a - c + 1, b - c + 1, 2 - c, x).$$

- Accordingly, when c is not an integer, we have two independent Frobenius series solutions and hence

$$y = c_1 F(a, b, c, x) + c_2 x^{1-c} F(a - c + 1, b - c + 1, 2 - c, x) \quad (6)$$

is the general solution of the hypergeometric equation (1) near the singular point $x = 0$.

- Note that the above solution is only valid near the origin. We now solve GHE (1) near another singular point $x = 1$.

General solution of GHE at the regular point $x = 1$

- The simplest procedure is to obtain this solution from the one already found, by introducing a new independent variable $t = 1 - x$. ($x = 1 - t$, $dy/dx = -dy/dt$ and $d^2y/dx^2 = d^2y/dt^2$).
- This makes $x = 1$ correspond to $t = 0$ and transforms (1) into

$$t(1-t)y'' + [(a+b-c+1) - (a+b+1)t]y' - aby = 0$$

where the primes denotes the derivatives with respect to t .

- Since the above is a hypergeometric equation, its general solution near $t = 0$ can be written down at once from (6) by replacing x by t and c by $a+b-c+1$ and then we replace t by $1-x$ to get the general solution of GHE (1) near $x = 1$:

$$y = c_1 F(a, b, a+b-c+1, 1-x) + c_2 (1-x)^{c-a-b} F(c-b, c-a, c-a-b+1, 1-x). \quad (7)$$

(In this case it is necessary to assume that $c-a-b$ is not an integer).

Some comments, remarks and applications of hypergeometric functions F

- Formulas (6) and (7) show that the adaptability of the constants in equation GHE (1) makes it possible to express the general solution of this equation near each of its singular points in terms of the single function F .
- Moreover, the general solution of a wide class of differential equations can be written in terms of hypergeometric functions F .
- Any differential equation in which the coefficients of y'' , y' and y are polynomials of degree 2, 1 and 0 respectively and also the first of these polynomials has distinct real roots can be brought into hypergeometric form by a linear change of the independent variable, and hence can be solved near its singular point in terms of the hypergeometric function.
- The above remarks can be explained as follows. We consider the class of the equation of this type:

$$(x - A)(x - B)y'' + (C + Dx)y' + Ey = 0 \quad (8)$$

where $A \neq B$.

- If we change the independent variable from x to t linearly by means of $(x - A)(x - B) = kt(t - 1)$ for some constant k i.e.

$$t = \frac{x - A}{B - A}$$

then $x = A$ corresponds to $t = 0$ and $x = B$ to $t = 1$.

- After changing the independent variable from x to t the equation (8) assumes the form

$$t(1 - t)y'' + (F + Gt)y' + Hy = 0,$$

where F , G and H are certain combinations of the constants in (8) and the primes indicate derivatives with respect to t .

- This is a **hypergeometric equation** with constants a , b and c defined by $F = c$, $G = -(a + b + 1)$ and $H = -ab$ and can therefore be solved near $t = 0$ and $t = 1$ in terms of the hypergeometric function.
- This means that (8) can be solved in terms of the same function near $x = A$ and $x = B$.

Problems

Find the general solution of each of the following differential equations near the indicated singular point in terms of hypergeometric function.

$$\textcircled{1} \quad x(1-x)y'' + \left(\frac{3}{2} - 2x\right)y' + 2y = 0, \quad x = 0;$$

$$\textcircled{2} \quad (2x^2 + 2x)y'' + (1 + 5x)y' + y = 0, \quad x = 0;$$

$$\textcircled{3} \quad (x^2 - 1)y'' + (5x + 4)y' + 4y = 0, \quad x = -1;$$

$$\textcircled{4} \quad (x^2 - x - 6)y'' + (5 + 3x)y' + y = 0, \quad x = 3.$$

Hint: Convert the equation into hypergeometric equation with a suitable change of independent variable as explained in the previous slides.

Before we start to solve these problems, we recall hypergeometric equation and its general solutions near its regular singular points.

Hypergeometric equation and solutions

Hypergeometric equation:

$$x(1-x)y'' + [c - (a+b+1)x]y' - aby = 0$$

General solution at $x = 0$:

$$y = c_1 F(a, b, c, x) + c_2 x^{1-c} F(a-c+1, b-c+1, c-2, x)$$

General solution at $x = 1$:

$$y = c_1 F(a, b, a+b-c+1, 1-x) \\ + c_2 (1-x)^{c-a-b} F(c-b, c-a, c-a-b+1, 1-x)$$

Hypergeometric function:

$$F(a, b, c, x) = 1 + \sum_{n=1}^{\infty} \frac{a(a+1) \cdots (a+n-1) b(b+1) \cdots (b+n-1)}{n! c(c+1) \cdots (c+n-1)} x^n$$

Solution of Problem 1

$$x(1-x)y'' + \left(\frac{3}{2} - 2x\right)y' + 2y = 0, \quad x = 0.$$

If we compare this equation with the hypergeometric equation we have the coefficients

$$c = \frac{3}{2}; \quad a + b + 1 = 2; \quad ab = -2.$$

Or $c = \frac{3}{2}$; $a + b = 1$; $ab = -2$. And then $a - b$ can be found by using the formula

$$a - b = \sqrt{(a+b)^2 - 4ab} = \sqrt{1 - 4(-2)} = 3.$$

By solving these equations we will get $a = 2$, $b = -1$, $c = \frac{3}{2}$.

Therefore the General solution at $x = 0$ is given by

$$\begin{aligned} y &= c_1 F(a, b, c, x) + c_2 x^{1-c} F(a - c + 1, b - c + 1, c - 2, x) \\ &= c_1 F\left(2, -1, \frac{3}{2}, x\right) + c_2 x^{-1/2} F\left(\frac{3}{2}, -\frac{3}{2}, -\frac{1}{2}, x\right) \\ &= c_1 \left(1 - \frac{4}{3}x\right) + c_2 x^{-1/2} F\left(\frac{3}{2}, -\frac{3}{2}, -\frac{1}{2}, x\right). \end{aligned}$$

Solution of Problem 4

$$(x^2 - x - 6)y'' + (5 + 3x)y' + y = 0, \quad x = 3.$$

We use the change of independent variable from x to t by $t = \frac{x - A}{B - A}$, where A and B are roots of the polynomial $x^2 - x - 6$ which is a coefficient of y'' in given equation.

We see that $A = 3$ and $B = -2$ so that $t = \frac{3 - x}{5}$ or $x = -5t + 3$.

$$\frac{dy}{dx} = \frac{dy}{dt} \cdot \frac{dt}{dx} = -\frac{1}{5} \frac{dy}{dx}; \quad \frac{d^2y}{dx^2} = \frac{d}{dt} \left(-\frac{1}{5} \frac{dy}{dx} \right) = \frac{1}{25} \frac{d^2y}{dt^2}.$$

This transformation reduces the equation into

$$(-5t)(-5t + 5) \left(\frac{1}{25} \right) y'' + (5 - 15t + 9) \left(-\frac{1}{5} \right) y' + y = 0$$

$$t(1 - t)y'' + \left(\frac{14}{5} - 3t \right) y' - y = 0.$$

If we compare this equation with the hypergeometric equation we have the coefficients

$$c = \frac{14}{5}; \quad a + b + 1 = 3; \quad ab = 1.$$

$a + b = 2$; $ab = 1$ and hence $a - b = \sqrt{4 - 4} = 0$. Therefore

$$a = 1, \quad b = 1, \quad c = \frac{14}{5}.$$

The General solution at $t = 0$ or $x = 3$ is given by

$$\begin{aligned} y &= c_1 F(a, b, c, t) + c_2 t^{1-c} F(a - c + 1, b - c + 1, c - 2, t) \\ &= c_1 F\left(1, 1, \frac{14}{5}, t\right) + c_2 t^{-9/5} F\left(-\frac{4}{5}, -\frac{4}{5}, -\frac{4}{5}, t\right) \quad \left(\text{since } t = \frac{3-x}{5}\right) \\ &= c_1 F\left(1, 1, \frac{14}{5}, \frac{3-x}{5}\right) + c_2 \left(\frac{3-x}{5}\right)^{-9/5} F\left(-\frac{4}{5}, -\frac{4}{5}, -\frac{4}{5}, \frac{3-x}{5}\right). \end{aligned}$$

Problem 5

Some differential equations are of the hypergeometric type even though they may not appear to be so. Find the general solution of

$$(1 - e^x)y'' + \frac{1}{2}y' + e^xy = 0$$

near $x = 0$ by changing the independent variable to $t = e^x$.

Sol. The transformation $t = e^x$ implies $dt/dx = e^x = t$ and

$$\frac{dy}{dx} = \frac{dy}{dt} \cdot \frac{dt}{dx} = t \frac{dy}{dt}; \quad \frac{d^2y}{dx^2} = t^2 \frac{d^2y}{dt^2} + t \frac{dy}{dt}$$

We substitute all these in given equation to get

$$t^2(1 - t) \frac{d^2y}{dt^2} + t \left(\frac{3}{2} - t \right) \frac{dy}{dt} + ty = 0.$$

Or

$$t(1 - t) \frac{d^2y}{dt^2} + \left(\frac{3}{2} - t \right) \frac{dy}{dt} + y = 0.$$

After comparing with standard hypergeometric equation we get the constants $c = \frac{3}{2}$, $a + b + 1 = 1$ and $ab = -1$. If solve these equations. then we will get $a = 1$, $b = -1$, $c = \frac{3}{2}$.

The General solution at $t = 1$ or $x = 0$ is given by

$$\begin{aligned}
 y &= c_1 F(a, b, a + b - c + 1, 1 - t) + c_2 (1 - t)^{c-a-b} F(c - b, c - a, c - a - b + 1, 1 - t) \\
 &= c_1 F\left(1, -1, -\frac{1}{2}, 1 - t\right) + c_2 (1 - t)^{3/2} F\left(\frac{5}{2}, \frac{1}{2}, \frac{5}{2}, 1 - t\right) \\
 &= c_1 F\left(1, -1, -\frac{1}{2}, 1 - e^x\right) + c_2 (1 - e^x)^{3/2} F\left(\frac{5}{2}, \frac{1}{2}, \frac{5}{2}, 1 - e^x\right).
 \end{aligned}$$

Problem 6

Consider the Chebyshev's equation

$$(1 - x^2)y'' - xy' + p^2y = 0,$$

where p is a nonnegative constant. Transform it into a hypergeometric equation by changing the independent variable from x to $t = \frac{1}{2}(1 - x)$, and show that its general solution near $x = 1$ is

$$y = c_1 F\left(p, -p, \frac{1}{2}, \frac{1-x}{2}\right) + c_2 \left(\frac{1-x}{2}\right)^{1/2} F\left(p + \frac{1}{2}, -p + \frac{1}{2}, \frac{3}{2}, \frac{1-x}{2}\right).$$

Problem 7

Consider the differential equation

$$x(1-x)y'' + [p - (p+2)x]y' - py = 0$$

where p is a constant.

- (a) If p is not an integer, find the general solution near $x = 0$ in terms of hypergeometric functions.
- (b) Write the general solution found in (a) in terms of elementary functions.
- (c) When $p = 1$, the differential equation becomes

$$x(x-1)y'' + (1-3x)y' - y = 0$$

and the solution in (b) is no longer the general solution. Find the general solution in this case by the method of the use of known solution to find another.

Problem 8. Consider the Legendre's equation

$$(1 - x^2)y'' - 2xy' + n(n + 1)y = 0 \quad (9)$$

where n is understood to be nonnegative integer.

We have found analytic solutions near ordinary point $x = 0$. However, the solutions most useful in the applications are those bounded near $x = 1$.

(a) Find the general solution near $x = 1$ by changing the independent variable from x to $\frac{1}{2}(1 - x)$

(b) Find the solutions from (a) that are bounded near $x = 1$.

Sol. The change of independent variable x to $\frac{1}{2}(1 - x)$ makes $x = 1$ correspond to $t = 0$ and transforms the equation (9) into

$$t(1 - t)y'' + (1 - 2t)y' + n(n + 1)y = 0 \quad (10)$$

where the primes denotes derivatives with respect to t .

This is hypergeometric equation with $a = -n$, $b = n + 1$, and $c = 1$, so it has the following polynomial solution near $t = 0$:

$$y_1 = F(-n, n + 1, 1, t). \quad (11)$$

Since the exponents of the equation (10) at the origin $t = 0$ are both zero ($m_1 = 0$ and $m_2 = 1 - c = 0$), we seek a second solution via the formula $y_2(t) = y_1(t)v(t)$ with $v(t)$ is given by

$$v(t) = \int \frac{1}{y_1^2} e^{-\int P(t) dt} dt$$

where $P(t) = \frac{1 - 2t}{t(1 - t)}$. Note that

$$P(t) = \frac{1 - 2t}{t(1 - t)} = \frac{1}{t} - \frac{1}{1 - t}$$

therefore

$$e^{-\int P(t) dt} = e^{-\ln t - \ln(1-t)} = \frac{1}{t(1 - t)}.$$

So we have that

$$v(t) = \int \frac{1}{y_1^2} \frac{1}{t(1-t)} dt = \int \frac{1}{t} \left[\frac{1}{y_1^2} (1-t) \right] dt$$

Since y_1 is a polynomial with constant term 1, the bracket expression on right is an analytic function of the form $1 + a_1 t + a_2 t^2 + \cdots$. This yields

$v(t) = \log t + a_1 t + \cdots$, so

$$y_2 = y_1(\log t + a_1 t + \cdots).$$

and the general solution is given by

$$y = c_1 y_1 + c_2 y_2. \quad (12)$$

If we replace t in (12) by $\frac{1}{2}(1-x)$, we will get the general solution of Legendre's equation.

Because of the presence of the term $\log t$ in y_2 , it is clear that (12) is bounded if and only if $c_2 = 0$.

If we replace t in (11) by $\frac{1}{2}(1-x)$, it follows that the solutions of Legendre's equation (9) are bounded near $x = 1$ are precisely constant multiple of the polynomial

$$F\left[-n, n+1, 1, \frac{1}{2}(1-x)\right].$$

Here the polynomial

$$F\left[-n, n+1, 1, \frac{1}{2}(1-x)\right]$$

is called n -th Legendre polynomial.

Problem 9. Verify the following identities

$$(a) \quad (1+x)^p = F(-p, b, b, -x)$$

$$(b) \quad e^x = \lim_{b \rightarrow \infty} F\left(a, b, a, \frac{x}{b}\right)$$

$$(c) \quad \log(1+x) = xF(1, 1, 2, -x)$$

$$(d) \quad \cos x = \lim_{a \rightarrow \infty} F\left(a, a, \frac{1}{2}, -\frac{x^2}{4a^2}\right)$$

Legendre's equation

Legendre Polynomials

- The Legendre Equation is

$$(1 - x^2) y'' - 2x y' + n(n + 1) y = 0,$$

where n is a constant.

- **How will you try to find its solution at $x = 0$?**
- **Series Solution at an Ordinary Point:** Here $x = 0$ is an ordinary point and the general solution is

$$y(x) = a_0 \left[1 - \frac{n(n+1)}{2!} x + \frac{n(n-2)(n+1)(n+3)}{3!} x^4 - \dots \right] \\ + a_1 \left[x - \frac{(n-1)(n+2)}{3!} x^3 + \frac{(n-1)(n-3)(n+2)(n+4)}{5!} x^5 - \dots \right].$$

Note:

- If n is not an integer, then each series is convergent in $-1 < x < 1$.
- If n is a non-negative integer, one of the series terminates and thus is a polynomial.

Solution near $x = 1$

- However, the solutions, most useful in the applications, are those bounded near $x = 1$.
- So our interest is to solve the Legendre equation near $x = 1$.
- For this we proceed by the approach of Hypergeometric equation.

The Legendre Equation is

$$(1 - x^2) y'' - 2x y' + n(n + 1) y = 0,$$

where n is a constant.

- What are its regular singular points?

Solution near $x = 1$

The Legendre Equation is

$$(1 - x^2) y'' - 2x y' + n(n + 1) y = 0,$$

where n is a constant.

- Looking at coefficients of y'' , y' , y we notice that we can transform this into a hypergeometric equation in standard form.
- What transformation will be helpful to find the solution near $x = 1$?
- Can we find the transformed Differential equation?

Solution near $x = 1$

The Legendre Equation is

$$(1 - x^2) y'' - 2x y' + n(n + 1) y = 0,$$

where n is a constant.

- We change the independent variable from x to t by

$$t = \frac{x - 1}{(-1) - 1} = \frac{1}{2}(1 - x).$$

- This makes $x = 1$ correspond to $t = 0$ and transforms the equation into

$$t(1 - t)y'' + (1 - 2t)y' + n(n + 1)y = 0,$$

where the primes represent derivatives with respect to t .

What are its solutions?

Solution near $x = 1$

The Transformed Equation is

$$t(1-t)y'' + (1-2t)y' + n(n+1)y = 0,$$

where n is a constant.

- This is hypergeometric equation with $a = -n, b = n+1, c = 1$.
- When $1-c$ is a real number but not an integer, then the general solution of the above equation can be found as

$$y = c_1 F(a, b, c, x) + c_2 x^{1-c} F(a-c+1, b-c+1, 2-c, x) \quad (6)$$

- But in our case we have $1-c = 0$. Infact both exponents are 0 ($m_1 = 0, m_2 = 0$.)
- So we need an alternative approach.
- Note, the first solution is $y_1 = F(-n, n+1, 1, t)$.

The Transformed Equation is

$$t(1-t)y'' + (1-2t)y' + n(n+1)y = 0,$$

where n is a constant. Note that we have $P(x) = \frac{(1-2t)}{t(1-t)}$.

- We seek a second solution by the method of Section 16. This second solution is $y_2 = v y_1$, where

$$v' = \frac{1}{y_1^2} e^{-\int p dt} = \frac{1}{y_1^2} e^{\int \frac{(2t-1)}{t(1-t)} dt} = \frac{1}{y_1^2 t(1-t)} = \frac{1}{t} \left[\frac{1}{y_1^2(1-t)} \right].$$

- On solving, we get

$$y_2 = y_1 (\log t + a_1 t + \dots)$$

and the general solution of (2) near the origin is

$$y = c_1 y_1 + c_2 y_2.$$

Bounded Solutions Near $x = 1$

- Because of the presence of the term $\log t$ in y_2 , it is clear that the general solution

$$y = c_1 y_1 + c_2 y_2$$

is bounded near $t = 0$ if and only if $c_2 = 0$.

- Replace t by

$$t = \frac{1}{2}(1 - x),$$

- The solutions of Legendre's equation that are bounded near $x = 1$ are precisely constant multiples of the polynomial

$$F(-n, n+1, 1, \frac{1}{2}(1-x)).$$

- This brings us to the fundamental definition. The n^{th} Legendre polynomial is denoted by $P_n(x)$ and defined by

$$P_n(x) = F\left(-n, n+1, 1, \frac{1}{2}(1-x)\right)$$

Definition of $P_n(x)$ via F

Definition (Legendre polynomial). The n^{th} Legendre polynomial $P_n(x)$ is defined by

$$P_n(x) = F\left(-n, n+1, 1, \frac{1-x}{2}\right).$$

$$\begin{aligned} P_n(x) &= F\left(-n, n+1, 1, \frac{1-x}{2}\right) \\ &= 1 + \frac{(-n)(n+1)}{(1!)^2} \left(\frac{1-x}{2}\right) + \frac{(-n)(-n+1)(n+1)(n+2)}{(2!)^2} \left(\frac{1-x}{2}\right)^2 + \cdots \\ &\quad + \frac{(-n)(-n+1)(-n+(n-1))(n+1)(n+2) \cdots (2n)}{(n!)^2} \left(\frac{1-x}{2}\right)^n \end{aligned}$$

$$P_n(x) = 1 + \frac{n(n+1)}{(1!)^2 2} (x-1) + \frac{n(n-1)(n+1)(n+2)}{(2!)^2 2^2} (x-1)^2 + \cdots + \frac{(2n)!}{(n!)^2 2^n} (x-1)^n.$$

Remark: The above formula is a very inconvenient tool to use in studying $P_n(x)$, so we look for something simpler.

Rodrigues' formula

The n^{th} Legendre polynomial can equivalently be defined by Rodrigues' formula:

$$P_n(x) = \frac{1}{2^n n!} \frac{d^n}{dx^n} [(x^2 - 1)^n], \quad n = 0, 1, 2, \dots$$

Remark. The above provides a relatively easy method for computing successive Legendre polynomials.

- **First few polynomials (by Rodrigues):**

$$P_0(x) = 1, \quad P_1(x) = x,$$

$$P_2(x) = \frac{1}{2}(3x^2 - 1), \quad P_3(x) = \frac{1}{2}(5x^3 - 3x),$$

$$P_4(x) = \frac{1}{8}(35x^4 - 30x^2 + 3), \quad P_5(x) = \frac{1}{8}(63x^5 - 70x^3 + 15x), \dots$$

Orthogonality

On $[-1, 1]$, the sequence $\{P_n\}_{n \geq 0}$ is orthogonal:

$$\int_{-1}^1 P_m(x) P_n(x) dx = \begin{cases} 0, & m \neq n, \\ \frac{2}{2n+1}, & m = n. \end{cases}$$

Problem. State and prove the orthogonality property of Legendre polynomials.

Thank you for your attention