

Introduction

This document is an amalgam of notes-to-self, useful formulas, and matrix derivative computations for Gauss-Turan Quadrature. It is based on the BTG model due to Oliveira, Kedem, and Short.

-Leo Huang

Notation and Equations

$\{Z(s), s \in D, D \subseteq \mathbb{R}^d\}$: random field

$Z = [Z(s_1), \dots, Z(s_n)]^T$: observations

$Z_0 = [Z(s_{01}), \dots, Z(s_{0k})]^T$: unobserved random vector (to be predicted)

$z = [z_1, \dots, z_n]^T, z_i = Z(s_i)$

$\mathcal{G} = \{g_\lambda(\cdot) | \lambda \in \Lambda\}$

$\eta = [\beta, \tau, \theta, \lambda]^T$: regression, random field, structural, and transform parameters

$L(\eta; z) = \left(\frac{\tau}{2\pi}\right)^{n/2} |\sigma_\theta|^{-1/2} \exp \left\{ -\frac{\tau}{2} (g_\lambda(z) - X\beta)^T \sigma_\theta^{-1} (g_\lambda(z) - X\beta) \right\} \prod_{i=1}^n |g'_\lambda(z_i)|$: likelihood

2.1: Model Description

One feature of BTG is that it considers a range of sampling distributions instead of a single one, thereby lending itself to "more robust predictive inference" (4.2)

$$Y(s) = g_\lambda(z)$$
$$[g_\lambda(Z_0), g_\lambda(Z)]^T \sim \mathcal{N}_{n+k} \left(\begin{bmatrix} X_0\beta \\ X\beta \end{bmatrix}, \frac{1}{\tau} \begin{bmatrix} E_\theta & B_\theta \\ B_\theta^T & \Sigma_\theta \end{bmatrix} \right)$$

Prior specification:

$$p(\beta, \tau, \theta, \lambda) \propto \frac{p(\theta)p(\lambda)}{\tau J_\lambda^{p/n}}$$

$p(\theta)$ and (λ) are prior marginals of θ and λ . If we assume for the time being that λ is fixed, then this simplifies to

$$p(\beta, \tau, \theta) \propto \frac{p(\theta)p(\lambda)}{\tau}$$

2.2: Posterior of Model Parameters

The treatment of β is that it is an improper distribution, i.e. does not integrate to unity. Rather, the paper allows β to span an unbounded space and take on all values with equal probability.

A least squares problem:

$$\min \|X\beta - g_\lambda(z)\|_{\Sigma_\theta^{-1}}^2$$
$$\hat{\beta}_{\theta, \lambda} = (X^T \Sigma_\theta^{-1} X)^{-1} X^T \Sigma_\theta^{-1} g_\lambda(z)$$

Next for the quadratic form,

$$\tilde{q}_{\theta, \lambda} = (g_\lambda(z) - X\hat{\beta}_{\theta, \lambda}) \Sigma_\theta^{-1} (g_\lambda(z) - X\hat{\beta}_{\theta, \lambda})$$
$$\int_{\Omega} p(z|\eta) p(\eta) d\eta = \int_{\Lambda} \int_{\Theta} |\Sigma_\theta|^{-1/2} |X^T \Sigma_\theta^{-1} X|^{-1/2} \tilde{q}_{\theta, \lambda}^{-(n-p)/2} J_\lambda^{1-(p/n)} p(\theta) p(\lambda) d\theta d\lambda$$

It is true that $p(\beta, \tau|\theta, \lambda, z) = p(\beta|\tau, \theta, \lambda, z)p(\tau|\theta, \lambda, z)$ is normal gamma, i.e.

$$(\beta|\tau, \theta, \lambda, z) \sim \mathcal{N}_p \left(\hat{\beta}_{\theta, \lambda}, \frac{1}{\tau} (X^T \Sigma_\theta^{-1} X)^{-1} \right)$$

$$(\tau|\theta, \lambda, z) \sim \text{Gamma}(\frac{n-p}{2}, \frac{2}{\tilde{q}_{\theta, \lambda}})$$

In addition,

$$p(\theta, \lambda|z) = p(\beta, \tau, \theta, \lambda|z)p(\beta, \tau|\theta, \lambda, z)$$

Applying Bayes theorem to the numerator gives

$$p(\theta, \lambda|z) \propto |\Sigma_\theta|^{-1/2} |X^T \Sigma_\theta^{-1} X|^{-1/2} \tilde{q}_{\theta, \lambda}^{-(n-p)/2} J_\lambda^{1-(p/n)} p(\theta) p(\lambda)$$

The posterior distribution is determined up to a multiplicative constant:

$$p(\beta, \tau, \theta, \lambda|z) = p(\beta, \tau|\theta, \lambda, z)p(\theta, \lambda|z)$$

2.3: Prediction of Z_0

Bayesian predictive density function:

$$p(z_0|z) = \int_{\Omega} p(z_0|\eta, z) p(\eta|z) d\eta$$

Fixing λ :

Appendix

Computing Derivatives

We are interested in computing the derivatives of $p(\theta, \lambda|z)$ for maximum a posteriori estimation of θ, λ . We are also interested in the derivatives of $p(z_0|\theta, \lambda, z)$ so we can compute statistics of interest of the CDF $\Phi(z_0|z)$ and the PDF $p(z_0|z)$, such as the median and narrowest 95% confidence interval.

Some Facts

- $\frac{\partial}{\partial \theta_i} \det(K(\theta)) = \det(K(\theta)) \text{tr} \left(K^{-1}(\theta) \frac{\partial K(\theta)}{\partial \theta_i} \right)$
- $\frac{\partial}{\partial \theta_i} (K(\theta))^{-1} = -K^{-1}(\theta) \frac{\partial K(\theta)}{\partial \theta_i} K^{-1}(\theta)$
- $\frac{\partial}{\partial \theta_i} A(\theta) B(\theta) = A'(\theta) B(\theta) + A(\theta) B'(\theta)$
- $\frac{\partial}{\partial \theta_i} A(\theta) X B(\theta) = A'(\theta) X B(\theta) + A(\theta) X B'(\theta)$

From equation 8 in the paper, we know that

$$p(\theta, \lambda|z) = C \det(\Sigma_\theta)^{-1/2} \det(X^T \Sigma_\theta^{-1} X)^{-1/2} \tilde{q}_{\theta, \lambda}^{-\frac{n-p}{2}} J_\lambda^{1-\frac{p}{n}} p(\theta) p(\lambda)$$

where

$$\tilde{q}_{\theta, \lambda} = (g_\lambda(z) - X \hat{\beta}_{\theta, \lambda}) \Sigma_\theta^{-1} (g_\lambda(z) - X \hat{\beta}_{\theta, \lambda})$$

and

$$\hat{\beta}_{\theta, \lambda} = (X^T \Sigma_\theta^{-1} X)^{-1} X^T \Sigma_\theta^{-1} g_\lambda(z)$$

The convention here is that $\theta \in \mathbb{R}^q$. We seek to compute $\frac{\partial}{\partial \theta_i} p(\theta, \lambda|z)$ and $\frac{\partial}{\partial \lambda} p(\theta, \lambda|z)$.

0.1 Computing $\frac{\partial}{\partial\theta}p(\theta, \lambda|z)$

By the product rule,

$$\frac{\partial}{\partial\theta}p(\theta, \lambda|z) = C \left(\frac{\partial}{\partial\theta}(\det(\Sigma_\theta)^{-1/2}) \right) \det(X^T \Sigma_\theta^{-1} X)^{-1/2} \tilde{q}_{\theta, \lambda}^{-\frac{n-p}{2}} J_\lambda^{1-\frac{p}{n}} p(\theta) p(\lambda) \quad (1)$$

$$+ C \det(\Sigma_\theta)^{-1/2} \left(\frac{\partial}{\partial\theta}(\det(X^T \Sigma_\theta^{-1} X)^{-1/2}) \right) \tilde{q}_{\theta, \lambda}^{-\frac{n-p}{2}} J_\lambda^{1-\frac{p}{n}} p(\theta) p(\lambda) \quad (2)$$

$$+ C \det(\Sigma_\theta)^{-1/2} \det(X^T \Sigma_\theta^{-1} X)^{-1/2} \left(\frac{\partial}{\partial\theta}(\tilde{q}_{\theta, \lambda}^{-\frac{n-p}{2}}) \right) J_\lambda^{1-\frac{p}{n}} p(\theta) p(\lambda) \quad (3)$$

$$+ C \det(\Sigma_\theta)^{-1/2} \det(X^T \Sigma_\theta^{-1} X)^{-1/2} \tilde{q}_{\theta, \lambda}^{-\frac{n-p}{2}} J_\lambda^{1-\frac{p}{n}} \left(\frac{\partial}{\partial\theta} p(\theta) \right) p(\lambda) \quad (4)$$

Each of the derivatives enclosed in large parentheses are computed explicitly below.

Expression (1)

$$\begin{aligned} \frac{\partial}{\partial\theta}(\det(\Sigma_\theta)^{-1/2}) &= -\frac{1}{2} \det(\Sigma_\theta)^{-3/2} \det(\Sigma_\theta) \operatorname{tr}(\Sigma_\theta^{-1} \frac{\partial}{\partial\theta} \Sigma_\theta) \\ &= -\frac{1}{2} \det(\Sigma_\theta)^{-1/2} \operatorname{tr}(\Sigma_\theta^{-1} \frac{\partial}{\partial\theta} \Sigma_\theta) \end{aligned}$$

Expression (2)

$$\begin{aligned} \frac{\partial}{\partial\theta}(\det(X^T \Sigma_\theta^{-1} X)^{-1/2}) &= -\frac{1}{2}(\det(X^T \Sigma_\theta^{-1} X)^{-3/2} \det(X^T \Sigma_\theta^{-1} X) \operatorname{tr}((X^T \Sigma_\theta^{-1} X)^{-1} \frac{\partial}{\partial\theta}(X^T \Sigma_\theta^{-1} X))) \\ &= -\frac{1}{2}(\det(X^T \Sigma_\theta^{-1} X)^{-1/2} \operatorname{tr}((X^T \Sigma_\theta^{-1} X)^{-1} X^T (-\Sigma_\theta^{-1} \frac{\partial \Sigma_\theta}{\partial\theta} \Sigma_\theta^{-1}) X)) \end{aligned}$$

Expression (3)

$$\frac{\partial}{\partial\theta}(\tilde{q}_{\theta, \lambda}^{-\frac{n-p}{2}}) = -\left(\frac{n-p}{2}\right) \tilde{q}_{\theta, \lambda}^{-\frac{n-p+2}{2}} \frac{\partial}{\partial\theta} \tilde{q}_{\theta, \lambda}$$

where

$$\begin{aligned} \frac{\partial}{\partial\theta} \tilde{q}_{\theta, \lambda} &= \frac{\partial}{\partial\theta} \left((g_\lambda(z) - X \hat{\beta}_{\theta, \lambda})^T \Sigma_\theta^{-1} (g_\lambda(z) - X \hat{\beta}_{\theta, \lambda}) \right) \\ &= \left(-X \frac{\partial}{\partial\theta} \hat{\beta}_{\theta, \lambda} \right)^T \Sigma_\theta^{-1} (g_\lambda(z) - X \hat{\beta}_{\theta, \lambda}) \\ &\quad + (g_\lambda(z) - X \hat{\beta}_{\theta, \lambda})^T \left(-\Sigma_\theta^{-1} \frac{\partial \Sigma_\theta}{\partial\theta} \Sigma_\theta^{-1} \right) (g_\lambda(z) - X \hat{\beta}_{\theta, \lambda}) \\ &\quad + (g_\lambda(z) - X \hat{\beta}_{\theta, \lambda})^T \Sigma_\theta^{-1} \left(-X \frac{\partial}{\partial\theta} \hat{\beta}_{\theta, \lambda} \right) \end{aligned}$$

and

$$\begin{aligned} \frac{\partial}{\partial\theta} \hat{\beta}_{\theta, \lambda} &= \frac{\partial}{\partial\theta} \left((X^T \Sigma_\theta^{-1} X)^{-1} X^T \Sigma_\theta^{-1} g_\lambda(z) \right) \\ &= -(X^T \Sigma_\theta^{-1} X)^{-1} \left(-X^T \Sigma_\theta^{-1} \frac{\partial \Sigma_\theta}{\partial\theta} \Sigma_\theta^{-1} X \right) (X^T \Sigma_\theta^{-1} X)^{-1} X^T \Sigma_\theta^{-1} g_\lambda(z) \\ &\quad + (X^T \Sigma_\theta^{-1} X)^{-1} X^T \left(-\Sigma_\theta^{-1} \frac{\partial \Sigma_\theta}{\partial\theta} \Sigma_\theta^{-1} \right) g_\lambda(z) \end{aligned}$$

Expression (4)

Depends on $p(\theta)$, the prior distribution over θ .

0.2 Computing $\frac{\partial}{\partial \lambda} p(\theta, \lambda | z)$

Recall that $g : \mathbb{R}_1 \mapsto \mathbb{R}^1$ and that $g_\lambda(\mathbf{z}) = [g_\lambda(z_1), g_\lambda(z_2), \dots, g_\lambda(z_n)]^T$. By assumption (section 2.1), $g'_\lambda(x) = \frac{\partial}{\partial x} g_\lambda$ exists.

By the product rule,

$$\frac{\partial}{\partial \lambda} p(\theta, \lambda | z) = C \det(\Sigma_\theta)^{-1/2} \det(X^T \Sigma_\theta^{-1} X)^{-1/2} \left(\frac{\partial}{\partial \lambda} \tilde{q}_{\theta, \lambda}^{-\frac{n-p}{2}} \right) J_\lambda^{1-\frac{p}{n}} p(\theta) p(\lambda) \quad (5)$$

$$+ C \det(\Sigma_\theta)^{-1/2} \det(X^T \Sigma_\theta^{-1} X)^{-1/2} \tilde{q}_{\theta, \lambda}^{-\frac{n-p}{2}} \left(\frac{\partial}{\partial \lambda} J_\lambda^{1-\frac{p}{n}} \right) p(\theta) p(\lambda) \quad (6)$$

$$+ C \det(\Sigma_\theta)^{-1/2} \det(X^T \Sigma_\theta^{-1} X)^{-1/2} \frac{\partial}{\partial \lambda} \tilde{q}_{\theta, \lambda}^{-\frac{n-p}{2}} J_\lambda^{1-\frac{p}{n}} p(\theta) \left(\frac{\partial}{\partial \lambda} p(\lambda) \right) \quad (7)$$

The expressions in parentheses are evaluated below.

Computing $\frac{\partial}{\partial \lambda} \tilde{q}_{\theta, \lambda}^{-\frac{n-p}{2}}$

$$\frac{\partial}{\partial \lambda} \tilde{q}_{\theta, \lambda}^{-\frac{n-p}{2}} = - \left(\frac{n-p}{2} \right) \tilde{q}_{\theta, \lambda}^{-\frac{n-p+2}{2}} \frac{\partial}{\partial \lambda} \tilde{q}_{\theta, \lambda}$$

where

$$\begin{aligned} \frac{\partial}{\partial \lambda} \tilde{q}_{\theta, \lambda} &= \left(\frac{\partial}{\partial \lambda} g_\lambda(z) - X \frac{\partial}{\partial \lambda} \hat{\beta}_{\theta, \lambda} \right)^T \Sigma_\theta^{-1} (g_\lambda(z) - X \hat{\beta}_{\theta, \lambda}) \\ &\quad + (g_\lambda(z) - X \hat{\beta}_{\theta, \lambda})^T \Sigma_\theta^{-1} \left(\frac{\partial}{\partial \lambda} g_\lambda(z) - X \frac{\partial}{\partial \lambda} \hat{\beta}_{\theta, \lambda} \right) \end{aligned}$$

and

$$\frac{\partial}{\partial \lambda} \hat{\beta}_{\theta, \lambda} = (X^T \Sigma_\theta^{-1} X)^{-1} X^T \Sigma_\theta^{-1} \frac{\partial}{\partial \lambda} g_\lambda(z)$$

Computing $\frac{\partial}{\partial \lambda} J_\lambda^{1-\frac{p}{n}}$

By definition, $J_\lambda = \prod_{i=1}^n |g'_\lambda(z_i)|$. By assumption (section 2.1), g_λ is assumed to be monotone, so

$$J_\lambda = \begin{cases} (-1)^n \prod_{i=1}^n g'_\lambda(z_i) & g(\cdot) \text{ monotone decreasing} \\ \prod_{i=1}^n g'_\lambda(z_i) & g(\cdot) \text{ monotone increasing} \end{cases}$$

Now the derivative of the expression is equal to

$$\frac{\partial}{\partial \lambda} J_\lambda^{1-\frac{p}{n}} = \left(1 - \frac{p}{n} \right) J_\lambda^{-\frac{p}{n}} \frac{\partial}{\partial \lambda} J_\lambda$$

where

$$\frac{\partial}{\partial \lambda} J_\lambda = \begin{cases} (-1)^n \sum_{j=1}^n \frac{\partial}{\partial \lambda} g'_\lambda(z_j) \prod_{i \neq j, i=1}^n g'_\lambda(z_i) \\ \sum_{j=1}^n \frac{\partial}{\partial \lambda} g'_\lambda(z_j) \prod_{i \neq j, i=1}^n g'_\lambda(z_i) \end{cases}$$

Computing $\frac{\partial}{\partial \lambda} p(\lambda)$

Depends on $p(\lambda)$

0.3 Computing $\frac{\partial}{\partial z_0} p(z_0 | \theta, \lambda, \mathbf{z})$

From equation (12),

$$p(z_0 | \theta, \lambda, \mathbf{z}) = \frac{\Gamma\left(\frac{n-p+k}{2}\right) \prod_{j=1}^k |g'_\lambda(z_{0j})|}{\Gamma\left(\frac{n-p}{2}\right) \pi^{k/2} |\tilde{q}_{\theta,\lambda} \mathbf{C}_\theta|^{1/2}} [1 + (g_\lambda(z_0) - \mathbf{m}_{\theta,\lambda})^T (\tilde{q}_{\theta,\lambda} \mathbf{C}_\theta)^{-1} (g_\lambda(z_0) - \mathbf{m}_{\theta,\lambda})]^{-\frac{n-p+k}{2}}$$

where we write for convenience:

$$\mathbf{m}_{\theta,\lambda} = \mathbf{B}_\theta \Sigma_\theta^{-1} g_\lambda(\mathbf{z}) + \mathbf{H}_\theta \hat{\beta}_{\theta,\lambda}$$

$$\mathbf{H}_\theta = \mathbf{X}_0 - \mathbf{B}_\theta \Sigma_\theta^{-1} \mathbf{X}$$

$$\mathbf{C}_\theta = \mathbf{D}_\theta + \mathbf{H}_\theta (\mathbf{X}^T \Sigma_\theta^{-1} \mathbf{X})^{-1} \mathbf{H}_\theta^T$$

$$\mathbf{D}_\theta = \mathbf{E}_\theta - \mathbf{B}_\theta \Sigma_\theta^{-1} \mathbf{B}_\theta^T$$

and \mathbf{B}_θ and \mathbf{E}_θ are covariances matrices from

$$(g_\lambda(Z_0), g_\lambda(Z) | \beta, \tau, \theta, \lambda) \sim \left(\begin{pmatrix} X_0 \beta \\ X \beta \end{pmatrix}, \frac{1}{\tau} \begin{pmatrix} \mathbf{E}_\theta & \mathbf{B}_\theta \\ \mathbf{B}_\theta^T & \Sigma_\theta \end{pmatrix} \right)$$

The function $p(z_0 | \theta, \lambda, \mathbf{z})$ is a function from \mathbb{R}^n to \mathbb{R}^1 . The derivative should be equal to the Jacobian under continuity assumptions. The i th entry of the Jacobian is $\frac{\partial}{\partial z_{0i}} p(z_0 | \theta, \lambda, \mathbf{z})$.

Ignoring the constant terms, we have that

$$\begin{aligned} \frac{\partial}{\partial z_{0i}} p(z_0 | \theta, \lambda, \mathbf{z}) &= \left(\frac{\partial}{\partial z_{0i}} J_\lambda(z_0) \right) [1 + (g_\lambda(z_0) - \mathbf{m}_{\theta,\lambda})^T (\tilde{q}_{\theta,\lambda} \mathbf{C}_\theta)^{-1} (g_\lambda(z_0) - \mathbf{m}_{\theta,\lambda})]^{-\frac{n-p+k}{2}} \\ &+ J_\lambda(z_0) \left(-\frac{n-p+k}{2} \right) (1 + (g_\lambda(z_0) - \mathbf{m}_{\theta,\lambda})^T (\tilde{q}_{\theta,\lambda} \mathbf{C}_\theta)^{-1} (g_\lambda(z_0) - \mathbf{m}_{\theta,\lambda}))^{-\frac{n-p+k+2}{2}} \\ &\left\{ \left(\frac{\partial}{\partial z_{0i}} g_\lambda(z_0) \right)^T (\tilde{q}_{\theta,\lambda} \mathbf{C}_\theta)^{-1} (g_\lambda(z_0) - \mathbf{m}_{\theta,\lambda}) + (g_\lambda(z_0) - \mathbf{m}_{\theta,\lambda})^T (\tilde{q}_{\theta,\lambda} \mathbf{C}_\theta)^{-1} \left(\frac{\partial}{\partial z_{0i}} g_\lambda(z_0) \right) \right\} \end{aligned}$$

Since

$$J_\lambda(z_0) = \begin{cases} (-1)^n \prod_{i=1}^n g'_\lambda(z_{i0}) & g(\cdot) \text{ monotone decreasing} \\ \prod_{i=1}^n g'_\lambda(z_{i0}) & g(\cdot) \text{ monotone increasing} \end{cases}$$

we have

$$\frac{\partial}{\partial z_{0i}} J_\lambda = \begin{cases} (-1)^n \frac{\partial}{\partial z_{0i}} g''_\lambda(z_{i0}) \prod_{j=1, j \neq i}^n g'_\lambda(z_{j0}) & g(\cdot) \text{ monotone decreasing} \\ \frac{\partial}{\partial z_{0i}} g''_\lambda(z_{i,0}) \prod_{j=1, j \neq i}^n g'_\lambda(z_{j0}) & g(\cdot) \text{ monotone increasing} \end{cases}$$

Also, be definition,

$$\frac{\partial}{\partial z_{0i}} g_\lambda(z_0) = g'_\lambda(z_{0i})$$

0.4 Computing $\frac{\partial}{\partial \theta} p(z_0 | \theta, \lambda, \mathbf{z})$

We first compile some useful derivatives:

$$\frac{\partial}{\partial \theta} \mathbf{m}_{\theta, \lambda} = \frac{\partial}{\partial \theta} \mathbf{B}_{\theta} \Sigma_{\theta}^{-1} g_{\lambda}(z) - \mathbf{B}_{\theta} \Sigma_{\theta}^{-1} \frac{\partial \Sigma_{\theta}}{\partial \theta} \Sigma_{\theta}^{-1} g_{\lambda}(z) + \frac{\partial}{\partial \theta} \mathbf{H}_{\theta} \hat{\beta}_{\theta, \lambda} + \mathbf{H}_{\theta} \frac{\partial}{\partial \theta} \hat{\beta}_{\theta, \lambda}$$

$$\frac{\partial}{\partial \theta} \mathbf{H}_{\theta} = -\frac{\partial}{\partial \theta} \mathbf{B}_{\theta} \Sigma_{\theta}^{-1} X + \mathbf{B}_{\theta} \Sigma_{\theta}^{-1} \frac{\partial \Sigma_{\theta}}{\partial \theta} \Sigma_{\theta}^{-1} X$$

$$\begin{aligned} \frac{\partial}{\partial \theta} \mathbf{C}_{\theta} &= \frac{\partial}{\partial \theta} \mathbf{D}_{\theta} + \frac{\partial}{\partial \theta} \mathbf{H}_{\theta} (\mathbf{X}^T \Sigma_{\theta}^{-1} \mathbf{X})^{-1} \mathbf{H}_{\theta}^T + \mathbf{H}_{\theta} (\mathbf{X}^T \Sigma_{\theta}^{-1} \mathbf{X})^{-1} \left(\mathbf{X}^T \Sigma_{\theta}^{-1} \frac{\partial \Sigma_{\theta}}{\partial \theta} \Sigma_{\theta}^{-1} \mathbf{X} \right) (\mathbf{X}^T \Sigma_{\theta}^{-1} \mathbf{X})^{-1} \mathbf{H}_{\theta}^T \\ &\quad + \mathbf{H}_{\theta} (\mathbf{X}^T \Sigma_{\theta}^{-1} \mathbf{X})^{-1} \frac{\partial}{\partial \theta} \mathbf{H}_{\theta}^T \end{aligned}$$

$$\frac{\partial}{\partial \theta} \mathbf{D}_{\theta} = \frac{\partial}{\partial \theta} \mathbf{E}_{\theta} - \frac{\partial}{\partial \theta} \mathbf{B}_{\theta} \Sigma_{\theta}^{-1} \mathbf{B}_{\theta}^T + \mathbf{B}_{\theta} \Sigma_{\theta}^{-1} \frac{\partial \Sigma_{\theta}}{\partial \theta} \Sigma_{\theta}^{-1} \mathbf{B}_{\theta}^T - \mathbf{B}_{\theta} \Sigma_{\theta}^{-1} \frac{\partial}{\partial \theta} \mathbf{B}_{\theta}^T$$

Note that $\frac{\partial}{\partial \theta} \hat{\beta}_{\theta, \lambda}$ were computed previously. Abstracting the parts of equation (12) which don't depend on θ , we are left with

$$g(\theta) := \det(\tilde{q}_{\theta, \lambda} \mathbf{C}_{\theta})^{-1/2} [1 + (g_{\lambda}(z_0) - \mathbf{m}_{\theta, \lambda})^T (\tilde{q}_{\theta, \lambda} \mathbf{C}_{\theta})^{-1} (g_{\lambda}(z_0) - \mathbf{m}_{\theta, \lambda})]^{-\frac{n-p+k}{2}}$$

We have

$$\begin{aligned} \frac{\partial}{\partial \theta} g(\theta) &= \left(\frac{\partial}{\partial \theta} \det(\tilde{q}_{\theta, \lambda} \mathbf{C}_{\theta})^{-1/2} \right) [1 + (g_{\lambda}(z_0) - \mathbf{m}_{\theta, \lambda})^T (\tilde{q}_{\theta, \lambda} \mathbf{C}_{\theta})^{-1} (g_{\lambda}(z_0) - \mathbf{m}_{\theta, \lambda})]^{-\frac{n-p+k}{2}} \\ &\quad + \det(\tilde{q}_{\theta, \lambda} \mathbf{C}_{\theta})^{-1/2} \left(-\frac{n-p+k}{2} \right) (1 + (g_{\lambda}(z_0) - \mathbf{m}_{\theta, \lambda})^T (\tilde{q}_{\theta, \lambda} \mathbf{C}_{\theta})^{-1} (g_{\lambda}(z_0) - \mathbf{m}_{\theta, \lambda}))^{-\frac{n-p+k+2}{2}} \\ &\quad \left(\frac{\partial}{\partial \theta} ((g_{\lambda}(z_0) - \mathbf{m}_{\theta, \lambda})^T (\tilde{q}_{\theta, \lambda} \mathbf{C}_{\theta})^{-1} (g_{\lambda}(z_0) - \mathbf{m}_{\theta, \lambda})) \right) \end{aligned}$$

For the derivative of the determinant, we get

$$\begin{aligned} \frac{\partial}{\partial \theta} \det(\tilde{q}_{\theta, \lambda} \mathbf{C}_{\theta})^{-1/2} &= -\frac{1}{2} \det(\tilde{q}_{\theta, \lambda} \mathbf{C}_{\theta})^{-3/2} \det(\tilde{q}_{\theta, \lambda} \mathbf{C}_{\theta}) \operatorname{tr} \left((\tilde{q}_{\theta, \lambda} \mathbf{C}_{\theta})^{-1} \left(\left(\frac{\partial}{\partial \theta} \tilde{q}_{\theta, \lambda} \right) \mathbf{C}_{\theta} + \tilde{q}_{\theta, \lambda} \frac{\partial}{\partial \theta} \mathbf{C}_{\theta} \right) \right) \\ &= -\frac{1}{2} \det(\tilde{q}_{\theta, \lambda} \mathbf{C}_{\theta})^{-1/2} \operatorname{tr} \left((\tilde{q}_{\theta, \lambda} \mathbf{C}_{\theta})^{-1} \left(\left(\frac{\partial}{\partial \theta} \tilde{q}_{\theta, \lambda} \right) \mathbf{C}_{\theta} + \tilde{q}_{\theta, \lambda} \frac{\partial}{\partial \theta} \mathbf{C}_{\theta} \right) \right) \end{aligned}$$

where $\frac{\partial}{\partial \theta} \tilde{q}_{\theta, \lambda}$ was computed previously. For the second derivative, we have

$$\begin{aligned} \frac{\partial}{\partial \theta} ((g_{\lambda}(z_0) - \mathbf{m}_{\theta, \lambda})^T (\tilde{q}_{\theta, \lambda} \mathbf{C}_{\theta})^{-1} (g_{\lambda}(z_0) - \mathbf{m}_{\theta, \lambda})) &= - \left(\frac{\partial}{\partial \theta} \mathbf{m}_{\theta, \lambda} \right)^T (\tilde{q}_{\theta, \lambda} \mathbf{C}_{\theta})^{-1} (g_{\lambda}(z_0) - \mathbf{m}_{\theta, \lambda}) \\ &\quad + (g_{\lambda}(z_0) - \mathbf{m}_{\theta, \lambda})^T \left(\frac{\partial}{\partial \theta} (\tilde{q}_{\theta, \lambda} \mathbf{C}_{\theta})^{-1} \right) (g_{\lambda}(z_0) - \mathbf{m}_{\theta, \lambda}) \\ &\quad - (g_{\lambda}(z_0) - \mathbf{m}_{\theta, \lambda})^T (\tilde{q}_{\theta, \lambda} \mathbf{C}_{\theta})^{-1} \left(\frac{\partial}{\partial \theta} \mathbf{m}_{\theta, \lambda} \right) \end{aligned}$$

where

$$\frac{\partial}{\partial \theta} (\tilde{q}_{\theta, \lambda} \mathbf{C}_{\theta})^{-1} = -(\tilde{q}_{\theta, \lambda} \mathbf{C}_{\theta})^{-1} \left(\frac{\partial}{\partial \theta} \tilde{q}_{\theta, \lambda} \mathbf{C}_{\theta} + \tilde{q}_{\theta, \lambda} \frac{\partial}{\partial \theta} \mathbf{C}_{\theta} \right) (\tilde{q}_{\theta, \lambda} \mathbf{C}_{\theta})^{-1}$$

0.5 Computing $\frac{\partial}{\partial \lambda} p(z_0 | \theta, \lambda, \mathbf{z})$

The part of equation (12) that depends on λ is

$$h(\lambda) := \underbrace{\left(\prod_{j=1}^k |g'_\lambda(z_{0j})| \right)}_{\text{Expr. 1}} \underbrace{\det(\tilde{q}_{\theta,\lambda} \mathbf{C}_\theta)^{-1/2}}_{\text{Expr. 2}} \underbrace{[1 + (g_\lambda(z_0) - \mathbf{m}_{\theta,\lambda})^T (\tilde{q}_{\theta,\lambda} \mathbf{C}_\theta)^{-1} (g_\lambda(z_0) - \mathbf{m}_{\theta,\lambda})]^{-\frac{n-p+k}{2}}}_{\text{Expr. 3}}$$

For expression 1, we have

$$\frac{\partial}{\partial \lambda} \prod_{j=1}^k |g'_\lambda(z_{0j})| = \begin{cases} (-1)^n \sum_{j=1}^n \frac{\partial}{\partial \lambda} g'(z_{j0}) \prod_{i \neq j, i=1}^n g'_\lambda(z_{i0}) & g(\cdot) \text{ monotone decreasing} \\ \sum_{j=1}^n \frac{\partial}{\partial \lambda} g'(z_{j0}) \prod_{i \neq j, i=1}^n g'_\lambda(z_{i0}) & g(\cdot) \text{ monotone increasing} \end{cases}$$

For expression 2, we have

$$\frac{\partial}{\partial \lambda} \det(\tilde{q}_{\theta,\lambda} \mathbf{C}_\theta)^{-1/2} = -\frac{1}{2} \det(\tilde{q}_{\theta,\lambda} \mathbf{C}_\theta)^{-1/2} \text{tr}((\tilde{q}_{\theta,\lambda} \mathbf{C}_\theta)^{-1} \frac{\partial}{\partial \lambda} (\tilde{q}_{\theta,\lambda} \mathbf{C}_\theta))$$

where $\frac{\partial}{\partial \lambda} \tilde{q}_{\theta,\lambda}$ was computed previously. For expression 3, we obtain

$$\begin{aligned} & \left(-\frac{n-p+k}{2} \right) [1 + (g_\lambda(z_0) - \mathbf{m}_{\theta,\lambda})^T (\tilde{q}_{\theta,\lambda} \mathbf{C}_\theta)^{-1} (g_\lambda(z_0) - \mathbf{m}_{\theta,\lambda})]^{-\frac{n-p+k+2}{2}} \\ & \frac{\partial}{\partial \lambda} ((g_\lambda(z_0) - \mathbf{m}_{\theta,\lambda})^T (\tilde{q}_{\theta,\lambda} \mathbf{C}_\theta)^{-1} (g_\lambda(z_0) - \mathbf{m}_{\theta,\lambda})) \end{aligned}$$

where

$$\begin{aligned} \frac{\partial}{\partial \lambda} ((g_\lambda(z_0) - \mathbf{m}_{\theta,\lambda})^T (\tilde{q}_{\theta,\lambda} \mathbf{C}_\theta)^{-1} (g_\lambda(z_0) - \mathbf{m}_{\theta,\lambda})) &= \left(\frac{\partial}{\partial \lambda} g_\lambda(z_0) - \frac{\partial}{\partial \lambda} \mathbf{m}_{\theta,\lambda} \right)^T (\tilde{q}_{\theta,\lambda} \mathbf{C}_\theta)^{-1} (g_\lambda(z_0) - \mathbf{m}_{\theta,\lambda}) \\ &+ (g_\lambda(z_0) - \mathbf{m}_{\theta,\lambda})^T \left(\frac{\partial}{\partial \lambda} (\tilde{q}_{\theta,\lambda} \mathbf{C}_\theta)^{-1} \right) (g_\lambda(z_0) - \mathbf{m}_{\theta,\lambda}) \\ &+ (g_\lambda(z_0) - \mathbf{m}_{\theta,\lambda})^T (\tilde{q}_{\theta,\lambda} \mathbf{C}_\theta)^{-1} \left(\frac{\partial}{\partial \lambda} g_\lambda(z_0) - \frac{\partial}{\partial \lambda} \mathbf{m}_{\theta,\lambda} \right) \end{aligned}$$

where

$$\frac{\partial}{\partial \lambda} (\tilde{q}_{\theta,\lambda} \mathbf{C}_\theta)^{-1} = -(\tilde{q}_{\theta,\lambda} \mathbf{C}_\theta)^{-1} \left(\frac{\partial}{\partial \lambda} \tilde{q}_{\theta,\lambda} \mathbf{C}_\theta \right) (\tilde{q}_{\theta,\lambda} \mathbf{C}_\theta)^{-1}$$

and

$$\frac{\partial}{\partial \lambda} \mathbf{m}_{\theta,\lambda} = \mathbf{B}_\theta \Sigma_\theta^{-1} \frac{\partial}{\partial \lambda} g_\lambda(z)$$

For convenience, partials of $\tilde{q}_{\lambda,\theta}$ and $\hat{\beta}_{\theta,\lambda}$ are repeated here:

$$\frac{\partial}{\partial \lambda} \tilde{q}_{\theta,\lambda} = \left(\frac{\partial}{\partial \lambda} g_\lambda(z) - X \frac{\partial}{\partial \lambda} \hat{\beta}_{\theta,\lambda} \right)^T \Sigma_\theta^{-1} (g_\lambda(z) - X \hat{\beta}_{\theta,\lambda}) + (g_\lambda(z) - X \hat{\beta}_{\theta,\lambda})^T \Sigma_\theta^{-1} \left(\frac{\partial}{\partial \lambda} g_\lambda(z) - X \frac{\partial}{\partial \lambda} \hat{\beta}_{\theta,\lambda} \right)$$

$$\frac{\partial}{\partial \lambda} \hat{\beta}_{\theta,\lambda} = (X^T \Sigma_\theta^{-1} X)^{-1} X^T \Sigma_\theta^{-1} \frac{\partial}{\partial \lambda} g_\lambda(z)$$

0.5.1 Some Derivatives of the Box Cox Transform

Note that the derivative of the Box Cox Transform w.r.t λ is

$$\frac{\partial}{\partial \lambda} g_{\lambda}(z) = \begin{cases} \frac{\lambda x^{\lambda} \ln x - x^{\lambda} + 1}{\lambda^2} & \lambda \neq 0 \\ 0 & \lambda = 0 \end{cases}$$

The second derivative is given by

$$\frac{\partial^2}{\partial \lambda^2} g_{\lambda}(z) = \begin{cases} \frac{x^{\lambda} (\ln x)^2}{\lambda} - \frac{2x^{\lambda} \ln x}{\lambda^2} + \frac{2(x^{\lambda} - 1)}{\lambda^3} & \lambda \neq 0 \\ 0 & \lambda = 0 \end{cases}$$

References

- Bayesian Transformed Gaussian (De Oliveria, Kedem, and Short)
- Orthogonal Polynomials: Computation and Approximation by Gautschi