

# Hierarchical effects of contrast and motion coherence in early visual cortex

Dan Birman, Justin Gardner

Department of Psychology, Stanford University  
danbirman@stanford.edu

## 1. Introduction



An existing model of **contrast** discrimination suggests early visual cortex is sufficient to explain behavioral performance<sup>1</sup>.




Can that approach be used to jointly explain **motion coherence** discrimination?

## 3. Discrimination task


We collected data on how well participants could discriminate small increments in contrast and motion coherence.




## 2. Model: neural responses





**Contrast** and **motion coherence** response functions constrained by behavior (n=21)



**V1 contrast** and **MT motion coherence** response functions constrained by fMRI (n=11)




## 5. Response functions




## 6. Behavior to fMRI

We fit a neural noise<sup>5</sup> parameter to scale response models into the same space. Based on previous work<sup>1</sup> we fit this parameter using the functions constrained on V1 and contrast discrimination.




Model comparison of behavioral **contrast** and **motion coherence** response models to V1 and MT, respectively




## 4. Cortical measurements (fMRI)


Cortical responses were measured during discrimination and passive observation while performing a fixation task.



V1



MT



All data shows bootstrapped mean  $\pm$  95% CI across subjects. Non-linear response functions were modeled using an exponential that can interpolate between linear and non-linear responses:  $Response(s) = \alpha - \alpha e^{-\kappa s}$

1. Boynton, G. M., Demb, J. B., Glover, G. H., & Heeger, D. J. *Vision Research* (1999).  
3. V5/MT is thought to respond linearly to increasing motion coherence (see also 4). Rees, G., Friston, K., & Koch, C. *Nature neuroscience* (2000).

4. Simoncelli, E. P., & Heeger, D. J. *Vision Research* (1998).  
5. Previous reported values for neural noise in a similar model of contrast discrimination were 0.064% and 0.016% for distributed and focal attention. Pestilli, F., Carrasco, M., Heeger, D. J., & Gardner, J. L. *Neuron* (2011).