

ler : LVK (LIGO-Virgo-KAGRA collaboration) Event (compact-binary mergers) Rate calculator and simulator

Phurailatpam Hemantakumar¹, Anupreeta More^{2,3}, Harsh Narola^{4,5}, Ng Chung Yin (Leo)¹, Justin Janquart^{4,5}, Chris Van Den Broeck^{4,5}, Otto Akseli HANNUKSELA¹, Neha Singh⁶, and David Keitel⁶

1 Department of Physics, The Chinese University of Hong Kong, Shatin, New Territories, Hong Kong
2 The Inter-University Centre for Astronomy and Astrophysics (IUCAA), Post Bag 4, Ganeshkhind, Pune 411007, India
3 Kavli Institute for the Physics and Mathematics of the Universe (IPMU), 5-1-5 Kashiwanoha, Kashiwa-shi, Chiba 277-8583, Japan
4 Department of Physics, Utrecht University, Princetonplein 1, 3584 CC Utrecht, The Netherlands
5 Nikhef – National Institute for Subatomic Physics, Science Park, 1098 XG Amsterdam, The Netherlands
6 Departament de Física, Universitat de les Illes Balears, IAC3-IEEC, Crta. Valldemossa km 7.5, E-07122 Palma, Spain

DOI: [10.21105/joss.03000](https://doi.org/10.21105/joss.03000)

Software

- [Review](#) ↗
- [Repository](#) ↗
- [Archive](#) ↗

Editor: [Daniel S. Katz](#) ↗

Reviewers:

- [@khanx169](#)
- [@RobertRosca](#)

Submitted: 26 January 2021

Published: 06 April 2021

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

Summary

Gravitational waves (GWs) are ripples in the fabric of space and time caused by the acceleration of unevenly distributed mass or masses. Observable GWs are created especially during the violent cosmic events of merging compact binaries, such as ‘binary black holes’ (BBH), ‘binary neutron stars’ (BNS), and ‘neutron star and black hole pair’ (NSBH). GWs emitted by these events can be distorted or magnified by the gravitational fields of massive objects such as galaxies or galaxy clusters, a phenomenon known as gravitational lensing. Profound comprehension of gravitational lensing’s impact on GW signals is imperative to their accurate interpretation and the extraction of astrophysical insights therein. For this purpose, statistical modelling of GWs lensing can provide valuable insights into the properties of the lensing objects and GW sources. Such statistics require accurate and efficient means to calculate the detectable lensing rates, which depend on up-to-date modeling and implementation of lens and source properties and their distribution. The outcomes of these computational analyses not only contribute to generating dependable forecasts but also play an important role in validating forthcoming lensing events (Janquart et al. 2023).

Obtaining precise outcomes in statistical analyses of this nature necessitates the utilization of large-scale sampling, often numbering in the millions. However, this process is computationally demanding. The *ler* framework addresses this by employing innovative techniques to optimize the workflow and computation efficiency required for handling large-scale statistical analyses, essential for modeling detectable events and calculating rates. Its integration of modular statistical components enhances the framework’s adaptability and extendability, thus proving to be an invaluable asset in the evolving field of gravitational wave research. Detailed description, source code, and examples are available in *ler* [documentation](#).

Statement of need

ler is a statistical-based Python package specifically designed for computing detectable rates of both lensed and unlensed gravitational wave (GW) events, catering to the requirements of the LIGO-Virgo-KAGRA Scientific Collaboration and astrophysics research scholars. The core functionality of *ler* intricately hinges upon the interplay of various components which include sampling the properties of compact-binary sources, lens galaxies characteristics, solving lens equations to derive properties of resultant images, and computing detectable GW rates. This comprehensive functionality builds on the leveraging of array operations and linear algebra from the *numpy* (NumPy Community 2022) library, enhanced by interpolation methods from

scipy (Virtanen et al. 2020) and Python’s *multiprocessing* capabilities. Efficiency is further boosted by the *numba* (Lam, Pitrou, and Seibert 2022) library’s Just-In-Time (*njit*) compilation, optimizing extensive numerical computations and employing the inverse transform sampling method to replace more cumbersome rejection sampling. The modular design of *ler* not only optimizes speed and functionality but also ensures adaptability and upgradability, supporting the integration of additional statistics as research evolves. Currently, *ler* is an important tool in generating simulated gravitational wave (GW) events—both lensed and unlensed—and provides astrophysically accurate distributions of event-related parameters for both detectable and non-detectable events. This functionality aids in event validation and enhances the forecasting of detection capabilities across various GW detectors to study such events. The architecture of the *ler* API facilitates seamless compatibility with other software packages, allowing researchers to integrate and utilize its functionalities based on specific scientific requirements.

Design and Structure

The architecture of the *ler* API is deliberately organized such that each distinct functionality holds its own significance in scientific research. Simultaneously, these functionalities seamlessly integrate and can be employed collectively to accommodate diverse scientific objectives. Key features of *ler* and its dependencies can be summarized as follows:

- Sampling Gravitational Wave (GW) Source Properties:
 - For the unlensed events, The sampling distribution ($R_m(z_s)$) for the source’s redshift (z_s) is derived from the merger rate density of compact binaries, which, in turn, is based on the star formation rate. The code is meticulously designed to enable the straightforward integration of future updates or user-specified distributions of these sources.
 - Intrinsic and extrinsic parameters (θ) of GW sources are sampled using prior distributions ($P(\theta)$) from the *gwcosmo* (gwcosmo Contributors 2022) and *bilby* (Ashton2019?) packages, with options for users to input custom distributions.
- Sampling of lens galaxies attributes and source red-shifts:
 - For lensed case, the source redshift (z_s) is sampled under the strong lensing condition (SL), based on the precomputed probability of strong lensing with source at z_s (optical depth: $P(\text{SL}|z_s)$ or $\tau(z_s)$). This probability can be recalculated for specified configurations of lens galaxies, leveraging *multiprocessing* and *njit* functionalities for enhanced efficiency.
 - The package utilizes the Elliptical Power Law with external shear (EPL+Shear) model (Wempe et al. 2022) for galaxy parameters (θ_L) sampling, following Wierda et al. (2021). Rejection sampling is applied on the above samples on condition that whether the event is strongly lensed or not, $P(\text{SL}|z_s, \theta_L)$.
- Generation of image properties:
 - Source position (β) is sampled from the caustic in the source plane.
 - Sampled lens properties and source position is fed in *lenstronomy* (Birrer et al. 2021) to generate properties of the images. This is the slowest part of the entire simulation, which *ler* tackles through parallelization with multiprocessing.
 - Image properties like magnification (μ_i) and time delay (Δt_i) modifies the original source signal strength, changing the signal-to-noise ratio SNR and our ability to detect.
- Calculation of Detectable Merger Rates Per Year:
 - The calculation of rates necessitates integration over simulated events that meet specific detection criteria. This process includes computing SNRs (ρ) for each event or its lensed images, followed by an assessment against a predetermined threshold(s) (ρ_{th}).
 - SNR calculations are optimized using *gwsnr*, leveraging interpolation and multiprocessing for accuracy and speed.
 - Simulated events and rate results, along with input configurations, are systematically

archived for easy access and future analysis. Additionally, all interpolators used in the process are preserved for future applications.

- Most cosmology-related calculations within the *ler* package are performed using the *astropy* library (Astropy Collaboration 2013). The default cosmological model is LambdaCDM ($H_0 = 70$, $\Omega_m = 0.3$, $\Omega_\Lambda = 0.7$); however, users have the flexibility to employ any cosmology available in *astropy*. All internal calculations in *ler* will then be based on the user-selected cosmological model.

Equations

Detectable Unlensed rates:

$$R_U = \int dz_s \frac{dV_c}{dz_s} \frac{R_m(z_s)}{1+z_s} \{ \Theta[\rho(z_s, \theta) - \rho_{th}] P(\theta) d\theta \}$$

z_s : GW source redshift, $\frac{dV_c}{dz_s}$: Differential co-moving volume, $\frac{1}{1+z_s}$: Time dilation correction factor, $R_m(z_s)$: source frame merger rate density, θ : GW source parameters, P : probability distribution, ρ : SNR, ρ_{th} : SNR threshold, Θ : Heaviside function to select detectable GW events.

Detectable Lensed rates:

$$R_L = \int dz_s \frac{dV_c}{dz_s} \tau(z_s) \frac{R_m(z_s)}{1+z_s} \mathcal{O}_{images}(z_s, \theta, \mu_i, \Delta t_i, \rho_{th}) P(\theta) P(\theta_L | SL, z_s) P(\beta | SL) d\theta d\beta d\theta_L dz_s$$

$\tau(z_s)$: Optical-depth of strong lensing, θ_L : lens parameters, β : source position, μ : image magnification, Δt : image time delay, \mathcal{O} : operator to select detectable lensed GW events, i : index of images of a lensed event, SL: strong lensing condition.

Acknowledgements

The authors express their sincere appreciation for the significant contributions that have been instrumental in completing this research. Special thanks are extended to the academic advisors for their invaluable guidance and steadfast support. The collaborative efforts and enriching discussions with research colleagues significantly enhanced the study's quality. Acknowledgement is given to the Department of Physics, The Chinese University of Hong Kong, for the Postgraduate Studentship that facilitated this research. Further gratitude is extended to the Netherlands Organisation for Scientific Research (NWO) for their support. The authors also recognize the contributions of individuals who added empirical depth to this work. Appreciation is conveyed for the computational resources provided by the LIGO Laboratory, supported by National Science Foundation Grants No. PHY-0757058 and No. PHY-0823459.

References

- [ref-astropy] Astropy Collaboration. 2013. "Astropy: A community Python package for astronomy." *Astronomy and Astrophysics* 558 (October). <https://doi.org/10.1051/0004-6361/201322068>.

Birrer, Simon, Anowar J. Shajib, Daniel Gilman, Aymeric Galan, Jelle Aalbers, Martin Millon, Robert Morgan, et al. 2021. "Lenstronomy II: A Gravitational Lensing Software Ecosystem." *Journal of Open Source Software* 6 (62): 3283. <https://doi.org/10.21105/joss.03283>.

gwcosmo Contributors. 2022. "gwcosmo: A Python package for gravitational-wave cosmology." *GitHub Repository*. GitHub. <https://github.com/gwcosmo/gwcosmo>.

Janquart, J, M Wright, S Goyal, J C L Chan, A Ganguly, Á Garrón, D Keitel, et al. 2023. "Follow-up analyses to the O3 LIGO–Virgo–KAGRA lensing searches." *Monthly Notices of the Royal Astronomical Society* 526 (3): 3832–60. <https://doi.org/10.1093/mnras/stad2909>.

Lam, Stan, Stéphane Pitrou, and Mark Seibert. 2022. "Numba: A High Performance Python Compiler." *Numba Documentation*. Anaconda, Inc. <https://numba.pydata.org/>.

NumPy Community. 2022. "NumPy: A Fundamental Package for Scientific Computing with Python." *NumPy Website*. NumPy. <https://numpy.org/>.

Virtanen, Pauli, Ralf Gommers, Travis E. Oliphant, Matt Haberland, Tyler Reddy, David Cournapeau, Evgeni Burovski, et al. 2020. "SciPy 1.0: Fundamental Algorithms for Scientific Computing in Python." *Nature Methods*. SciPy. <https://www.scipy.org/>.

Wempe, Ewoud, Léon V. E. Koopmans, A. Renske A. C. Wierda, Otto Akseli Hannuksela, Alberto Agnello, Cyril Bonvin, Bendetta Bucciarelli, et al. 2022. "A Lensing Multi-Messenger Channel: Combining LIGO-Virgo-Kagra Lensed Gravitational-Wave Measurements with Euclid Observations." <https://arxiv.org/abs/2204.08732>.

Wierda, A. Renske A. C., Ewoud Wempe, Otto A. Hannuksela, Léon V. E. Koopmans, Alberto Agnello, Cyril Bonvin, Bendetta Bucciarelli, et al. 2021. "Beyond the Detector Horizon: Forecasting Gravitational-Wave Strong Lensing." *The Astrophysical Journal* 921 (1): 154. <https://doi.org/10.3847/1538-4357/ac1bb4>.