

- tion and relapse of childhood B-lineage acute lymphoblastic leukemia. *Blood* 83:1355-1362, 1994
41. Storb R, Etzioni R, Anasetti C, et al: Cyclophosphamide combined with antithymocyte globulin in preparation of allogeneic marrow transplants in patients with aplastic anemia. *Blood* 84:941-949, 1994
 42. Suttorp M, Schmitz N, Dreger P, et al: Monitoring of chimerism after allogeneic bone marrow transplantation with unmanipulated marrow by use of DNA polymorphisms. *Leukemia* 7:679-687, 1993
 43. Tanaka J, Kasai M, Imamura M, et al: Evaluation of mixed chimerism by two-step polymerase chain reaction amplification of hypervariable region MCT118 after allogeneic bone marrow transplantation. *Ann Hematol* 68:189-193, 1994
 44. Ugozzoli L, Yam P, Petz LD, et al: Amplification by the polymerase chain reaction of hypervariable regions of the human genome for evaluation of chimerism after bone marrow transplantation. *Blood* 77:1607-1615, 1991
 45. van Leeuwen JEM, van Tol MJD, Joosten AM, et al: Mixed T-lymphoid chimerism after allogeneic bone marrow transplantation for hematologic malignancies of children is not correlated with relapse. *Blood* 82:1921-1928, 1993
 46. van Leeuwen JEM, van Tol MJD, Joosten AM, et al: Persistence of host-type hematopoiesis after allogeneic bone marrow transplantation for leukemia is significantly related to the recipient's age and/or the conditioning regimen, but it is not associated with an increased risk of relapse. *Blood* 83:3059-3067, 1994
 47. Vermeylen C, Cornu G: Bone marrow transplantation for sickle cell disease—the European experience. *Am J Pediatr Hematol Oncol* 16:18-21, 1994
 48. Viard F, Merel P, Bilhou-Nabera C, et al: Mixed chimerism after sex-mismatched allogeneic BMT: Evaluation of two molecular techniques. *Bone Marrow Transplant* 11:27-31, 1993
 49. Wagner JE, Zahurak M, Piantadosi S, et al: Bone marrow transplantation of chronic myelogenous leukemia in chronic phase: Evaluation of risks and benefits. *J Clin Oncol* 10:779-789, 1992
 50. Yam PY, Petz LD, Knowlton RG, et al: Use of DNA restriction fragment length polymorphisms to document marrow engraftment and mixed hematopoietic chimerism following bone marrow transplantation. *Transplant* 43:399-407, 1987

Address reprint requests to

Paul G. Rothberg, PhD
Molecular Genetics Laboratory
Children's Mercy Hospital
2401 Gillham Road
Kansas City, MO 64108

DETECTION OF MYCOBACTERIUM TUBERCULOSIS BY MOLECULAR METHODS

Vivian Jonas, MS, and Mathew Longiaru, PhD

Molecular methods for identifying *Mycobacterium tuberculosis* in culture have been routinely used in the clinical laboratory for many years. Nucleic acid probes are now commonly used to specifically identify *M. tuberculosis* grown from both solid media and liquid media such as BACTEC (Becton Dickinson Diagnostic Instrument Systems, Sparks, Maryland) broth. The Gen-Probe AccuProbe Mycobacterium Tuberculosis Complex Culture Identification Test demonstrates a sensitivity and specificity of 99.2% and 99.9%, respectively, prior to discrepant resolution (Gen-Probe AMTD, package insert). Evans et al¹⁴ have described the use of the AccuProbe tests directly from BACTEC. They concluded that the identification of *M. tuberculosis* occurred more rapidly with AccuProbe than with conventional identification methods, but that a sensitive and rapid method for detecting *M. tuberculosis* directly from clinical specimens was required. The sensitivity of nucleic acid probes for the identification of *M. tuberculosis* using the AccuProbe acridinium ester technology³⁴ is approximately 500,000 to 1 million organisms (unpublished data), which explains the need for more sensitive methods. Probes are available for *M. tuberculosis*, *M. avium*, *M. intracellulare*, *M. gordonae*, *M. kansasii*, and *M. avium* complex.

High-pressure liquid chromatography (HPLC) is also used to identify mycobacteria and can identify any known species. This method has

From Gen-Probe Incorporated, San Diego, California

CLINICS IN LABORATORY MEDICINE

VOLUME 17 • NUMBER 1 • MARCH 1997

119

been used routinely in the clinical laboratory with isolates from solid media as described by Butler and Kilburn.⁶ More recently, Jost et al²⁰ have applied this method to growth from BACTEC medium and to clinical specimens. They reported 99.0% sensitivity, relative to culture, with HPLC applied to BACTEC growth; however, sensitivity determined directly from clinical specimens was only 56.8%. The specificity of both applications of HPLC was 100%. The sensitivity and specificity reported for HPLC from the BACTEC medium is adequate for use in the clinical laboratory as a tool for diagnosing tuberculosis; however, HPLC applied to clinical specimens lacks adequate sensitivity.

Because molecular methods such as hybridization with nucleic acid probes and HPLC are not sufficiently sensitive for use directly from clinical specimens and require culturing the organism, several weeks to 2 months or more are still needed to detect and identify *M. tuberculosis*. Thus, several groups have developed nucleic acid target amplification technologies for more rapid diagnosis of tuberculosis.

Various nucleic acid target amplification technologies have been applied to detection of *M. tuberculosis*.^{1-5, 7-19, 21-44} These include polymerase chain reaction (PCR) and transcription-mediated amplification (TMA) as well as strand displacement amplification (SDA) and ligase chain reaction (LCR).^{34, 38, 39} All have been used with patient specimens; however, because the latter two methods have not been routinely used in the clinical laboratory to date, this review will focus on PCR and TMA. The primary amplification targets for PCR are the IS6110 insertion sequence and the 16S rRNA gene. TMA's target is the 16S rRNA.

TMA is a transcription-based target amplification system whose primary product is RNA. The amplification can use either RNA or DNA as a target. For detecting mycobacteria, the preferred target is rRNA. Ribosomal RNA is present at about 2000 copies per mycobacterial cell, which provides an increase in sensitivity compared with targets such as IS6110, which are present in 1 to 10 copies per cell.¹³ As shown in Figure 1, two primers are utilized for TMA. The first primer contains a promoter sequence for RNA polymerase as well as a region complementary to the 16S rRNA target sequence. Reverse transcriptase extends this primer to make a cDNA copy of the rRNA target. RNase H activity, associated with reverse transcriptase, then makes the cDNA available to the second primer. The second primer is complementary to a portion of the cDNA copy. Reverse transcriptase then extends this primer, resulting in a double-stranded DNA intermediate. A part of the DNA intermediate contains a double-stranded promoter region that is recognized by RNA polymerase. The RNA polymerase generates multiple copies of RNA. This RNA is available for amplification using the same primers and enzymes. A billion-fold amplification is achieved using TMA with *M. tuberculosis* rRNA as a target in less than 2 hours.




Figure 1. Transcription-mediated amplification. Two primers are used for TMA (see text for further details). RT = reverse transcriptase; Pol = RNA polymerase. Black bar = DNA extension product of RT; Dark tone bar = promoter primer; Medium tone bar = RNA amplicon; Light tone bar = primer; Vertical lines = base-pairing.

PCR uses two primers, one enzyme, and repeated thermal cycling between denaturation, reannealing, and extension steps. Each cycle of PCR begins with denaturation of the double-stranded DNA amplicon followed by primer annealing to target and extension of PCR products by *Taq* polymerase. Each cycle theoretically doubles the number of

target nucleic acid molecules; therefore, after 20 cycles a million-fold amplification is achieved, and after 30 cycles, a billion-fold.

PCR has been applied to direct detection of *M. tuberculosis* using a number of different target nucleic acids. Owing to the lack of standardization of reagents and protocols with "home brew" PCR protocols, there have been inconsistent results in the clinical performance from laboratory to laboratory, and sensitivity and specificity have varied greatly.²⁷ More recently, the literature has indicated that results with manufactured kit reagents are more consistent. Specificities of PCR have been in the high range (exceeding 95%), whereas sensitivities, compared with culture, have ranged between 67% and 100%.^{2, 3, 8, 11, 22, 24, 25, 33, 42} This variation in sensitivity is probably due to multiple factors including sampling error and specimen inhibition of the PCR. Amplification-based assays from Gen-Probe and Roche Molecular Systems have shown comparable levels of sensitivity and specificity in clinical evaluations.^{1, 18, 37, 44}

TMA has been combined with the Hybridization Protection Assay (HPA) for amplicon detection to develop a single-tube clinical diagnostic test for the detection of *M. tuberculosis* in respiratory sediments. The test has recently been approved by the US Food and Drug Administration (FDA) for use in acid-fast bacilli (AFB) smear-positive specimens. The Gen-Probe Amplified Mycobacterium Tuberculosis Direct Test (AMTD) was evaluated in a six-site clinical trial in the United States. The results of AFB smear, culture, and AMTD were evaluated for 6079 specimens from 2609 patients. Of the total specimens tested, 198 specimens were AFB smear-positive from 78 patients not on therapy for tuberculosis. The sensitivity of the AMTD test for these samples was 95.5% (147 of 154) and the specificity was 100% (44 of 44).

Table 1. REPORTED CLINICAL SENSITIVITY AND SPECIFICITY OF THE GEN-PROBE AMPLIFIED MYCOBACTERIUM TUBERCULOSIS DIRECT (AMTD) TEST IN SEVERAL LABORATORIES

Reference	No. Specimens	AMTD	
		Sensitivity (%)	Specificity (%)
Abe et al ¹	135	91.9	100
Bodmer et al ⁴	617	71.4	99
Bradley et al ⁵	844	93.6	97.8
Ehlers et al ¹²	71	96.2	97.7
Jonas et al ¹⁹	758	82.4	99.4
LaRocco et al ²³	760	95	98
Miller et al ²⁴	750	91	98.5
Pfyffer et al ²⁸	938	97.8	97.1
Pfyffer et al ³⁰	1117	86.6	96.4
Vlasopolder et al ³⁶	412	98.4	98.9
Welch et al ⁴¹	339	100	99.7

Table 2. REPORTED CLINICAL SENSITIVITY AND SPECIFICITY OF AMPLICOR/PCR FOR *M. TUBERCULOSIS* IN SEVERAL LABORATORIES

Reference	No. Specimens	PCR	
		Sensitivity (%)	Specificity (%)
Beavis et al ²	532	95	96
Bergmann et al ³	956	79.4	99.6
D'Amato et al ⁸	985	66.7	99.6
DeVallois et al ¹¹	784	90.9	100
Kirschner et al ²²	729	84.5	99.5
Moore et al ²⁵	1009	85	99.6
Schirm et al ³³	504	70.4	>98
Wobeser et al ⁴²	1480	79	99

Elsewhere in the world the test has been evaluated with a wide variety of samples for the diagnosis of tuberculosis. Sensitivities and specificities of AMTD and PCR on AFB smear-positive and -negative respiratory specimens have ranged from 71% to 100%. Tables 1 to 3 summarize these data. Researchers have reported that the test allows the reliable and rapid detection of tuberculosis in patients suspected of disease.

Vlasopolder et al³⁶ have applied the AMTD test to 138 nonrespiratory specimens. They report sensitivity, specificity, and positive and negative predictive values of 100%, 95.0%, 85.0%, and 100%, respectively, on the samples they tested, with the exception of pleural exudates, which were analyzed separately. One of five culture-positive samples and two culture-negative samples from pleural exudates were detected by AMTD. Vlasopolder et al³⁶ suggest the possibility of using alternative specimen-processing procedures to improve the performance of AMTD with pleural exudates, which may have inhibitors not present in respiratory samples.

Pfyffer et al³⁰ performed the largest study to date on nonrespiratory specimens. They tested 322 samples for *M. tuberculosis* using the ampli-

Table 3. COMPARISON OF GEN-PROBE AMTD AND AMPLICOR/PCR PERFORMANCE IN SEVERAL LABORATORIES

Reference	No. Specimens	AMTD		PCR	
		Sensitivity (%)	Specificity (%)	Sensitivity (%)	Specificity (%)
Ichiyama et al ¹⁸	422	100	99.3	97.8	98.9
Miller et al ²⁴	750	91	98.5	91	92.3
Vuorinen et al ³⁷	256	86.2	100	82.8	100
Zolnir-Dove et al ⁴⁴	281	94.0	94.4	96.0	94.8

fied AMTD test. Sensitivity and specificity compared with culture were 93.1% and 97.7%, respectively. Of special interest is their adaptation of the AMTD test for use with cerebrospinal fluid. They used a 500 μ sample instead of the 50 μ recommended for respiratory specimens, pretreated the sample with sodium dodecyl sulfate (SDS), and increased the amplification time to 3 hours. Six cases of tuberculous meningitis were diagnosed by AMTD. Five of the six cases were detected by culture; the sixth was culture-negative, but the patient had disseminated tuberculosis documented by culture-positive specimens from other sites in the body. Pfyffer et al³⁰ also tested 65 pleural fluids, of which two were positive by AMTD and culture, three were positive by AMTD only, and two were positive by culture only. Again it was demonstrated that pretreatment of the sample with SDS improved the results. SDS pretreatment might improve the performance of the test by eliminating the potential inhibitors present in pleural fluids.

Folgueira et al¹⁵ describe the use of PCR for the rapid diagnosis of *M. tuberculosis* bacteremia by PCR. *M. tuberculosis* DNA was present in peripheral blood mononuclear cell samples from 9 of 11 HIV-infected patients and in 7 of 21 non-HIV infected patients with different forms of tuberculosis. All samples from HIV-infected patients with extrapulmonary tuberculosis were PCR-positive, as were 5 of 13 samples from non-HIV infected patients. Samples from 12 control patients with positive PPD tests were PCR-negative. These researchers concluded that only patients with active tuberculosis will be PCR-positive, and that PCR applied to blood samples will aid in the rapid diagnosis of tuberculosis, especially in HIV-infected patients.

Several researchers have questioned the relevance of a positive nucleic acid target amplification result on specimens from patients on therapy when culture is negative for *M. tuberculosis*.^{7, 16, 21, 26, 43} It has been shown that viability of *M. smegmatis* can be assessed by amplification of rRNA using nucleic acid sequence-based amplification (another transcription based amplification system).³⁵ *M. smegmatis* that had been treated with antituberculosis drugs and shown to be nonviable were still detected by PCR, and not by NASBA. The results of the study by Van Der Vliet et al³⁵ imply that the decay of rRNA occurs more rapidly after cell death than does the degradation of genomic DNA. It should, therefore, be possible to determine the viability of *M. tuberculosis* by using another amplification method such as TMA to detect rRNA. A study was undertaken with the public health laboratory in Santa Ana, California (Orange County) to evaluate the persistence of rRNA in successfully treated patients.²⁶ Patients whose samples were AFB smear-positive within the first three samples submitted were enrolled in the study. All specimens submitted for analysis while patients were under care were included in the study and were tested by AFB smear, routine culture methods, and

AMTD. Results of smear and culture were unknown by the person performing the AMTD test. The AFB smear result was the last positive test result in 4% of cases, culture was last in 22%, and the AMTD test was last in 52% of cases. Fifty-six percent of patients had a period of shedding noncultivable organisms that were detected by AMTD. The noncultivable period lasted from 7 to 245 days. All three tests became reproducibly negative and remained negative during follow-up of up to 1 year. These data suggest that with time (and more experience), TMA may be applied to determine whether therapy is efficacious.

Schluger et al³² evaluated the clinical utility of IS6110 PCR for the diagnosis of tuberculosis and concluded it is not useful in a routine setting based on its lack of specificity for diagnosis of active TB. As discussed previously, however, Moore et al believe that nucleic acid amplification tests will play a role in the routine diagnosis of tuberculosis. Kennedy et al²¹ also believe that "... PCR is a promising method for assessing treatment response in pulmonary tuberculosis," and Yuen et al⁴³ agreed with this assessment of clinical utility of PCR. Hellyer et al¹⁶ have concluded that the detection of *M. tuberculosis* DNA by target amplification will not be useful in therapeutic monitoring based on the inability of PCR to differentiate between live and dead organisms. This area is being further explored as researchers become more familiar with the power of nucleic acid amplification tests.

Nucleic acid amplification techniques such as TMA and PCR have been shown to be effective in the detection of *M. tuberculosis* from a variety of specimens. Preliminary data suggest that nucleic acid amplification methods that detect rRNA can be used on samples from patients undergoing therapy for tuberculosis. Much more work remains to be done in this area. Clearly, TMA and PCR aid in the rapid detection of tuberculosis in patients suspected of having the disease.

Obviously these methods are currently being used to detect tuberculosis. Nucleic acid amplification tests already aid in patient management and in making public health decisions. Improvements in sensitivity will render these tests even more useful than they are today. Ultimately the tests will be automated for greater ease of use. It is clear that improved detection and monitoring of tuberculosis will help in the fight to eradicate this disease and effectively manage the infected patient.

References

1. Abe C, Hirano K, Wada M, et al: Detection of *Mycobacterium tuberculosis* in clinical specimens by polymerase chain reaction and Gen-Probe Amplified Mycobacterium Tuberculosis Direct Test. *J Clin Microbiol* 31:3270-3274, 1993
2. Beavis KG, Lichty MB, Jungkind DJ, et al: Evaluation of Amplicor PCR for the

- direct detection of *Mycobacterium tuberculosis* from sputum specimens. *J Clin Microbiol* 33:2582-2586, 1995
3. Bergmann JS, Woods GL: Clinical evaluation of the Roche AMPLICOR PCR *Mycobacterium tuberculosis* test for detection of *M. tuberculosis* in respiratory specimens. *J Clin Microbiol* 34:1083-1085, 1996
 4. Bodmer T, Gurtner A, Schopfer K, et al: Screening of respiratory tract specimens for the presence of *Mycobacterium tuberculosis* by using the Gen-Probe Amplified *Mycobacterium Tuberculosis* Direct Test. *J Clin Microbiol* 32:1483-1487, 1994
 5. Bradley SP, Reed SI, Catanzaro A: Clinical efficacy of the Amplified *Mycobacterium Tuberculosis* Direct Test for the diagnosis of pulmonary tuberculosis. *Am Rev Respir Dis* 153:1606-1610, 1996
 6. Butler WR, Kilburn JO: Identification of major slowly growing pathogenic mycobacteria and *Mycobacterium gordonae* by high-performance liquid chromatography of their mycolic acids. *J Clin Microbiol* 26:50-53, 1988
 7. Chin DP, Yajko DM, Hadley WK, et al: Clinical utility of a commercial test based on the polymerase chain reaction for detecting *Mycobacterium tuberculosis* in respiratory specimens. *Am J Respir Crit Care Med* 151:1872-1877, 1995
 8. D'Amato RF, Wallman AA, Hochstein LH, et al: Rapid diagnosis of pulmonary tuberculosis by using Roche AMPLICOR *Mycobacterium tuberculosis* PCR test. *J Clin Microbiol* 33:1832-1834, 1995
 9. Delacourt C, Poveda J-D, Chureau C, et al: Use of polymerase chain reaction for improved diagnosis of tuberculosis in children. *J Pediatrics* 126:703-709, 1995
 10. Del Portillo P, Thomas MC, Martinez E, et al: Multiprimer system for differential identification of mycobacteria in clinical samples. *J Clin Microbiol* 34:324-328, 1996
 11. DeVallois A, Legrand E, Rastogi N: Evaluation of Amplicor MTB test as adjunct to smears and culture for direct detection of *Mycobacterium tuberculosis* in the French Caribbean. *J Clin Microbiol* 34:1065-1068, 1996
 12. Ehlers S, Pirmann M, Zaki W, et al: Evaluation of a commercial rRNA target amplification assay for detection of *Mycobacterium tuberculosis* complex in respiratory specimens. *Eur J Clin Microbiol Infect Dis* 13:827-829, 1994
 13. Eisenach KD, Cave MO, Bates JH, et al: Polymerase chain reaction amplification of a repetitive DNA sequence specific to *Mycobacterium tuberculosis*. *J Infect Dis* 161:977-981, 1990
 14. Evans KD, Nakasone AS, Sutherland PA, et al: Identification of *Mycobacterium tuberculosis* and *Mycobacterium avium-M. intracellulare* directly from primary BACTEC cultures by using acridinium ester labeled DNA probes. *J Clin Microbiol* 30:2427-2431, 1992
 15. Folgueira L, Delgado R, Palenque E, et al: Rapid diagnosis of *Mycobacterium tuberculosis* bacteremia by PCR. *J Clin Microbiol* 34:512-515, 1996
 16. Hellyer TJ, Fletcher TW, Bates JH, et al: Strand displacement amplification and the polymerase chain reaction for monitoring response to treatment in patients with pulmonary tuberculosis. *J Infect Dis* 173:934-941, 1996
 17. Hermans PWM, Schuitema ARJ, Van Soolingen D, et al: Specific detection of *Mycobacterium tuberculosis* complex strains by polymerase chain reaction. *J Clin Microbiol* 28:1204-1213, 1990
 18. Ichiyama S, Iinuma Y, Tawada Y, et al: Evaluation of Gen-Probe Amplified *Mycobacterium Tuberculosis* Direct Test and Roche PCR-Microwell Plate Hybridization Method (Amplicor *Mycobacterium*) for direct detection of mycobacteria. *J Clin Microbiol* 34:130-133, 1996
 19. Jonas V, Alden MJ, Curry JL, et al: Detection and identification of *Mycobacterium tuberculosis* directly from sputum sediments by amplification of rRNA. *J Clin Microbiol* 31:2410-2416, 1993
 20. Jost KC Jr, Dunbar DF, Barth SS, et al: Identification of *Mycobacterium tuberculosis* and *M. avium* complex directly from smear-positive sputum specimens and BACTEC 12B cultures by high performance liquid chromatography with fluorescence detection and computer-driven pattern recognition models. *J Clin Microbiol* 33:1270-1277, 1995
 21. Kennedy N, Gillespie SH, Saruni AOS, et al: Polymerase chain reaction for assessing treatment response in patients with pulmonary tuberculosis. *J Infect Dis* 170:713-716, 1994
 22. Kirschner P, Rosenau J, Springer B, et al: Diagnosis of mycobacterial infections by nucleic acid amplification: 18-month prospective study. *J Clin Microbiol* 34:304-312, 1996
 23. LaRocco MT, Wanger A, Ocera H, et al: Evaluation of a commercial rRNA amplification assay for direct detection of *Mycobacterium tuberculosis* in processed sputum. *Eur J Clin Microbiol Infect Dis* 13:726-731, 1994
 24. Miller N, Hernandez SG, Cleary TJ: Evaluation of Gen-Probe Amplified *Mycobacterium Tuberculosis* Direct Test and PCR for direct detection of *Mycobacterium tuberculosis* in clinical specimens. *J Clin Microbiol* 32:393-397, 1994
 25. Moore DF, Curry JL: Detection and identification of *Mycobacterium tuberculosis* directly from sputum sediments by Amplicor PCR. *J Clin Microbiol* 33:2686-2691, 1995
 26. Moore DF, Curry JL, Knott CA, et al: Amplification of ribosomal RNA for assessing treatment response of pulmonary tuberculosis patients during antimicrobial therapy. *J Clin Microbiol*, in press
 27. Noordhoek GT, Kolk AHG, Bjune G, et al: Sensitivity and specificity of PCR for detection of *Mycobacterium tuberculosis*: A blind comparison study among seven laboratories. *J Clin Microbiol* 32:277-284, 1994
 28. Pfyffer GE: Amplification techniques: Hope or illusion in the direct detection of tuberculosis? *Medical Microbiology Letters* 3:335-347, 1994
 29. Pfyffer GE, Kissling P, Wirth R, et al: Direct detection of *Mycobacterium tuberculosis* complex in respiratory specimens by a target-amplified test system. *J Clin Microbiol* 32:918-923, 1994
 30. Pfyffer GE, Kissling P, Jahn EMI, et al: Diagnostic performance of amplified *Mycobacterium tuberculosis* direct test with cerebrospinal fluid, other non-respiratory, and respiratory specimens. *J Clin Microbiol* 34:834-841, 1996
 31. Richeldi L, Barnini S, Saltini C: Molecular diagnosis of tuberculosis. *Eur Respir J* 8(suppl):689s-700s, 1995
 32. Schluger NW, Kinney D, Harkin TJ, et al: Clinical utility of the polymerase chain reaction in the diagnosis of infections due to *Mycobacterium tuberculosis*. *Chest* 105:1116-1121, 1993
 33. Schirm J, Oostendorp LAB, Mulder JG: Comparison of Amplicor, in-house PCR, and conventional culture for detection of *Mycobacterium tuberculosis* in clinical samples. *J Clin Microbiol* 33:3221-3224, 1995
 34. Shinnick TM, Jonas V: Molecular approaches to the diagnosis of tuberculosis. In Bloom BR (ed): *Tuberculosis: Pathogenesis and Control*. Washington, DC, American Society for Microbiology, 1994
 35. Van Der Vliet GME, Schepers P, Schukink RAF, et al: Assessment of mycobacterial viability by RNA amplification. *J Clin Microbiol* 38:1959-1965, 1994
 36. Vlasopolder F, Singer P, Roggeveen C: Diagnostic value of an amplification method (Gen-Probe) compared with that of culture for the diagnosis of tuberculosis. *J Clin Microbiol* 33:2699-2703, 1995
 37. Vuorinen P, Miettinen A, Vuento R, et al: Direct detection of *Mycobacterium tuberculosis* complex in respiratory specimens by Gen-Probe Amplified *Mycobacterium Tuberculosis* Direct Test and Roche Amplicor *Mycobacterium Tuberculosis* Test. *J Clin Microbiol* 33:1856-1859, 1995
 38. Walker DA, Taylor IK, Mitchell DM, et al: Comparison of polymerase chain reaction amplification of two mycobacterial DNA sequences, IS6110 and the 65kDa antigen gene, in the diagnosis of tuberculosis. *Thorax* 47:690-694, 1992
 39. Walker GT, Fraiser MS, Schram JL, et al: Strand displacement amplification—an isothermal, *in vitro* DNA amplification technique. *Nucleic Acids Res* 20:1691-1696, 1992
 40. Walker GT, Nadeau JG, Linn CP, et al: Strand displacement amplification (SDA) and transient-state fluorescence polarization detection of *Mycobacterium tuberculosis* DNA. *Clin Chem* 42:9-13, 1996
 41. Welch K, Brown G, Jonas V, et al: Performance of the Gen-Probe Amplified *Mycobacterium Tuberculosis* Direct Test in a laboratory that infrequently isolates *Mycobacterium tuberculosis*. *Diagn Microbiol Infect Dis* 22:297-299, 1995
 42. Wobeser WL, Krajden M, Conly J, et al: Evaluation of Roche Amplicor PCR assay for *Mycobacterium tuberculosis*. *J Clin Microbiol* 34:134-139, 1996

43. Yuen KY, Chan KS, Chan CM, et al: Use of PCR in routine diagnosis of treated and untreated pulmonary tuberculosis. *J Clin Pathol* 46:318-322, 1992
44. Zolnir-Dove M, Poljak M, Seme K, et al: Evaluation of two commercial amplification assays for the detection of *Mycobacterium tuberculosis* complex in respiratory specimens. *Infection* 23:216-221, 1995

Address reprint requests to
Vivian Jonas, MS
Gen-Probe Incorporated
9880 Campus Point Drive
San Diego, CA 92121

SIMULTANEOUS MULTIANALYTE NUCLEIC ACID DETECTION FOR GASTROINTESTINAL BACTERIAL PATHOGENS USING GENESTAR TECHNOLOGY

Linxian Wu, PhD, Jana Coombs, BS, Sharon Malmstrom, BS,
and Michael Glass, PhD

Acute diarrheal illnesses are very common among young children and infants.^{5, 11} In developing countries, approximately 4 to 5 million deaths each year have been reported as the result of gastroenteritis.⁵ In developed countries, gastrointestinal (GI) bacterial infections are also a major cause of morbidity and mortality.⁵ Five bacterial species have been determined to play a significant role in bacterial gastroenteritis: *Salmonella* spp., *Shigella* spp. and enteroinvasive *Escherichia coli* (EIEC), *Campylobacter* spp., enterohemorrhagic *E. coli* (EHEC), especially 0157:H7, and *Yersinia enterocolitica*. Differentiation of these bacterial species is critical for physicians to select an adequate treatment regimen. The wrong treatment regimen may worsen the illness or even result in death.

Classical culture techniques are typically used in clinical laboratories to identify and differentiate the GI bacterial pathogens.¹¹ Such methods usually take a minimum of 18 hours to several days. It is evident that rapid identification methods are needed for clinical diagnosis of GI bacterial pathogens.

From the Department of Research and Development, Gull Laboratories, Inc., Salt Lake City, Utah

CLINICS IN LABORATORY MEDICINE