

Chapter 1 Exercises

David Piper

September 7, 2020

1.1 D and e are sets as they are in curly braces. The other examples have some items just listed by themselves.

- 1.2 (a) $\{x \in S \mid x \in \mathbb{N}\}$
(b) $\{x \in S \mid x \geq 0\}$
(c) $\{x \in S \mid x < 0\}$
(d) $\{x \in S \mid |x| > 1\}$

- 1.3 (a) $|A| = 5$
(b) $|B| = 21$
(c) $|C| = 50$
(d) $|D| = 2$
(e) $|E| = 1$
(f) $|E| = 2$

- 1.4 (a) $A = \{-3, -2, -1, 0, 1, 2, 3, 4\}$
(b) $B = \{-2, -1, 0, 1, 2\}$
(c) $C = \{1, 2, 3, 4, 5, 6, 7, 8, 9\}$
(d) $D = \{0, 1\}$
(e) $E = \emptyset$

- 1.5 (a) $A = \{x \in \mathbb{Z} \mid x < 0\}$
(b) $B = \{x \in \mathbb{Z} \mid x^2 < 10\}$
(c) $C = \{x \in \mathbb{Z} \mid 0 < x^2 < 5\}$

- 1.6 (a) $A = \{2x + 1 \mid x \in \mathbb{Z}\} = \{\dots, -3, -1, 1, 3, 5, \dots\}$
 (b) $B = \{4n \mid n \in \mathbb{Z}\} = \{\dots, -16, -8, -4, 0, 4, 8, 16, \dots\}$
 (c) $C = \{3q + 1 \mid q \in \mathbb{Z}\} = \{\dots, -7, -4, 1, 4, 7, \dots\}$
- 1.7 (a) $A = \{3x + 2 \mid x \in \mathbb{Z}\}$
 (b) $B = \{5x \mid x \in \mathbb{Z}\}$
 (c) $C = \{x^3 \mid x \in \mathbb{Z}\}$
- 1.8 (a) $A = \{-4, -3, -2, 2, 3, 4\}$
 (b) $\frac{9}{4}, \frac{10}{4}$, and $\frac{21}{8}$
 (c) $C = \{2, \sqrt{2}\}$
 (d) $D = \{2\}$
 (e) $|A| = 6, |C| = 2$, and $|D| = 1$
- 1.9 $B = \{5, 7, 8, 10, 13\}$ so $C = \{5, 8\}$
- 1.10 (a) $A = \{1\}, B = \{1\}$, and $C = \{1, 2\}$
 (b) $A = \{1\}, B = \{\{1\}, 2\}$, and $C = \{\{\{1\}, 2\}\}$
 (c) $A = \{1\}, B = \{\{1\}, 2\}$, and $C = \{1, 2\}$
- 1.11 The interval I can just be (a, b) since the two intervals would be equal and thus subsets. If we let $c = \frac{b+a}{2}$ then the interval I would be centered at c ; as it would be the center of the original interval.
- 1.12 $A = B = C = D = E$
- 1.13 Skipping this because the problem is pretty straightforward and it's really complicated to draw Venn diagrams in LaTeX (aka the benefits are not worth the effort).
- 1.14 (a) $\mathcal{P}(A) = \{\emptyset, \{1\}, \{2\}, \{1, 2\}\}$ and $|\mathcal{P}(A)| = 4$
 (b) $\mathcal{P}(A) = \{\emptyset, \{\emptyset\}, \{1\}, \{\emptyset, 1\}, \{\{a\}\}, \{\emptyset, \{a\}\}\}, \{1, \{a\}\}, \{\emptyset, 1, \{a\}\}\}$
 and $|\mathcal{P}(A)| = 8$
- 1.15 $\mathcal{P}(A) = \{\emptyset, \{0\}, \{\{0\}\}, \{\emptyset, \{0\}\}, \{0, \{0\}\}, \{\emptyset, 0, \{0\}\}\}$

- 1.16 Find $\mathcal{P}(\mathcal{P}(\{1\}))$ and its cardinality. $\mathcal{P}(\{1\}) = \{\emptyset, \{1\}\}$
 $\mathcal{P}(\mathcal{P}(\{1\})) = \{\emptyset, \{\emptyset\}, \{\{1\}\}, \{\emptyset, \{1\}\}\}$
 $|\mathcal{P}(\mathcal{P}(\{1\}))| = 4$
- 1.17 $\mathcal{P}(A) = \{\emptyset, \{0\}, \{\emptyset\}, \{\{\emptyset\}\}, \{0, \emptyset\}, \{\emptyset, \{\emptyset\}\}, \{0, \{\emptyset\}\}, \{0, \emptyset, \{\emptyset\}\}\}$
 $|\mathcal{P}(A)| = 8$
- 1.18 $A = \{\emptyset, 0, \{\emptyset\}, \{0\}\}$
 $\mathcal{P}(A) = \{\emptyset, \{\emptyset\}, \{0\}, \{\{\emptyset\}\}, \{\{0\}\}, \{\emptyset, 0\},$
 $\{\emptyset, \{\emptyset\}\}, \{\emptyset, \{0\}\}, \{\{0\}, \{\emptyset, 0, \{\emptyset\}\}\},$
 $\{\emptyset, 0, \{\emptyset\}\}, \{\emptyset, 0, \{0\}\}, \{\emptyset\}\}$
 // I am stopping here this is really tedious, I know I am missing a few sets, but it isn't valuable enough to spend more time on. May come back with pen and paper.
- 1.19 (a) $S = \emptyset$
 (b) $S = \{1\}$
 (c) $\{\{1\}, \{2\}, \{3\}, \{4\}, \{5\}\}$
 (d) $\{1, 2, 3, 4, 5\}$
- 1.20 (a) This statement could be either true or false; we don't have enough information. Specifically, if the *only* element of A is 1, then it would be true. However if $A = \{1, \{1\}\}$ then it would be false.
- (b) *Proof.* Since we know that $A \subset \mathcal{P}(B) \subset C$ and that $|A| = 2$ then we know that $|\mathcal{P}(B)|$ must be less than $|C|$; since a proper subset must have less elements than its proper superset. So we have the following inequality $|A| < |\mathcal{P}(B)| < |C|$.
- Since the cardinality of a set must be a nonzero positive integer we know that the smallest possible cardinality for $\mathcal{P}(B)$ is 3. However, we also know that the cardinality of a power set $\mathcal{P}(K)$ must be $2^{|K|}$ for some set K . Thus, the cardinality of $\mathcal{P}(B)$ cannot be 3 (the next largest integer after 2; which is $|A|$) because $\log_2 3$ is not an integer. However, it *can* be 4. So the smallest possible cardinality of $\mathcal{P}(B) = 4$. Since $|\mathcal{P}(B)| < |C|$ the smallest possible cardinality of $C = 5$. Q.E.D.
- (c) *Proof.* We are given that $|B| = |A| + 1$. We also know that the cardinality of a power set of a set K must be $2^{|K|}$. So in this case,

$|\mathcal{P}(A)| = 2^{|A|}$ and $|\mathcal{P}(B)| = 2^{|B|} = 2^{|A+1|}$. From this equation we can observe that regardless of the cardinality of B it must always be at least 2^1 or simply 2 more than the cardinality of A . Q.E.D.

- (d) *Proof.* We have four sets: A, B, C , and D which are subsets of $\{1, 2, 3\}$. Furthermore, we know that $|A| = |B| = |C| = |D| = 2$. There are only 3 subsets of $\{1, 2, 3\}$ with a cardinality of 2, namely $\{1, 2\}, \{1, 3\}$, and $\{2, 3\}$. Since we have four sets all of which are subsets of $\{1, 2, 3\}$ with a cardinality of 2 we must conclude that at least one of them are the same set. Q.E.D.

- 1.21 One approach we can take to find a solution that satisfies all of the conditions is attempting to minimize the sums of the elements in each set. We know from (a) that A must contain 1, from (b) that A and C must contain 2, and that A must contain 3 from (c). So we know that the minimum sum of A is 6. If we choose 3 to be in C from (c) then the difference in the sums of A and C is exactly 1 which satisfies (f). So we'd like to maintain that if we can. Observing that (d) requires 4 to be in an even number of any of the sets we can choose to add it to both A and C , thus maintaining the difference of 1 for their sums. If we then choose to satisfy (e) by adding 5 to B we have satisfied all of the conditions leaving us with $B = \{1, 5\}$.