

Using n -dimensional Hypervolumes to
measure Ecosystem Stability

David Bridgwood

August 2018

Word Count: 3516

A thesis submitted in part fulfilment of the requirements for the degree of
Master of Research at Imperial College London

Formatted in the Journal style of Ecology Letters

Submitted for the MRes in Computational Methods in Ecology and
Evolution

Declaration of Authorship

Data Used

All data from this thesis was provided by projects based at SAFE, a large-scale forest fragmentation experiment (?). Data on trees came from Terhi Riutta (?), small-mammals from Philip Chapman (?) and beetles from Adam Sharp (https://www.safeproject.net/datasets/view_dataset?id=123).

Analysis

All code and analysis was undertaken by myself with the use of Python and R (?). The R package 'hypervolumes' (?) was utilised in the construction of hypervolumes. All code is available at https://github.com/dbridg15/masters_project.

Supervision

Prof. Rob Ewers and Dr. David Orme provided regular input on my analysis, suggesting possible methods and extensions.

Acknowledgements

I would like to thank the following people for their contribution to the thesis:

- My Supervisor Rob Ewers, for his guidance throughout the project.
- David Orme, for his help in developing my analysis.
- Samraat Pawar and James Rosindell for useful discussions about the project over the months.

Table of Contents

1 Abstract

Continued demand for resources has seen the conversion of natural landscapes to urban and agricultural uses increase. These land use changes, which are particularly prevalent in the highly biodiverse tropical rainforests of Southeast Asia impact the structure of ecosystems and their resilience to environmental perturbation. The debate surrounding the relationship of biodiversity with stability remains contentious with both theoretical and empirical studies providing evidence on either side. Stability has traditionally been measured as the variability of communities through time, ‘community temporal stability’, though a plethora of definitions exist. Advances in the delineation of n -dimensional hypervolumes offers a new approach to studying ecosystem stability. Here I use species abundance data from the SAFE project in Sabah, Malaysia to construct ‘community composition hypervolumes’ for the taxonomic groups of trees, mammals and beetles at plots across a gradient of forest quality and logging history. I assess ecosystem stability using measures of hypervolume overlap between censuses, demonstrating a new approach to measuring ecosystem stability.

Keywords:

ecosystem stability, n -dimensional hypervolumes, land-use change, SAFE.