

Large values of Dirichlet Polynomials and Zero Density Results

Honors Thesis Presentation

Chi Li

May 1, 2025

1. Motivation
2. Zeta function and PNT
3. Primes in short intervals and Zero density
4. Large Values Estimates results

Motivation

- We want to estimate the distribution of primes

$$\pi(x) \stackrel{\text{def}}{=} \sum_{p \leq x} 1.$$

- The Prime Number Theorem gives

$$\pi(x) = (1 + o(1)) \frac{x}{\log x}.$$

- What about

$$\sum_{x \leq p \leq x+y} 1$$

for $y = o(x)$? Is it still $\sim y / \log x$?

Zeta function

- For $\Re(s) > 1$ define

$$\zeta(s) \stackrel{\text{def}}{=} \sum_{n \geq 1} n^{-s}.$$

- This also has a product representation

$$\zeta(s) = \prod_p (1 - p^{-1})^{-1}.$$

- We can also analytically continue this to the whole complex plane.

$$\pi^{-s/2} \Gamma(s/2) \zeta(s) = \pi^{-(1-s)/2} \Gamma((1-s)/2) \zeta(1-s).$$

Idea of proof: Use Integral representation of Γ , exchange summation and integration and apply Poisson summation.

Relation to PNT

1. We scale everything by $\log x$. Let

$$\Lambda(n) \stackrel{\text{def}}{=} \begin{cases} \log p, & \text{if } n = p^k, \\ 0, & \text{else,} \end{cases}$$

and

$$\psi(x) \stackrel{\text{def}}{=} \sum_{n \leq x} \Lambda(n).$$

2. Using product representation

$$\frac{\zeta'}{\zeta} = [\log \zeta]' = - \sum_n \frac{\Lambda(n)}{n^s}$$

Relation to PNT (cont.)

We can make this connection precise through the explicit formula

Theorem (Riemann-von Mangoldt explicit formula)

Let N be not a prime power. We have

$$\psi(N) = N - \lim_{T \rightarrow \infty} \sum_{|\Im(\rho)| \leq T, \zeta(\rho)=0} \frac{N^\rho}{\rho} - \frac{\zeta'}{\zeta}(0) + \frac{1}{2} \log(1 - N^{-2}).$$

Theorem (Truncated Riemann-von Mangoldt explicit formula)

We have

$$\psi(N) = N - \sum_{|\Im(\rho)| \leq T, \zeta(\rho)=0} \frac{N^\rho}{\rho} + O(N(\log NT)^2/T) + O(\log N).$$

Assume Riemann Hypothesis, then $|N^\rho| = N^{1/2}$, so we have $\psi(N) = N + O(N^{1/2+\epsilon})$.

Prime numbers in short intervals

What about $\sum_{x \leq n \leq x+y} \Lambda(n) \sim y$? If RH holds then we can take $y = x^{1/2+\epsilon}$.

Show enough zeros have 'small' real part. Use a zero counting function for zeros with large real part.

Definition (Zero density)

$$N(\sigma, T) \stackrel{\text{def}}{=} \#\{\rho : \zeta(\rho) = 0, |\Im(\rho)| \leq T\}.$$

Theorem (Hoheisel)

If $N(\sigma, T) \ll T^{a(1-\sigma)} \log^b T$ uniformly for $\sigma \in [1/2, T]$ then we can take $y = x^\theta$, for


$$\theta > 1 - \frac{1}{a + \frac{b}{A}},$$

and A is a (large) absolute constant.


Zero density

Before 2023: $N(\sigma, T) \ll T^{12(1-\sigma)/5+o(1)}$ (Ingham-Huxley)

2024: $N(\sigma, T) \ll T^{30(1-\sigma)/13+o(1)}$ (Guth-Maynard).



(a) $\sigma = 3/4$ is also the bottleneck when written in Hoheisel's form.



(b) The exponent is reduced around the bottleneck region.

Idea of Zero density proof

1. $\frac{1}{\zeta}(s)$ can be represented as

$$\sum_n \frac{\mu(n)}{n^s}.$$

2. Let $M_x(s) \stackrel{\text{def}}{=} \sum_{n \leq x} \frac{\mu(n)}{n^s}$, which is an approximation of $1/\zeta$. Then we have by Möbius inversion

$$\zeta(s)M_x(s) = 1 + \sum_{n > x} a_{n,x} n^{-s} \approx 1 + \sum_{n > x} a_{n,x} n^{-s} e^{-n/y} \approx 1 + \sum_{y^2 > n > x} a_{n,x} n^{-s} e^{-n/y}$$

with some error decreasing in y .

3. If s is a zero of ζ then the left hand side is zero, which means the Dirichlet series on the right hand side has magnitude close to 1.

Guth-Maynard zero density result

Theorem (Guth-Maynard)

Let $D_N(t) \stackrel{\text{def}}{=} \sum_{N \leq x < 2N} b_n n^{it}$. If $W \subset [0, T]$ is a set of 1-separated points such that

$$|D_N(t_i)| > V \quad \forall t_i \in W$$

then

$$|W| \leq T^{o(1)} \left(\max_{N \leq n < 2N} |b_n| \right) (N^2 V^{-2} + N^{18/5} V^{-4} + TN^{12/5} V^{-4}).$$

Hybrid Zero Density and Dirichlet Value Estimate

There are generalizations for zero densities of Dirichlet-L functions $L(s, \chi) \stackrel{\text{def}}{=} \sum_n \chi(n) n^{-s}$.

Theorem

Let $D_N(t, \chi) \stackrel{\text{def}}{=} \sum_{N \leq x < 2N} b_n \chi(n) n^{it}$. If $W = \{t_i, \chi_i\}$ such that χ_i is a primitive character mod q , $t_i \in [0, T]$ are 1-separated for the same character,

$$|D_N(t_i, \chi_i)| > V \quad \forall (t_i, \chi_i) \in W.$$

Then we have for $N \geq q^{5/6}$

$$|W| \leq (qT)^{o(1)} \left(\max_{N \leq n < 2N} |b_n| \right) (N^2 V^{-2} + N^{18/5} V^{-4} + qTN^{12/5} V^{-4}).$$