



Molecular origin and biological effects of exercise mimetics

Yuping Zhu, Gang Song^{*}

Key Lab of Physical Fitness Evaluation and Motor Function Monitoring, College of Physical Education, Southwest University, Chongqing, 400715, China

ARTICLE INFO

Keywords:

Exercise mimetics
Pharmacology research
Target

ABSTRACT

With the rapid development of sports science and molecular biology technology, academia refers to molecules or microorganisms that mimic or enhance the beneficial effects of exercise on the body, called “exercise mimetics.” This review aims to clarify the concept and development history of exercise mimetics, and to define the concept of exercise mimetics by summarizing its characteristics and functions. Candidate molecules and drug targets for exercise mimetics are summarized, and the relationship between exercise mimetics and exercise is explained, as well as the targeting system and function of exercise mimetics. The main targeting systems for exercise mimetics are the exercise system, circulatory system, endocrine system, endocrine system, and nervous system, while the immune system is potential targeting systems. Finally, future research directions for exercise mimetics are discussed.

1. Introduction

Currently, new drug development is becoming increasingly difficult and time-consuming. According to reports, the number of new drugs that make it to market for every \$1 billion invested is halved every nine years, and is stuck in the ‘anti-Moore’s Law’ dilemma.¹ The ‘anti-Moore’s Law’, signifying that the returns from drug development are significantly lower than the capital invested. Exercise mimetics, an avant-garde paradigm in drug development, harness the advantageous outcomes derived from established physical activities. Through the integration of artificial intelligence (AI) technology, this approach not only provides novel perspectives in drug discovery but also presents a promising trajectory to confront the limitations posed by the ‘anti-Moore’s Law’. It is well known that ‘exercise is medicine’ (EIM), and can be involved in the prevention and treatment of up to 26 diseases such as obesity, diabetes, and cognitive dysfunction.^{2–4} Suitable exercise can induce adaptive changes in the body, producing health benefits that involve numerous cellular and molecular changes, countless intricate biochemical reactions, and interactions between tissue and organs.^{5–7} These molecules or microorganisms involved in adaptation may provide inspiration for pharmaceuticals.

The concept of ‘Enviromimetics’ was introduced 20 years ago and refers to new therapies that are able to mimic or enhance beneficial environmental stimuli through complementary approaches based on gene-environment interactions and experience-dependent molecular mechanisms of plasticity, which can be broadly engaged in the

treatment of disease.^{8,9} Based on this, in June 2021, Gubert proposed a new drug development approach that designs adaptive changes in the health benefits induced by exercise as efficacy substances for preventing and treating certain specific diseases, called ‘Exercise Mimetics’.⁷ Therefore, it is presumed that ‘Exercise Mimetics’ is a subclass of ‘Enviromimetics’. Studies have designed a transmembrane protein 130 (GP130) ligand small molecule IC7Fc based on the characteristics of interleukin 6 (IL-6) and ciliary neurotrophic factor (CNTF), which removed one GP130 binding site in IL-6 and replaced it with a leukemia inhibitory factor (LIF) receptor binding site in CNTF, and then fused it with the immunoglobulin G (IgG) Fc domain. IC7Fc has CNTF morphology and IL-6 receptor-dependent features, and is used to treat type 2 diabetes, and has passed preclinical trials. This achievement was published in the journal Nature and recommended on its public account.¹⁰ This is the first potential drug developed and validated using the exercise mimetic development strategy. In addition, studies have found that aerobic exercise can upregulate the expression of CD8⁺ T cells and IL-15Rα sensitive to interleukin 15 (IL-15), leading to the enrichment of CD8⁺ T cells and IL-15Rα on pancreatic cancer cells and killing tumor cells. Novartis developed an IL-15 agonist, NIZ985, based on this mechanism, which can simulate the effect of exercise to enhance the IL-15/IL-15Rα signaling pathway, reduce potential inflammatory reactions, and produce sustained anti-tumor effects.¹¹ This is the first report of using exercise mimetics to treat cancer. It is imperative to recognize that the realm of exercise mimetics is currently in its nascent stages, marked by an absence of conclusive determinations regarding

^{*} Corresponding author.

E-mail address: songgang@aliyun.com (G. Song).

<https://doi.org/10.1016/j.jesf.2023.12.002>

Received 11 September 2023; Received in revised form 5 December 2023; Accepted 5 December 2023

Available online 8 December 2023

1728-869X/© 2023 The Society of Chinese Scholars on Exercise Physiology and Fitness. Published by Elsevier (Singapore) Pte Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

the cause-and-effect relationships between candidate molecules and established drug targets. Consequently, exercise mimetics currently represent a conceptual framework, and only the two studies mentioned earlier have conducted direct clinical trials under this guiding principle. Both of these studies have exclusively reported results from animal experiments, with no disclosed outcomes from human trials. Additionally, as of now, no exercise mimetic drugs have progressed through Phase III clinical trials to attain market approval. This underscores that research on exercise mimetics remains in its early developmental phase, with no definitive conclusions drawn regarding the efficacy and potential adverse effects of these compounds. It implies that the transition from exercise mimetics to “exercise pills” has not been fully actualized. Future endeavors should prioritize the expansion of clinical research to refine and authenticate this research paradigm.

In summary, ‘Exercise Mimetics’ is a subclass of ‘Enviromimetics’, the development strategy of exercise mimetics can provide new directions for new drug development, which can help break the ‘anti-Moore’s Law’ of drug discovery and accelerate the speed of drug development. This review aims to clarify the concept of exercise mimetics, trace its developmental history, and provide a comprehensive literature review on its candidate molecule sources, established drug targets, and biological mechanisms. By systematically collecting relevant literature in the field of exercise mimetics, we intend to integrate distinctive features from these sources to enhance the understanding of exercise mimetics. Finally, it will explore the application and trends of AI and bioinformatics tools in the development of exercise mimetics.

2. Overview of exercise mimetics

Exercise mimetics are a proposed class of therapeutics that specifically mimic or enhance the therapeutic effects of exercise.⁷ Exercise mimetics mainly benefit health through three ways: synthesizing myokines, regulating signaling pathways, and targeting small molecule targets (Table 1).

2.1. Myokines

The process by which certain factors produced in muscles during exercise act on various tissues such as the brain, liver, and adipose tissue through endocrine mechanisms, exerting beneficial effects on overall health, is referred to as the “work stimulus,” “work stimulus,” “work factor,” or “exercise factor”.^{31,32} Pedersen’s seminal work quantified a striking surge, approximately 100–150 times the baseline, in IL-6 concentrations within the bodies of marathon runners immediately post-race, predominantly originating from active musculature. Subsequent investigations unveiled the multifaceted impacts of IL-6, encompassing heightened glucose absorption and insulin responsiveness in the healthful populace, augmentation of skeletal muscle mass and strength, and facilitation of adipose tissue breakdown coupled with enhanced fatty acid oxidation. This groundbreaking revelation substantiates, for the first time, a substantial escalation of IL-6 within functioning muscles post-exercise, conclusively affirming the presence of muscle-derived IL-6.¹² Therefore, Pedersen boldly proposed that IL-6 produced by working muscles during exercise is considered a “myokine” and named it as such.^{31,33,34} In January 2012, Nature reported the discovery of a PGC1- α -dependent myokine called irisin, and confirmed that both mice and humans promote the secretion of irisin after exercise, acting on white adipose tissue, stimulating the expression of uncoupling protein 1 (UCP1) and brown fat-like development, improving glucose homeostasis and preventing obesity.¹⁵ This investigation unveiled escalated extracellular heat shock protein 90 α (eHsp90 α) levels in the interstitial fluid and plasma of murine muscle tissues post-exercise. Notably, it elucidated the binding capacity of eHsp90 α to integrin α V β 5, inducing conformational alterations in Irisin. The activated Irisin, in turn, engages integrin through high-affinity binding sites, initiating signal transmission via the Hsp90 α / α V β 5 complex. These findings offer crucial

insights into the functional mechanism of Irisin, thereby contributing to the advancement of exercise mimetics grounded in Irisin³⁵. Therefore, This is an extension and expansion of the “myokine” family. In October 2016, Febbraio referred to more than 200 myokines that mainly promote health as a myokinome.²⁰

2.2. Signaling pathways and small molecular targets

The skeletal muscle signaling pathway is an important target for exercise mimetics. Exercise training activates the interaction between AMPK and PPAR- δ agonists, participating in the regulation of muscle fiber metabolism, contraction, and control of the metabolic phenotype of muscle fibers. Moreover, even in the absence of exercise, oral administration of the AMPK agonist AICAR can target the AMPK-PPAR δ pathway to enhance training adaptability and increase endurance, making AMPK and PPAR- δ considered crucial candidate molecules for exercise mimetics.¹⁴ It is well-established that various forms of exercise have been demonstrated to increase adiponectin mRNA expression and monomeric protein levels of adiponectin, accompanied by an elevation in AdipoR1 protein levels.^{36–38} Adiponectin activates AMPK/SIRT1/PGC-1 α through its receptor AdipoR1 in skeletal muscle, enhancing insulin sensitivity and exercise endurance in obese diabetic mice. Additionally, muscle-specific upregulation of AdipoR1 leads to the restoration of mitochondrial-related gene expression, similar to the recovery observed in mice subjected to exercise under high-fat diet conditions. This suggests that upregulation of AdipoR1 is a potential candidate for exercise mimetics.²⁴ In addition, Li et al.¹⁸ proposed to use signaling pathway molecules activated by exercise as pharmacological targets for the development of “exercise pills” that can improve fatty acid oxidation, skeletal muscle fiber type transformation, mitochondrial biogenesis, angiogenesis, and exercise capacity. As is well known, exercise has the ability to increase the expression of mitochondrial PGC-1 α and promote mitochondrial biogenesis.^{39,40} Further research found that the PGC-1 gene can encode the PGC-1 α 4 protein, and found that in a mouse model of skeletal muscle-specific PGC-1 α 4 overexpression, it can specifically induce mouse IGF-1, inhibit myostatin, and resist cancer cachexia-induced muscle atrophy, while significantly improving skeletal muscle strength and mass.¹⁶ Nine years later, Ruas discovered that after eight weeks of voluntary wheel running exercise, skeletal muscle neurotrophin (NRTN) expression increased, and by constructing a mouse model of NRTN overexpression (human alpha-skeletal actin neurotrophin, HSA-NRTN), it was found that compared with wild-type mice, HSA-NRTN mice had higher integrity of neuromuscular junctions, more stable muscle nerve fiber structure, and optimized local energy metabolism function of muscles, significantly improving endurance, speed, and delaying degenerative motor neuron diseases. In addition, HSN-NRTN mice significantly reduced body weight and improved glucose tolerance by affecting lean body weight and reducing subcutaneous fat accumulation.²⁷ Therefore, the skeletal muscle PGC-1 gene and NRTN is also considered a potential candidate molecule for exercise mimetics. In addition, some specific small molecules that promote overall health are also considered as candidate molecular targets for exercise mimetics. For example, in March 2022, Leiter reported in Cell Metabolism that acute treadmill exercise for four days significantly upregulated the expression of selenoprotein P (SEPP1) in mouse plasma, promoting hippocampal neural progenitor cell proliferation and adult neurogenesis without affecting cell apoptosis. Further research found that exogenous selenium supplementation significantly improved hippocampus-dependent spatial learning and memory ability in mice, indicating that selenium may be one of the potential targets for treating hippocampus-related neurodegenerative and cognitive decline diseases.²⁹ In June 2022, Li reported in Nature that mice and humans after exercise showed increased levels of irisin, a myokine produced by skeletal muscles, in blood circulation. Irisin was found to stimulate beige adipocyte formation and improve metabolic health, suggesting that irisin could be a potential target for the treatment of metabolic

Table 1

The research history of the exercise mimetics.

Author/Year	Journal/Article type	Article	Research results	Target
Ostrowski et al. (1998) ¹²	The Journal of physiology; Article	Evidence that interleukin-6 is produced in human skeletal muscle during prolonged running	The concentrations of IL-6 were measured in marathon runners before, immediately after, and 2 h after the race. It was found that the IL-6 level increased up to 100–150 times higher immediately after the race compared to the pre-race level. The source of IL-6 was not white blood cells and was not related to exercise-induced damage, but mainly produced in the working muscles.	Myokines
Pedersen et al. (2001) ¹³	Current opinion in hematology; Article	Exercise and interleukin-6	The mechanism of IL-6 secretion from working muscle was reviewed, and it was suggested that the increase in muscle-derived IL-6 may be significantly associated with chronic diseases such as obesity and dyslipidemia.	Myokines
Narkar et al. (2008) ¹⁴	Cell; Article	AMPK and PPAR δ agonists are exercise mimetics	The study found that AMPK and PPAR- δ signaling interact with each other, inducing many exercise adaptations in skeletal muscle. Even sedentary mice, when treated with AICAR alone for 4 weeks, showed an upregulation of several protein genes involved in oxidative metabolism and a 44 % increase in running endurance. AMPK-PPAR- δ was reported for the first time as a targeted pathway for oral drug administration to enhance exercise adaptation and even increase endurance without exercise. AMPK and PPAR- δ considered crucial candidate molecules for exercise mimetics.	Signaling pathways
Bostrom et al. (2012) ¹⁵	Nature; Article	A PGC1- α -dependent myokine that drives brown-fat-like development of white fat and thermogenesis	For the first time, a PGC1- α -dependent myokine called irisin was discovered and its function was verified. Exercise induces the secretion of irisin, which acts on white adipose tissue, stimulating the expression of UCP1 and the development of brown fat-like tissue.	small molecular targets
Ruas et al. (2012) ¹⁶	Cell metabolism; Article	A PGC-1 α isoform induced by resistance training regulates skeletal muscle hypertrophy	The PGC-1 α 4, which is strongly expressed in working muscles. In a skeletal muscle-specific PGC-1 α 4 overexpression mouse model, it was found that specific induction of IGF-1 and suppression of myostatin could effectively resist cancer cachexia-induced muscle wasting, while significantly increasing muscle mass and strength. This suggests that PGC-1 α 4 is an important regulatory factor in the modulation of skeletal muscle mass and strength.	small molecular targets
Fan et al. (2013) ¹⁷	Journal of molecular endocrinology; Review	Road to exercise mimetics: targeting nuclear receptors in skeletal muscle	Nuclear receptors and their co-regulators play a key role in regulating skeletal muscle energy metabolism and exercise-induced muscle remodeling. This review provides an overview of the collaborative role of nuclear receptors in skeletal muscle oxidative metabolism and summarizes the latest advances in exercise mimetics targeting nuclear receptors and their co-regulators.	
(Li et al., 2015) ¹⁸	Trends in pharmacological sciences; Review	Exercise Pills: At the Starting Line	This article introduces the concept of “exercise pills” and how they simulate the effects produced by physical exercise. It provides an overview of the biological effects of “exercise pills,” including the transformation of oxidative muscle fiber types, mitochondrial biogenesis, increased fat oxidation, vascular generation, and improved exercise capacity. The article also compares the beneficial effects and molecular mechanisms of sports and candidate exercise drugs.	Review of candidate exercise mimetics
Wall et al. (2016) ¹⁹	Journal of molecular endocrinology; Review	Nuclear receptors and AMPK: can exercise mimetics cure diabetes?	This review summarizes the nuclear receptors and co-regulators that are potential targets for exercise mimetics and evaluates their therapeutic effects on diabetes.	small molecular targets
Whitham et al. (2016) ²⁰	Nature reviews. Drug discovery; Review	The ever-expanding myokinome: discovery challenges and therapeutic implications	The concept of “myokinome” is proposed, and substances that have been discovered and may serve as candidate exercise mimetics are summarized and classified. Techniques for discovering myokines, such as omics technologies, mass spectrometry, and proteomics, are introduced. The target of myokine therapy is elucidated, as are the current challenges of myokinome discovery.	Myokines
Fan et al. (2017) ²¹	Cell metabolism; Review	Exercise Mimetics: Impact on Health and Performance	Exercise mimetics are an alternative therapy for intervening in the decline of health caused by sedentary behavior, while further exploring the scope of application for existing exercise mimetics such as GW501516 and AICAR.	Signaling pathways/ASEM
Findeisen et al. (2019) ¹⁰	Nature; Article	Treatment of type 2 diabetes with the designer cytokine IC7Fc	The concept of exercise mimetics was first truly designed and put into practice for the treatment of T2DM. IL-6 and CNTF were found to regulate metabolic homeostasis, but their therapeutic effects on T2DM were not satisfactory. Therefore, the gp130 binding site on IL-6 was replaced by CNTF and then combined with immunoglobulin G to remove harmful groups and retain effective groups, forming	ASEM

(continued on next page)

Table 1 (continued)

Author/Year	Journal/Article type	Article	Research results	Target
Sanford et al. (2020) ²²	Cell; Article	Molecular Transducers of Physical Activity Consortium (MoTrPAC): Mapping the Dynamic Responses to Exercise	IC7Fc. It can effectively treat T2DM, prevent fat accumulation in the liver, and has minimal side effects. Propose that there are still errors in determining the detailed molecular signaling underlying the beneficial health effects and disease prevention induced by exercise, but establishing a molecular atlas and database of exercise and utilizing multi-omics analysis can greatly promote addressing this issue and treating diseases.	Signaling pathways
Contrepolis et al. (2020) ²³	Cell; Article	Molecular Choreography of Acute Exercise;	This project explores a multimodal and multiomics approach to investigate changes in the metabolome, lipidome, immunome, proteome, transcriptome, and microbiome during maximal oxygen uptake testing, providing new insights for the development of exercise mimetics.	Research ideas of multimethodology
Gubert et al. (2021) ⁷	Nature reviews. Drug discovery; Review	Exercise mimetics: harnessing the therapeutic effects of physical activity	Proposing the new concept of exercise mimetics, and advocating for it to become a new class of therapeutic intervention with a particular focus on its impact on enhancing brain function and cognition, especially in the field of central nervous system diseases.	The concept of exercise mimetics
Iwabu et al. (2021) ²⁴	Communications biology; Article	AdipoR agonist increases insulin sensitivity and exercise endurance in AdipoR-humanized mice	The adiponectin derived from adipocytes has been shown to activate the AMPK/SIRT1/PGC-1 α pathway in skeletal muscle through its receptor AdipoR1, and increase insulin sensitivity and exercise endurance in obese diabetic mice. Therefore, the AdipoR1 signal of adiponectin is considered an important signal for exercise mimetics.	Signaling pathways
Fasipe et al. (2021) ²⁵	Sports Medicine and Health Science; Review	Harnessing the cardiovascular benefits of exercise: Are Nrf2 activators useful?	Exercise promotes adaptive responses by activating Nrf2, thereby improving cardiovascular health, suppressing pathological cardiac remodeling, and alleviating hypertension. Additionally, Nrf2 can be activated through pharmacological means, providing new avenues for the design of exercise mimetics.	small molecular targets
Drake et al. (2021) ²⁶	Proceedings of the National Academy of Sciences of USA; Article	Mitochondria-localized AMPK responds to local energetics and contributes to exercise and energetic stress-induced mitophagy	In both mouse and human skeletal muscle, an energy sensor called "mitoAMPK" exists, and metformin activates mitoAMPK in skeletal muscle without activating AMPK in other parts of the cell. This suggests the importance of mitoAMPK in the treatment of chronic diseases. The team also developed an effective gene model to predict the key steps of mitoAMPK activation, which can be used to promote regular exercise, prevent diseases, and develop effective exercise mimetics.	Signaling pathways
Correia et al. (2021) ²⁷	Cell metabolism; Article	Muscle-secreted neurturin couples myofiber oxidative metabolism and slow motor neuron identity	After eight weeks of rotating exercise, the expression of the neurotrophic factor NRTN in mouse skeletal muscles was found to be elevated along with PGC-1. Additionally, the mRNA of NRTN in the gastrocnemius muscle of rats and humans increased significantly after exercise. A transgenic mouse model overexpressing NRTN (HSA-NRTN) was constructed, and the results showed that HSA-NRTN not only plays a key role in improving neuromuscular junctions, maintaining muscle fiber structure, and regulating local energy metabolism but also significantly improves mouse endurance and speed, as well as slowing down the degenerative disease of motor neurons.	Signaling pathways
De Miguel et al. (2021) ²⁸	Nature; Article	Exercise plasma boosts memory and dampens brain inflammation via clusterin	The study for the first time transferred plasma from active individuals to sedentary individuals in vitro and found that one of the important mechanisms by which exercise improves neurodegenerative diseases is that exercise can produce clusterin, a protein that reduces baseline expression of neuroinflammatory genes and experimentally induced brain inflammation, and has a significant therapeutic effect on neurodegenerative diseases.	small molecular targets
Leiter et al. (2022) ²⁹	Cell metabolism; Article	Selenium mediates exercise-induced adult neurogenesis and reverses learning deficits induced by hippocampal injury and aging	Treadmill exercise significantly upregulates the expression of SEPP1 in mice and promotes the proliferation of NPC and the potential for neuronal lineage differentiation in the hippocampus, without affecting cell apoptosis. Animal cognitive function tests have found that exogenous selenium supplementation can significantly improve mice's hippocampus-dependent spatial learning and memory abilities, indicating that selenium may be a key substance for restoring age-related hippocampal functional defects and effectively reversing hippocampal injury-related cognitive decline. These findings provide new ideas for developing exercise mimetics.	small molecular targets
Li et al. (2022) ³⁰	Nature; Article	An exercise-inducible metabolite that suppresses feeding and obesity	After treadmill exercise, the small molecule metabolite Lac-Phe significantly increased in the plasma of mice and horses. Lac-Phe is synthesized by CNDP2 through the condensation of lactate and phenylalanine. Lac-Phe can	small molecular targets

(continued on next page)

Table 1 (continued)

Author/Year	Journal/Article type	Article	Research results	Target
Kurz et al. (2022) ¹¹	Cancer Cell; Article	Exercise-induced engagement of the IL-15/IL-15Rα axis promotes anti-tumor immunity in pancreatic cancer	effectively inhibit the food intake of mice, improve glucose homeostasis, reduce fat mass and body weight, and prevent obesity. Human studies have found that the level of Lac-Phe in the plasma continues to increase after exercise, and it is one of the key metabolites regulating human exercise. Aerobic exercise can increase adrenaline secretion and upregulate the expression of CD8 ⁺ T cells and IL-15Rα, which are sensitive to IL-15, thereby leading to the accumulation of CD8 ⁺ T cells and IL-15Rα on pancreatic cancer cells and ultimately killing the tumor cells and inhibiting the development of pancreatic cancer. Based on this mechanism, Novartis has developed an IL-15 agonist called NIZ985, which can mimic the effect of exercise in enhancing the IL-15/IL-15Rα signaling pathway, reduce potential inflammatory reactions, and achieve sustained anti-tumor effects.	ASEM

Note. UCP1, uncoupling protein 1; PGC-1α4, PGC-1 gene encodes a protein; IGF-1, insulin-like growth factor-1; T2DM, type 2 diabetes; AMPK, AMP-activated protein kinase; SEPP1, selenium protein P; NPC, neural progenitor cells; Lac-Phe, N-lactoyl-phenylalanine; CNDP2, cytosolic non-specific dipeptidase 2; IL-15, interleukin 15; ASEM, Artificially synthesized exercise mimetics.

diseases.³⁰ Therefore, signaling pathways and small molecular targets related to skeletal muscles and exercise mimetics have been extensively studied, and may provide new insights and strategies for the development of exercise mimetics as pharmacological interventions for various health conditions.

2.3. Artificially synthesized exercise mimetics

In October 2019, Mark Febbraio of Gavin Institute of Medicine, Sydney, Australia reported in Nature a method for artificially synthesizing exercise mimetics and designed a small molecule compound, IC7Fc,¹⁰ which has therapeutic effects on type 2 diabetes. This provides a new strategy for exercise mimetics in the field of metabolic disease treatment. In July 2022, Professor Dafna Bar-Sagi of Grossman Medical College of new york University reported the anti-tumor effect of exercise in Cancer Cell and developed exercise mimetics. Research has revealed that the IL-15 agonist NIZ985 can simulate exercise-enhanced IL-15/IL-15Rα signaling pathways, thereby promoting the destruction of tumor cells.¹¹ This presents a novel avenue for the application of exercise mimetics in the field of cancer therapy.

It should be pointed out that in the field of exercise mimetics research, breakthroughs and explorations are still ongoing. It is urgent to promote the development of exercise mimetics through the interdisciplinary integration of sports science, pharmacy, and other fields.

3. The relationship between exercise mimetics and exercise

Exercise mimetics use the health benefits produced by exercise as a source of drug design to seek potential new methods for disease treatment. This method of treatment, which simulates or enhances the therapeutic effects or health benefits of exercise, is known as exercise mimetics. In recent years, exercise mimetics have gradually emerged as a research focal point. The changing lifestyles and working habits of individuals have resulted in a lack of physical activity and an increase in sedentary behavior. Consequently, a global estimate indicates that 31.1 % of adults fail to meet the minimum recommended exercise standards for maintaining health.⁴¹ This indirectly leads to a dramatic increase in the incidence of “modern civilization diseases” (including obesity, type 2 diabetes,⁴² and cardiovascular disease, etc.⁴³). Appropriate exercise training is one of the important means to prevent and treat these diseases, Including moderate-intensity aerobic training,⁴⁴ high-intensity aerobic training,⁴⁵ and lower limb resistance training.⁴⁶ For example, adults with type 2 diabetes and metabolic syndrome who underwent three months of aerobic exercise training and high-intensity exercise training observed a decrease in insulin resistance and systemic

inflammation markers, which appeared to be related to an increase in peak oxygen uptake rather than weight loss.⁴⁷ Furthermore, during exercise, immune cells in the tumor infiltrate in large numbers, leading to a reduction of over 60 % in the incidence and growth of tumors in several mouse models, as well as the prevention of tumor or cancer-related death.⁴⁸ Exercise training is also an effective method for treating most neurological disorders.⁴⁹ By injecting circulating blood factors from mice after exercise into sedentary mice, baseline neuro-inflammatory gene expression and experimentally induced brain inflammation can be reduced, and cognitive function can be improved.²⁸

Considering the insufficient amount of human exercise prescriptions and the human and economic burden caused by these diseases, and taking into account the health benefits of exercise, including neurological and psychiatric disorders,^{50,51} cardiovascular diseases,⁵² metabolism and inflammation,⁵³ and tumors,⁵⁴ the academic community has gradually focused on alternative therapies using exercise mimetics.⁵⁵ Exercise mimetics are closely related to human physical activity. Different types and intensities of exercise have different health benefits. In theory, the greater the intensity of exercise, the more significant the improvement in various aspects of heart rate, metabolism, and physiological function.^{56,57} In addition, different genetic characteristics, environments, cells, molecules, or system levels of individuals can also lead to differences in the exercise benefits produced by different types of exercise (aerobic or resistance), including improvement or inhibition of organismal health benefits.^{58–60} Based on the design concept of exercise mimetics, this study will focus on known forms of exercise that can produce therapeutic effects, and summarize and determine the exercise molecular targets that are related to therapeutic effects as effective sources of exercise mimetic molecules.

It is worth noting that although exercise mimetics may not be able to fully mimic the wide range of benefits of exercise, the emerging candidate exercise mimetics^{14,61,62} are a favorable potential choice for people who cannot exercise regularly due to obesity, amputation, spinal cord injury, metabolic diseases, and musculoskeletal or cardiovascular diseases. The design concept of these exercise mimetics is to activate signal molecules related to exercise, which are logically considered to be effective pharmacological targets for such exercise mimetics.^{7,63,64} Therefore, the development of exercise mimetics is feasible.

4. Candidate molecular sources of exercise mimetics

The human body is a highly complex system that maintains homeostasis through dynamic interactions among multiple levels of entities, ranging from cells, tissues, organs to systems, through various means such as neural, endocrine and exocrine pathways. Exercise, as an

external stimulus, can disrupt and modulate this complex system, and appropriate exercise can have beneficial effects on human health. These changes include the metabolic, proteomic, genomic, immunological, and gut microbiota profiles, which can serve as potential sources of exercise mimetics. Exercise mimetics can be roughly classified into nine categories based on their sources and effects (Fig. 1).

- ① Neurotrophins, such as brain-derived neurotrophic factor (BDNF). Exercise can upregulate BDNF expression, which plays a critical role in neuronal survival, proliferation, maturation, and growth in the brain (including the hippocampus, hypothalamus, and cortex).⁶⁵ Clinical studies have found that BDNF has a positive therapeutic effect on clinical models of Alzheimer's disease (AD), Huntington's disease (HD), and other brain disorders.^{20,66,67} It is worth noting that BDNF is not only produced in the brain but also observed to increase in skeletal muscles after exercise.⁶⁸ Further research has confirmed that exercise can upregulate BDNF in skeletal muscles and participate in exercise-induced muscle regeneration.⁶⁹ In addition, BDNF can also affect tissue metabolism via autocrine or paracrine signaling, including fat oxidation,⁷⁰ insulin resistance, and playing a crucial role in angiogenesis, cardiovascular development, and cardiac protection.⁷¹
- ② Neurotransmitters and neuropeptides, including glutamate (GLU), serotonin (5-HT), and neuropeptide Y (NPY). Exercise can regulate the level of GLU,⁷² which can stimulate the production of BDNF as a neurotransmitter, thereby changing the sensitivity of neurons to GLU and Ca²⁺ stability and ultimately changing the plasticity of neurons.⁷³ 5-HT is an important neurotransmitter for exercise-induced adult neurogenesis⁷⁴ and a key factor in hippocampal neurogenesis.⁷⁵ In addition, exercise can upregulate the plasma NPY level in the rat hypothalamus,⁷⁶ which is closely related to increased anxiety, cognitive impairment, and changes in hippocampal synaptic plasticity.⁷⁷
- ③ Intestinal flora and their metabolites, such as the phylum Firmicutes and short-chain fatty acids (SCFA), etc. With the development of related technologies such as genomics and bioinformatics, the relationship between fecal intestinal flora and their metabolites and health has gradually been discovered.




Fig. 1. Molecular mediators and drug-making targets of exercise mimetics.

Exercise can change the composition and abundance of intestinal flora, and then improve overall gut health and promote organismal health.^{78–80} In addition, the intestinal flora has a bidirectional communication with various organs and tissues of the body, such as the cardiovascular system (gut-heart axis), the immune system (gut-immune axis), the skeletal muscle (gut-muscle axis), and the brain (gut-brain axis),⁸¹ making it a potential target for disease treatment.

- ④ Myokines, including AMPK or PPAR agonists such as AICAR and GW501516, muscle-derived neurotrophic factors (BDNF (muscle)), as well as myokines such as irisin, insulin-like growth factor 1 (IGF1), and cathepsin B. Exercise can induce skeletal muscle remodeling and related metabolic pathways through the AMPK-sirtuin 1 (SIRT1)-PPAR δ pathway, which is crucial for muscle energy metabolism and mitochondrial biogenesis, and is one of the main targets for developing exercise mimetics.^{21,82} AICAR and GW501516 (a selective PPAR δ agonist) have been shown to indirectly improve memory function and neurogenesis (possibly through the IL-6-mediated increase in BDNF⁸³ and cathepsin B⁸⁴).⁸⁵ In addition, metformin (targeting AMPK) has a positive effect on cognitive function in an Alzheimer's disease mouse model.⁸⁶ These are potential molecules for developing exercise mimetics.
- ⑤ Vascular growth factors, exercise-induced increases in vascular endothelial growth factor (VEGF), VEGFR2, Ang2, and CD34⁺ cells (vascular generation markers) can effectively prevent the occurrence and development of stroke and related diseases.⁸⁷ In addition, studies have found that the use of SU1498 (a VEGF receptor FLK1 inhibitor) prevents exercise-induced improvement of depression-like behavior, indicating that VEGF is a key factor in the exercise anti-depression effect.⁸⁸ Moreover, VEGF is also a key substance for exercise-induced adult hippocampal neurogenesis.⁸⁹ Moreover, recent advancements in research suggest that exercise triggers adaptive alterations in early lymphatic behavior, inducing an upregulation of lymphangiogenesis. Key regulatory factors in muscle lymphangiogenesis include vascular endothelial growth factor receptor-3 (VEGFR-3) and its ligands, VEGF-C and VEGF-D.⁹⁰ Notably, recent investigations emphasize the involvement of lymphangiogenesis in exercise-induced physiological cardiac hypertrophy. This process entails the elevation of VEGFR3 levels through exercise, and the initiation of cardiac lymphangiogenesis requires the activation of VEGFR3.⁹¹ Additionally, a growing body of evidence highlights the role of lymphangiogenesis in tissue repair and regeneration.⁹² In summary, this underscores the potential contributions of lymphangiogenesis and its associated receptor proteins in facilitating tissue recovery and regeneration through exercise. Consequently, lymphangiogenesis and its related secretory receptor proteins emerge as promising therapeutic targets.
- ⑥ Extracellular vesicles, which refer to molecular structures that can contain and transport biological materials (including DNA, RNA, proteins, and lipid molecules) between cells, are connecting tissues throughout the body.⁹³ The number of extracellular vesicles in circulation increases 2–4 times after exercise, which is considered one of the important pathways for inter-tissue signaling during exercise and has positive promoting effects on health.^{94,95}
- ⑦ Non-coding RNA, such as microRNAs (miRNAs). Exercise can regulate the expression of miRNAs.⁹⁶ miRNAs are the most extensively studied type of non-coding RNA among small non-coding RNAs (sncRNAs) and can act as “master regulators” by regulating gene expression,⁹⁷ with positive therapeutic effects.⁹⁸ It has also been shown that long non-coding RNA (lncRNA) is responsive to physical exercise.⁹⁹ lncRNAs play a crucial role in regulating signaling pathways associated with DNA, mRNA, and protein, thereby influencing gene expression in




Fig. 3. Schematic diagram of exercise mimetics regulating endocrine system.

response to metabolic disorders caused by fasting response, etc.^{126,127} Intense exercise can induce FGF21 expression and increase FGF21 levels in circulation.¹²⁸ In addition, FGF21 can stimulate glucose uptake and fatty acid oxidation in metabolic tissues such as muscle and fat, effectively preventing the body from entering a state of hunger.^{129,130} Animal experiments have also confirmed that in FGF21 overexpression mouse models, high doses of FGF21 significantly improved insulin sensitivity and resistance to metabolic syndrome.¹³¹ This suggests that FGF21 could be a potential candidate exercise mimetic for improving metabolic diseases. In the future, the focus should be on exploring the normal physiological metabolic response induced by FGF21 production (not caused by intense exercise or high-intensity exercise) or researching corresponding effective activators; at the same time, the activation process of PPAR- α should be avoided (should not be produced by hunger), and corresponding blocking methods should be studied.

PPAR- γ is also one of the important targets for treating metabolic diseases. PPAR- γ plays a crucial role in the development of adipose tissue by regulating the expression of lipid metabolism genes and adipokines with key metabolic functions, such as adiponectin¹³². There are many factors that influence PPAR- γ activity and the expression of target genes in adipose tissue, including exercise and other stressors, suggesting that PPAR- γ is a potential target for the development of novel exercise mimetics.¹³³ Thiazolidinediones (TZDs) are a classic class of drugs that activate PPAR- γ and counteract diabetes.¹³⁴ TZDs enhance the development and lipid handling of adipocytes by activating PPAR- γ , increasing the uptake of circulating fatty acids into adipose tissue, and reduce insulin resistance.¹³⁵ Recent studies have found that TZDs inhibit the phosphorylation of the S273 residue of PPAR- γ by cyclin-dependent kinase 5 (CDK5), resulting in changes in the expression pattern of PPAR- γ target genes in white adipose tissue, which may be a potential factor preventing obesity.¹³⁶ Phosphorylation of S273 can bind to thyroid hormone receptor-associated protein 3 (Thrap3), which is crucial for PPAR- γ activity in obesity.¹³⁷ Unlike TZD agonist drugs, studies have found that non-agonist PPAR- γ ligands (UHC1) effectively block the S273 phosphorylation of PPAR- γ and improve insulin sensitivity and diet-induced obesity in the treatment of T2D.¹³⁸ These studies suggest that PPAR- γ can be used as a therapeutic target for diabetes (with Thrap3 and CDK5 as key influencing factors) and as a candidate molecule for exercise mimetics.

In addition, the adiponectin receptor AdipoR1 signaling pathway is considered an important signal for exercise mimetics. Studies have found that in AdipoR1 gene knockout mouse models, the use of AdipoR agonists (AdipoRon) can exert beneficial effects through AdipoR in muscles, increasing insulin sensitivity and exercise endurance in AdipoR mice.²⁴ This suggests that adiponectin from adipocytes displays its beneficial effects in skeletal muscles through its receptor AdipoR1, activating AMPK/SIRT1/PGC-1 α , and increasing insulin sensitivity and

exercise endurance in obese diabetic mice.²⁴ Adiponectin and its receptor can be potential exercise mimetic molecules for treating metabolic diseases.

5.3. Circulatory system

The circulatory system, especially the cardiovascular system, is also one of the future research directions of exercise mimetics materials. Its main mechanism is shown in Fig. 4.

The IGF1/PI3K/AKT signaling pathway is a key mechanism in exercise-induced physiological myocardial hypertrophy and cardiac protection. Exercise induces the secretion of IGF1, activates IGF1R tyrosine kinase (TK),^{139,140} and recruits PI3K.¹⁴¹ PI3K further converts PIP2 to PIP3 in the plasma membrane,¹⁴² which recruits PDK1 and AKT to the plasma membrane and then activates and phosphorylates PDK1 and AKT,¹⁴³ which is the IGF1/PI3K/AKT signaling pathway. Activation of this pathway can effectively promote myocardial cell contraction, which is crucial in regulating exercise-induced physiological myocardial hypertrophy. It plays a protective role in the heart by promoting myocardial cell contraction, improving survival rate, reducing myocardial cell apoptosis, inhibiting pathological hypertrophy, promoting angiogenesis, and so on.^{144–146}

Noncoding RNAs also play an important role in the cardiovascular system. Exercise upregulates miR-222 and LncRNA CPhar in the heart, while downregulating lncExACT1.¹⁴⁷ LncRNA CPhar binds to DDX17, sequestering C/EBP β and thereby inhibiting ATF7 transcription.¹⁴⁸ Additionally, lncExACT1 can bind to miR-222, and exercise downregulates lncExACT1, leading to the release of more miR-222, which helps to induce exercise-induced physiological cardiac hypertrophy.¹⁴⁹ lncExACT1 also actively regulates the most similar protein-coding gene, DCHS2, and reduces lncExACT1 activation of Yes-associated protein (YAP).¹⁴⁸ These noncoding RNAs are involved in regulating exercise-induced physiological cardiac hypertrophy and have a protective effect on myocardial injury and pathological cardiac hypertrophy.¹⁵⁰ Furthermore, exercise can regulate miR-34a and miR-126, inducing endothelial cell-derived small extracellular vesicles (sEVs) containing miR-342-5p and cardiac sEVs derived from brown adipose tissue containing miR-125b-5p, miR-128-3p, and miR-30d-5p to




Fig. 4. Schematic diagram of the effects of exercise mimetics on the regulation of the circulatory system.

prevent myocardial injury and cardiac remodeling.¹⁵¹ These findings suggest that the IGF1/PI3K/AKT signaling pathway and noncoding RNAs and their regulatory signaling pathways can serve as potential targets for exercise mimetics, as well as potential molecular targets for treating cardiovascular diseases.

5.4. Neurological system

The neurological system is currently a hot research topic in the field of exercise mimetics. The main mechanism of its action is shown in Fig. 5.

Another key target system for exercise mimetics is the neurological system, which enhances neuronal plasticity, particularly in adult neurogenesis and synaptic plasticity, improving cognitive function and preventing the occurrence of neurological disorders.^{51,82} Different molecular processes and signaling pathways are associated with the enhancement of adult neurogenesis and synaptic activity after exercise, including signal transduction pathways such as BDNF, tyrosine receptor kinase B (TrkB), glutamatergic, dopaminergic, and adrenergic systems.^{77,152–155} In addition, at the cellular level, the effects of exercise on glial generation, neurogenesis, synaptic genesis, and angiogenesis lead to structural and functional changes, ultimately enhancing brain function, including cognition.¹⁵⁶ Studies have found that exercise can increase the volume of the hippocampal dentate gyrus/CA3 and enhance memory in young people.¹⁵⁷ Furthermore, extensive evidence supports the enhancement of adult neurogenesis and synaptic plasticity induced by exercise.^{158–160} Exercise has also been shown to reverse learning deficits caused by hippocampal injury by promoting adult neurogenesis.¹⁶¹ It is evident that exercise is one of the important factors that affect the neurological system. The main pathways by which exercise affects the neurological system include: (1) exercise can affect molecules such as BDNF, neurotransmitters, neuropeptides, and non-coding RNAs, promoting adult neurogenesis and enhancing hippocampal synaptic

plasticity⁷; (2) exercise can induce the expression of clusterin and selenoprotein P (SEPP1). Clusterin can reduce baseline expression of inflammatory genes and experimentally induced brain inflammation, thus preventing the occurrence of neurodegenerative diseases.²⁸ SEPP1 can promote proliferation of hippocampal NPCs and their neuronal lineage potential, effectively reversing cognitive decline related to hippocampal injury.²⁹ These findings suggest that exercise mimetics can mimic the expression of brain-derived substances or induce the generation of certain key factors, thereby enhancing the neuroprotective effect and improving brain function.

5.5. Other targeted organs and functions of exercise mimetics

There has been less research on exercise mimetics in other systems of the human body, such as the reproductive, respiratory, digestive, and urinary systems. The immune system is a promising molecular target for exercise mimetics, and may play a role in cancer treatment. It is well known that the mechanism of cancer development is complex, involving multi-factorial and multi-step reactive processes that are closely related to lifestyle, environment, genetics, and other factors. Treatment is difficult, and the main diagnostic and therapeutic methods currently available are drug therapy and surgery.¹⁶² However, with the development of exercise science and drug research, scientists have discovered that exercise is a potential approach for cancer treatment. For example, the IL-15 agonist NIZ985 has demonstrated the ability to mimic the effects of exercise by enhancing the IL-15/IL-15R α signaling pathway, leading to the inhibition of pancreatic cancer.¹¹ NIZ985 may become a new exercise mimetic for cancer treatment. This suggests that exercise mimetics are likely to be one of the potential directions for cancer treatment, and in the future, more exercise targets for the treatment of other cancers need to be explored, providing new directions for cancer treatment.

6. Future research trends of exercise mimetics

Exercise mimetics have enormous potential for the prevention and treatment of human diseases. However, the current number of developed and utilized exercise mimetics is extremely limited, due to the complex physiological, metabolic, and endocrine changes and adaptations that occur during exercise. It is difficult for a single exercise mimetic to summarize all such systemic changes. However, it should be noted that exercise mimetics only need to mimic or enhance a subset of therapeutic effects of exercise to have clinical impact, which is the main research goal of current exercise mimetics. In addition, the development of exercise mimetics requires a deeper understanding of the molecular and cellular mechanisms of beneficial therapeutic effects induced by exercise, including changes at the molecular, cellular, tissue, organ, and systemic levels. However, understanding this mechanism is challenging, as exercise itself directly or indirectly affects changes at various levels of the body.¹⁶³ Despite the difficulty in elucidating this complex mechanism of action, current research on exercise mimetics has enabled us to begin utilizing some of their therapeutic potential.¹⁶⁴

Possible research directions in the future include utilizing interdisciplinary technologies such as bioinformatics and artificial intelligence to investigate the characteristics of existing motion simulation agents. These studies can help identify common structures of potential artificial motion simulation agents. For example, AlphaFold2 is a technique used to predict the key conformational changes of drug aggregation and the associated protein subunits, enabling precise prediction of the three-dimensional structural features of drugs.¹⁶⁵ There are many applications of artificial intelligence in the field of medicine, including: 1) Identification of Biomarkers or Biological Targets: AI accelerates the screening process of candidate molecules for exercise mimetics by simulating their structure and properties, leading to the identification of optimal candidates. 2) Simulation of Clinical Experiments and Outcome Prediction: AI, by integrating candidate exercise mimetic molecules,




Fig. 5. Schematic diagram of effects of exercise mimetics on brain structure and function.

simulates potential exercise mimetics, predicting their possible clinical trial results and effects, and provides prior feedback and adjustment suggestions. 3) Prediction and Monitoring of Drug Side Effects: AI analyzes medical records and drug databases to forecast potential side effects and adverse reactions associated with exercise mimetics. This capability ensures the timely detection and adjustment of safety issues related to these drugs. In general, the application of artificial intelligence in the field of medicine can greatly improve the efficiency and accuracy of drug development, enhance the success rate and effectiveness of clinical trials, help doctors make personalized treatment decisions, and ensure patient drug safety, which can also be used to guide the development of motion simulation drugs. Furthermore, motion simulation may also be applied in preventive medicine, where genomics and selective biomarker arrays can be used to identify high-risk populations who can be targeted for specific treatments to delay or prevent the onset of specific diseases. Additionally, further exploration of the targeted systems of motion simulation agents is necessary to enrich the study of the biological improvement effects of these agents.

7. Summary

Exercise mimetics are still in the early stages of research. The beneficial effects of exercise-induced cytokines and microbiota on the body have led to the development of exercise mimetics, a term that encompasses both such substances and living organisms, including new compounds synthesized through biotechnology. Exercise mimetics include small molecules, peptides, antibodies, non-coding RNA, gut microbiota, and epigenetic editing constructs, among others. Additionally, research on exercise mimetics is complementary to that on exercise itself, with the goal of using the features and functional properties of the health benefits induced by simulated exercise to design new potential methods for treating diseases. Currently, it has been found that the target systems of exercise mimetics include the exercise system, circulatory system, and nervous system, and there is likely a biological improvement effect in the immune and endocrine metabolism systems as well. In the future, the development of exercise mimetics should be aided by technologies such as bioinformatics and artificial intelligence.

Author contributions

Yuping Zhu performed the literature search and data analysis, and Gang Song drafted and critically revised the manuscript.

Funding

This work was supported by the National Social Science Fund of China [grant numbers 21BTY092], and the Graduate Students Innovation Program of Chongqing [grant numbers CYS22175].

Availability of data and materials

Data sharing is not applicable to this article as no datasets were generated or analysed during the current study.

Ethics approval and consent to participate

Not applicable.

Consent for publication

All authors agreed on the publication of the current version of manuscript.

Declaration of competing interest

The authors declare no conflict of interest.

Acknowledgements

Not applicable.

Abbreviations

EIM	exercise is medicine
GP130	glycoprotein 130
IL-6	interleukin 6
CNTF	ciliary neurotrophic factor
LIF	leukemia inhibitory factor
IgG	immunoglobulin G
UCP1	uncoupling protein 1
AMPK	AMP-activated protein kinase
IGF-1	insulin-like growth factor-1
SEPP1	selenium protein P-1
NPC	neural progenitor cells
Lac-Phe	N-lactoyl-phenylalanine
CNDP2	cytosolic non-specific dipeptidase 2
NRTN	neurotrophin
HSA-NRTN	human alpha-skeletal actin neurotrophin
BDNF	brain-derived neurotrophic factor
AD	Alzheimer's disease
HD	Huntington's disease
GLU	glutamate
NPY	neuropeptide Y
SCFA	short-chain fatty acids
SIRT1	sirtuin 1
VEGFs	vascular endothelial growth factors
HDAC	histone deacetylase
HAT	histone acetylation transferase
ROS	reactive oxygen species
FGF21	fibroblast growth factor 21
CDK5	cyclin-dependent kinase 5
TK	tyrosine kinase
sEVs	small extracellular vesicles
TrkB	tyrosine receptor kinase B

References

1. Van Norman GA. Overcoming the declining trends in innovation and investment in cardiovascular therapeutics. *JACC (J Am Coll Cardiol): Basic to Translational Science*. 2017;2:613–625. <https://doi.org/10.1016/j.jacbs.2017.09.002>.
2. Luan X, Tian X, Zhang H, Huang R, Li N, et al. Exercise as a prescription for patients with various diseases. *J Sport Health Sci*. 2019;8:422–441. <https://doi.org/10.1016/j.jshs.2019.04.002>.
3. Pedersen BK, Saltin B. Exercise as medicine - evidence for prescribing exercise as therapy in 26 different chronic diseases. *Scand J Med Sci Sports*. 2015;25:1–72. <https://doi.org/10.1111/sms.12581>.
4. Timmons JA, Baar K, Daviden PK, Atherton PJ. Is irisin a human exercise gene? *Nature*. 2012;488:E9–E10. <https://doi.org/10.1038/nature11364>.
5. Cento AS, Leigheb M, Caretti G, Penna F. Exercise and exercise mimetics for the treatment of musculoskeletal disorders. *Curr Osteoporos Rep*. 2022;20:249–259. <https://doi.org/10.1007/s11914-022-00739-6>.
6. Chen J, Zhou R, Feng Y, Cheng L. Molecular mechanisms of exercise contributing to tissue regeneration. *Signal Transduct Targeted Ther*. 2022;7:383. <https://doi.org/10.1038/s41392-022-01233-2>.
7. Gubert C, Hannan AJ. Exercise mimetics: harnessing the therapeutic effects of physical activity. *Nat Rev Drug Discov*. 2021;20:862–879. <https://doi.org/10.1038/s41573-021-00217-1>.
8. Mcomish CE, Hannan AJ. Enviromimetics: exploring gene environment interactions to identify therapeutic targets for brain disorders. *Expert Opin Ther Targets*. 2007;11:899–913. <https://doi.org/10.1517/14728222.11.7.899>.
9. Hannan AJ. Molecular mediators, environmental modulators and experience-dependent synaptic dysfunction in huntington's disease. *Acta Biochim Pol*. 2004;51:415–430.
10. Findeisen M, Allen TL, Henstridge DC, Kammoun H, Brandon AE, et al. Treatment of type 2 diabetes with the designer cytokine ic7fc. *Nature*. 2019;574:63–68. <https://doi.org/10.1038/s41586-019-1601-9>.
11. Kurz E, Hirsch CA, Dalton T, Shadaloe SA, Khodadadi-Jamayran A, et al. Exercise-induced engagement of the il-15/il-15 α axis promotes anti-tumor immunity in pancreatic cancer. *Cancer Cell*. 2022;40:720–737. <https://doi.org/10.1016/j.ccell.2022.05.006>.

12. Ostrowski K, Rohde T, Zacho M, Asp S, Pedersen BK. Evidence that interleukin-6 is produced in human skeletal muscle during prolonged running. *J Physiol.* 1998;508: 949–953. <https://doi.org/10.1111/j.1469-7793.1998.949bp.x>.
13. Pedersen BK, Steensberg A, Schjerling P. Exercise and interleukin-6. *Curr Opin Hematol.* 2001;8:137–141. <https://doi.org/10.1097/00062752-200105000-00002>.
14. Narkar VA, Downes M, Yu RT, Embler E, Wang Y, et al. Ampk and ppar δ agonists are exercise mimetics. *Cell.* 2008;134:405–415. <https://doi.org/10.1016/j.cell.2008.06.051>.
15. Boström P, Wu J, Jedrychowski MP, Korde A, Ye L, et al. A pgc1- α -dependent myokine that drives brown-fat-like development of white fat and thermogenesis. *Nature.* 2012;481:463–468. <https://doi.org/10.1038/nature10777>.
16. Ruas JL, White JP, Rao RR, Kleiner S, Brannan KT, et al. A pgc-1 α isoform induced by resistance training regulates skeletal muscle hypertrophy. *Cell.* 2012;151: 1319–1331. <https://doi.org/10.1016/j.cell.2012.10.050>.
17. Fan W, Atkins AR, Yu RT, Downes M, Evans RM. Road to exercise mimetics: targeting nuclear receptors in skeletal muscle. *J Mol Endocrinol.* 2013;51: T87–T100. <https://doi.org/10.1530/JME-13-0258>.
18. Li S, Laher I. Exercise pills: at the starting line. *Trends Pharmacol Sci.* 2015;36: 906–917. <https://doi.org/10.1016/j.tips.2015.08.014>.
19. Wall CE, Yu RT, Atkins AR, Downes M, Evans RM. Nuclear receptors and ampk: can exercise mimetics cure diabetes? *J Mol Endocrinol.* 2016;57:R49–R58. <https://doi.org/10.1530/JME-16-0073>.
20. Whitham M, Febbraio MA. The ever-expanding myokinome: discovery challenges and therapeutic implications. *Nat Rev Drug Discov.* 2016;15:719–729. <https://doi.org/10.1038/nrd.2016.153>.
21. Fan W, Evans RM. Exercise mimetics: impact on health and performance. *Cell Metabol.* 2017;25:242–247. <https://doi.org/10.1016/j.cmet.2016.10.022>.
22. Sanford JA, Nogiec CD, Lindholm ME, Adkins JN, Amar D, et al. Molecular transducers of physical activity consortium (motrap): mapping the dynamic responses to exercise. *Cell.* 2020;181:1464–1474. <https://doi.org/10.1016/j.cell.2020.06.004>.
23. Contrepois K, Wu S, Moneghetti KJ, Hornburg D, Ahadi S, et al. Molecular choreography of acute exercise. *Cell.* 2020;181:1112–1130. <https://doi.org/10.1016/j.cell.2020.04.043>.
24. Iwabu M, Okada-Iwabu M, Tanabe H, Ohuchi N, Miyata K, et al. Adipor agonist increases insulin sensitivity and exercise endurance in adipor-humanized mice. *Commun Biol.* 2021;4:45. <https://doi.org/10.1038/s42003-020-01579-9>.
25. Fasipe B, Li S, Laher I. Harnessing the cardiovascular benefits of exercise: are nrf2 activators useful? *Sports Medicine and Health Science.* 2021;3:70–79. <https://doi.org/10.1016/j.smhs.2021.04.002>.
26. Drake JC, Wilson RJ, Laker RC, Guan Y, Spaulding HR, et al. Mitochondria-localized ampk responds to local energetics and contributes to exercise and energetic stress-induced mitophagy. *Proc Natl Acad Sci USA.* 2021;118. <https://doi.org/10.1073/pnas.2025932118>.
27. Correia JC, Kelahmetoglu Y, Jannig PR, Schweingruber C, Shvaikovskaya D, et al. Muscle-secreted neurturin couples myofiber oxidative metabolism and slow motor neuron identity. *Cell Metabol.* 2021;33:2215–2230. <https://doi.org/10.1016/j.cmet.2021.09.003>.
28. De Miguel Z, Khoury N, Betley MJ, Lehallier B, Willoughby D, et al. Exercise plasma boosts memory and dampens brain inflammation via clusterin. *Nature.* 2021;600:494–499. <https://doi.org/10.1038/s41586-021-04183-x>.
29. Leiter O, Zhuo Z, Rust R, Wasielewska JM, Grönnert L, et al. Selenium mediates exercise-induced adult neurogenesis and reverses learning deficits induced by hippocampal injury and aging. *Cell Metabol.* 2022;34:408–423. <https://doi.org/10.1016/j.cmet.2022.01.005>.
30. Li VL, He Y, Contrepois K, Liu H, Kim JT, et al. An exercise-inducible metabolite that suppresses feeding and obesity. *Nature.* 2022;606:785–790. <https://doi.org/10.1038/s41586-022-04828-5>.
31. Pedersen BK, Steensberg A, Fischer C, Keller C, Keller P, et al. Searching for the exercise factor: is il-6 a candidate? *J Muscle Res Cell Motil.* 2003;24:113–119. <https://doi.org/10.1023/a:1026070911202>.
32. Winocour PH, Durrington PN, Bhatnagar D, Mbewu AD, Ishola M, et al. A cross-sectional evaluation of cardiovascular risk factors in coronary heart disease associated with type 1 (insulin-dependent) diabetes mellitus. *Diabetes Res Clin Pract.* 1992;18:173–184. [https://doi.org/10.1016/0168-8227\(92\)90143-f](https://doi.org/10.1016/0168-8227(92)90143-f).
33. Febbraio MA, Hiscock N, Sacchetti M, Fischer CP, Pedersen BK. Interleukin-6 is a novel factor mediating glucose homeostasis during skeletal muscle contraction. *Diabetes.* 2004;53:1643–1648. <https://doi.org/10.2337/diabetes.53.7.1643>.
34. Pedersen BK, Febbraio M. Muscle-derived interleukin-6—a possible link between skeletal muscle, adipose tissue, liver, and brain. *Brain Behav Immun.* 2005;19: 371–376. <https://doi.org/10.1016/j.bbi.2005.04.008>.
35. A M, Wallas TE, Zhou H, Draga-Coleta SV, Gorgulla C, et al. Irisin acts through its integrin receptor in a two-step process involving extracellular hsp90 α . *Mol Cell.* 2023;83:1903–1920. <https://doi.org/10.1016/j.molcel.2023.05.008>.
36. Martinez-Huenchullan SF, Maharjan BR, Williams PF, Tam CS, McLennan SV, Twigg SM. Skeletal muscle adiponectin induction depends on diet, muscle type/activity, and exercise modality in c57bl/6 mice. *Phys Rep.* 2018;6, e13848. <https://doi.org/10.14814/phy2.13848>.
37. Dai Y, Pang J, Gong H, Fan W, Zhang TM. Roles and tissue source of adiponectin involved in lifestyle modifications. *J Gerontol Ser A-Biol Sci Med Sci.* 2013;68: 117–128. <https://doi.org/10.1093/gerona/gls131>.
38. Goto A, Ohno Y, Ikuta A, Suzuki M, Ohira T, et al. Up-regulation of adiponectin expression in antigravitational soleus muscle in response to unloading followed by reloading, and functional overloading in mice. *PLoS One.* 2013;8, e81929. <https://doi.org/10.1371/journal.pone.0081929>.
39. Safdar A, Little JP, Stokl AJ, Hettinga BP, Akhtar M, Tarnopolsky MA. Exercise increases mitochondrial pgc-1 α content and promotes nuclear-mitochondrial cross-talk to coordinate mitochondrial biogenesis. *J Biol Chem.* 2018;293:4953. <https://doi.org/10.1074/jbc.EC118.002682>.
40. Sutherland LN, Bomhof MR, Capozzi LC, Basaraba SA, Wright DC. Exercise and adrenaline increase pgc-1 α mRNA expression in rat adipose tissue. *J. Physiol.-London.* 2009;587:1607–1617. <https://doi.org/10.1113/jphysiol.2008.165464>.
41. Hallal PC, Andersen LB, Bull FC, Guthold R, Haskell W, Ekelund U. Global physical activity levels: surveillance progress, pitfalls, and prospects. *Lancet.* 2012;380: 247–257. [https://doi.org/10.1016/S0140-6736\(12\)60646-1](https://doi.org/10.1016/S0140-6736(12)60646-1).
42. Mellett LH, Bousquet G. Heart-healthy exercise. *Circulation.* 2013;127:e571–e572. <https://doi.org/10.1161/CIRCULATIONAHA.112.000880>.
43. Chakravarthy MV, Booth FW. Eating, exercise, and “thrifty” genotypes: connecting the dots toward an evolutionary understanding of modern chronic diseases. *J Appl Physiol.* 2004;96:3–10. <https://doi.org/10.1152/japplphysiol.00757.2003>.
44. Wallberg-Henriksson H, Zierath JR. Exercise remodels subcutaneous fat tissue and improves metabolism. *Nat Rev Endocrinol.* 2015;11:198–200. <https://doi.org/10.1038/nrendo.2015.24>.
45. Ross R, Hudson R, Stotz PJ, Lam M. Effects of exercise amount and intensity on abdominal obesity and glucose tolerance in obese adults. *Ann Intern Med.* 2015; 162:325–334. <https://doi.org/10.7326/M14-1189>.
46. Ahtaiainen JP, Walker S, Silvenoinen M, Kyröläinen H, Nindl BC, et al. Exercise type and volume alter signaling pathways regulating skeletal muscle glucose uptake and protein synthesis. *Eur J Appl Physiol.* 2015;115:1835–1845. <https://doi.org/10.1007/s00421-015-3155-3>.
47. Balducci S, Zanuso S, Nicolucci A, Fernando F, Cavallo S, et al. Anti-inflammatory effect of exercise training in subjects with type 2 diabetes and the metabolic syndrome is dependent on exercise modalities and independent of weight loss. *Nutr Metabol Cardiovasc Dis.* 2010;20:608–617. <https://doi.org/10.1016/j.numecd.2009.04.015>.
48. Idorn M, Thor Straten P. Exercise and cancer: from “healthy” to “therapeutic”. *Cancer Immunol Immunother.* 2017;66:667–671. <https://doi.org/10.1007/s00262-017-1985-z>.
49. Streckmann F, Balke M, Cavaletti G, Toscanelli A, Bloch W, et al. Exercise and neuropathy: systematic review with meta-analysis. *Sports Med.* 2022;52: 1043–1065. <https://doi.org/10.1007/s40279-021-01596-6>.
50. Firth J, Cotter J, Elliott R, French P, Yung AR. A systematic review and meta-analysis of exercise interventions in schizophrenia patients. *Psychol Med.* 2015;45: 1343–1361. <https://doi.org/10.1017/S003329714003110>.
51. Alkadhi KA. Exercise as a positive modulator of brain function. *Mol Neurobiol.* 2018;55:3112–3130. <https://doi.org/10.1007/s12035-017-0516-4>.
52. Boa BCS, Yudkin JS, van Hinsbergh VWM, Bouskela E, Eringa EC. Exercise effects on perivascular adipose tissue: endocrine and paracrine determinants of vascular function. *Br J Pharmacol.* 2017;174:3466–3481. <https://doi.org/10.1111/bph.13732>.
53. Benatti FB, Pedersen BK. Exercise as an anti-inflammatory therapy for rheumatic diseases—myokine regulation. *Nat Rev Rheumatol.* 2015;11:86–97. <https://doi.org/10.1038/nrrheum.2014.193>.
54. Ruiz-Casado A, Martín-Ruiz A, Pérez LM, Provencio M, Fiuza-Luces C, Lucia A. Exercise and the hallmarks of cancer. *Trends Cancer.* 2017;3:423–441. <https://doi.org/10.1016/j.trecan.2017.04.007>.
55. Vina J, Sanchis-Gomar F, Martinez-Bello V, Gomez-Cabrera MC. Exercise acts as a drug: the pharmacological benefits of exercise. *Br J Pharmacol.* 2012;167:1–12. <https://doi.org/10.1111/j.1476-5381.2012.01970.x>.
56. Fletcher GF, Balady GJ, Amsterdam EA, Chaitman B, Eckel R, et al. Exercise standards for testing and training. *Circulation.* 2001;104:1694–1740. <https://doi.org/10.1161/hc3901.095960>.
57. Piercy KL, Troiano RP, Ballard RM, Carlson SA, Fulton JE, et al. The physical activity guidelines for americans. *JAMA, J Am Med Assoc.* 2018;320:2020. <https://doi.org/10.1001/jama.2018.14854>.
58. Sarzynski MA, Ghosh S, Bouchard C. Genomic and transcriptomic predictors of response levels to endurance exercise training. *J Physiol.* 2017;595:2931–2939. <https://doi.org/10.1113/JP272559>.
59. Hoffman NJ. Omics and exercise: global approaches for mapping exercise biological networks. *Cold Spring Harb. Perspect. Med.* 2017;7, a29884. <https://doi.org/10.1101/cshperspect.a029884>.
60. Al-Khelaifi F, Diboun I, Donati F, Botré F, Abraham D, et al. Metabolic gwas of elite athletes reveals novel genetically-influenced metabolites associated with athletic performance. *Sci Rep.* 2019;9, 19889. <https://doi.org/10.1038/s41598-019-56496-7>.
61. Rangwala SM, Wang X, Calvo JA, Lindsley L, Zhang Y, et al. Estrogen-related receptor γ is a key regulator of muscle mitochondrial activity and oxidative capacity. *J Biol Chem.* 2010;285:22619–22629. <https://doi.org/10.1074/jbc.M110.125401>.
62. Nogueira L, Ramirez-Sanchez I, Perkins GA, Murphy A, Taub PR, et al. (-)-epicatechin enhances fatigue resistance and oxidative capacity in mouse muscle. *J Physiol.* 2011;589:4615–4631. <https://doi.org/10.1113/jphysiol.2011.209924>.
63. Jang YJ, Byun S. Molecular targets of exercise mimetics and their natural activators. *Bmb Rep.* 2021;54:581–591. <https://doi.org/10.5483/BMBRep.2021.54.12.151>.
64. Hawley JA, Joyner MJ, Green DJ. Mimicking exercise: what matters most and where to next? *J Physiol-London.* 2021;599:791–802. <https://doi.org/10.1113/JP278761>.

65. Tapia-Arancibia L, Aliaga E, Silhol M, Arancibia S. New insights into brain bdnf function in normal aging and alzheimer disease. *Brain Res Rev.* 2008;59:201–220. <https://doi.org/10.1016/j.brainresrev.2008.07.007>.
66. Phillips C, Baktir MA, Srivatsan M, Salehi A. Neuroprotective effects of physical activity on the brain: a closer look at trophic factor signaling. *Front Cell Neurosci.* 2014;8:170. <https://doi.org/10.3389/fncel.2014.00170>.
67. Wang R, Holsinger RMD. Exercise-induced brain-derived neurotrophic factor expression: therapeutic implications for alzheimer's dementia. *Ageing Res Rev.* 2018;48:109–121. <https://doi.org/10.1016/j.arr.2018.10.002>.
68. Bishop-Bailey D. Mechanisms governing the health and performance benefits of exercise. *Br J Pharmacol.* 2013;170:1153–1166. <https://doi.org/10.1111/bph.12399>.
69. Yu T, Chang Y, Gao X, Li H, Zhao P. Dynamic expression and the role of bdnf in exercise-induced skeletal muscle regeneration. *Int J Sports Med.* 2017;38:959–966. <https://doi.org/10.1055/s-0043-118343>.
70. So B, Kim H, Kim J, Song W. Exercise-induced myokines in health and metabolic diseases. *Integr. Med. Res.* 2014;3:172–179. <https://doi.org/10.1016/j.imr.2014.09.007>.
71. Motamed S, Karimi I, Jafari F. The interrelationship of metabolic syndrome and neurodegenerative diseases with focus on brain-derived neurotrophic factor (bdnf): kill two birds with one stone. *Metab Brain Dis.* 2017;32:651–665. <https://doi.org/10.1007/s11011-017-9997-0>.
72. Shi K, Liu X, Hou L, Qiao D, Lin X. Effects of exercise on mglur-mediated glutamatergic transmission in the striatum of hemiparkinsonian rats. *Neurosci Lett.* 2019;705:143–150. <https://doi.org/10.1016/j.neulet.2019.04.052>.
73. Mattson MP. Glutamate and neurotrophic factors in neuronal plasticity and disease. *Ann N Y Acad Sci.* 2008;1144:97–112. <https://doi.org/10.1196/annals.1418.005>.
74. Klempin F, Beis D, Mosienko V, Kempermann G, Bader M, Alenina N. Serotonin is required for exercise-induced adult hippocampal neurogenesis. *J Neurosci.* 2013;33:8270–8275. <https://doi.org/10.1523/JNEUROSCI.5855-12.2013>.
75. Alenina N, Klempin F. The role of serotonin in adult hippocampal neurogenesis. *Behav Brain Res.* 2015;277:49–57. <https://doi.org/10.1016/j.bbr.2014.07.038>.
76. Chen J, Zhao X, Yue G, Wang Z. Influence of acute and chronic treadmill exercise on rat plasma lactate and brain npy, l-enk, dyn a1–13. *Cell Mol Neurobiol.* 2007;27:1–10. <https://doi.org/10.1007/s10571-006-9110-4>.
77. Reichmann F, Wegerer V, Jain P, Mayerhofer R, Hassan AM, et al. Environmental enrichment induces behavioural disturbances in neuropeptide y knockout mice. *Sci Rep.* 2016;6, 28182. <https://doi.org/10.1038/srep28182>.
78. Cerdá B, Pérez M, Pérez-Santiago JD, Tornero-Aguilera JF, González-Soltero R, Larrosa M. Gut microbiota modification: another piece in the puzzle of the benefits of physical exercise in health? *Front Physiol.* 2016;7:51. <https://doi.org/10.3389/fphys.2016.00051>.
79. Gubert C, Kong G, Renoir T, Hannan AJ. Exercise, diet and stress as modulators of gut microbiota: implications for neurodegenerative diseases. *Neurobiol Dis.* 2020;134, 104621. <https://doi.org/10.1016/j.nbd.2019.104621>.
80. Yu J, Youngson NA, Laybutt DR, Morris MJ, Leigh SJ. Complementary yet divergent effects of exercise and an exercise mimetic on microbiome in high-fat diet-induced obesity. *Physiol Genom.* 2023. <https://doi.org/10.1152/physiolgenomics.00066.2023>.
81. Cryan JF, O'Riordan KJ, Cowan CSM, Sandhu KV, Bastiaansen TFS, et al. The microbiota-gut-brain axis. *Physiol Rev.* 2019;99:1877–2013. <https://doi.org/10.1152/physrev.00018.2018>.
82. Guerrieri D, Moon HY, van Praag H. Exercise in a pill: the latest on exercise-mimetics. *Brain Plast.* 2017;2:153–169. <https://doi.org/10.3233/BPL-160043>.
83. Lauritzen HPMM, Brandauer J, Schjerling P, Koh H, Treebak JT, et al. Contraction and aicar stimulate il-6 vesicle depletion from skeletal muscle fibers in vivo. *Diabetes.* 2013;62:3081–3092. <https://doi.org/10.2337/db12-1261>.
84. Moon HY, Becke A, Berron D, Becker B, Sah N, et al. Running-induced systemic cathepsin b secretion is associated with memory function. *Cell Metabol.* 2016;24:332–340. <https://doi.org/10.1016/j.cmet.2016.05.025>.
85. Kobilo T, Yuan C, van Praag H. Endurance factors improve hippocampal neurogenesis and spatial memory in mice. *Learn Mem.* 2011;18:103–107. <https://doi.org/10.1101/lm.2001611>.
86. Ditacchio KA, Heinemann SF, Dziewczapolski G. Metformin treatment alters memory function in a mouse model of alzheimer's disease. *J Alzheim Dis.* 2015;44:43–48. <https://doi.org/10.3233/JAD-141332>.
87. Pianta S, Lee JY, Tuazon JP, Castelli V, Mantohac LM, et al. A short bout of exercise prior to stroke improves functional outcomes by enhancing angiogenesis. *Neuromol. Med.* 2019;21:517–528. <https://doi.org/10.1007/s12017-019-08533-x>.
88. Kiuchi T, Lee H, Mikami T. Regular exercise cures depression-like behavior via vegf-flk-1 signaling in chronically stressed mice. *Neuroscience.* 2012;207:208–217. <https://doi.org/10.1016/j.neuroscience.2012.01.023>.
89. Fabel K, Fabel K, Tam B, Kauffer D, Baiker A, et al. Vegf is necessary for exercise-induced adult hippocampal neurogenesis. *Eur J Neurosci.* 2003;18:2803–2812. <https://doi.org/10.1111/j.1460-9568.2003.03041.x>.
90. Ji RC. Recent advances and new insights into muscular lymphangiogenesis in health and disease. *Life Sci.* 2018;211:261–269. <https://doi.org/10.1016/j.lfs.2018.09.043>.
91. Bei Y, Huang Z, Feng X, Li L, Wei M, et al. Lymphangiogenesis contributes to exercise-induced physiological cardiac growth. *J Sport Health Sci.* 2022;11:466–478. <https://doi.org/10.1016/j.jshs.2022.02.005>.
92. Bei Y, Liu J, Xiao J. Lymphatic regulation in tissue repair and regeneration: recent advances and future perspective. *Curr Stem Cell Res Ther.* 2023;18:730–732. <https://doi.org/10.2174/1574888X17666220607122742>.
93. Veziroglu EM, Mias GI. Characterizing extracellular vesicles and their diverse rna contents. *Front Genet.* 2020;11:700. <https://doi.org/10.3389/fgene.2020.00700>.
94. Frühbeis C, Helmig S, Tug S, Simon P, Krämer-Albers E. Physical exercise induces rapid release of small extracellular vesicles into the circulation. *J Extracell Vesicles.* 2015;4, 28239. <https://doi.org/10.3402/jev.v4.28239>.
95. Whitham M, Parker BL, Friedrichsen M, Hingst JR, Hjorth M, et al. Extracellular vesicles provide a means for tissue crosstalk during exercise. *Cell Metabol.* 2018;27:237–251. <https://doi.org/10.1016/j.cmet.2017.12.001>.
96. Gomes CPC, de Gonzalo-Calvo D, Toro R, Fernandes T, Theisen D, et al. Non-coding rnas and exercise: pathophysiological role and clinical application in the cardiovascular system. *Clin Sci.* 2018;132:925–942. <https://doi.org/10.1042/CS20171463>.
97. Bartel DP. MicroRNAs: target recognition and regulatory functions. *Cell.* 2009;136:215–233. <https://doi.org/10.1016/j.cell.2009.01.002>.
98. Ling H, Fabbri M, Calin GA. MicroRNAs and other non-coding rnas as targets for anticancer drug development. *Nat Rev Drug Discov.* 2013;12:847–865. <https://doi.org/10.1038/nrd4140>.
99. Bonilauri B, Dallagiovanna B. Long non-coding rnas are differentially expressed after different exercise training programs. *Front Physiol.* 2020;11, 567614. <https://doi.org/10.3389/fphys.2020.567614>.
100. Statello L, Guo CJ, Chen LL, Huarte M. Gene regulation by long non-coding rnas and its biological functions. *Nat Rev Mol Cell Biol.* 2021;22:96–118. <https://doi.org/10.1038/s41580-020-00315-9>.
101. Urbano A, Smith J, Weeks RJ, Chatterjee A. Gene-specific targeting of dna methylation in the mammalian genome. *Cancers.* 2019;11. <https://doi.org/10.3390/cancers11101515>.
102. Moore LD, Le T, Fan G. Dna methylation and its basic function. *Neuropsychopharmacology.* 2013;38:23–38. <https://doi.org/10.1038/npp.2012.112>.
103. Kulis M, Esteller M. Dna methylation and cancer. *Adv Genet.* 2010;70:27–56. <https://doi.org/10.1016/B978-0-12-380866-0.60002-2>.
104. Park S, Kim J. A short guide to histone deacetylases including recent progress on class ii enzymes. *Exp Mol Med.* 2020;52:204–212. <https://doi.org/10.1038/s12276-020-0382-4>.
105. Fernandes J, Arida RM, Gomez-Pinilla F. Physical exercise as an epigenetic modulator of brain plasticity and cognition. *Neurosci Biobehav Rev.* 2017;80:443–456. <https://doi.org/10.1016/j.neubiorev.2017.06.012>.
106. McGowan PO, Roth TL. Epigenetic pathways through which experiences become linked with biology. *Dev Psychopathol.* 2015;27:637–648. <https://doi.org/10.1017/S0954579415000206>.
107. Widmann M, Nieß AM, Munz B. Physical exercise and epigenetic modifications in skeletal muscle. *Sports Med.* 2019;49:509–523. <https://doi.org/10.1007/s40279-019-01070-4>.
108. Guerrieri JL, Sotayo A, Ponichtera HE, Castrillon JA, Pourzia AL, et al. Class iia hdac inhibition reduces breast tumours and metastases through anti-tumour macrophages. *Nature.* 2017;543:428–432. <https://doi.org/10.1038/nature21409>.
109. Bagchi RA, Weeks KL. Histone deacetylases in cardiovascular and metabolic diseases. *J Mol Cell Cardiol.* 2019;130:151–159. <https://doi.org/10.1016/j.yjmcc.2019.04.003>.
110. Falkenberg KJ, Johnstone RW. Histone deacetylases and their inhibitors in cancer, neurological diseases and immune disorders. *Nat Rev Drug Discov.* 2014;13:673–691. <https://doi.org/10.1038/nrd4360>.
111. Kraemer WJ, Ratamess NA, Nindl BC. Recovery responses of testosterone, growth hormone, and igf-1 after resistance exercise. *J Appl Physiol.* 2017;122:549–558. <https://doi.org/10.1152/jappphysiol.00599.2016>.
112. Bouchard C, Rankinen T, Timmons JA. Genomics and genetics in the biology of adaptation to exercise. *Compr Physiol.* 2011;1:1603. <https://doi.org/10.1002/cphy.c100059>.
113. Lee C, Zeng J, Drew BG, Sallam T, Martin-Montalvo A, et al. The mitochondrial-derived peptide mots-c promotes metabolic homeostasis and reduces obesity and insulin resistance. *Cell Metabol.* 2015;21:443–454. <https://doi.org/10.1016/j.cmet.2015.02.009>.
114. Sprecher DL, Massien C, Pearce G, Billin AN, Perlestein I, et al. Triglyceride:high-density lipoprotein cholesterol effects in healthy subjects administered a peroxisome proliferator activated receptor δ agonist. *Arterioscler Thromb Vasc Biol.* 2007;27:359–365. <https://doi.org/10.1161/01.ATV.0000252790.70572.0c>.
115. Ismael A, Holmes M, Papoutsis E, Pantou L, Koutakis P. Resistance training, antioxidant status, and antioxidant supplementation. *Int J Sport Nutr Exerc Metabol.* 2019;29:539–547. <https://doi.org/10.1123/ijnsnem.2018.0339>.
116. Han J, Yang K, An J, Jiang N, Fu S, Tang X. The role of nrf2 in bone metabolism – friend or foe? *Front Endocrinol.* 2022;13, 813057. <https://doi.org/10.3389/fendo.2022.813057>.
117. Staurengo-Ferrari L, Badaro-Garcia S, Hohmann MSN, Manchope MF, Zaninelli TH, et al. Contribution of nrf2 modulation to the mechanism of action of analgesic and anti-inflammatory drugs in pre-clinical and clinical stages. *Front Pharmacol.* 2019;9:1536. <https://doi.org/10.3389/fphar.2018.01536>.
118. El Assar M, Angulo J, Rodríguez-Mañas L. Frailty as a phenotypic manifestation of underlying oxidative stress. *Free Radic Biol Med.* 2020;149:72–77. <https://doi.org/10.1016/j.freeradbiomed.2019.08.011>.
119. Miller KN, Clark JP, Anderson RM. Mitochondrial regulator pgc-1 α —modulating the modulator. *Current Opinion in Endocrine and Metabolic Research.* 2019;5:37–44. <https://doi.org/10.1016/j.coemr.2019.02.002>.
120. Belli R, Bonato A, De Angelis L, Mirabilli S, Ricciardi MR, et al. Corrigendum: metabolic reprogramming promotes myogenesis during aging. *Front Physiol.* 2021;12, 674698. <https://doi.org/10.3389/fphys.2021.674698>.

121. Zerlotin R, Oranger A, Pignataro P, Dicarlo M, Maselli F, et al. Irisin and secondary osteoporosis in humans. *Int J Mol Sci.* 2022;23:690. <https://doi.org/10.3390/ijms23020690>.
122. Estell EG, Le PT, Vegting Y, Kim H, Wrann C, et al. Irisin directly stimulates osteoclastogenesis and bone resorption in vitro and in vivo. *Elife.* 2020;9. <https://doi.org/10.7554/eLife.58172>.
123. Contreras AV, Torres N, Tovar AR. Ppar- α as a key nutritional and environmental sensor for metabolic adaptation. *Adv Nutr.* 2013;4:439–452. <https://doi.org/10.3945/an.113.003798>.
124. Dreyer C, Keller H, Mahfoudi A, Laudet V, Krey G, Wahli W. Positive regulation of the peroxisomal beta-oxidation pathway by fatty acids through activation of peroxisome proliferator-activated receptors (ppar). *Biol Cell.* 1993;77:67–76. [https://doi.org/10.1016/s0248-4900\(05\)80176-5](https://doi.org/10.1016/s0248-4900(05)80176-5).
125. Kersten S, Seydoux J, Peters JM, Gonzalez FJ, Desvergne B, Wahli W. Peroxisome proliferator-activated receptor α mediates the adaptive response to fasting. *J Clin Invest.* 1999;103:1489–1498. <https://doi.org/10.1172/JCI6223>.
126. Inagaki T, Dutchak P, Zhao G, Ding X, Gautron L, et al. Endocrine regulation of the fasting response by ppar α -mediated induction of fibroblast growth factor 21. *Cell Metabol.* 2007;5:415–425. <https://doi.org/10.1016/j.cmet.2007.05.003>.
127. Lundäsen T, Hunt MC, Nilsson L, Sanyal S, Angelin B, et al. Ppar α is a key regulator of hepatic fgf21. *Biochem Biophys Res Commun.* 2007;360:437–440. <https://doi.org/10.1016/j.bbrc.2007.06.068>.
128. Cuevas-Ramos D, Almeda-Valdés P, Meza-Arana CE, Brito-Córdova G, Gómez-Pérez FJ, et al. Exercise increases serum fibroblast growth factor 21 (fgf21) levels. *PLoS One.* 2012;7, e38022. <https://doi.org/10.1371/journal.pone.0038022>.
129. Mashili FL, Austin RL, Deshmukh AS, Fritz T, Caidahl K, et al. Direct effects of fgf21 on glucose uptake in human skeletal muscle: implications for type 2 diabetes and obesity. *Diabetes Metabol Res Rev.* 2011;27:286–297. <https://doi.org/10.1002/dmrr.1177>.
130. Potthoff MJ, Inagaki T, Satapati S, Ding X, He T, et al. Fgf21 induces pgc-1 α and regulates carbohydrate and fatty acid metabolism during the adaptive starvation response. *Proc Natl Acad Sci USA.* 2009;106:10853–10858. <https://doi.org/10.1073/pnas.0904187106>.
131. Zhang Y, Xie Y, Berglund ED, Coate KC, He TT, et al. The starvation hormone, fibroblast growth factor-21, extends lifespan in mice. *Elife.* 2012;1:e65. <https://doi.org/10.7554/eLife.00065>.
132. Berger JP, Akiyama TE, Meinke PT. Ppars: therapeutic targets for metabolic disease. *Trends Pharmacol Sci.* 2005;26:244–251. <https://doi.org/10.1016/j.tips.2005.03.003>.
133. Butcher LR, Thomas A, Backx K, Roberts A, Webb R, Morris K. Low-intensity exercise exerts beneficial effects on plasma lipids via ppar. *Med Sci Sports Exerc.* 2008;40:1263–1270. <https://doi.org/10.1249/MSS.0b013e31816c091d>.
134. Spiegelman BM. Ppar-gamma: adipogenic regulator and thiazolidinedione receptor. *Diabetes.* 1998;47:507–514. <https://doi.org/10.2337/diabetes.47.4.507>.
135. Lehrke M, Lazar MA. The many faces of ppar. *Cell.* 2005;123:993–999. <https://doi.org/10.1016/j.cell.2005.11.026>.
136. Choi JH, Banks AS, Estall JL, Kajimura S, Boström P, et al. Anti-diabetic drugs inhibit obesity-linked phosphorylation of ppar γ by cdk5. *Nature.* 2010;466:451–456. <https://doi.org/10.1038/nature09291>.
137. Choi JH, Choi S, Kim ES, Jedrychowski MP, Yang YR, et al. Thrap3 docks on phosphoserine 273 of ppar and controls diabetic gene programming. *Genes Dev.* 2014;28:2361–2369. <https://doi.org/10.1101/gad.249367.114>.
138. Choi S, Kim ES, Koh M, Lee S, Lim D, et al. A novel non-agonist peroxisome proliferator-activated receptor γ (ppar) ligand uhc1 blocks ppar phosphorylation by cyclin-dependent kinase 5 (cdk5) and improves insulin sensitivity. *J Biol Chem.* 2014;289:26618–26629. <https://doi.org/10.1074/jbc.M114.566794>.
139. Zebrowska A, Gasior Z, Langfort J. Serum igf-i and hormonal responses to incremental exercise in athletes with and without left ventricular hypertrophy. *J Sports Sci Med.* 2009;8:67–76.
140. Neri Serneri GG, Boddi M, Modesti PA, Cecioni I, Coppo M, et al. Increased cardiac sympathetic activity and insulin-like growth factor-i formation are associated with physiological hypertrophy in athletes. *Circ Res.* 2001;89:977–982. <https://doi.org/10.1161/hh2301.100982>.
141. Vanhaesebroeck B, Guillermet-Guibert J, Graupera M, Bilanges B. The emerging mechanisms of isoform-specific pi3k signalling. *Nat Rev Mol Cell Biol.* 2010;11:329–341. <https://doi.org/10.1038/nrm2882>.
142. Bass-Stringer S, Tai CMK, McMullen JR. Igf1-pi3k-induced physiological cardiac hypertrophy: implications for new heart failure therapies, biomarkers, and predicting cardiotoxicity. *J Sport Health Sci.* 2021;10:637–647. <https://doi.org/10.1016/j.jshs.2020.11.009>.
143. Ghigo A, Li M. Phosphoinositide 3-kinase: friend and foe in cardiovascular disease. *Front Pharmacol.* 2015;6:169. <https://doi.org/10.3389/fphar.2015.00169>.
144. McMullen JR, Shioi T, Zhang L, Tarnavski O, Sherwood MC, et al. Phosphoinositide 3-kinase(p110 α) plays a critical role for the induction of physiological, but not pathological, cardiac hypertrophy. *Proc Natl Acad Sci USA.* 2003;100:12355–12360. <https://doi.org/10.1073/pnas.1934654100>.
145. Shioi T, McMullen JR, Kang PM, Douglas PS, Obata T, et al. Akt/protein kinase b promotes organ growth in transgenic mice. *Mol Cell Biol.* 2002;22:2799–2809. <https://doi.org/10.1128/MCB.22.8.2799-2809.2002>.
146. Debusch B, Treskov I, Lupu TS, Weinheimer C, Kovacs A, et al. Akt1 is required for physiological cardiac growth. *Circulation.* 2006;113:2097–2104. <https://doi.org/10.1161/CIRCULATIONAHA.105.595231>.
147. Hou Z, Qin X, Hu Y, Zhang X, Li G, et al. Longterm exercise-derived exosomal mir-342-5p. *Circ Res.* 2019;124:1386–1400. <https://doi.org/10.1161/CIRCRESAHA.118.314635>.
148. Li H, Trager LE, Liu X, Hastings MH, Xiao C, et al. Lncxact1 and dchs2 regulate physiological and pathological cardiac growth. *Circulation.* 2022;145:1218–1233. <https://doi.org/10.1161/CIRCULATIONAHA.121.056850>.
149. Liu X, Xiao J, Zhu H, Wei X, Platt C, et al. Mir-222 is necessary for exercise-induced cardiac growth and protects against pathological cardiac remodeling. *Cell Metabol.* 2015;21:584–595. <https://doi.org/10.1016/j.cmet.2015.02.014>.
150. Gao R, Wang L, Bei Y, Wu X, Wang J, et al. Long noncoding rna cardiac physiological hypertrophy-associated regulator induces cardiac physiological hypertrophy and promotes functional recovery after myocardial ischemia-reperfusion injury. *Circulation.* 2021;144:303–317. <https://doi.org/10.1161/CIRCULATIONAHA.120.050446>.
151. Zhao H, Chen X, Hu G, Li C, Guo L, et al. Small extracellular vesicles from brown adipose tissue mediate exercise cardioprotection. *Circ Res.* 2022;130:1490–1506. <https://doi.org/10.1161/CIRCRESAHA.121.320458>.
152. Anacker C, Hen R. Adult hippocampal neurogenesis and cognitive flexibility — linking memory and mood. *Nat Rev Neurosci.* 2017;18:335–346. <https://doi.org/10.1038/nrn.2017.45>.
153. Garcia PC, Real CC, Britto LR. The impact of short and long-term exercise on the expression of arc and ampa receptors during evolution of the 6-hydroxy-dopamine animal model of Parkinson's disease. *J Mol Neurosci.* 2017;61:542–552. <https://doi.org/10.1007/s12031-017-0896-y>.
154. Graff RM, Kunz HE, Agha NH, Baker FL, Laughlin M, et al. B2-adrenergic receptor signaling mediates the preferential mobilization of differentiated subsets of cd8+ t-cells, nk-cells and non-classical monocytes in response to acute exercise in humans. *Brain Behav Immun.* 2018;74:143–153. <https://doi.org/10.1016/j.bbi.2018.08.017>.
155. Parrini M, Ghezzi D, Deidda G, Medrihan L, Castorflorio E, et al. Aerobic exercise and a bdnf-mimetic therapy rescue learning and memory in a mouse model of down syndrome. *Sci Rep.* 2017;7, 16825. <https://doi.org/10.1038/s41598-017-17201-8>.
156. El-Sayes J, Harasym D, Turco CV, Locke MB, Nelson AJ. Exercise-induced neuroplasticity: a mechanistic model and prospects for promoting plasticity. *Neuroscientist.* 2019;25:65–85. <https://doi.org/10.1177/1073858418771538>.
157. Nauer RK, Dunne MF, Stern CE, Storer TW, Schon K. Improving fitness increases dentate gyrus/ca3 volume in the hippocampal head and enhances memory in young adults. *Hippocampus.* 2020;30:488–504. <https://doi.org/10.1002/hipo.23166>.
158. Sarauli D, Costanzi M, Mastroianni V, Farioli-Vecchioli S. The long run: neuroprotective effects of physical exercise on adult neurogenesis from youth to old age. *Curr Neuropharmacol.* 2017;15:519–533. <https://doi.org/10.2174/1570159X14666160412150223>.
159. van Praag H, Kempermann G, Gage FH. Running increases cell proliferation and neurogenesis in the adult mouse dentate gyrus. *Nat Neurosci.* 1999;2:266–270. <https://doi.org/10.1038/6368>.
160. Belzung C, Wigmore P. *Springer Berlin/Heidelberg.* 2013:189–210. Germany.
161. Codd LN, Blackmore DG, Vukovic J, Bartlett PF. Exercise reverses learning deficits induced by hippocampal injury by promoting neurogenesis. *Sci Rep.* 2020;10, 19269. <https://doi.org/10.1038/s41598-020-76176-1>.
162. Mullard A. Addressing cancer's grand challenges. *Nat Rev Drug Discov.* 2020;19:825–826. <https://doi.org/10.1038/d41573-020-00202-0>.
163. Hawley JA, Hargreaves M, Joyner MJ, Zierath JR. Integrative biology of exercise. *Cell.* 2014;159:738–749. <https://doi.org/10.1016/j.cell.2014.10.029>.
164. Catoire M, Kersten S. The search for exercise factors in humans. *Faseb J.* 2015;29:1615–1628. <https://doi.org/10.1096/fj.14.263699>.
165. Goulet A, Cambillau C. Structure and topology prediction of phage adhesion devices using alphafold2: the case of two oenococcus oeni phages. *Microorganisms.* 2021;9:2151. <https://doi.org/10.3390/microorganisms9102151>.