

Account

Dashboard

Courses

Groups

Calendar

Inbox

History

Course Material Website

Commons

Help

Resources

Lecture 19 Activity Results for Test Student

Score for this attempt: 1 out of 1
Submitted Mar 12 at 8:27am
This attempt took 1 minute.

Correct!

Question 1

1 / 1 pts

Consider the second order differential operator L :

$$L(u) \equiv a(x,y)u_{xx} + 2b(x,y)u_{xy} + c(x,y)u_{yy} + d(x,y)u_x + e(x,y)u_y + f(x,y)u$$

Its discriminant is $\delta(x,y) = (b(x,y))^2 - a(x,y)c(x,y)$.

Suppose $\delta(x,y) > 0$ at (x,y) . Which statement below is true?

☒ Operator L is hyperbolic at (x,y) .

☐ Operator L is parabolic at (x,y) .

☐ Operator L is elliptic at (x,y) .

☐ There is not enough information to determine the classification of operator L at (x,y) . The classification is affected by coefficients $d(x,y)$ and $e(x,y)$.

☐ Operator L is degenerate at (x,y) .

Additional Comments:

Correct!

Question 2

0 / 0 pts

Consider the second order differential operator L :

$$L(u) \equiv a(x,y)u_{xx} + 2b(x,y)u_{xy} + c(x,y)u_{yy} + d(x,y)u_x + e(x,y)u_y + f(x,y)u$$

Its discriminant is $\delta(x,y) = (b(x,y))^2 - a(x,y)c(x,y)$.

Suppose $\delta(x,y) = 0$ at (x,y) and $a(x,y)$, $b(x,y)$ and $c(x,y)$ are not all zeros. Which statement below is true?

☐ Operator L is hyperbolic at (x,y) .

☒ Operator L is parabolic at (x,y) .

☐ Operator L is elliptic at (x,y) .

☐ There is not enough information to determine the classification of operator L at (x,y) . The classification is affected by coefficients $d(x,y)$ and $e(x,y)$.

☐ Operator L is degenerate at (x,y) .

Additional Comments:

Correct!

Correct!

Question 3

0 / 0 pts

Which of the following is true? **Select all that apply.**

☐ $\partial_x(a(x,y)\partial_x u) = a(x,y)\partial_x^2 u$

☐ $\partial_x(a(x,y)\partial_x u) = \partial_x^2(a(x,y)u)$

☒ $\partial_x(a(x,y)\partial_x u)$ and $a(x,y)\partial_x^2 u$ have the same principal part.

☒ $\partial_x(a(x,y)\partial_x u)$ and $\partial_x^2(a(x,y)u)$ have the same principal part.

☐ $a(x,y)\partial_x^2 u = \partial_x^2(a(x,y)u)$

Additional Comments:

Correct!

Question 4

0 / 0 pts

Consider the second order differential operator L :

$$L(u) \equiv a(x,y)u_{xx} + 2b(x,y)u_{xy} + c(x,y)u_{yy} + d(x,y)u_x + e(x,y)u_y + f(x,y)u$$

The coefficient matrix is $A(x,y) = \begin{pmatrix} a(x,y) & b(x,y) \\ b(x,y) & c(x,y) \end{pmatrix}$.

Which of the following is true? **Select all that apply.**

☐ The principal part of $L(u)$ is $\begin{pmatrix} \partial_x \\ \partial_y \end{pmatrix} A(x,y) \begin{pmatrix} \partial_x \\ \partial_y \end{pmatrix} u$.

☐ The principal part of $L(u)$ is $\begin{pmatrix} \partial_x \\ \partial_y \end{pmatrix}^T A(x,y) \begin{pmatrix} \partial_x \\ \partial_y \end{pmatrix} u$.

☐ $L(u)$ and $\begin{pmatrix} \partial_x \\ \partial_y \end{pmatrix} A(x,y) \begin{pmatrix} \partial_x \\ \partial_y \end{pmatrix} u$ have the same principal part.

☒ $L(u)$ and $\begin{pmatrix} \partial_x \\ \partial_y \end{pmatrix}^T A(x,y) \begin{pmatrix} \partial_x \\ \partial_y \end{pmatrix} u$ have the same principal part.

☐ $L(u)$ and $A(x,y) \begin{pmatrix} \partial_x \\ \partial_y \end{pmatrix}^T \begin{pmatrix} \partial_x \\ \partial_y \end{pmatrix} u$ have the same principal part.

☐ $L(u)$ and $\begin{pmatrix} \partial_x \\ \partial_y \end{pmatrix}^T \begin{pmatrix} \partial_x \\ \partial_y \end{pmatrix} (A(x,y)u)$ have the same principal part.

Additional Comments:

Correct!

Question 5

0 / 0 pts

Consider the second order differential operator $L(u)$ with coefficient matrix $A(x,y)$. $L(u)$ and $\begin{pmatrix} \partial_x \\ \partial_y \end{pmatrix}^T A(x,y) \begin{pmatrix} \partial_x \\ \partial_y \end{pmatrix} u$ have the same principal part.

Consider the change of variable $\begin{cases} \xi = \xi(x,y) \\ \eta = \eta(x,y) \end{cases}$. We have

$$\begin{pmatrix} \partial_x \\ \partial_y \end{pmatrix} u = \begin{pmatrix} \xi_x & \eta_x \\ \xi_y & \eta_y \end{pmatrix} \begin{pmatrix} \partial_\xi \\ \partial_\eta \end{pmatrix} u = \begin{pmatrix} \partial(\xi,\eta) \\ \partial(x,y) \end{pmatrix}^T \begin{pmatrix} \partial_\xi \\ \partial_\eta \end{pmatrix} u$$

Let $\tilde{L}(u)$ denote operator $L(u)$ in terms of new variables (ξ,η) .

What is the coefficient matrix of $\tilde{L}(u)$?

☐ $\tilde{A}(\xi,\eta) = A(x,y)$

☐ $\tilde{A}(\xi,\eta) = \begin{pmatrix} \partial(\xi,\eta) \\ \partial(x,y) \end{pmatrix}^T A(x,y) \frac{\partial(\xi,\eta)}{\partial(x,y)}$

☒ $\tilde{A}(\xi,\eta) = \frac{\partial(\xi,\eta)}{\partial(x,y)} A(x,y) \begin{pmatrix} \partial(\xi,\eta) \\ \partial(x,y) \end{pmatrix}^T$

☐ $\tilde{A}(\xi,\eta) = A(x,y) \begin{pmatrix} \partial(\xi,\eta) \\ \partial(x,y) \end{pmatrix}^T$

☐ $\tilde{A}(\xi,\eta) = \frac{\partial(\xi,\eta)}{\partial(x,y)} A(x,y)$

☐ $\tilde{A}(\xi,\eta)$ is affected by the first derivative terms in $L(u)$. There is not enough information to determine $\tilde{A}(\xi,\eta)$.

Additional Comments:

Correct!

Question 6

0 / 0 pts

Consider the second order differential operator $L(u)$ with coefficient matrix $A(x,y)$. Consider the change of variable $\begin{cases} \xi = \xi(x,y) \\ \eta = \eta(x,y) \end{cases}$ with non-singular Jacobian $\frac{\partial(\xi,\eta)}{\partial(x,y)}$. Let $\tilde{L}(u)$ denote operator $L(u)$ in terms of new variables (ξ,η) .

Which statement below is true?

☐ $\tilde{L}(u)$ and $L(u)$ have the same coefficient matrix.

☒ $\tilde{L}(u)$ and $L(u)$ have the same classification.

☐ $\tilde{L}(u)$ and $L(u)$ have the same classification only if the non-singular Jacobian $\frac{\partial(\xi,\eta)}{\partial(x,y)}$ is an orthogonal matrix.

☐ $\tilde{L}(u)$ and $L(u)$ have the same classification only if the non-singular Jacobian $\frac{\partial(\xi,\eta)}{\partial(x,y)}$ is a positive definite matrix.

☐ $\tilde{L}(u)$ and $L(u)$ have the same classification only if $\det\left(\frac{\partial(\xi,\eta)}{\partial(x,y)}\right) > 0$.

☐ $\tilde{L}(u)$ and $L(u)$ have the same classification only if the non-singular Jacobian $\frac{\partial(\xi,\eta)}{\partial(x,y)}$ is a diagonal matrix.

Additional Comments:

Correct!

Question 7

0 / 0 pts

Consider the second order differential operator L :

$$L(u) \equiv a(x,y)u_{xx} + 2b(x,y)u_{xy} + c(x,y)u_{yy} + d(x,y)u_x + e(x,y)u_y + f(x,y)u$$

Suppose $\delta(x,y) = -\det\begin{pmatrix} a(x,y) & b(x,y) \\ b(x,y) & c(x,y) \end{pmatrix} > 0$ in a region of (x,y) .

What is the coefficient matrix of the canonical form?

☐ $\tilde{A}(\xi,\eta) = \alpha(\xi,\eta) \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$

☒ $\tilde{A}(\xi,\eta) = \alpha(\xi,\eta) \begin{pmatrix} 0 & \frac{1}{2} \\ \frac{1}{2} & 0 \end{pmatrix}$

☐ $\tilde{A}(\xi,\eta) = \alpha(\xi,\eta) \begin{pmatrix} 0 & \frac{-1}{2} \\ \frac{1}{2} & 0 \end{pmatrix}$

☐ $\tilde{A}(\xi,\eta) = \alpha(\xi,\eta) \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$

☐ $\tilde{A}(\xi,\eta) = \alpha(\xi,\eta) \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$

Additional Comments:

Correct!

Question 8

0 / 0 pts

Consider the second order differential operator L :

$$L(u) \equiv a(x,y)u_{xx} + 2b(x,y)u_{xy} + c(x,y)u_{yy} + d(x,y)u_x + e(x,y)u_y + f(x,y)u$$

Suppose $\delta(x,y) = -\det\begin{pmatrix} a(x,y) & b(x,y) \\ b(x,y) & c(x,y) \end{pmatrix} < 0$ in a region of (x,y) .

What is the coefficient matrix of the canonical form?

☐ $\tilde{A}(\xi,\eta) = \alpha(\xi,\eta) \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$

☐ $\tilde{A}(\xi,\eta) = \alpha(\xi,\eta) \begin{pmatrix} 0 & \frac{1}{2} \\ \frac{1}{2} & 0 \end{pmatrix}$

☐ $\tilde{A}(\xi,\eta) = \alpha(\xi,\eta) \begin{pmatrix} 0 & \frac{-1}{2} \\ \frac{1}{2} & 0 \end{pmatrix}$

☐ $\tilde{A}(\xi,\eta) = \alpha(\xi,\eta) \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$

☒ $\tilde{A}(\xi,\eta) = \alpha(\xi,\eta) \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$

Additional Comments:

Fudge Points: --

You can manually adjust the score by adding positive or negative points to this box.

Final Score: 1 out of 1

Update Scores

Here's the latest quiz results for Test Student. You can modify the points for any question and add more comments, then click "Update Scores" at the bottom of the page.

Quiz Submissions

Attempt 1: 1

Test Student has 1 attempt left

Allow this student an extra attempt