Have you tested for change in hysterantery (veg-flodates) in HF data over time flor w/ blimate? You should's

concordance=TRUE

1. Phenology, the study of seasonal life cycle events, has long been studied, and recently drawn increasing scientific attention. Recent work is beginning to show that it is not only individual phenophases that are important, but also the realtionship between them, even if they are seemingly disperate. Examples: Phenology shifts can change life history ie winter to 7 sumemr annual or others. But this work is really just beggining.

Good but the camples

2. One phenological sequence that has attracted some attention is the flower leaf sequence (FLS). Why do some species flower before leaves while other produce leaves first?

(it's common ...

3. It has been suggest that flowering first, also known as hysteranthy, proteranthy, or precicious flowering, may be an important mediator of reproductive or physiological functioning, and several hypotheses have emerged.

- (a) Here are the hypotheses:
- (b) Wind Pollination effeciency: This is the dominant temperate hypothesis. It has a corrilary in the tropics: insect visability.
- (c) Hydration issues, the primary idea emerging from research in dry
- (d) Several other hypotheses of hysteranthy are actually just hypotheses for early flowering, ie. seed size, cold tolerance, others mentioned in "the paper". These would imply the FLS is a a byproduct of differential selection regimes on leaves and flower.
- (e) or phylogenetic conservatism.
- (f) It is clear that climate change is altering phenology. This could alter FLS. If FLS is truly impotant, which we assume it is, alterations might be bad for plants, and we should know more about this.

4. Despite the rich theory begind it, there is limited, quality, empirical data about FLS.

- (a) Why? Flowering and leafing are often observed seperately, we don't have good phenological records.
- (b) The best data are verbal description from guide books like "flowers before leaves" etc.
- (c) Sadly, these verbal descriptions are incompatible with the quantitative phenology we use. There are three major sources of uncertainty in these data, limiting their use in a meaningful way-1 natural variation 2 Definitional ambiguity with the observer, and 3 ambiguity in researcher interpretation. Each of which we will discuss in detail

SubHEADER - Sources of uncertainty

(A) and . oh, you could re-work the author into a short intro have a subsection on hypotheses before Sources of Variation. I

Somewhere around here (as areader) law

(a) We do not have a great sense about the range of interannual variation, or interpopulation difference in FLS which could be really important. Perhaps species with short lag times in their FLS may switch between year, or life stages, or have different FLS across their range. None of this can be captured with the data sources we have—we don't know if "or" statements like "Flowers before or with leaves" reflect interanual variation, variation between individuals, branches or over a region or overlap?

Why might there be variability? *Think more about this*? fernt cues are more reliable for differnt phenophases, some year cues converges or diverges and the patterns respond.

(c) adaptive reasons?

- (d) We dont know, but understanding this would be huge for actually testing hypotheses. You could ask questions like in years of more extreme FLS seperation, do we see more pollination success, or are they droughtier?
- (e) To get a baseline sense of this, we looked at a few species from harvard forest again. And look there is considerable variation between trees and years.
- (f) It is considerable. [FIGURE 2]
- (g) We also don't know if differences between sources can be attributed to biological differences or definitional ones which we will discuss below.

6. Ambiguity in definition

- (a) What does an author mean when they say "flowers before leaves?". There are different ways to characterize this as we seem from other studies (cite some or talk about BBCH).
- (b) Using harvard forest data we demonstate this matters. If an author means flower bud burst before leaf budburst you get a very different species list than flowers open before leaf expansion. [FIG-URE OR TABLE 1] This could explain differences between sources, but does rule out there are biological differences mentioned above, and this ambiguity limits our ability to understand this trait across space and time and it makes it difficult to validate these descriptions with other phenological observation that speficially specify budburst, leafout etc.

7. Ambiguity in interpretation and modeling choices

(a) For modeling, these descriptions allow for the FLS to be characterized binary or at best, catagorical approximations only, when in fact, it is a continuous trait. Where to draw the line between catagories is up to researcher interpretation, and these choices could affect downstream analysis.

(b) These choices might be influenced by our bias. If I favor the wind pollination hypothesis, I would be biologically justified for choosing the more expansive definition, but if I think it hysteranthy is a physiological constraint, a more conservative physiological definition would be appropriate.

8. Do these sources of error matter to how we understand FLS? We illustrate the implications of all of this uncertaintly through modeling. We show the support for the various hypotheses is sensative to researcher choices using our analysis from MTSV and Silvics.

(a) Look! The phylogenetic signal changes with our choices and between data sources [FIGURE 3 phylogeny figures]. This is also discuss in "the paper's" random forest suppliment models. Not only that, we see that the result reinforce the model choices and data sources, ie the more expaisive definition strengthens wind pollination effect [FIGURE 4].

- (b) Our analysis shows that generally early flowering, pollination snydrome and seed characteristics are decent predictors of hysteranthy, though the strength the of these effects, and the phylogenetic signals, are sensative to choices made during analysis that could be entagled in the three levels of data ambiguity discussed above.
- (c) This makes its hard to conclusively favor a single hypothesis from the ones laid out at the beginning
- (d) But that's okay, because it is likely hysteranthy didn't evolve for a single function but may serve different functions for different life histories, biogeographic origins etc (Lizzie had a better way of saying this and some citations I think).
- (e) What we can say right now is that there is good eveidence that FLS and other phenological sequences are impotant beyound the phases they contain, and study of these patterns and the relationship between them should continue and be improved.
- 9. We have a few suggestions for how this area of study should progress:
 - (a) More observation, so we can treat FLS as continuous rather than discrete variables. Do this over multiple years and locations
 - (b) Mechaism. We have to identify the variability in the cues
 - (c) View this in the context of the phenological cycle as a whole especcially budset.

The sound of the s

(a) More discretion (b) Mecha (c) View cially data what days that days that

see email

underground trees

(geo xylexfree...)

protect from front

at high elevations

of fire elsewhere.