

ABSTRACT

SOIL SEED BANK GERMINATION AND UNDERSTORY DIVERSITY IN OAK SAVANNA RESTORATIONS

Michael L Saxton, M.S.
Department of Geography
Northern Illinois University, 2012
David Goldblum and Lesley S. Rigg, Co-Directors

The cumulative effect of widespread conversion of natural areas to agricultural land use, invasion of non-native species and 20th century fire suppression has increased habitat fragmentation and has led to the overall decline of species diversity and ecosystem functionality of oak savannas across the North American prairie-forest ecotone region. This research was conducted in a 30ha oak savanna restoration project in north central Illinois, USA. We analyzed differences in restoration management history and practices by comparing a five-year-old restoration effort against a degraded unmanaged unit. We use multivariate analyses to determine the influence of variation in understory light levels using hemispherical lens photography and variation in soil N, C, pH, and percent sand/silt/clay on herbaceous species diversity in random sampling plots. Results indicate a significant difference in species richness, cover and stem counts between the managed and unmanaged units. Additionally, we collected soil samples from both units and germinated the soil seed banks using three different treatment methods: watering with aqueous smoke extract, heating samples to 100 °C, and watering samples with distilled water. Significant differences were found in species per tray and germinants per tray between treatments.

PREVIEW

NORTHERN ILLINOIS UNIVERSITY
DE KALB, ILLINOIS

AUGUST 2012

SOIL SEED BANK GERMINATION AND UNDERSTORY
DIVERSITY IN OAK SAVANNA RESTORATIONS

BY

MICHAEL L. SAXTON
©2012 Michael L. Saxton

A THESIS SUBMITTED TO THE GRADUATE SCHOOL
IN PARTIAL FULFILLMENT OF THE REQUIREMENTS
FOR THE DEGREE
MASTER OF SCIENCE

DEPARTMENT OF GEOGRAPHY

Thesis Co-Directors:
David Goldblum
Lesley S. Rigg

UMI Number: 1519738

All rights reserved

INFORMATION TO ALL USERS

The quality of this reproduction is dependent upon the quality of the copy submitted.

In the unlikely event that the author did not send a complete manuscript and there are missing pages, these will be noted. Also, if material had to be removed, a note will indicate the deletion.



UMI 1519738

Published by ProQuest LLC (2012). Copyright in the Dissertation held by the Author.

Microform Edition © ProQuest LLC.

All rights reserved. This work is protected against unauthorized copying under Title 17, United States Code



ProQuest LLC.
789 East Eisenhower Parkway
P.O. Box 1346
Ann Arbor, MI 48106 - 1346

ACKNOWLEDGEMENTS

I am deeply indebted to many people for the support I have been fortunate to receive. My family has always been steadfast in their love and support. Thanks, Mom and Dad. My five years spent working at Nachusa were the most formative years of my life where I made many lasting friendships and learned more than I ever could have imagined. Much love to the Nachusa community. I am also grateful to the Friends of Nachusa Grasslands for supporting my research with a generous grant.

I was privileged to have two stellar advisors that have been supportive and generous with their time and advice. Thanks Drs. Lesley Rigg and David Goldblum for everything. A special thank you to Shannon McCarragher for our partnership on fieldwork and data collection. To my fellow grad students Michaela, Joe, Mike, Kris and many others, thanks for the memories.

TABLE OF CONTENTS

	Page
LIST OF TABLES	vi
LIST OF FIGURES	viii
LIST OF APPENDECIES	x
Chapter	
1. INTRODUCTION	1
Oak Savanna	1
Ecosystem Origin.....	1
Anthropocene.....	3
Fragmentation	5
<i>Lonicera maackii</i>	6
Site Description.....	7
Management Units.....	10
Reference for Restoration	11
Objectives	12
2. METHODS	14
Point Centered Quarter Sampling	14
Germination	14
Sample Collection.....	15

Chapter	Page
Smoky Water	15
Baked Soil	16
Sample Trays and Growth Chamber	16
Vegetation Sampling and Environmental Measurement	17
Vegetation Sampling	17
Floristic Quality Index	18
Soil Lab Analysis	19
Moisture Factor Analysis	19
Soil pH Analysis	20
Particle Size Analysis	20
Carbon and Nitrogen Analysis	22
Hemispherical Photography	23
3. RESULTS	24
Point Centered Quarter Sampling	24
Germination	28
Viability Testing	28
Overview	28
Smoky Water	29
Distilled Water	32
Baked Soil	34
All Treatments	35

Chapter	Page
Units and Treatments Comparison.....	37
Vegetation Sampling.....	39
Environmental Sampling	40
Regression.....	41
Ordination	41
4. DISCUSSION.....	44
Point Centered Quarter Sampling.....	44
Germination	47
Sampling	49
5. APPLICATION FOR RESTORATION.....	53
6. CONCLUSION.....	56
REFERENCES	58
APPENDICES.....	65

LIST OF TABLES

Table	Page
1. <u>P</u> oint <u>C</u> entered <u>Q</u> uarter Sampling Results from Brandt.....	25
2. Point Centered Quarter Sampling Results from Bennett	26
3. Total Germinants, Mean Germinants per Tray, Mean Species per Tray, Total Species and Mean C-value per Management Unit in Smoky Water Trial	29
4. Prevalent Species per Unit in Smoky Water Trial.....	31
5. Total Germinants, Mean Species per Tray, Mean Germinants per Tray and Total Species in Distilled Water Trial	32
6. Prevalent Species per Unit in Distilled Water Trial.....	34
7. Total Germinants, Species per Tray and Total Species in Baked Soil Trial.....	34
8. Prevalent Species from Brandt in Baked Soil Trial	34
9. Total Germinants, Mean Species per Tray, Mean Germinants per Tray and Total Species for All Treatment Trial	35
10. Prevalent Species per Unit in All Treatments Trial	37
11. Per Management Unit Total Germinants, Mean Germinants per Tray, Total Species and Mean Species per Tray Comparing Management Units	37
12. Per Treatment Total Germinants, Mean Germinants per Tray, Total Species and Mean Species per Tray Comparing Treatments	38
13. Most Prevalent Species in Total Sampling by Stem Count	39
14. Most Prevalent Species per Unit by Stem Count in Total Sampling.....	39
15. Per Plot Averages per Management Units from Vegetation Sampling	40

Table

Page

16. Per Plot Average of Environmental Variables per Management Unit.....	40
17. Eigenvalues and Variance Explained in Canonical Correspondence Analysis	42

PREVIEW

LIST OF FIGURES

Figure	Page
1. Nachusa Grasslands Map Indicating County Lines and Location Within Illinois.....	8
2. 1840 Survey Map with Notes that Indicate Prairie-Forest Boundary	9
3. 1939 and 2012 Aerial Photographs of the Bennett and Brandt Management Units	12
4. Management Units with Vegetation Sampling Plot Locations Indicated.....	18
5. Size Structure Graph of Brandt Based on Point Centered Quarter Sampling.....	27
6. Size structure Graph of Bennett based on Point Centered Quarter Sampling.....	27
7. 12-Week Germination Curve for Bennett and Brandt in Smoky Water Trial	30
8. Number of Germinants Added per Week per Management unit in Smoky Water Trial	30
9. 12-Week Germination Curve for Bennett and Brandt in Distilled Water trial.....	33
10. Number of Germinants Added per Week per Management Unit in Distilled Water Trial.....	33
11. 12-Week Germination Curve for Bennett and Brandt All Treatments	36
12. Number of Germinants Added per Week for All Treatments.....	36
13. Mean Germinants per Tray per Management Unit.....	38
14. Mean Species per Tray per Management Unit	38

Figure	Page
15. Canonical Correspondence Analysis of Vegetation Plot Data by Management Unit with Environmental Measures.....	42
16. Canonical Correspondence Analysis of Species Correspondence to Environmental Variables.....	43

PREVIEW

LIST OF APPENDICES

Appendix	Page
A. SPECIES GERMINATED IN SMOKY TRIAL	65
B. SPECIES GERMINATED IN DISTILLED TRIAL	67
C. VEGETATION SAMPLING SPECIES LIST	69

PREVIEW

CHAPTER 1

INTRODUCTION

Oak Savanna

Typified by fluid and shifting boundaries resulting from interactions between climate, topography, and fire, Midwestern savannas formed a unique ecotone between the eastern deciduous forests and the tall grass prairies of the Plains states (Nuzzo 1986; Anderson and Bowles 1999). Savanna ecosystems are characterized as having a generally scattered, open-grown oak-dominated overstory structure with a dense herbaceous layer of forbs and grasses whose distribution pattern is heavily influenced by disturbance events such as wildfires and droughts (Anderson 1998; Karnitz 2006). Savannas are exceedingly species diverse due in part to the highly varied mosaic of soil and light gradients caused by their broken canopy and spatially distinct structure (Leach and Givnish 1999).

Ecosystem Origin

The genesis of what is considered the oak savanna ecotone dates back to a warming period nearly 8,000 – 3,000 years before present during the late Pleistocene period, an era marked by glaciations, inter-glacial periods, and species migration, both flora and fauna,

due to changes in global climate (Anderson 2006). This warming period is frequently referred to as the Hypsithermal period, characterized by a generally warmer and drier climate that was very conducive to frequent and intense fires (Camill et al. 2003). The millennia following the end of the Wisconsin Glaciation saw a constantly changing boundary between woodlands and prairie. This transitional zone is known as an ecotone where the spatial extent of either ecotype is directly controlled by disturbance factors such as fire and grazing pressure and also by climatic factors such as precipitation rates and droughts (Harrington 1991).

Prairie forbs and grasses are evolved to withstand drought conditions more successfully than most tree species. This adaption, coupled with tree's inability to withstand frequent intense fire, allowed prairies to expand their range after the last glaciation. Additionally, Native American populations increased substantially during this time period, which in turn had a dramatic effect on ecosystem function and species compositions. Native Americans were the primary source of ignition in most regions. By managing their surroundings with fire for thousands of years, Native Americans were essentially "keystone species" (Nowacki 2008). The indigenous populations were using fire as a tool to herd game species including elk and bison, to clear land for agricultural purposes, to promote vegetation growth, and to maintain a relatively open space free from thickets and trees (Pyne 1982).

Climate, fire and topography were major driving forces behind the patchy mosaic characteristic of savannas. Fires played a pivotal role in defining vegetation types and tree distribution across the oak savanna ecotone. Fires, driven by the prevailing west-to-east winds, would race across flat prairies and up dry south-facing slopes. The lee sides

of hills, however, were protected from fires and thus maintained trees and thickets (Abrams 1992). Rivers, streams, wetlands, and sandy uplands worked as fire breaks which helped to maintain a patchy mosaic of burned and unburned areas.

The rain shadow created by the Rocky Mountains, in conjunction with changing air current patterns of the Hypsithermal period (Clark et al. 2001), had a direct effect on precipitation rates across the Great Plains. This in turn caused a west-to-east gradient of short grass prairies, mixed grass prairies, tallgrass prairies, and savannas progressing into the Eastern deciduous forest, each with its own suite of flora and fauna. At the heart of the tallgrass prairie region are two of the longest rivers in the world; the Missouri and the Mississippi. This interconnected network of waterways created a matrix of fire breaks, wetlands, floodplains, and riparian areas that contributed immensely to the overall heterogeneity of the tallgrass prairie and oak savanna ecotone region.

Anthropocene

Prior to European settlement in the 1840s, oak savanna covered 11-13 million hectares of the Midwest extending over an eight-state region. By 1985, however, approximately only 0.02% of high-quality oak savanna remained intact due to the cumulative effect of widespread conversion of natural areas to agricultural land use, invasion of non-native species and 20th-century fire suppression (Nuzzo 1986). Today, these anthropocentric factors have resulted in increased habitat fragmentation and have led to the overall decline of species diversity and ecosystem functionality of oak savannas across the region (McCune and Cottam 1985).

Fires, frequently ignited by Native Americans as a tool to clear land for agriculture and drive bison herds, were prevalent in the pre-settlement Midwest and played a vital role in the formation and maintenance of oak savanna ecosystems (Abrams 1992; Whitney 1994; Nowacki and Abrams 2008). Compared to many fire-intolerant species such as elm (*Ulmus sp.*), walnut (*Juglans sp.*), and cherry (*Prunus sp.*), oaks (*Quercus sp.*) thrived in this frequent fire environment due in part to their thick, fire-resistant bark and their ability to re-sprout after being top-killed by fire (Stearns 1991; Abrams 1992). Typically, oak savannas burned on a 4.5-year interval cycle which maintained oak dominance by keeping fire-intolerant tree species populations in check (Wolf 2004). The sparse spatial distribution of oaks created a patchy mosaic of available resources which enabled tallgrass prairie species and more shade-tolerant herbs to thrive in the transitional savanna community (Leach and Givnish 1999).

Anthropogenic processes, namely fire suppression associated with the onset of European settlement of the Midwest in the early part of the 19th century, pushed natural systems beyond ecological thresholds and along trajectories bound for radically different ecosystem composition, form, and structure (Rayburn and Major 2008). What little oak savanna that remained after the initial logging and agricultural conversion was further degraded by cessation of wildfire. Many of these small oak savanna remnants transitioned into closed canopy woodlands with increased proportions of mesophytic, fire-tolerant tree species (Apfelbaum and Haney 1991; Anderson 1998; Anderson et al. 2000). In addition, the greater stand density caused a marked increase in humidity, decrease in solar radiation and wind speeds, and a cooler and moister forest floor which drove down species diversity and favored more shade-tolerant herbs (Nowacki and