



Dear Dr. Pinfield-Wells,

We propose a “Viewpoint” about the drivers of flower-leaf phenological sequences (FLSs), with implications for future plant communities. Evolutionary theory predicts that both flowering and leaf phenology are critical fitness components of woody plants, and a century of empirical research supports this assertion (Munguia-Rosas *et al.*, 2011; Forrest & Miller-Rushing, 2010). In recent decades, this work has been extended to predict that it is not only individual phenophases but also the relationship between them that determines woody plant fitness (Menzel & Fabian, 1999; Ettinger *et al.*, 2018). Many deciduous woody plants flower before leafing, yet sustained research efforts have yet to yield a well-supported explanation for this. These unresolved hypotheses are critically important now as climate change is shifting FLSs—which may exacerbate fitness differences between species and reshape eco-systems of the future. Our “Viewpoint” shows how progress in this area has been stalled by our conceptual framework for FLSs; we detail a new approach built on continuous measures of FLS and intra-specific and within-individual variation to rapidly advance progress.

What hypotheses or questions does this work address?

Research suggests FLSs are under strong selection and critical to fitness. FLS variation may be an adaptation for wind-pollination (Rathcke & Lacey, 1985), reducing water stress (Reich & Borchert, 1984), or early season flowering (Primack, 1987), but these conflicting hypotheses remain unresolved. A novel approach focusing on intra-specific FLS variation and quantitative comparisons is necessary to accurately evaluate these hypotheses.

How does this work advance our current understanding of plant science?

We show: 1) The current framework fails to capture the inter- and intra- specific FLS variation in nature impeding hypothesis testing. 2) Variation provides novel insights about the function of FLSs, revealing consistencies and anomalies in support for the hypotheses. 3) Leveraging intra-specific variation will advance our understanding of FLS.

Why is this work important and timely?

We show that climate change is altering FLSs, but effects vary across species (Fig. 1). Shifts could be beneficial or adverse, and predicting this outcome requires researchers to effectively evaluate the current hypotheses. Our framework is the first that can do this accurately, making it critical to fundamental and applied research.

We expect this manuscript will be titled “Reconciling competing hypotheses regarding flower-leaf sequences in temperate forests for fundamental and global change biology”. It will be co-authored by I. Morales-Castilla, and E.M. Wolkovich. This proposed manuscript is not under consideration anywhere else. Thank you for your consideration.

Sincerely,

Daniel Buonaiuto

Abstract:

Phenology is a major component of an organism's fitness. While individual phenological events affect fitness, growing evidence suggests that the relationship between events may be equally or more important. This may explain why deciduous woody plants exhibit considerable variation in the order of reproductive and vegetative events, or flower-leaf sequences (FLSs). Research suggests that FLSs are adaptive, with several competing hypotheses to explain their function. Reconciling these hypotheses has been impeded by how FLS patterns are described and defined. Here, we advance the existing hypotheses to account for the FLS variation in nature and evaluate them with four case studies. Our inquiry provides three major insights towards a new framework for understanding FLSs. First, we find concurrent support for multiple hypotheses, suggesting progress can come from studies addressing overlapping hypotheses. Second, support for FLS hypotheses is sensitive to how FLSs are defined, with quantitative definitions proving most useful. Finally, we highlight the limits of trait-based hypothesis testing. Researchers should adopt an intra-specific approach and evaluate fitness consequences of FLS variation to test the FLS hypotheses and to predict how climate-related alterations to FLSs will affect plant communities and re-shape eco-systems in the future.

Selected Figures:




Figure 1: **FLS across Europe for three tree species from 1960 to 2015 suggests climate change has generally increased the time between flowering and leafing**, but the direction and rate of change differs across species, which may exacerbate fitness differences within forest communities. To detect the effect of climate change on average FLS, we used models that allow for shifts in FLS after 1980. Lines represent the mean trend in FLS per species, and the highlighted regions indicate historic range of FLS variability (upper and lower 95% credible intervals of the pre-1980 average) from the PEP725 database (Templ *et al.*, 2018).

References

- Ettinger, A., Gee, S. & M. Wolkovich, E. (2018) Phenological sequences: how early season events define those that follow. *American Journal of Botany* **105**.
- Forrest, J. & Miller-Rushing, A.J. (2010) Toward a synthetic understanding of the role of phenology in ecology and evolution. *Philosophical Transactions of the Royal Society B: Biological Sciences* **365**, 3101–3112.
- Menzel, A. & Fabian, P. (1999) Growing season extended in Europe. *Nature* **397**, 659.
- Munguia-Rosas, M.A., Ollerton, J., Parra-Tabla, V. & Arturo De-Nova, J. (2011) Meta-analysis of phenotypic selection on flowering phenology suggests that early flowering plants are favoured. *Ecology Letters* **14**, 511–521.
- Primack, R.B. (1987) Relationships among flowers, fruits, and seeds. *Annual Review of Ecology and Systematics* **18**, 409–430.
- Rathcke, B. & Lacey, E.P. (1985) Phenological patterns of terrestrial plants. *Annual Review of Ecology and Systematics* **16**, 179–214.
- Reich, P. & Borchert, R. (1984) Water-stress and tree phenology in a tropical dry forest in the lowlands of costa-rica. *Journal of Ecology* **72**, 61–74.
- Templ, B., Koch, E., K.Bolmgren, Ungersböck, M., Paul, A., Scheifinger, H. & et al. (2018) Pan european phenological database (pep725): a single point of access for european data. *Int. J. Biome-teorology* .