

# Reconciling historic hypotheses regarding flower-leaf sequences in temperate forests for fundamental and global change biology

Daniel Buonaiuto, Nacho, Lizzie

August 13, 2019

## Abstract

As the study of phenology progresses as a discipline, it is increasingly clear that it is not only individual phenological events that affect organism fitness and ecological functioning, but also the relationship between stages. Deciduous woody plants exhibit considerable variation in the relative order of their reproductive and vegetative events, or their flower-leaf sequence (FLS). A century of research suggests that FLS patterns are adaptive, and several competing hypotheses stand to explain their evolution and function.

Reconciling these hypotheses has been impeded by our conceptual orientation towards them. Classically, FLSs are treated as discrete categories and described at the species level. However, this framework obscures important inter-specific differences and masks substantial intra-specific variation ignored by the hypotheses. Here, we review and modify the existing hypotheses to account for the high levels of both inter- and intra-specific FLS variation seen in nature. We then evaluate these hypotheses with four case studies from temperate forest species.

Our review and case studies provide three major insights towards a new conceptual framework for understanding FLSs. First, we find concurrent support for multiple hypotheses. Future research should accommodate this coincidence by allowing for overlapping hypotheses in large community models and by testing individual hypotheses in smaller sub-groupings that control for variation in other traits. Second, we show that the support for FLS hypotheses is highly sensitive to how FLSs are defined. Researchers should, when possible, move away from the classic categorization scheme and use continuous measures of FLS, such as number of days between flowering and leafing. Finally, researchers should use these intra-specific, continuous data to test for fitness consequences in FLS variation. This will advance our understanding of the fundamental biology of FLSs, and help us to predict how climate-related alterations to FLSs will affect tree communities in the changing future.

## Introduction

Phenology, the timing of seasonal life cycle events, allows organisms to synchronize important life history transitions with optimum environmental conditions (?), and is a critical component of ecosystem structure and function (??). Recent work in woody plant phenology has shown that it is not only individual phenological stages that affect these processes, but also the relationships between them (?).

One phenological relationship that has long received scientific interest (see ?), and recently, increased attention in the literature (??) is the flower-leaf phenological sequence (FLS) of deciduous woody plants. In a typical model of plant life-history, vegetative growth precedes reproduction. However, for many species in the forests of Eastern North America, it is not the green tips of new shoots that mark the commencement of the growing season, but the subtle reds and yellows of their flowers. This flowering-first FLS is common in these regions, and its prevalence suggests that this FLS has adaptive significance (?).

A deep inquiry into the nature of this phenological pattern is necessary and timely because anthropogenic climate change is altering FLSs (Fig. ??). For the three European tree species we examined, the number of days between flowering and leafout have increased as a result of climate change, but the rate of change differs among them. If, as suggested, optimum FLS patterning is an important component of fitness, this differential FLS sensitivity to climate change may influence which species will persist under altered climate conditions.

Despite recent advances in characterizing the evolution and underlying physiology of FLS (??), a major challenge to predicting how FLS patterns will shift in response to climate change is that we don't have a solid baseline understanding of variability in FLS. While some authors present general correlations between flowering and leafing phenology (??), fine scale FLS variability has never been evaluated. We suggest that characterizing FLS variation among individuals and populations will not only improve our ability to predict how FLS patterns will change in the future, but also allow for a more biologically relevant evaluation of the current FLS hypotheses, revealing avenues for future, direct hypothesis testing.

Here we 1) Review the hypotheses of woody plant FLSs and their respective predictions, 2) Evaluate variation in FLSs, and explore how FLS variation within species, populations and individuals alters the predictions of the hypotheses, 3) Show how the incorporation of variation reveals consistencies and anomalies in support for FLS hypotheses using several case studies from temperate forests, and 4) make recommendations for future study of FLSs.

## Defining FLS

Flower-leaf sequences have traditionally been classified into qualitative categories that are almost always defined at the species level. However, as we will demonstrate using a long term phenological data set from Harvard Forest in Petersham, MA (?), accurately classifying species into these categories is not a simple task.

The terms hysternanthy, protanthy, proteranthy or precocious flowering describe plants that produce flowers before their leaves (??). A classic example of this FLS is *Populus tremuloides*, which reaches peak flowering weeks before any sign of leaf development (Fig. ??). These species tend to exhibit a degree of physiological specialization, such as the separation of flower and leaf buds.

Seranthy describes species like *Nyssa sylvatica* in which flowers open as leaves approach their full size (Fig ??). These species can still differentiate flower buds in the previous season, but may rely less on stored energy than flowering-first taxa.

But what about species whose FLS separation is less clear? It is possible to describe all species whose flowering period overlaps their leaf development as synanthous (?), but this third category may obscure important inter-specific differences.

Take *Betula alleghaniensis* for example: One would be justified in classifying this species as hysternanthous because its flower buds tend to burst before its leaf buds, or as synanthous because its open flowers overlap the beginning of leaf growth (Fig. ??). Neither category is descriptive enough, because no matter the choice, it is clear that the FLS of this species is fundamentally different from *Acer rubrum*, whose flowers open weeks before the leaves and *Acer pensylvanicum* whose flowers do not open until leaves are well along in their expansion (Fig. ??).

In addition to obscuring species differences within FLS categories, the subjectivity of categorization makes it difficult to synthesize information across datasets. If a species is called hysternanthous in one data set and synanthous in another, should we interpret this as temporal or geographic variability in FLS or as an artifact of observer decision-making? This uncertainty hampers our ability to accurately test the existing FLS hypotheses because any statistical relationship between FLS and other traits is biased by the original observer.

However, we can extend our thinking regarding the hypotheses to overcome this bias. By examining the biological mechanisms underlying each hypotheses and incorporating their predictions regarding variability and overlapping phenophases in FLSs into analyses, we can arrive at better ways to utilize existing data and identify a new framework for treating FLS moving forward.

# Hypotheses of FLS and their predictions about flower-leaf overlap

Behind each prevailing FLS hypothesis there are evolutionary drivers that yield the phenological sequences observed today, and these mechanisms inherently accommodate different degrees of overlap between vegetative and floral phenophases. Here, we review the current FLS hypotheses, identify the underlying biology of each, and clarify how much overlap between flowering and vegetative growth they predict.

## Wind pollination

The most prevalent FLS hypothesis suggests that hysteresis is an adaptation critical for effective wind pollination, with leafless flowering allowing for more efficient pollen dispersal and transfer (???). This hypothesis hinges on the fact that leaves create a substantial physical disruption to pollen transfer, a premise that we would not necessarily expect to be true for the early stages of leaf expansion when tiny leaf primordia would have little impact on environmental structure. In this framework, we expect that trees that flower during the early stages of leaf expansion would gain similar mechanical advantage to those who complete their flowering before any leaf activity. We see that in Harvard Forest, while wind-pollinated species flower both before and after bud burst, they all flower before their leaves reach 75% of their final size (Fig. ??). This hypothesis predicts that wind pollinated species should flower before or with their leaves, while in animal pollinated species, FLS should be random or co-vary with pollinator activities.

## Water dynamics

Another hypothesis, emerging from the dry-deciduous tropics where flowering during the leafless season is also common (?), suggests that flowering before leaf development is an adaptation to reduce water stress associated with maintaining floral hydration while leaves are transpiring (?). This hypothesis asserts there is a significant cost to maintaining floral structures during any stage of leaf activity, and therefore only species whose flowering occurs before any leaf expansion would gain this drought advantage. This hypothesis predicts that species that are drought tolerant should flower before leafing out, with minimal overlap between the floral and vegetative phenophases. Species that are not drought tolerant gain no advantage from flowering first, so in these species FLSs should be random.

## Early flowering

A third possibility is that the flowering-first FLS is a physiological byproduct of selection for early flowering (?). Within this framework, there is no advantage to a species being hysteresis vs. seranthous, as long as the absolute flowering time of

the contrasting FLSs is the same. However, recent work from ? has demonstrated that floral hydration is primarily maintained by the phloem rather than the xylem. With leaf phenology constrained by the timing of xylem re-genesi, selection for early-season reproduction could drive flowering into the early season when xylem function is still suppressed, producing the hysternanthous FLS. This might explain why hysternanthous species tend to be the earliest species to flower (Fig. ??). Here, we expect longer time between flowering and leafing to be associated with earlier flowering phenology in general, and we expect more phenological overlap or a switch to seranth in later flowering species. For hysternanth, we might also expect to see strong associations with other early flowering traits such as seed mass, dispersal season or cold tolerance, but the hypothesis does not exclusively require the selective driver of early flowering to be one of these traits (?). This hypothesis predicts that early flowering times should be strongly associated with flowering-first FLSs. It also is likely there would be a relationship between this FLS and other early flowering traits, but the absence of these associations does not invalidate the hypothesis.

## Phylogenetics

Finally, it is also possible that FLSs are highly conserved traits, and the preponderance of hysternanth in the temperate zone is a product of phylogenetic representation of the region rather than an adaptive quality to the trait. In this framework, a species' FLS is under very weak or no selection so there are no expectations regarding the degree of overlap between flower and leaf phenological activity. This hypothesis predicts strong phylogenetic patterning in the FLS with no correlation with other traits.

*Need help with a short conclusions section hereon why figuring out these hypotheses NOW is most important for fundamental science, and for climate change, which has made this variation more obvious ... transition to variation. cite that it isn't really addressed anywhere else*

## Variation in FLS

. Intra-specific variation is the engine of natural selection, and if it is substantial in FLS patterns, we can infer much about the origins of this trait, and its trajectory as the climate changes. We investigated individual FLS variation in the Harvard Forest data (?), and found that the time between flowering and leaf activity varied by as much as several weeks for most species. This variability is lost completely in the classic framework of categorization. For example, for *Q. rubra*, a species classically listed as flowering and leafing in synanth, there are some years in which flower budburst is more than a week before leaf budburst, and other years in which leaf buds burst weeks prior to floral budburst (Fig. ??). We also found significant population-level variation in FLS, using the Pan European phenological database PEP725 (?), with the average time between

flowering and leafing varying between sites by a week or more.

Given the variability of FLSs at the individual and population level, it is clear that considering FLS variability at only higher taxonomic levels obscures important realities about the biology of this phenological trait. Below, we discuss how the observed variation below the species level may alter the existing FLS hypotheses.

## How FLS variation alters predictions

### Wind pollination

Pollination syndrome is generally treated as a species-level trait, considered to be fairly immutable across ecological time and space. Because of this, we would not expect significant variation in FLS across populations or individuals because we would not expect variation in pollination syndrome. However, as discussed above, a tree with no overlap between flowering and leafing phenology does not necessarily gain a significant pollen transfer advantage over an individual with some overlap. The pollination efficiency advantage from flowering-first diminishes as the canopy fills in, but we do not know at what point during leaf expansion pollination would become significantly encumbered. It is possible that interannual and population-level variation in the timing between flowering and leaf out for hysteranthous and synanthous individuals could maintain a wind pollination advantage, as long as the overlap did not cross a certain unknown threshold. Therefore, based on the wind pollination efficiency hypothesis, we would not expect high levels of population or individual variation in FLS, but the detection of some FLS variability at these levels does not inherently challenge this hypothesis.

### Water dynamics

If FLS's are driven by water dynamics, we would expect there to be significant population-level variation in FLSs. Populations growing in drier habitats should flower earlier relative to their leaf activity than their counterparts growing in wetter areas that experience weaker selection for minimizing phenological overlap. Therefore, increased time between flowering and leafing should be negatively correlated with average soil moisture. Water availability may also drive interannual FLS variation, with drought years increasing hysteranthous, and wetter years permitting more FLS overlap. Because plants have many other physiological mechanisms for dealing with occasional drought, we only expect to see a signal for the association between drought tolerance and hysteranthous for observations of populations in consistently dry regions.

### Early flowering

This hypothesis predicts some variation on the population level based on local adaptation. We would expect populations in which selection for earlier phenology is stronger,

perhaps those in regions with shorter growing seasons, to flower earlier relative to their leaf development. At the individual level, FLS variability could be driven by interannual variability in spring conditions. Both flowering and leaf phenology are strongly cued to temperature and photoperiod (??), but with leaf phenology constrained by xylem activity and flowering phenology relatively independent of it, we would expect a more sensitive response to environment in flowering time resulting in FLS variation. This hypothesis predicts that early flowering years or populations should be associated with an increase in the time between flowering and leafing for hysteranthous species. It also predicts a tighter temporal correlation between flowering and leafing for seranthous species or those with mixed buds in which flower timing is constrained by leaf budburst.

## Phylogenetics

With the lack of treatment of intra-specific FLS variability in the literature, we have no strong basis for asserting whether the apparent variability in FLSs is a product of genetic or environmental controls. If there is a strong genetic component to FLS as has been show for other phenophases (?), some population-level variation could be driven by reproductive isolation. With strong genetic control of FLS, we might also see consistent genotypic differences in FLS among individuals within a population, but would not predict high levels of interannual variation.

## Available evidence for FLS hypotheses in temperate woody species

Direct tests of these hypotheses are relatively rare in the literature, and—when tested—support for them is mixed. Many studies only test a single hypothesis, making comparison between them difficult. For example, the primary evidence for the wind pollination hypotheses comes from pollen diffusion studies, e.g., particle movement through closed and open canopies (???), which provide no framework for comparatively evaluating the other hypotheses. We are aware of no direct tests that have tried and distinguish selection for hysternathy from selection for early flowering, but ? notes that hysternanthous, wind-pollinated species tend to also have large seed mass, and lack primary seed dormancy for germination, traits associated with early flowering in general. This raises the distinct possibility that hysternathy may simply be one component of a larger suite of early flowering traits. We are also unaware of any studies that have mechanistically evaluated the water dynamics hypothesis, though observations of flowering in the dry tropics by ?? suggest that the timing of flowering in hysternanthous taxa is associated with a plant water status recovery due to leaf drop. Only recently has it even been suggested that this hypothesis might be relevant in the temperate zone as well, as we wouldn't expect that water status

would limit biological activity in the wet springs of the temperate zone (?).

In contrast, studies testing multiple hypotheses have generally found support for more than one evolutionary driver of hysteresis. One study by ? showed that wind-pollinated species tend to also be earlier flowering than their biotocally-pollinated sister taxa, suggesting a relationship between the early flowering and wind pollination hypotheses. A recent study by ? tested multiple hypotheses by modeling associations between species' trait and FLS patterns in the Great Lakes region. They found strong support for both the water dynamics and early flowering (flower timing and seed characteristics) hypotheses along with strong phylogenetic clustering.

In all of these cases, variability in FLS below the species level was not addressed. Yet, there are datasets widely available that allow for testing these several hysteresis hypotheses concurrently, and at multiple taxonomic levels. To address this gap, we supplement our literature review by re-testing some previously-used datasets to examine all hypotheses, and leverage several widely-available datasets to test how support for these hypotheses varies across the inter- to intra-specific levels.

We evaluated hysteresis in four phenological datasets. Michigan Trees and its companion volume Michigan Shrubs and Vines (??) (MTSV) contains categorical FLS information for 195 woody plant species. The USFS Silvics manual volume II (?) contains categorical FLS descriptions for 81 woody species. These data can be used to test inter-specific FLS variation. Within these datasets, we applied two alternative FLS classification schemes; physiological hysteresis, which allowed for no overlap between floral and leaf phenophases, and functional hysteresis, which allowed for a degree of overlap as predicted by the wind pollination hypotheses. The Harvard Forest data set (HF) contains quantitative flowering and leaf phenology measurements for individuals of 24 woody species over a 15 year period, allowing for both inter- and intra-specific comparisons (?). In this data set, we approximated the two hysteresis classification schemes mentioned above by measuring the time between several different floral and leaf phenophases. From the Pan European Phenological Database (PEP725) (?) we obtained spatially and temporally explicit, quantitative flowering and leaf phenology for four common European tree species. This allows us to evaluate FLSs only at the intra-specific level, but permits us to address variability among populations.

In considering all data sets in tandem two clear trends emerge: First, in accordance with the recent literature, we found support for multiple hypotheses (figure ??). There was generally strong support for the early flowering and wind pollination hypotheses, poor support for the water dynamics hypothesis, and the phylogenetic signal was variable (figure ??). But we also found that relative importance of each predictor, and therefore, the strength of the support for each hypothesis, changed significantly depending on how we defined hysteresis in the data set. Using continuous measures of FLS stabilized this modeling artifact, but increased the uncertainty around the estimates, suggesting categorical data may be over-simplifying trait relationships.

We used our intra-specific datasets to test some of the predictions we made about



intra-specific variability in the water dynamics and early flower hypotheses. Contrary to our prediction, we found that dry years correlate with a decrease in time between flowering and leafing for hysteroanthous species, largely due to delayed flowering. When we examined the relationship between 30 year soil moisture records (?) and population level variation in FLS timing across Germany, we found a weak negative association between average soil moisture levels and time between flowering and leafing as predicted by the water dynamics hypothesis. However, when we incorporate other predictors like flowering time into our analysis, the association disappeared, and early flowering was the only predictor of hysteroanth (Fig. ??, PEP725 estimates).

In accordance with our predictions for the early flowering hypothesis, we found that for hysteroanthous species, FLS variation is much more tightly correlated with variation in flowering timing than in leafing timing, but this contrast is far less stark in seranthous *Aesculus hippocastum* (figure ??). Though our intra-specific data set is species limited, we can refine our prediction to say that plasticity in the first phenophase of the season (flowering for hysteroanthous species and leafing for seranthous species) seems to drive variability in FLSs, but this observation should be tested more rigorously and explicitly in future work.

While the inter- and intra-specific case studies are not perfectly comparable (ie the wind pollination hypothesis cannot be evaluated on the intra-specific level), the general insights from our intra-specific studies supports the relationships found in the inter-specific case studies and provide novel, higher resolution insights of their own.

## Future

Perhaps more important than the results of all of our specific case studies themselves, is that in considering them together, we gain a more comprehensive picture of where our understanding of this phenological trait is currently, and where it needs to go. Below we highlight five characteristics of FLS that should be incorporated into future research.

## Multiple hypotheses explain FLSs

Our results underscore other lines of evidence that show multiple hypotheses should be starting point for all future FLS research. While there is certainly value to broad taxonomic studies, and future large-scale analyses should continue, the consistent support for multiple hypotheses shows there are limits to the utility of these kinds of studies. We suggest that it is better to explore the evolutionary dynamics of hysteroanth with a more mechanistic approach, which may mean utilizing a more taxonomically-restricted focus. One option is to look within the hypotheses to address sub-grouping of taxa in which overlap between hypotheses could be controlled. For example, what drives hysteroanth among biotically-pollinated taxa? It certain isn't wind pollination efficiency. Or, what factors accounts for variability in hysteroanth among wind-pollinated taxa? Incorporat-

ing a more explicit phylo-biogeographic approach would be instructive at this level, for example: are their phylo-geographic commonalities between the few biotically-pollinated hysteroanthous species in Eastern flora?

Even with drilling down to sub-groupings, interspecific trait-association models can only can take us so far. One reality of these kinds of studies is that we never know that we are picking the right traits. For example, we used minimum precipitation across a species' range, one of the only available quantitative drought metrics at the scale of large inter-specific models, to represent the water dynamics hypothesis. Is this really a good proxy for drought tolerance? Further, species evolve a suite of traits for any function, and unmeasured traits might bias our results (?). For example, wind-pollinated species could compensate for pollen intercepted by a synanthous or seranthous FLS by over-producing pollen or through self-pollination. To really understand FLS across large taxonomic space, one would have to compare species across an unfeasibly large, N-dimensional trait space, suggesting we will need to utilize other, complementary approaches.

### **Intra-specific variation in FLS**

In this paper, we have shown that FLSs can be highly variable at the intra-specific level. This variation can be leveraged in research design to overcome many of the limitations of larger trait-correlation models. Unlike with inter-specific approaches, focusing on FLS variation within species holds most other traits relatively equal, avoiding the problem of tradeoffs with latent unmeasured traits. Evolutionary theory predicts that intra-specific variation should follow the same trends as inter-specific variation, and consistent agreement between inter- and intra-specific, as we found in our analysis, will help narrow in on certain hypotheses.

### **The FLS is a quantitative trait**

Treating FLS observations as continuous variables are the most accurate way to describe these data. Our modeling work shows that this is an important step towards reducing observer bias and revealing important inter-specific differences that are masked by categorization. Quantitative measure of phenology, e.g. the BBCH scale (?), standardize data across time and space, observer, and analyst. Adopting such measurements in the study of phenological sequences would allow for FLS patterns to be compared across larger temporal, geographic and taxonomic scales, giving researchers more power to accurately address questions about FLS variation.

## FLS and fitness

While trait associations point to past selection, fitness is the driver of trait evolution, and at the core of each FLS hypothesis is a fitness prediction. By utilizing intra-specific comparisons and continuous measurements of FLS, we can move beyond trait associations and test the fitness consequences of FLS variation.

Variability in hysteresis should lead to variability in fitness outcome at the intra-specific level. For example, the wind pollination hypothesis predicts that with all else equal, years with increased time between flowering and leafing should correlate with more pollination success. The water dynamics hypothesis suggests hysteretic populations with a consistently large time between flowering and leafing should better tolerate drought. These predictions could be directly assessed through well-designed experiments and field studies.

## FLS and physiology

Decades of research suggests that both floral and vegetative phenological events are cued by temperature and photoperiod (??), suggesting they are under shared genetic and physiological control. But to yield the FLS variation seen in nature, there must be systematic differences in reproductive and vegetative phenological responses to the environment. Researchers can use intra-specific variation in FLS to identify which cues dominate each phenological process and better understand the underlying genetic and physiological constraints that structure phenological sequences.

With our proposed framework, we paint a picture of possibility for the study of FLSs in woody plants. Through examining FLS variation in more targeted taxonomic assemblages and using quantitative data with mechanistic metrics, we can refine the existing FLS hypotheses and better comprehend the causes and consequences of FLS variation at multiple taxonomic scales. This is an essential step towards a more complete understanding of the fundamental biology of temperate woody plants, and for predicting the fate of these species as global climate continues to change.

## References

- (2019) Dwd climate data center (cdc): Multi-annual grids of soil moisture in 5cm depth under grass and sandy loam, version 0.x., digital.
- Barnes, B.V. & Wagner, W.H.J. (1981,2004) *Michigan Trees: A guide to the Trees of the Great Lakes Region*. University of Michigan Press.
- Bolmgren, K., Eriksson, O. & Linder, H.P. (2003) Contrasting flowering phenology and

species richness in abiotically and biotically pollinated angiosperms. *Evolution* **57**, 2001–2011.

Borchert, R. (1983) PHENOLOGY AND CONTROL OF FLOWERING IN TROPICAL TREES. *BIOTROPICA* **15**, 81–89.

Burns, R.M., Honkala, B.H. & coordinators], T. (1990) Silvics of north america: Volume 2. hardwoods. Tech. rep., United States Department of Agriculture (USDA), Forest Service.

Burton V. Barnes, Christopher W. Dick, M.E.G. (2016) *Michigan Shrubs Vines: A guide to species of the Great Lakes Region*. University of Michigan Press.

Cleland, E.E., Chuine, I., Menzel, A., Mooney, H.A. & Schwartz, M.D. (2007) Shifting plant phenology in response to global change. *Trends in Ecology Evolution* **22**, 357 – 365.

Davies, T.J., Regetz, J., Wolkovich, E.M. & McGill, B.J. (2019) Phylogenetically weighted regression: A method for modelling non-stationarity on evolutionary trees. *Global Ecology and Biogeography* **28**, 275–285.

Ettinger, A., Gee, S. & M. Wolkovich, E. (2018) Phenological sequences: how earlyseason events define those that follow. *American Journal of Botany* **105**.

Finn, G.A., Straszewski, A.E. & Peterson, V. (2007) A general growth stage key for describing trees and woody plants. *Annals of Applied Biology* **151**, 127–131.

Flynn, D.F.B. & Wolkovich, E.M. (2018) Temperature and photoperiod drive spring phenology across all species in a temperate forest community. *New Phytologist* **219**, 1353–1362.

Forrest, J. & Miller-Rushing, A.J. (2010) Toward a synthetic understanding of the role of phenology in ecology and evolution. *Philosophical Transactions of the Royal Society B: Biological Sciences* **365**, 3101–3112.

Franklin, D.C. (2016) Flowering while leafless in the seasonal tropics need not be cued by leaf drop: evidence from the woody genus *Brachychiton* (malvaceae). *Plant Ecology and Evolution* **149**, 272–279.

Friedman, J. & Barrett, S.C.H. (2009) Wind of change: new insights on the ecology and evolution of pollination and mating in wind-pollinated plants. *Annals of Botany* **103**, 1515–1527.

Gougherty, A.V. & Gougherty, S.W. (2018) Sequence of flower and leaf emergence in deciduous trees is linked to ecological traits, phylogenetics, and climate. *New Phytologist* **220**, 121–131.

- 426 Heinig, R. (1899) Glossary of the botanic terms used in describing flowering plants. Cal-  
427 cutta, India.
- 428 Janzen, D.H. (1967) Synchronization of sexual reproduction of trees within the dry season  
429 in central america **21**, 620–637.
- 430 Lamont, B.B. & Downes, K.S. (2011) Fire-stimulated flowering among resprouters and  
431 geophytes in australia and south africa. *Plant Ecology* **212**, 2111–2125.
- 432 Lechowicz, M.J. (1995) Seasonality of flowering and fruiting in temperate forest trees.  
433 *Canadian Journal of Botany* **73**, 175–182.
- 434 Milleron, M., Lopez de Heredia, U., Lorenzo, Z., Perea, R., Dounavi, A., Alonso, J., Gil,  
435 L. & Nanos, N. (2012) Effect of canopy closure on pollen dispersal in a wind-pollinated  
436 species (*Fagus sylvatica* L.). *PLANT ECOLOGY* **213**, 1715–1728.
- 437 Nathan, R. & Katul, G. (2005) Foliage shedding in deciduous forests lifts up long-distance  
438 seed dispersal by wind. *PROCEEDINGS OF THE NATIONAL ACADEMY OF SCI-*  
439 *ENCES OF THE UNITED STATES OF AMERICA* **102**, 8251–8256.
- 440 Niklas, K.J. (1985) The aerodynamics of wind pollination **51**, 328–386.
- 441 O’Keefe, J. (2015) Phenology of woody species at harvard forest since 1990.
- 442 Piao, S., Friedlingstein, P., Ciais, P., Viovy, N. & Demarty, J. (2007) Growing season  
443 extension and its impact on terrestrial carbon cycle in the northern hemisphere over  
444 the past 2 decades. *Global Biogeochemical Cycles* **21**.
- 445 Primack, R.B. (1987) Relationships among flowers, fruits, and seeds. *Annual Review of*  
446 *Ecology and Systematics* **18**, 409–430.
- 447 Rathcke, B. & Lacey, E.P. (1985) Phenological patterns of terrestrial plants. *Annual*  
448 *Review of Ecology and Systematics* **16**, 179–214.
- 449 Reich, P. & Borchert, R. (1984) WATER-STRESS AND TREE PHENOLOGY IN A  
450 TROPICAL DRY FOREST IN THE LOWLANDS OF COSTA-RICA. *JOURNAL OF*  
451 *ECOLOGY* **72**, 61–74.
- 452 Robertson, C. (1895) The philosophy of flower seasons, and the phaenological relations of  
453 the entomophilous flora and the anthophilous insect fauna **29**, 97–117.
- 454 Savage, J.A. (2019) A temporal shift in resource allocation facilitates flowering before leaf  
455 out and spring vessel maturation in precocious species. *AMERICAN JOURNAL OF*  
456 *BOTANY* **106**, 113–122.
- 457 Stephen H. Spurr, B.V.B. (1980) *Forest Ecology*. John Wiley Sons, 3rd edn.

- 458 Templ, B., Koch, E., K.Bolmgren, Ungersböck, M., Paul, A., Scheifinger, H. & et al.  
459 (2018) Pan european phenological database (pep725): a single point of access for euro-  
460 pean data. *Int. J. Biometeorology* .
- 461 Whitehead, D.R. (1969) Wind pollination in the angiosperms: Evolutionary and environ-  
462 mental considerations **23**, 28–35.
- 463 Wilczek, A.M., Burghardt, L.T., Cobb, A.R., Cooper, M.D., Welch, S.M. & Schmitt,  
464 J. (2010) Genetic and physiological bases for phenological responses to current and  
465 predicted climates. *Philosophical transactions of the Royal Society of London. Series*  
466 *B, Biological sciences* **365**, 3129–3147.




Figure 1: Trends in average FLS offset across Europe for three tree species from 1960 to 2015. Highlighted regions indicate historic range of FLS variability. For all species, the time between flowering and leafing is increasing but the rate of change differs between species and sites




Figure 2: Average day of phenological events for woody plant species at Harvard Forest in Petersham, MA from 1990-2015. The community displays all major FLS patterns, but because of overlapping floral and vegetative development and interannual variability in phenology, it is difficult to neatly assign all species to a FLS category. Other notable patterns relevant to the FLS hypotheses can be seen. For example, as predicted by the early flowering hypothesis, the earliest species to initiate spring phenology are hysteroanthous and as predicted by the pollination syndrome hypothesis, wind-pollinated species (indicated with a \*) may vary in whether their flowers or vegetative buds break first, but all open their flowers before leaves expand to 75%.






Figure 3: FLS variability among years and within a population of *Quercus rubra* at Harvard Forest. While this species is typically reported as flowering and leafing in tandem, the order of flower and leaf bud break, and the time between them varies considerably within the community in a give year, and within an individual across year.




Figure 4: Estimated effects of water dynamics (minimum precipitation across species range or average soil moisture), pollination syndrome, and flowering time on FLS patterns across four case studies. We used phylogenetic adjustments and standardized units to make a rough comparison of four datasets of different taxonomic scopes (intra-vs. interspecific variation) data types (categorical and continuous) and definitions of FLS. While absolute parameter estimates should not be directly compared due to scaling inconsistencies between models and different modeling approaches for differing data structure, all models support the consensus that wind pollination and early flowering is associated with a flowering first FLS, and there is little effect of measures of water dynamics. Lines represent 95% bootstrap or credible intervals depending on the modeling framework.