

1 Phenological responses to climate mediate seedling competition with
2 an invasive woodland herb

3 D.M. Buonaiuto ^{1,2,3a}, E.M. Wolkovich^{2,3,4}

In review at *Journal of Ecology*

4 *Author affiliations:*

5 ¹Department of Environmental Conservation, University of Massachusetts, Amherst, Massachusetts, USA.

6 ORCID: 0000-0003-4022-2591

7 ²Arnold Arboretum of Harvard University, Boston, Massachusetts, USA.

8 ³Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, Massachusetts, USA

9 ⁴Forest & Conservation Sciences, Faculty of Forestry, University of British Columbia, Vancouver, British
10 Columbia, Canada

11 ^aCorresponding author: 617.823.0687; dbuonaiuto@umass.edu

Abstract

1. Invasive plants are often characterized by rapid germination and precocious phenology. Theory suggests that early germination may provide invaders with competitive advantage over slower germinating natives, but the relative contribution of rapid germination vs. other intrinsic competitive traits to the success of invaders is poorly understood. Depending on the relationship between germination and competition, shifts in germination phenology due to climate change may increase the dominance of invaders or buffer communities against their impacts.
2. We investigated the link between temperature variation, germination phenology and competitive interactions with a sequence of controlled environment experiments. First, we evaluated the relationships between temperature variation and germination phenology for two North American herbaceous species, the invasive *Hesperis matronalis* and native *Cryptotaenia canadensis*. We then leveraged temperature-response differences to manipulate the relative germination phenology of these taxa and quantified the effects of their phenological differences on competition.
3. Seeds of the invasive *H. matronalis* germinated rapidly, reaching 50% germination in under ten days in all treatment combinations. *C. canadensis* did not reach 50% germination with less than seven weeks of cold stratification. However, with more than 10 weeks of cold stratification and low (20/10°C) incubation temperatures, the germination phenology of *C. canadensis* was well matched to that of *H. matronalis*. When grown together, we found that precocious germination phenology doubled the competitive impact of *H. matronalis* relative to its other intrinsic competitive traits. Phenological advantage of just two-three days relative to *C. canadensis* was enough to secure competitive dominance at the seedling stage.
4. *Synthesis.* This study revealed that the mechanistic link between the germination phenology and competitive success of an invasive plant can be strongly mediated by climate sensitivity differences between introduced and native species. Climate change will likely exacerbate these differences, especially in regions where warming reduces the cold stratification. Our findings suggest that phenological diversity

in native plant communities is an important property of invasion resistance. The relationship between environmental variation, germination dynamics and competition provide a path forward for forecasting climate change impacts on seasonal community assembly, and highlights the need to incorporate phenological diversity in restorations.

Keywords: competition, climate change, germination, invasion, phenology, priority effects, stratification

Introduction

A central tenet of community assembly theory is that the order of arrival of species mediates inter-specific interactions and can dictate the trajectory of community structure (Fukami, 2015). These historical contingencies, known as priority effects, alter the structure and function of communities, driving communities to long-term alternate stable states (Fukami & Nakajima, 2011). Yet in many ecosystems, plant communities must re-assemble each year after a period of dormancy. In these communities, priority effects are the products of phenology, the timing of seasonal life cycle events, rather than the timing of the arrival of propagules, which occurs prior to the dormant season in many cases (Rudolf, 2019; Howe & Smallwood, 1982; Baskin & Baskin, 1988a).

Invasive plants are often characterized by rapid germination and precocious phenology under a wide variety of environmental conditions (Gioria *et al.*, 2018; Gioria & Pyšek, 2017; Wolkovich & Cleland, 2011; Smith, 2013). By contrast, native plants tend to exhibit more constrained germination cues (Marushia *et al.*, 2010; Wainwright & Cleland, 2013; Van Clef & Stiles, 2001). In many temperate systems, seeds of native plants are dispersed with deep physiological dormancy, requiring prolonged exposure to specific environmental conditions, such as cold stratification (cool temperatures of 0-10° C), to break dormancy and stimulate germination (ten Brink *et al.*, 2013; Cavieres & Sierra-Almeida, 2017; Bradford & Nonogaki, 2007).

These differences in germination physiology can yield strong differences in the relative germination phenology of invasive and native plants, with invaders germinating well before their native competitors (Gioria & Pyšek, 2017). This difference in relative germination timing among species, which we refer to as **phenological advantage**, can contribute significantly to the competitive abilities, and ultimately invasion success, of invasive plants. By allowing them to begin drawing down seasonal resources and modifying their environment before their native competitors emerge (Kardol *et al.*, 2013), invaders gain a competitive advantage through a **seasonal priority effect** (Wainwright *et al.*, 2011).

Despite the growing interest in seasonal priority effects, it has been difficult to quantify their overall contri-

bution to the competitive success of invaders. Germination is notoriously difficult to monitor in the field, and rapid phenology often co-varies with other competitive traits (Dickson *et al.*, 2012; Milbau *et al.*, 2003; Hao *et al.*, 2009). Because of these difficulties, many experiments vary phenological advantage by sowing competing seeds at different time intervals (Young *et al.*, 2017). While these experiments have provided strong evidence that phenological advantage—on the order of just days to weeks—can yield substantial priority effects (Weidlich *et al.*, 2020), their experimental set-up is difficult to translate into natural communities in which priority effects are mediated by climate, and difficult to use for forecasting. Understanding the role that phenological advantage plays in mediating the dynamics of interspecific competition is critical for predicting and managing the structure and function of plant communities in the face of anthropogenic climate change. Due to interspecific differences in responses to environmental variation, sustained alterations to environmental conditions are already shifting community-wide patterns of germination (Walck *et al.*, 2011). If patterns of germination are indeed tightly linked to the competitive dynamics of communities, then phenological reorganization is likely to shift the strength of species' interactions, change patterns of invasion, and strongly influence biological filtering of plant communities.

In this study, we generated contrasting levels of phenological advantage among two woodland herbaceous species (the North American invasive *Hesperis matronalis* and native *Cryptotaenia canadensis*) by leveraging their differences in germination timing in response to environmental variation. First, we performed a series of germination assays in controlled environments under varying temperature regimes to estimate a realistic range of climate-driven variation in phenological advantage. We then used competition trials under contrasting environmental conditions to indirectly manipulate the phenological advantage between these two taxa and quantify the contribution of seasonal priority effects to their competitive dynamics. By linking climate variation, phenological advantage, and seasonal priority effects, our study has important implications for how anthropogenic climate change will alter phenological assembly and, in turn, plant community interactions in the decades to come.

Materials and Methods

Focal species

For this study, we focused on a pair of woodland herbaceous species. Dames Rocket (*Hesperis matronalis*) is a herbaceous biennial/perennial species in the *Brassicaceae* family, originally from Eurasia, and introduced to North America in the 19th century (Francis *et al.*, 2009). It can rapidly invade meadows, forest edges and woodlands, forming thick stands and excluding native vegetation (Francis *et al.*, 2009). It is currently listed as a noxious or invasive weed in several states and provinces in the United States and Canada (Susko & Hussein, 2008). Honewort (*Cryptotania canadensis*) is a herbaceous perennial in the *Apiaceae* family, native to forests and woodlands of eastern North America (Hawkins *et al.*, 2007). The habitat overlap of these two species suggests that they may compete in nature. While their habitat requirements may be similar, the two species display a substantially different germination niche, making them a suitable model for our study. *C. canadensis* seeds are classified with deep physiological dormancy and require a substantial period of cold moist stratification to release dormancy and initiate germination (Baskin & Baskin, 1988b). While some reports suggest that cold stratification enhances germination in *H. matronalis* at low incubation temperatures, several studies have demonstrated that fresh and after-ripened (dry-stored) seeds of *H. matronalis* are capable of rapid and complete germination at a wide range of temperatures (Susko & Hussein, 2008).

Experiment I: Germination Assays

To investigate the relationship between environmental variation and relative germination timing among species, we obtained seeds of *C. canadensis* from Prairie Moon Nursery (Winona, MN) and seeds of *H. matronalis* from American Meadows (Shelburne, VT). We performed germination assays in the growth facilities of the Arnold Arboretum in Boston, Massachusetts, USA (42.3074° N, 71.1208° W). We assigned seeds to a fully-crossed set of twenty experimental treatments; 10 levels of cold stratification duration (0,2,4,5,6,7,8,9,11,13 weeks at 4°C) and two levels of incubation temperature (warm— 25°C:15°C (day/night),

cool— 20°C:10°C (day/night)).

Prior to applying experimental treatments we performed a “float test” in which all seeds were placed in distilled water, and unfilled seeds (floating) were removed from the experiment (Baskin & Baskin, 2014). We imbibed the remaining seeds in distilled water for 24 hours and then placed 20 seeds for each species/treatment combination in petri dishes on moist pool-filter sand, with three replicates per treatment. For the cold stratification treatments, we wrapped petri dishes in aluminum foil to prevent light exposure and placed them in a growth chamber at 4°C. After each stratification interval, we transferred the petri dishes to their assigned incubation chamber for 25 days, moistening the germination substrate as necessary to maintain maximum saturation of the medium without flooding the seeds. We checked for new germinates every two days, defining a seed as germinated when its radical or cotyledon tissue was visible (Baskin & Baskin, 2014). We assessed the viability of any seeds that did not germinate in the 25 day incubation period by performing a “crush test” in which we applied pressure to the intact seed to evaluate its condition (Baskin & Baskin, 2014). We excluded any seeds deemed unviable from all subsequent analyses. Due to the staggering of our stratification treatments the experiment took place between 27 August - 12 December 2018.

Experiment II: Competition Trials

To quantify the contribution of seasonal priority effects to interspecific competition dynamics we performed competition trials under controlled conditions in a research greenhouse at the Arnold Arboretum from October 2020 - February 2021. We planted seeds of *C. canadensis* and *H. matronalis* into 8.9 cm square pots, employing a response surface design where we varied both the overall density of seeds and proportion of each species in each pot (Inouye, 2001). High and low density treatments consisted of 14 and 8 seeds respectively. Proportion treatments were 100:0%, 25:75%, 50:50%, 75:25%, 0:100% (*C. canadensis*:*H. matronalis*). Each density by proportion treatment was replicated six times.

We randomly assigned half of the pots to low (45 days) and high (72 days) cold stratification treatments in dark growth chambers at 4°C. We staggered the start of the treatments, so that at the conclusion of the

cold stratification, all pots were transferred to a heated greenhouse maintained at 15-25°C with 14 hours of supplemental light. Germination was observed daily from 24 December 2020 - 13 January 2021 and every two days from 15 January 2021 - 01 February 2021. The locations of each pot in the greenhouse were randomly reassigned every three days to minimize any blocking effects on germination or growth.

After 35 days we added 1 tsp per gallon of water of Peters 20-10-20 liquid feed fertilizer to all pots. After 62 days, we harvested the above-ground biomass from all pots, dried it for 48 hours at 60°C, and recorded the dry weight of each species/pot.

Statistical analysis

Germination Assays

To assess interspecific differences in the relationship between germination rate and temperature variability, we fit a Bayesian mixed-effect accelerated failure time model (AFT, Onofri *et al.*, 2010) using a Weibull distribution for the likelihood function. We included weeks of stratification and incubation temperature and their interaction with species as fixed effects. The model written below is modified from Onofri *et al.* (2010).

$$t_{50} = t_0 \phi$$

where t_{50} is time to 50% germination and corresponds to the germination times of a references seed lot (t_0) multiplied by and “acceleration factor” (ϕ). The acceleration factor is a product of the experimental treatments of our study through the equation:

$$\phi = \exp(\beta_{sp}X_{sp} + \beta_{strat}X_{strat} + \beta_{inc}X_{inc} + \beta_{stratXspecies}X_{strat} * X_{sp} + \beta_{incXspecies}X_{inc} * X_{sp})$$

where X_{sp} , X_{strat} and X_{inc} are the species, stratification and incubation treatment levels in our experiment, and β_{sp} , β_{strat} , β_{inc} , $\beta_{stratXspecies}$ and $\beta_{incXspecies}$ are the estimated effects on ϕ for adding an additional week or stratification or degree of incubation for each species respectively.

The AFT modeling framework let us robustly compare germination timing (t_{50} , time to 50% germination)

even among treatments with different final germination percentages by accounting for viable seeds that did not germinate during our incubation window (Soltani *et al.*, 2015; Onofri *et al.*, 2010). One drawback of this approach is that this class of models assumes that all viable seeds will eventually germinate, which we would not expect in nature. For this reason, we considered any estimated t50 values greater than 40 days to indicate that seeds would not reach 50% germination under those conditions.

Competition trials

We quantified phenological advantage between species by subtracting the mean germination time of *H. matronalis* from that of *C. canadensis* in each pot. This allowed us to evaluate the effect phenological advantage with a regression design (Cottingham *et al.*, 2005), with advantage values ranging from -1.3 to 9.5 (*C. canadensis* mean germination time 1.3 days earlier to 9.5 days later than *H. matronalis*).

For each pot, we calculated the relative growth rate difference (RGRD) among species using the equation below, modified from Connolly & Wayne (2005).

$$\text{RGRD} = \ln\left(\frac{Y_{Cc}}{y_{Cc}}\right) - \ln\left(\frac{Y_{Hm}}{y_{Hm}}\right)$$

where Y_{Hm} and Y_{Cc} are the final biomass of the species at the end of the experiment and y_{Hm} and y_{Cc} are the initial biomass of the seeds planted at the outset of the experiment. For this calculation we obtained estimates of seed mass for our focal species from the Kew Gardens Seed Information Database (Kew, 2022).

We then modeled the effect of seedling density of *C. canadensis*, *H. matronalis* and phenological advantage using on RGRD. The model is written below:

$$\text{RGRD}_i = N(\alpha + \beta_{Hm}n_{Hm} + \beta_{Cc}n_{Cc} + \beta_{pri}MGT, \sigma_{RGRD}^2)$$

where β_{Hm} and β_{Cc} are known as the species influence parameters—representing the intrinsic competitive ability of each species—or the estimated effect of increasing the seedling density of each species by one individual on the RGRD (Connolly & Wayne, 2005). n_{Hm} and n_{Cc} are the number of germinated individuals of *H. matronalis* and *C. canadensis* respectively. The relationship between the species influence parameters and n_{Hm} and n_{Cc} indicate that these parameter estimates are dependent on both the number of seeds planted (i.e. the seed bank in each pot) and the fraction of those seeds that germinated. This formulation allows our model to partition the effects of interspecific differences in germination phenology from the effects of interspecific differences in germination fraction under each environmental treatment. β_{pri} is the priority effect, or the effect of increasing the difference in mean germination time between *H. matronalis* and *C. canadensis* by one day. In this formulation, α is an un-interpretable intercept (Connolly & Wayne, 2005).

Model Implementation

We fit all models using the R package “brms” (Bürkner, 2018). We ran the model on four chains with 4000 iterations and a 3000 iteration warm-up for a total of 4000 posterior draws for each parameter, using weakly informative priors. We validated model performance by obtaining \hat{R} values between 1 and 1.01, high effective sample sizes and no divergent transitions. For all models we report the mean posterior estimate along with 90% uncertainty intervals (I_{90}).

Results

Germination advantage

H. matronalis reached 50% germination in under ten days for all environmental treatments, always exceeding 75% germination regardless of environmental conditions (Fig. 1, 2, Tab. S1). Increasing cold stratification duration and incubation temperature only marginally enhanced the germination rate of this species (Fig. 2). By contrast, increasing incubation temperature had a delaying effect on the germination rate of *C.*

canadensis, suggesting that the mean 20°C temperature of our warm incubation treatment is supra-optimal for the species (Fig. 2). Without sufficient cold stratification (<7 weeks for low incubation and <10 weeks for high incubation temperatures), seeds of *C. canadensis* did not reach 50% germination during the duration of our experiment (Fig. 1, 2, Tab. S1). However, under high levels of cold stratification germination rates of *C. canadensis* began to converge on those of *H. matronialis*, and at levels of stratification >10 weeks and low incubation, the germination rate and fraction of *C. canadensis* was well matched to that of *H. matronialis* (Fig. 1, 2, Tab. S1).

Germination priority effects

In the absence of phenological advantage, the influence of adding one seedling of *H. matronialis* to a pot community was almost 4X less than adding one *C. canadensis* seedling on the pot-level RGRD (represented by the species' influence parameters—*H. matronialis* (β_{Hm}): 0.13, I_{90} : 0.08, 0.17, *C. canadensis* (β_{Cc}): -0.40, I_{90} : -0.46, -0.35). Each day increase in the phenological advantage of *H. matronialis* had approximately the same influence on shifting the community biomass composition towards *H. matronialis* as adding an individual of that species to the community (seasonal priority effect (β_{pri}): 0.15, I_{90} : 0.09, 0.20, Fig. 3, Tab. S2). Together, these results suggest that *H. matronialis* will come to dominate the community biomass composition unless *C. canadensis* is at high relative abundance or the phenological advantage of *H. matronialis* is small.

Discussion

Environmental drivers of seasonal priority effects

Our results identify a natural mechanism—species' differential sensitivity to temperature—that can generate seasonal priority effects, and thus reshape the competitive landscape between native and invasive species. In the absence of phenological advantage, the intrinsic competitive abilities of each species in our study suggest that the native species, *C. canadensis*, is the stronger competitor (Fig. 3). However, the influence of

one day of phenological advantage for *H. matronalis* virtually doubled its influence on the final community composition, suggesting that seasonal priority effects play a major role in the competitive success of *H. matronalis* (Fig. 3). Our results indicate that *C. canadensis* can compete with the invasive *H. matronalis* at high relative abundance levels and/or when phenological advantage is low, joining a growing body of experiments demonstrating that relative germination phenology can function as a seasonal priority effect, enhancing the performance of the earliest germinating species at the expense of later germinants (Koerner *et al.*, 2008; Dickson *et al.*, 2012; Ross & Harper, 1972).

Our findings advance decades of work on seasonal priority effects by connecting them directly to environmental variation and extending them to forest systems. While research on the importance of seasonal priority effects has increased in recent years, the vast majority of studies have focused on dry grassland communities, and typically introduce priority effects artificially — by staggered planting of competitors (Young *et al.*, 2017; Weidlich *et al.*, 2020). In contrast, our experiment employed a more ecologically realistic mechanism—generating variation in phenological advantage indirectly by leveraging interspecific differences in germination responses to cold stratification. Our results suggest that seasonal priority effects can strongly influence seedling competition even in environment where germination phenology is primarily under temperature control. The results of our pair-wise competition trial suggests that seasonal priority effects, manifested through rapid germination phenology and propagule pressure, are mechanistically related to the competitive dominance, and—ultimately—invasion success of *H. matronalis*.

Seasonal priority effects and anthropogenic climate change

The implications of our study for the role of climate variability in mediating seasonal priority effects is two-fold. First, our results suggest that interannual climate variability should generate both among- and within-season variation in competition strength among species, potentially driving species coexistence via the storage effect (Chesson, 2003). Second, the key role we observed of climate in generating germination advantage and therefore seasonal priority effects suggests that sustained alteration to historic patterns of climate variability,

like those driven by anthropogenic climate change, are likely to alter the dynamics of competing seedlings. Changing patterns of phenological assembly have downstream effects on the structure and function of plant communities.

In our study, the phenological advantage of *H. matronalis* was maximized under lower stratification treatments and warmer incubation temperatures. This suggests that the warming temperatures associated with anthropogenic climate change may increase the magnitude of seasonal priority effects, largely due to the delay of germination in more climate-sensitive native species like *C. canadensis*. Interestingly, the difference in phenological advantage among our focal species was much higher in our germination assays than in our competition trials even at comparable levels of stratification. There are likely several explanations for these differences.

First, we used different metrics of germination speed, time to 50% germination (t50) and mean germination time in each experiment. While the metrics are related and often confused, there are important differences between them that make one or the other more appropriate for the two types of experiments we ran (see Supporting Information: “Measures of germination speed”). Second, the incubation temperatures in our greenhouse competition trials were more variable than in our growth chamber germination assays. The lower germination fractions we observed in *C. canadensis* under greenhouse conditions suggest that the temperature range was likely supra-optimal for this species, and the lower germination fraction increased the difference between t50 estimates and mean germination time measurements. Finally, we conducted germination assays and competition trials in different growth media (filter sand vs. potting soil), which have different moisture retention and light transmissible capacities. Germination media can affect germination rates (Baskin & Baskin, 2014), which may further explain differences among our two experiments.

Despite these differences, in both experiments increasing cold stratification advanced the germination phenology of *C. canadensis* and weakly that of *H. matronalis*, resulting in weaker phenological advantage for *H. matronalis* at higher stratification levels. (Fig. 2, Fig. S1) suggesting that the relationships between cold stratification and germination phenology we observed were robust.

Climate change may also increase the risk of precocious phenology (Inouye, 2000). While in our experiment, there was no cost to germinating too early, in natural systems, selection usually limits early activity. Across environments, optimum germination phenology is likely driven by a trade-off between maximizing the length of the growing season and the risk of exposure to damaging environmental episodes when germinating too early (Augspurger & Salk, 2017). For example, in dry grassland ecosystems, the precocious germination of invasives has a substantial cost if water availability is too low (Wainwright *et al.*, 2011). In temperate forest ecosystems, the primary risk of early phenology is damage from late season frost (Kollas *et al.*, 2014) and climate change is also altering the timing and frequency of frost events (Ma *et al.*, 2019). Understanding how climate change will reshape this tradeoff between seasonal priority effects and frost risk is a critical next step for understanding plant community interactions in an era of global change.

While we found that seasonal priority effects impacted competition among seedlings, our experiment was not able to quantify the role of seasonal priority effects in influencing the long-term, among-year dynamics of these perennial species. Many studies suggest that these short term priority effects may be transient, though several studies that used staggered planting methods at similar scale to the phenological advantage we observed in our trials saw the influence of these initial priority effects on community composition maintained several seasons later (Vaughn & Young, 2015; Young *et al.*, 2017; Torrez *et al.*, 2017). In perennial communities, these long term dynamics are even more difficult to assess. Many perennial herbs, *C. canadensis* included, rely heavily on vegetative reproduction (Hawkins *et al.*, 2005), and competition between ramets, and between ramets and seedlings, may also impact species interactions in the long-term. Understanding how phenological differences across life stages of long-lived perennial plants affects competition is an important need for predicting how communities may be impacted by interannual environmental variation and climate change.

Because we found that seasonal priority effects mediated competition for early ontological stages (germination, seedling), our findings may be most relevant to global change biology in the context of native plant establishment; whether colonizing new areas due to range shifts, recovering from novel disturbances or eco-

logical restoration. In fact, there has been a growing call to increase phenological diversity in restoration planning (Hess *et al.*, 2019). Studies have found that including early active species in plantings can suppress the abundance of invaders in both grassland (Cleland *et al.*, 2013) and forest ecosystems (Schuster *et al.*, 2020). At the same time, restoration mixes tend to lack species which fill the early season phenological niche (Havens & Vitt, 2016). The results of our study suggest that minimizing the priority effect conferred to invasive species due to rapid germination and early phenology by including species with similar, early phenological traits could be a powerful tool for managing plant invasions and restoring native ecosystems in an era of global change.

Conclusion

By leveraging the differential germination sensitivities to environmental cues of two competing species to manipulate phenological advantage between them, we were able to quantify the contribution of seasonal priority effects gained through rapid phenology on the competitive ability of the invasive species *H. matronalis*. We found that priority effects were approximately as strong as the intrinsic competitive traits of *H. matronalis* in influencing its competitive dominance over the native forest herb *Cryptotaenia canadensis*, suggesting seasonal priority effects mechanistically increase the invasion success of *H. matronalis*. Variation in germination phenology was strongly mediated by differences in how species respond to temperature cues, suggesting that sustained climate change will alter patterns of phenological advantage, potentially strengthening the seasonal priority effects of invaders as climate warms. Our findings highlight the important role of phenological diversity in the invasion resistance of native plant communities, implying that measures of phenological diversity should be incorporated into plant community assessments and ecological restoration.

Author Contributions

DMB and EMW conceived of this manuscript; DMB designed and executed the experiments, collected the data and performed the analyses; DMB led the writing of the manuscript. All authors contributed critically to drafts and gave final approval for submission.

Acknowledgements

Special thanks to K.J Woodruff, L. Toomey and the rest of the Arnold Arboretum research greenhouses staff for helping to manage the growth facilities as well as maintain and repeat these experiments throughout lab closures, quarantines and all other Covid19-related challenges.

Conflict of Interest

The authors declare no conflict of interest.

Data Availability

Data from the germination assays and competition trials, and associated modeling code will be made available at the time of publication at KNB (<https://knb.ecoinformatics.org/>).

References

- Augsburger, C.K. & Salk, C.F. (2017) Constraints of cold and shade on the phenology of spring ephemeral herb species. *Journal of Ecology* **105**, 246–254.
- Baskin, C. & Baskin, J. (2014) *Seeds: Ecology, Biogeography, and Evolution of Dormancy and Germination*. Elsevier Inc.
- Baskin, C.C. & Baskin, J.M. (1988a) Germination ecophysiology of herbaceous plant species in a temperate region. *American Journal of Botany* **75**, 286–305.

- Baskin, J.M. & Baskin, C.C. (1988b) The ecological life cycle of *cryptotaenia canadensis* (L.) DC. (Umbelliferae), a woodland herb with monocarpic ramets. *The American Midland Naturalist* **119**, 165–173.
- Bradford, K.J.K.J. & Nonogaki, H. (2007) *Seed development, dormancy, and germination*. Blackwell Pub., Oxford, OX, UK; Ames, Iowa, USA.
- Bürkner, P.C. (2018) Advanced bayesian multilevel modeling with the R package brms. *R Journal* **10**, 395–411.
- Cavieres, L. & Sierra-Almeida, A. (2017) Assessing the importance of cold-stratification for seed germination in alpine plant species of the high-andes of central Chile. *Perspectives in Plant Ecology, Evolution and Systematics* **30**.
- Chesson, P. (2003) Quantifying and testing coexistence mechanisms arising from recruitment fluctuations. *Theoretical Population Biology* **64**, 345–357.
- Cleland, E.E., Larios, L. & Suding, K.N. (2013) Strengthening invasion filters to reassemble native plant communities: Soil resources and phenological overlap. *Restoration Ecology* **21**, 390–398.
- Connolly, J. & Wayne, P. (2005) Assessing determinants of community biomass composition in two-species plant competition studies. *Oecologia* **142**, 450–457.
- Cottingham, K.L., Lennon, J.T. & Brown, B.L. (2005) Knowing when to draw the line: designing more informative ecological experiments. *Frontiers in Ecology and the Environment* **3**, 145–152.
- Dickson, T.L., Hopwood, J.L. & Wilsey, B.J. (2012) Do priority effects benefit invasive plants more than native plants? an experiment with six grassland species. *Biological Invasions* **14**, 2617–2624.
- Francis, A., Cavers, P.B. & Warwick, S.I. (2009) The biology of Canadian weeds. 140. *hesperis matronalis* L. *Canadian Journal of Plant Science* **89**, 191–206.
- Fukami, T. (2015) Historical contingency in community assembly: Integrating niches, species pools, and priority effects. *Annual Review of Ecology, Evolution, and Systematics* **46**, 1–23.
- Fukami, T. & Nakajima, M. (2011) Community assembly: alternative stable states or alternative transient states? *Ecology Letters* **14**, 973–984.
- Gioria, M. & Pyšek, P. (2017) Early bird catches the worm: germination as a critical step in plant invasion. *Biological Invasions* **19**, 1055–1080.
- Gioria, M., Pyšek, P. & Osborne, B.A. (2018) Timing is everything: does early and late germination favor invasions by herbaceous alien plants? *Journal of Plant Ecology* **11**, 4–16.
- Hao, J.H., Qiang, S., Liu, Q.Q. & Cao, F. (2009) Reproductive traits associated with invasiveness in *conyza sumatrensis*. *Journal of Systematics and Evolution* **47**, 245–254.
- Havens, K. & Vitt, P. (2016) The importance of phenological diversity in seed mixes for pollinator restoration. *Natural Areas Journal* **36**, 531–537.
- Hawkins, T.S., Baskin, J.M. & Baskin, C.C. (2005) Life cycles and biomass allocation in seed- and ramet-derived plants of *cryptotaenia canadensis* (apiaceae), a monocarpic species of eastern North America. *Canadian Journal of Botany* **83**, 518–528.
- Hawkins, T.S., Baskin, J.M. & Baskin, C.C. (2007) Seed morphology, germination phenology, and capacity to form a seed bank in six herbaceous layer apiaceae species of the eastern deciduous forest. *Castanea* **72**, 8–14.
- Hess, M.C.M., Mesléard, F. & Buisson, E. (2019) Priority effects: Emerging principles for invasive plant species management. *Ecological Engineering* **127**, 48–57.

- Howe, H.F. & Smallwood, J. (1982) Ecology of seed dispersal **13**, 201–228.
- Inouye, B. (2001) Response surface experimental designs for investigating interspecific competition. *Ecology* **82**, 2696–2706.
- Inouye, D.W. (2000) The ecological and evolutionary significance of frost in the context of climate change. *Ecology Letters* **3**, 457–463.
- Kardol, P., Souza, L. & Classen, A.T. (2013) Resource availability mediates the importance of priority effects in plant community assembly and ecosystem function. *Oikos* **122**, 84–94.
- Kew, R.B.G. (2022) Seed information database (sid). version 7.1.
- Koerner, C., Stoecklin, J., Reuther-Thiebaud, L. & Pelaez-Riedl, S. (2008) Small differences in arrival time influence composition and productivity of plant communities. *New Phytologist* **177**, 698–705.
- Kollas, C., Körner, C. & Randin, C.F. (2014) Spring frost and growing season length co-control the cold range limits of broad-leaved trees. *Journal of Biogeography* **41**, 773–783.
- Ma, Q., Huang, J.G., Hänninen, H. & Berninger, F. (2019) Divergent trends in the risk of spring frost damage to trees in europe with recent warming. *Global Change Biology* **25**, 351–360.
- Marushia, R.G., Cadotte, M.W. & Holt, J.S. (2010) Phenology as a basis for management of exotic annual plants in desert invasions. *Journal of Applied Ecology* **47**, 1290–1299.
- Milbau, A., Nijs, I., Van Peer, L., Reheul, D. & De Cauwer, B. (2003) Disentangling invasiveness and invasibility during invasion in synthesized grassland communities. *New Phytologist* **159**, 657–667.
- Onofri, A., Gresta, F. & Tei, F. (2010) A new method for the analysis of germination and emergence data of weed species. *Weed Research* **50**, 187–198.
- Ross, M.A. & Harper, J.L. (1972) Occupation of biological space during seedling establishment **60**, 77–88.
- Rudolf, V.H.W. (2019) The role of seasonal timing and phenological shifts for species coexistence. *Ecology Letters* **22**, 1324–1338.
- Schuster, M.J., Wragg, P.D., Williams, L.J., Butler, E.E., Stefanski, A. & Reich, P.B. (2020) Phenology matters: Extended spring and autumn canopy cover increases biotic resistance of forests to invasion by common buckthorn (*rhamnus cathartica*). *Forest Ecology and Management* **464**, 118067.
- Smith, L.M. (2013) Extended leaf phenology in deciduous forest invaders: mechanisms of impact on native communities. *Journal of Vegetation Science* **24**, 979–987.
- Soltani, E., Ghaderi-Far, F., Baskin, C.C. & Baskin, J.M. (2015) Problems with using mean germination time to calculate rate of seed germination. *Australian Journal of Botany* **63**, 631–635.
- Susko, D.J. & Hussein, Y. (2008) Factors affecting germination and emergence of dame’s rocket (*hesperis matronalis*). *Weed Science* **56**, 389–393.
- ten Brink, D.J., Hendriksma, H.P. & Bruun, H.H. (2013) Habitat specialization through germination cueing: a comparative study of herbs from forests and open habitats. *Annals of Botany* **111**, 283–292.
- Torrez, V., Mergeay, J., Meester, L.D., Honnay, O., Helsen, K. & Standish, R. (2017) Differential effects of dominant and subordinate plant species on the establishment success of target species in a grassland restoration experiment. *Applied vegetation science* **20**, 363–375.
- Van Clef, M. & Stiles, E.W. (2001) Seed longevity in three pairs of native and non-native congeners: Assessing invasive potential. *Northeastern Naturalist* **8**, 301–310.

- 415 Vaughn, K.J. & Young, T.P. (2015) Short-term priority over exotic annuals increases the initial density and
416 longer-term cover of native perennial grasses. *Ecological Applications* **25**, 791–799.
- 417 Wainwright, C.E. & Cleland, E.E. (2013) Exotic species display greater germination plasticity and higher
418 germination rates than native species across multiple cues. *Biological Invasions* **15**, 2253–2264.
- 419 Wainwright, C.E., Wolkovich, E.M. & Cleland, E.E. (2011) Seasonal priority effects: implications for invasion
420 and restoration in a semi-arid system. *Journal of Applied Ecology* **49**, 234–241.
- 421 Walck, J.L., Hidayati, S.N., Dixon, K.W., Thompson, K. & Poschlod, P. (2011) Climate change and plant
422 regeneration from seed. *Global Change Biology* **17**, 2145–2161.
- 423 Weidlich, E.W.A., Nelson, C.R., Maron, J.L., Callaway, R.M., Delory, B.M. & Temperton, V.M. (2020)
424 Priority effects and ecological restoration. *Restoration Ecology* **n/a**, e13317.
- 425 Wolkovich, E.M. & Cleland, E.E. (2011) The phenology of plant invasions: a community ecology perspective.
426 *Frontiers in Ecology and the Environment* **9**, 287–294.
- 427 Young, T.P., Stuble, K.L., Balachowski, J.A. & Werner, C.M. (2017) Using priority effects to manipulate
428 competitive relationships in restoration. *Restoration Ecology* **25**, S114–S123.

429 **Figures**




Figure 1: Cumulative germination rates of *H. matronalis* and *C. canadensis* under cool (20/10° C day/night) and warm (25/15° C day/night) incubation temperatures for each stratification treatment (number of weeks at 4° C, right axis) in our germination assays. Points depict the mean cumulative germination at each observation time, and vertical bars depict the standard error across replicates. Lines represent a generalized trend for cumulative germination, fit with a loess function. See Fig. 2 for model estimates from these data.




Figure 2: The effects of weeks of cold stratification at 4° C on the time to 50% germination of *Cryptotaenia canadensis* and *Hesperis matronalis* under warm (20/10° C day/night) and cool (25/15° C day/night) incubation conditions, estimated with accelerated failure time model. We show here only stratification treatment levels which allowed both species to reach 50% germination in less than 40 days. The solid lines depict the mean estimate, while lighter lines depict uncertainty.




Figure 3: Estimated effects of species' abundance (intrinsic competitive ability parameters) and phenological advantage (seasonal priority effects) on the relative growth rate difference between *H. matronalis* and *C. canadensis*. Negative parameter estimates indicate the community biomass composition shifts to favor *C. canadensis* while positive estimate towards dominance by *H. matronalis*. The points indicate the mean estimated effect of each parameter and bars the 90% uncertainty intervals. The full posterior distribution for each parameter is also depicted as an additional measure of uncertainty.