

A Collider Search for Dark Matter Produced in Association
with a Higgs Boson with the CMS Detector at the 13 TeV LHC

By

DUSTIN RAY BURNS

B.S. Physics (Georgia Institute of Technology) 2011

B.S. Applied Mathematics (Georgia Institute of Technology) 2011

M.S. Physics (University of California, Davis) 2012

DISSERTATION

Submitted in partial satisfaction of the requirements for the degree of

DOCTOR OF PHILOSOPHY

in

Physics

in the

OFFICE OF GRADUATE STUDIES

of the

UNIVERSITY OF CALIFORNIA

DAVIS

Approved:

Chair Michael Mulhearn

Robin Erbacher

Albert De Roeck

Committee in Charge

2017

Copyright © 2017 by
Dustin Ray Burns
All rights reserved.

*To someone very important . . .
a nice dedication.*

CONTENTS

List of Figures	iv
List of Tables	v
Abstract	vi
Acknowledgments	vii
1 Motivation and theoretical background	1
1.1 The Standard Model	1
1.1.1 Symmetries of nature	1
1.1.2 Particle content	8
1.2 Dark matter	11
1.2.1 Background	11
1.2.2 Beyond the Standard Model	12

LIST OF FIGURES

- 1.1 4l invariant mass distribution showing H discovery in the ZZ^* decay channel. The red line shows the signal distribution for $m_H = 126$ GeV. Figure taken from <https://twiki.cern.ch/twiki/bin/view/CMSPublic/Hig12041TWiki> 11

LIST OF TABLES

1.1	The particles of the Standard Model.	9
-----	----------------------------------------------	---

ABSTRACT

A Collider Search for Dark Matter Produced in Association with a Higgs Boson with the CMS Detector at the 13 TeV LHC

The abstract submitted as part of your dissertation, in the introductory pages, does not have a word limit. It should follow the same format as the rest of your dissertation (1.5 inch left margin, double-spaced, consecutive page numbering, etc.).

ACKNOWLEDGMENTS

Acknowledgements to those who helped you get to this point. They should be listed by chapter when appropriate.

Chapter 1

Motivation and theoretical background

The first section of this chapter lays the theoretical framework of the Standard Model (SM) of particle physics, including the historical discovery timeline of symmetries of nature, each corresponding to a leap forward in our physical understanding, and the model's particle content and their properties. The second section extends the symmetry group of the SM to include candidates for the observed cosmic dark matter (DM) and the particles that could mediate the DM-SM interactions.

1.1 The Standard Model

1.1.1 Symmetries of nature

Each major advance in the history of physics has corresponded to the discovery and mathematical implementation of a new global space-time, global discrete, or local gauge symmetry. In this section, I will review these discoveries, informally developing the mathematical framework needed to understand the symmetry groups of the SM. Along the way, two important subplots will play out: the development of our understanding of physics at smaller distance scales and higher energies, and the unification of previously separate physical sectors.

Our ancient ancestors were aware that certain geometrical shapes, e.g. the Platonic solids, possessed the quality of symmetry, and were driven to understand the composition

of physical substances by breaking them down into fundamental, indivisible units. These units, known as atoms by the ancient Greeks, interact and rearrange themselves according to physical laws to account for the variety of substances and physical phenomena we observe. The attraction to applying symmetry to nature is evidenced by the centuries-long belief that the Earth lie at the center of the universe, with the celestial bodies orbiting in perfect, divine circles.

The end of the scientifically repressive middle ages brought along an improvement in astronomical observations and the growth of the pseudo-scientific field of alchemy, which attempted to reduce, understand, and manipulate the fundamental elements of physical substances. When Kepler discovered three laws of planetary motion, he unified the description of the motion of the planets. For the first time, the conservation of a physical quantity, what we now know as angular momentum, was associated with a general set of physical objects. At the burgeoning of the scientific revolution, Newton championed the idea that the same physical laws can be applied to all physical events, and that properties of these laws can be abstracted to apply to nature at a fundamental level.

Newton defined an inertial reference frame implicitly as one where his first law held, that is, that an object remains at constant motion unless acted on by an outside force. Since his laws were the same in all inertial reference frames, a new symmetry of nature was discovered, now called symmetry under Euclidean transformations. Since Euclidean transformations form a mathematical group, it is said that classical mechanics is invariant under the Euclidean group. The invariance under the Euclidean transformations can be used to derive conservation laws: Newton's laws don't depend on spatial translations or rotations, implying the conservation of linear and angular momentum, respectively. These relationships foreshadow Noether's abstraction of the connection between continuous symmetries and conserved quantities. She proved that there is a conserved quantity, or current, associated with every symmetry of a physical system [Noether(1918a)]. This famous theorem facilitates the derivation of conserved quantities and will be used extensively in the theories that follow. The extension of the Euclidean transformations to

include time translations and motion at constant velocity (boosts) forms the Galilean group.

The next symmetry of nature to be discovered came when Lorentz found that Maxwell's equations, which unified the classical electricity and magnetism sectors, were invariant under a set of transformations that generalized the classical Galilean translations, now called Lorentz transformations, which form the Lorentz group [Lorentz(1898)]. The Lorentz symmetry corresponds to the conservation law of total electric charge [Noether(1918b)]. Soon after, Einstein derived the Lorentz symmetry as a property of space-time itself through his special theory of relativity, assuming two simple postulates: the laws of physics and the speed of light are the same in all inertial reference frames [Einstein(1905a)]. From these simple assumptions, Einstein was able to extend the classical laws of physics to the high velocity, high energy sector. Before Einstein, the symmetries of nature were thought to be consequences of the physical laws themselves, but Einstein's major paradigm shift was to view the symmetry itself as the more fundamental property, an insight key to his formulation of the general theory of relativity for the gravitational interaction [Einstein(1914)], which is not discussed further in this thesis. Combining the Lorentz and Euclidean transformations yields the Poincare group, the final global space-time symmetry of the SM.

Just as relativists were probing physics at higher speeds and energies, other physicists were investigating the behavior of systems at smaller distance scales, conducting experiments to explore phenomena such as the Compton effect [Compton(1923)] and photoelectric effect [Einstein(1905b)], which showed the quantized, particle-like behavior of light, and electron beam diffraction [Davisson and Germer(1927)], which showed the wave-like behavior of electrons. Quantum theory was developed to consolidate the wave-like and particle-like behaviors of systems, and extended the validity of classical physics to microscopic scales. Where in classical physics particles states were described by their absolute position and momentum, quantum theory describes the state of a system by a probabilistic wavefunction. The classical symmetries and conservation laws carried over to quantum theory, with the concept of angular momentum generalized to include spin. Particles with zero or integer spin are called Bosons, and were shown to obey Bose-Einstein statistics,

where the wavefunction is symmetric under all permutations of particles. Particles with half-integer spin, called Fermions, obey Fermi-Dirac statistics, where the wavefunction is symmetric under even permutations and changes sign under odd permutations, implying that no two particles can occupy the same state. These many-particle symmetries were applied to derive properties of materials and states of electrons in atomic and periodic systems, blackbody radiation, and many other properties of matter and radiation [Shankar(1994)].

Quantum theory was successfully applied to a myriad of low-energy systems. Dirac extended quantum theory and the Schrodinger equation, which describes the evolution of a quantum system in time, to the relativistic regime [Dirac(1967)]. His formulation of the wave equation for a spin-1/2 particle with mass m is inherently Lorentz invariant:

$$(i\gamma^\mu\partial_\mu - m)\psi = 0 \tag{1.1}$$

The solutions to the Dirac equation were found to have both positive and negative energy solutions, to the surprise of Dirac. His explanation was that the vacuum consisted of a "sea" of negative energy solutions, each in a distinct state due to the Pauli exclusion principle, and when a pair of electrons was produced, a positive energy state was filled and a negative energy state was vacated, creating a "hole" in the sea. The particle corresponding to this hole would have the same energy as the electron, but in order to conserve total charge, must have the opposite sign charge. This positively charged electron was not known to exist at the time, but was soon discovered in cosmic ray experiments and named the positron [Anderson(1933)]. Dirac's theoretical prediction of the positron and its subsequent discovery opened the door for the discovery that every particle has an oppositely charged antimatter partner.

In an attempt to apply relativistic quantum theory to the electromagnetic field and the spin-0 photon, Dirac formulated the theory of quantum electrodynamics (QED) [Dirac(1927)]. QED is the first example of a quantum field theory (QFT), where the physical dynamics apply to the quantum field associated with a particle rather than the particle itself, and particles/antiparticles interact as excitations of the field. This formu-

lation of creation and annihilation of particles and antiparticles gave a more physically intuitive explanation for the negative energy solutions of the Dirac equation than the Dirac sea. QED became the prototype for developing future relativistic quantum theories and Dirac's procedure the template for quantizing a general field theory.

With the discovery of antimatter and the apparatus of QFT in place, physicists continued searching for additional symmetries. If a particle state is an eigenvector of a symmetry operator, then the eigenvalue is an important quantum number of the state, since if the symmetry is conserved, the quantum number can be used to determine if a decay or interaction of this state is allowed. Three important discrete symmetries operations, and their products, have had significant importance: space coordinate parity inversion (P), particle-antiparticle charge conjugation (C), and time reversal (T) [Bettini(2014)]. Violation of P was observed for the weak interactions by Wu and others in the decays of Co-60 atoms [Wu et al.(1957)Wu, Ambler, Hayward, Hoppes, and Hudson]. Although it was long believed to be a symmetry of all the interactions, CP, the product of C and P was also found to be violated by the weak decays of K mesons [Christenson et al.(1964)Christenson, Cronin, Fitch, and T]. It is only the combination of P, C, and T that is now thought to be an exact symmetry of nature, obeyed by all interactions. The CPT symmetry was proven to be conserved in all relativistic QFTs [Schwinger(1951), Luders(1954)].

The mid-20th century saw an explosion of discovery in particle physics and was a golden age for the feedback between theoretical and experimental work. The discovery of additional particles such as the muon [Anderson and Neddermeyer(1936)], pion [Lattes et al.(1947)Lattes, Occhialini, and Powell], neutrino[Cowan et al.(1956)Cowan, Reines, Harrison, and many others inspired the theoretical development of the quark model [Gell-Mann(1964)], the refinement of QED [Schwinger(1948a), Schwinger(1948b), Feynman(1949a), Feynman(1949b), Feynman(1950), Tomonaga(1946), Dyson(1949a), Dyson(1949b)], and the formulation of theories to explain the strong [Gell-Mann(1964), Zweig(1964a), Zweig(1964b), Politzer(1973), Gross and Wilczek(1973)] and weak [Fermi(1933)] nuclear forces. Conversely, these new theories led to the prediction and subsequent discovery of new fundamental particles, such as the charm [Augustin et al.(1974)Augustin, Boyarski, Breidenbach, Bulos, Dakin, Feldman, Fischer, Fr

Aubert et al.(1974)Aubert, Becker, Biggs, Burger, Chen, Everhart, Goldhagen, Leong, McCorriston, Rh and bottom [Herb et al.(1977)Herb, Hom, Lederman, Sens, Snyder, Yoh, Appel, Brown, Brown, Innes, U quarks, and composite particles, such as the Ω^- [Barnes et al.(1964)].

The development of the theories of the strong and weak forces unveiled a new set of symmetries: local invariance under unitary gauge transformations. In keeping with the trend of the discovery of new symmetries being associated with the unification of physical sectors, the electromagnetic and weak nuclear interactions were found to be components of a single force, called the electroweak force [?, ?]. Again, with QED as the prototype, the Langrangians for the electroweak and strong forces were constructed to be invariant under unitary groups. Early work by Weyl [Weyl(1927)] showing the gauge invariance of electromagnetism was extended to QED, and lay the mathematical framework for describing the gauge invariance of the other forces. To demonstrate gauge invariance and how the gauge group generators are associated with the force carriers, take the QED terms of the SM Lagrangian as an example case [Michael E. Peskin(1995)]:

$$\mathcal{L} \supset \bar{\psi}(i\gamma^\mu D_\mu - m)\psi - \frac{1}{4}F^{\mu\nu}F_{\mu\nu} \quad (1.2)$$

where A_μ is the EM 4-potential and field corresponding to the photon, $F^{\mu\nu} = \partial^\mu A^\nu - \partial^\nu A^\mu$ is the electromagnetic field tensor, and $D_\mu = \partial_\mu + ieA_\mu$ is the gauge covariant derivative. These terms are invariant under the the U(1) transformations

$$\psi \rightarrow e^{i\theta(x)}\psi. \quad (1.3)$$

In general, the generators of an interaction's continuous symmetry group correspond to the gauge fields whose excitations are the gauge bosons that mediate that interaction. Just how A_μ is the field corresponding to the photon in QED and generator of the U(1) symmetry group of EM, the W and Z bosons and gluons are formed from the generators of the symmetry groups for the electroweak and strong interactions, SU(2)xU(1) and SU(3), respectively. Additionally, the covariant derivative is transformed in a way analagous to the QED covariant derivative, adding a factor for each generator with the corresponding charge in the coefficient. These charges (electric, color, weak isospin and hypercharge)

are conserved for all of the forces in all interactions[Bettini(2014)]. The charges of the SM particles are described in more detail in the next section.

Although the mathematical descriptions of their symmetries are similar, the EM, weak, and strong nuclear forces are quite different qualitatively. The EM force is the most familiar, being responsible for the interactions of matter and radiation at macroscopic scales, having only one type of charge which can be positive or negative. The strong force is mediated by eight gluons, corresponding to the eight generators of $SU(3)$, and carries three types of charge, known as red(R), green(G), and blue(B). The "negatives" of these charges are called anticolors: antired(\bar{R}), antigreen(\bar{G}), and antiblue(\bar{B}). Keeping with the visible color analogy, the theory of the strong force is called quantum chromodynamics (QCD). At low energies, the quarks and antiquarks are confined to form only color-neutral states. This is known as color confinement, and implies no free quarks exist, but only come in neutral combinations called hadrons. Since hadrons are strictly color-neutral, their states transform as the singlet representation under $SU(3)$. At high energies, the strong coupling decreases, and the quarks can be treated as effectively free particles with perturbation theory using Feynman diagrams. This property of the strong force is known as asymptotic freedom. The weak force is qualitatively different than either the EM or strong forces, as particles do not exchange its mediators to form bound states of any kind. The low-energy limit of the weak theory was developed by Fermi to explain beta decays [Fermi(1933)], but the full description was not developed until the electroweak force was formulated.

The unification of the electromagnetic and weak interactions into the electroweak force at about 100 GeV had one major shortcoming: the invariance of the Lagrangian required the gauge bosons, B from $U(1)$ and W^1, W^2, W^3 from $SU(2)$, to be massless. While the photon is indeed massless, the W and Z bosons have a nonzero mass []. Higgs and others [?, ?, ?, ?, ?, ?] proposed a solution by introducing a new scalar field whose excitations were called the Higgs boson (H). The scalar field is a complex doublet, meaning it has four total real components, and its vacuum expectation value (vev), or value throughout all of space, is one of many non-zero values in the bottom of its "Mexican hat" potential.

This spontaneous symmetry can be broken by choosing one of the values for the vev: $H = \frac{1}{\sqrt{2}}(0, \nu)$, where $\nu = 246$ GeV is called the H vev. At energies above $O(100)$ GeV, the electroweak symmetry is obeyed, the gauge bosons are massless, and the Higgs field has one of many values along the circle at the base of its potential. When the H field is expressed as a perturbation about this vev, the electroweak symmetry is broken into the weak and EM forces, and mass terms are generated for the weak force bosons, which are expressed as linear combinations of B, W^1, W^2, W^3 . The process of the Higgs acquiring a vev, spontaneously breaking the symmetry in its potential, and generating the masses of the weak bosons, is known as the Higgs mechanism, and results in the breaking of the electroweak symmetry. This mechanism can be carried out for a general operator depending on H with the substitution

$$H \rightarrow \frac{1}{\sqrt{2}}(\nu + h) \quad (1.4)$$

where h is the physical Higgs, corresponding to the leftover degree of freedom of the H doublet that is not absorbed by the three massive gauge bosons. H couples to the SM fermions, detailed in the next section, not by the same mixing as described for the gauge bosons, but via Yukawa interactions, direct couplings whose coefficients are related to the fermion masses. The Higgs mechanism was the final piece of the puzzle to understanding the fundamental laws of particle physics.

The final result of this saga is the standard model of particle physics, a relativistic gauge quantum field theory, globally invariant under the Poincare group, and locally invariant under the product of the strong and electroweak symmetry groups, $SU(3) \times SU(2) \times U(1)$, which describes all of the fundamental particles and their interactions. The particle content of the SM is detailed in the next section.

1.1.2 Particle content

The particle content of the SM is displayed in Table 1.1

The particles of the standard model consist of the spin-1/2 fermions, which interact to form regular matter, the spin-1 bosons, which mediate the interactions of the fermions, and the spin-0 scalar Higgs boson, which generates the masses of the bosons and fermions

u	c	t	...	γ	H
d	s	b		g	...
e	μ	τ		W	
ν_e	ν_μ	ν_τ		Z	

Table 1.1. The particles of the Standard Model.

via the Higgs mechanism. Unless otherwise labeled, the material in this section comes from [Bettini(2014)].

The fermions come in three sets of increasing masses, called generations, corresponding to the first three columns of Table 1.1. Across the rows, the particles have similar properties, and are abbreviated u^i, d^i, e^i, ν_e^i from top to bottom with $i = 1, 2, 3$ the generation index. Within each generation, the fermions are divided into two categories: the quarks, which are charged under the EM, weak, and strong forces, and the leptons, which are charged under the EM and weak forces. Each fermion has a corresponding antiparticle, whose mass is the same, but whose charges are opposite in sign. The quarks are SU(3) triplets, having a color charge of either R, G, or B. Since the weak force violates P, the fermions can be distinguished by their chirality, being labelled as either right handed (e_R^i) or left handed (e_L^i). The $(u^i, d^i)_L$ and $(e^i, \nu_e^i)_L$ pairs and their right handed anti-particle pairs are SU(2) doublets and interact via the weak force. The right handed fermions (left handed anti-fermions) are SU(2) singlets and do not interact via the weak force. The weak isospins (T_3) for the left handed fermions are: $(u^i, d^i)_L = (1/2, -1/2)$ and $(e^i, \nu_e^i)_L = (-1/2, 1/2)$. The EM charges are $(u^i, d^i)_L = (2/3e, -1/3e)$ and $(e^i, \nu_e^i)_L = (-1e, 0)$, and finally, the weak hypercharges are, from the relation $Y_W = 2(Q - T_3)$, $(u^i, d^i)_L = (1/3, 1/3)$ and $(e^i, \nu_e^i)_L = (-1, -1)$.

The fourth column of Table 1.1 lists the force mediators, or gauge bosons. g stands for the 8 gluons of QCD that mediate the strong nuclear force. The gluons are octets under SU(3), and correspond to linear combinations of the generator gauge fields G_μ^a . Gluons carry color charge themselves, but are electrically neutral. They are massless, consistent with the fact that they correspond to generators of a conserved symmetry, and may be

represented using the Gell-Mann matrices as the linearly independent set of states [?]:

$$\frac{1}{\sqrt{2}}(r\bar{b}+b\bar{r})\frac{1}{\sqrt{2}}(r\bar{g}+g\bar{r})\frac{1}{\sqrt{2}}(b\bar{g}+g\bar{b})\frac{1}{\sqrt{2}}(r\bar{r}-b\bar{b})-i\frac{1}{\sqrt{2}}(r\bar{b}-b\bar{r})-i\frac{1}{\sqrt{2}}(r\bar{g}-g\bar{r})-i\frac{1}{\sqrt{2}}(b\bar{g}-g\bar{b})\frac{1}{\sqrt{6}}(r\bar{r}+b\bar{b}-2g\bar{g}) \quad (1.5)$$

while the color singlet state that the colorless hadrons are in is:

$$\frac{1}{\sqrt{3}}(r\bar{r} + g\bar{g} + b\bar{b}). \quad (1.6)$$

The remaining gauge bosons mediate the electroweak force. Before electroweak symmetry breaking, the generators of $SU(2) \times U(1)$ correspond to the gauge fields B, W^1, W^2, W^3 , whose excitations are massless gauge bosons. After symmetry breaking via the Higgs mechanism, three of the bosons acquire mass and the electroweak gauge bosons are reparametrized as:

$$W^\pm = \frac{1}{\sqrt{2}}(W^1 \pm iW^2)Z = \cos\theta_w W^3 - \sin\theta_w B\gamma = \sin\theta_w W^3 + \cos\theta_w B \quad (1.7)$$

where θ_w is the weak mixing angle, the massive W^\pm and Z bosons mediate the weak force, and the massless γ is the photon which mediates EM. W^\pm have electric charge $\pm 1e$ while the Z and γ are neutral. The isospin of W^\pm is ± 1 and 0 for Z and γ , giving hypercharges of 0 for W^\pm and 0 for Z and γ .

The final particle of the SM is the scalar H. H is electrically neutral and constructed to be an $SU(2)$ doublet before electroweak symmetry breaking, with one component having weak isospin 1/2 (hypercharge -1), and the neutral component having isospin -1/2 (hypercharge 1), which includes the physical h. After electroweak symmetry breaking, three of the H components are absorbed by the gauge bosons, and the remaining physical h remains neutral. The parity of h is 1. Although H couples to all massive fermions and bosons, the decay channels that are relevant for collider searches are: $ZZ^* \rightarrow 4l, WW^* \rightarrow 2l2\nu, \gamma\gamma, \tau\bar{\tau}$, and $b\bar{b}$. After a decades long search, the discovery and verification of quantum numbers of H was announced by the CMS and ATLAS experiments in 2012 [?, ?]. The four-lepton invariant mass distribution, showing the H peak at its observed mass of $m_H = 126$ GeV is shown in Figure 1.1.




Figure 1.1. $4l$ invariant mass distribution showing H discovery in the ZZ^* decay channel. The red line shows the signal distribution for $m_H = 126$ GeV. Figure taken from <https://twiki.cern.ch/twiki/bin/view/CMSPublic/Hig12041TWiki>

1.2 Dark matter

1.2.1 Background

This section gives an overview of the most compelling sources of observational evidence for the existence of DM, the potential particles candidates for DM, and the methods for detecting them.

1.2.1.1 Observational evidence

The earliest indication that there may be matter in the universe that cannot be detected by conventional optical observations, so called dark matter (DM), came from measurements of the orbital velocities of astronomical bodies in galaxy clusters [?, ?] and galaxies [?, ?]. Classical Newtonian gravity gives a galactic rotation curve, which shows how the velocity v of a massive object depends on its distance r from the center of a galaxy, as $vr \propto \sqrt{M(r)/r}$, where $M(r)$ is the total mass in the galaxy within a radius r . However, measurements of the orbital velocities of objects outside of the visible part of galaxies, where $v(r) \propto 1/\sqrt{r}$, show instead $v(r) \propto \text{const.}$, i.e. that the mass $M(r) \propto r$ instead of being constant [?]. The larger than expected velocities imply the existence of this dark halo of non-luminous matter in galaxies.

bullet cluster and lensing

CMB

structure formation

1.2.2 Beyond the Standard Model

1.2.2.1 Effective field theory

1.2.2.2 Simplified models

REFERENCES

- [Noether(1918a)] Emmy Noether. Invariant Variation Problems. *Gott. Nachr.*, 1918: 235–257, 1918a. doi: 10.1080/00411457108231446. [Transp. Theory Statist. Phys.1,186(1971)].
- [Lorentz(1898)] H. A. Lorentz. Simplified Theory of Electrical and Optical Phenomena in Moving Systems. *Koninklijke Nederlandse Akademie van Wetenschappen Proceedings Series B Physical Sciences*, 1:427–442, 1898.
- [Noether(1918b)] Emmy Noether. Invariant Variation Problems. *Gott. Nachr.*, 1918: 235–257, 1918b. doi: 10.1080/00411457108231446. [Transp. Theory Statist. Phys.1,186(1971)].
- [Einstein(1905a)] Albert Einstein. On the electrodynamics of moving bodies. *Annalen Phys.*, 17:891–921, 1905a. doi: 10.1002/andp.200590006. [Annalen Phys.14,194(2005)].
- [Einstein(1914)] Albert Einstein. On the Foundations of the Generalized Theory of Relativity and the Theory of Gravitation. *Phys. Z.*, 15:176–180, 1914.
- [Compton(1923)] Arthur H. Compton. A quantum theory of the scattering of x-rays by light elements. *Phys. Rev.*, 21:483–502, May 1923. doi: 10.1103/PhysRev.21.483. URL <http://link.aps.org/doi/10.1103/PhysRev.21.483>.
- [Einstein(1905b)] Albert Einstein. Concerning an heuristic point of view toward the emission and transformation of light. *Annalen Phys.*, 17:132–148, 1905b.
- [Davisson and Germer(1927)] C. Davisson and L. H. Germer. Diffraction of Electrons by a Crystal of Nickel. *Phys. Rev.*, 30:705–740, 1927. doi: 10.1103/PhysRev.30.705.
- [Shankar(1994)] R. Shankar. *Principles of Quantum Mechanics*. Plenum Press, New York, 2 edition, 1994.
- [Dirac(1967)] P.A.M. Dirac. *The Principles of Quantum Mechanics*. Oxford University Press, 4 edition, 1967.
- [Anderson(1933)] Carl D. Anderson. The positive electron. *Phys. Rev.*, 43:491–494, Mar 1933. doi: 10.1103/PhysRev.43.491. URL <http://link.aps.org/doi/10.1103/PhysRev.43.491>.
- [Dirac(1927)] P. A. M. Dirac. The quantum theory of the emission and absorption of radiation. *Proceedings of the Royal Society of London A: Mathematical, Physical and Engineering Sciences*, 114(767): 243–265, 1927. ISSN 0950-1207. doi: 10.1098/rspa.1927.0039. URL <http://rspa.royalsocietypublishing.org/content/114/767/243>.

- [Bettini(2014)] Alessandro Bettini. *Introduction to Elementary Particle Physics*. Cambridge University Press, 2 edition, 2014.
- [Wu et al.(1957)Wu, Ambler, Hayward, Hoppes, and Hudson] C. S. Wu, E. Ambler, R. W. Hayward, D. D. Hoppes, and R. P. Hudson. Experimental test of parity conservation in beta decay. *Phys. Rev.*, 105:1413–1415, Feb 1957. doi: 10.1103/PhysRev.105.1413. URL <http://link.aps.org/doi/10.1103/PhysRev.105.1413>.
- [Christenson et al.(1964)Christenson, Cronin, Fitch, and Turlay] J. H. Christenson, J. W. Cronin, V. L. Fitch, and R. Turlay. Evidence for the 2π decay of the k_2^0 meson. *Phys. Rev. Lett.*, 13:138–140, Jul 1964. doi: 10.1103/PhysRevLett.13.138. URL <http://link.aps.org/doi/10.1103/PhysRevLett.13.138>.
- [Schwinger(1951)] Julian Schwinger. The theory of quantized fields. i. *Phys. Rev.*, 82:914–927, Jun 1951. doi: 10.1103/PhysRev.82.914. URL <http://link.aps.org/doi/10.1103/PhysRev.82.914>.
- [Luders(1954)] Gerhart Luders. On the Equivalence of Invariance under Time Reversal and under Particle-Antiparticle Conjugation for Relativistic Field Theories. *Kong. Dan. Vid. Sel. Mat. Fys. Med.*, 28N5:1–17, 1954.
- [Anderson and Neddermeyer(1936)] Carl D. Anderson and Seth H. Neddermeyer. Cloud chamber observations of cosmic rays at 4300 meters elevation and near sea-level. *Phys. Rev.*, 50:263–271, Aug 1936. doi: 10.1103/PhysRev.50.263. URL <http://link.aps.org/doi/10.1103/PhysRev.50.263>.
- [Lattes et al.(1947)Lattes, Occhialini, and Powell] C. M. G. Lattes, G. P. S. Occhialini, and C. F. Powell. Observations on the Tracks of Slow Mesons in Photographic Emulsions. 1. *Nature*, 160:453–456, 1947. doi: 10.1038/160453a0.
- [Cowan et al.(1956)Cowan, Reines, Harrison, Kruse, and McGuire] C. L. Cowan, Jr., F. Reines, F. B. Harrison, H. W. Kruse, and A. D. McGuire. Detection of the Free Neutrino: A Confirmation. *Science*, 124:103–104, July 1956. doi: 10.1126/science.124.3212.103.
- [Gell-Mann(1964)] Murray Gell-Mann. A Schematic Model of Baryons and Mesons. *Phys. Lett.*, 8:214–215, 1964. doi: 10.1016/S0031-9163(64)92001-3.
- [Schwinger(1948a)] Julian Schwinger. On quantum-electrodynamics and the magnetic moment of the electron. *Phys. Rev.*, 73:416–417, Feb 1948a. doi: 10.1103/PhysRev.73.416. URL <http://link.aps.org/doi/10.1103/PhysRev.73.416>.
- [Schwinger(1948b)] Julian Schwinger. Quantum electrodynamics. i. a covariant formulation. *Phys. Rev.*, 74:1439–1461, Nov 1948b. doi: 10.1103/PhysRev.74.1439. URL <http://link.aps.org/doi/10.1103/PhysRev.74.1439>.

- [Feynman(1949a)] R. P. Feynman. The theory of positrons. *Phys. Rev.*, 76:749–759, Sep 1949a. doi: 10.1103/PhysRev.76.749. URL <http://link.aps.org/doi/10.1103/PhysRev.76.749>.
- [Feynman(1949b)] R. P. Feynman. Space-time approach to quantum electrodynamics. *Phys. Rev.*, 76:769–789, Sep 1949b. doi: 10.1103/PhysRev.76.769. URL <http://link.aps.org/doi/10.1103/PhysRev.76.769>.
- [Feynman(1950)] R. P. Feynman. Mathematical formulation of the quantum theory of electromagnetic interaction. *Phys. Rev.*, 80:440–457, Nov 1950. doi: 10.1103/PhysRev.80.440. URL <http://link.aps.org/doi/10.1103/PhysRev.80.440>.
- [Tomonaga(1946)] S. Tomonaga. On a relativistically invariant formulation of the quantum theory of wave fields. *Prog. Theor. Phys.*, 1:27–42, 1946. doi: 10.1143/PTP.1.27.
- [Dyson(1949a)] F. J. Dyson. The radiation theories of tomonaga, schwinger, and feynman. *Phys. Rev.*, 75:486–502, Feb 1949a. doi: 10.1103/PhysRev.75.486. URL <http://link.aps.org/doi/10.1103/PhysRev.75.486>.
- [Dyson(1949b)] F. J. Dyson. The s matrix in quantum electrodynamics. *Phys. Rev.*, 75:1736–1755, Jun 1949b. doi: 10.1103/PhysRev.75.1736. URL <http://link.aps.org/doi/10.1103/PhysRev.75.1736>.
- [Zweig(1964a)] G. Zweig. An $SU(3)$ model for strong interaction symmetry and its breaking. Version 1. 1964a.
- [Zweig(1964b)] G. Zweig. An $SU(3)$ model for strong interaction symmetry and its breaking. Version 2. In D.B. Lichtenberg and Simon Peter Rosen, editors, *DEVELOPMENTS IN THE QUARK THEORY OF HADRONS. VOL. 1. 1964 - 1978*, pages 22–101. 1964b. URL <https://inspirehep.net/record/4674/files/cern-th-412.pdf>.
- [Politzer(1973)] H. David Politzer. Reliable perturbative results for strong interactions? *Phys. Rev. Lett.*, 30:1346–1349, Jun 1973. doi: 10.1103/PhysRevLett.30.1346. URL <http://link.aps.org/doi/10.1103/PhysRevLett.30.1346>.
- [Gross and Wilczek(1973)] David J. Gross and Frank Wilczek. Asymptotically free gauge theories. i. *Phys. Rev. D*, 8:3633–3652, Nov 1973. doi: 10.1103/PhysRevD.8.3633. URL <http://link.aps.org/doi/10.1103/PhysRevD.8.3633>.
- [Fermi(1933)] Enrico Fermi. Tentativo di una teoria dell’emissione dei raggi beta. *Ric. Sci.*, 4:491–495, 1933.
- [Augustin et al.(1974)] Augustin, Boyarski, Breidenbach, Bulos, Dakin, Feldman, Fischer, Fryberger, Han J. E. Augustin, A. M. Boyarski, M. Breidenbach, F. Bulos, J. T. Dakin, G. J. Feldman, G. E. Fischer, D. Fryberger, G. Hanson, B. Jean-Marie, R. R. Larsen, V. Lüth, H. L. Lynch, D. Lyon, C. C. Morehouse, J. M. Paterson, M. L. Perl,

- B. Richter, P. Rapidis, R. F. Schwitters, W. M. Tanenbaum, F. Vannucci, G. S. Abrams, D. Briggs, W. Chinowsky, C. E. Friedberg, G. Goldhaber, R. J. Hollebeek, J. A. Kadyk, B. Lulu, F. Pierre, G. H. Trilling, J. S. Whitaker, J. Wiss, and J. E. Zipse. Discovery of a narrow resonance in e^+e^- annihilation. *Phys. Rev. Lett.*, 33:1406–1408, Dec 1974. doi: 10.1103/PhysRevLett.33.1406. URL <http://link.aps.org/doi/10.1103/PhysRevLett.33.1406>.
- [Aubert et al.(1974)] Aubert, Becker, Biggs, Burger, Chen, Everhart, Goldhagen, Leong, McCorriston, R. J. J. Aubert, U. Becker, P. J. Biggs, J. Burger, M. Chen, G. Everhart, P. Goldhagen, J. Leong, T. McCorriston, T. G. Rhoades, M. Rohde, Samuel C. C. Ting, Sau Lan Wu, and Y. Y. Lee. Experimental observation of a heavy particle j . *Phys. Rev. Lett.*, 33:1404–1406, Dec 1974. doi: 10.1103/PhysRevLett.33.1404. URL <http://link.aps.org/doi/10.1103/PhysRevLett.33.1404>.
- [Herb et al.(1977)] Herb, Hom, Lederman, Sens, Snyder, Yoh, Appel, Brown, Brown, Innes, Ueno, Yamanouchi, S. W. Herb, D. C. Hom, L. M. Lederman, J. C. Sens, H. D. Snyder, J. K. Yoh, J. A. Appel, B. C. Brown, C. N. Brown, W. R. Innes, K. Ueno, T. Yamanouchi, A. S. Ito, H. Jöstlein, D. M. Kaplan, and R. D. Kephart. Observation of a dimuon resonance at 9.5 gev in 400-gev proton-nucleus collisions. *Phys. Rev. Lett.*, 39:252–255, Aug 1977. doi: 10.1103/PhysRevLett.39.252. URL <http://link.aps.org/doi/10.1103/PhysRevLett.39.252>.
- [Barnes et al.(1964)] V. E. Barnes et al. Observation of a Hyperon with Strangeness -3. *Phys. Rev. Lett.*, 12:204–206, 1964. doi: 10.1103/PhysRevLett.12.204.
- [Weyl(1927)] H. Weyl. Quantum mechanics and group theory. *Z. Phys.*, 46:1, 1927. doi: 10.1007/BF02055756.
- [Michael E. Peskin(1995)] Daniel V. Schroeder Michael E. Peskin. *An Introduction to Quantum Field Theory*. Addison-Wesley Publishing Company, 1 edition, 1995.

Dustin Ray Burns
June 2017
Physics

A Collider Search for Dark Matter Produced in Association
with a Higgs Boson with the CMS Detector at the 13 TeV LHC

Abstract

The abstract that is submitted to UMI must be formatted as shown in the example here. The body of the abstract cannot exceed 350 words. It should be in typewritten form, double-spaced, and on bond paper. It is important to write an abstract that gives a clear description of the content and major divisions of the dissertation, since UMI will publish the abstract exactly as submitted. Students completing their requirements under Plan A should provide extra copies of the typed summary for use by the dissertation committee during the examination.

The abstract that is submitted to UMI must be formatted as shown in the example here. The body of the abstract cannot exceed 350 words. It should be in typewritten form, double-spaced, and on bond paper. It is important to write an abstract that gives a clear description of the content and major divisions of the dissertation, since UMI will publish the abstract exactly as submitted. Students completing their requirements under Plan A should provide extra copies of the typed summary for use by the dissertation committee during the examination.

The abstract that is submitted to UMI must be formatted as shown in the example here. The body of the abstract cannot exceed 350 words. It should be in typewritten form, double-spaced, and on bond paper. It is important to write an abstract that gives a clear description of the content and major divisions of the dissertation, since UMI will publish the abstract exactly as submitted. Students completing their requirements under Plan A should provide extra copies of the typed summary for use by the dissertation committee during the examination.

The abstract that is submitted to UMI must be formatted as shown in the example here. The body of the abstract cannot exceed 350 words. It should be in typewritten

form, double-spaced, and on bond paper. It is important to write an abstract that gives a clear description of the content and major divisions of the dissertation, since UMI will publish the abstract exactly as submitted. Students completing their requirements under Plan A should provide extra copies of the typed summary for use by the dissertation committee during the examination.