

1 Static Semantics

$\boxed{\Theta; \Delta; \Gamma \vdash e : t}$ Typing rules for expressions

$$\frac{}{\Theta; \Delta; \cdot, x : t \vdash x : t} \text{TY_VAR_LIN}$$

$$\frac{x : t \in \Delta}{\Theta; \Delta; \cdot \vdash x : t} \text{TY_VAR}$$

$$\frac{\begin{array}{l} \Theta; \Delta; \Gamma \vdash e : t \\ \Theta; \Delta; \Gamma', x : t \vdash e' : t' \end{array}}{\Theta; \Delta; \Gamma, \Gamma' \vdash \mathbf{let } x = e \mathbf{ in } e' : t'} \text{TY_LET}$$

$$\frac{}{\Theta; \Delta; \cdot \vdash () : \mathbf{unit}} \text{TY_UNIT_INTRO}$$

$$\frac{\begin{array}{l} \Theta; \Delta; \Gamma \vdash e : \mathbf{unit} \\ \Theta; \Delta; \Gamma' \vdash e' : t \end{array}}{\Theta; \Delta; \Gamma, \Gamma' \vdash \mathbf{let } () = e \mathbf{ in } e' : t} \text{TY_UNIT_ELIM}$$

$$\frac{}{\Theta; \Delta; \cdot \vdash \mathbf{true} : \mathbf{bool}} \text{TY_BOOL_TRUE}$$

$$\frac{}{\Theta; \Delta; \cdot \vdash \mathbf{false} : \mathbf{bool}} \text{TY_BOOL_FALSE}$$

$$\frac{\begin{array}{l} \Theta; \Delta; \Gamma \vdash e : !\mathbf{bool} \\ \Theta; \Delta; \Gamma' \vdash e_1 : t' \\ \Theta; \Delta; \Gamma' \vdash e_2 : t' \end{array}}{\Theta; \Delta; \Gamma, \Gamma' \vdash \mathbf{if } e \mathbf{ then } e_1 \mathbf{ else } e_2 : t} \text{TY_BOOL_ELIM}$$

$$\frac{}{\Theta; \Delta; \cdot \vdash k : \mathbf{int}} \text{TY_INT_INTRO}$$

$$\frac{}{\Theta; \Delta; \cdot \vdash el : \mathbf{elt}} \text{TY_ELT_INTRO}$$

$$\frac{\begin{array}{l} \Theta; \Delta; \cdot \vdash v : t \\ v \neq l \cdot f \end{array}}{\Theta; \Delta; \cdot \vdash \mathbf{Many } v : !t} \text{TY_BANG_INTRO}$$

$$\frac{\begin{array}{l} \Theta; \Delta; \Gamma \vdash e : !t \\ \Theta; \Delta, x : t; \Gamma' \vdash e' : t' \end{array}}{\Theta; \Delta; \Gamma, \Gamma' \vdash \mathbf{let Many } x = e \mathbf{ in } e' : t'} \text{TY_BANG_ELIM}$$

$$\frac{\begin{array}{l} \Theta; \Delta; \Gamma \vdash e : t \\ \Theta; \Delta; \Gamma' \vdash e' : t' \end{array}}{\Theta; \Delta; \Gamma, \Gamma' \vdash (e, e') : t \otimes t'} \text{TY_PAIR_INTRO}$$

$$\frac{\begin{array}{l} \Theta; \Delta; \Gamma \vdash e_{12} : t_1 \otimes t_2 \\ \Theta; \Delta; \Gamma', a : t_1, b : t_2 \vdash e : t \end{array}}{\Theta; \Delta; \Gamma, \Gamma' \vdash \mathbf{let } (a, b) = e_{12} \mathbf{ in } e : t} \text{TY_PAIR_ELIM}$$

$$\begin{array}{c}
\frac{\Theta \vdash t' \text{ Type} \quad \Theta; \Delta; \Gamma, x : t' \vdash e : t}{\Theta; \Delta; \Gamma \vdash \mathbf{fun} \, x : t' \rightarrow e : t' \multimap t} \text{ TY_LAMBDA} \\
\\
\frac{\Theta; \Delta; \Gamma \vdash e : t' \multimap t \quad \Theta; \Delta; \Gamma' \vdash e' : t'}{\Theta; \Delta; \Gamma, \Gamma' \vdash e \, e' : t} \text{ TY_APP} \\
\\
\frac{\Theta, fc; \Delta; \Gamma \vdash e : t}{\Theta; \Delta; \Gamma \vdash \mathbf{fun} \, fc \rightarrow e : \forall fc. t} \text{ TY_GEN} \\
\\
\frac{\Theta \vdash f \text{ Cap} \quad \Theta; \Delta; \Gamma \vdash e : \forall fc. t}{\Theta; \Delta; \Gamma \vdash e[f] : t[f/fc]} \text{ TY_SPC} \\
\\
\frac{\Theta; \Delta, g : t \multimap t'; \cdot, x : t \vdash e : t'}{\Theta; \Delta; \cdot \vdash \mathbf{fix} \, (g, x : t, e : t') : t \multimap t'} \text{ TY_FIX}
\end{array}$$

2 Dynamic Semantics

$$\boxed{\langle \sigma, e \rangle \rightarrow \text{StepsTo}} \quad \text{operational semantics}$$

$$\frac{}{\langle \sigma, \mathbf{let} \, () = () \mathbf{in} \, e \rangle \rightarrow \langle \sigma, e \rangle} \text{ OP_LET_UNIT}$$

$$\frac{}{\langle \sigma, \mathbf{let} \, x = v \mathbf{in} \, e \rangle \rightarrow \langle \sigma, e[x/v] \rangle} \text{ OP_LET_VAR}$$

$$\frac{}{\langle \sigma, \mathbf{if} \, (\mathbf{Many} \, \mathbf{true}) \mathbf{then} \, e_1 \mathbf{else} \, e_2 \rangle \rightarrow \langle \sigma, e_1 \rangle} \text{ OP_IF_TRUE}$$

$$\frac{}{\langle \sigma, \mathbf{if} \, (\mathbf{Many} \, \mathbf{false}) \mathbf{then} \, e_1 \mathbf{else} \, e_2 \rangle \rightarrow \langle \sigma, e_2 \rangle} \text{ OP_IF_FALSE}$$

$$\frac{}{\langle \sigma, \mathbf{let} \, \mathbf{Many} \, x = \mathbf{Many} \, v \mathbf{in} \, e \rangle \rightarrow \langle \sigma, e[x/v] \rangle} \text{ OP_LET_MANY}$$

$$\frac{}{\langle \sigma, \mathbf{let} \, (a, b) = (v_1, v_2) \mathbf{in} \, e \rangle \rightarrow \langle \sigma, e[a/v_1][b/v_2] \rangle} \text{ OP_LET_PAIR}$$

$$\frac{}{\langle \sigma, (\mathbf{fun} \, fc \rightarrow v)[f] \rangle \rightarrow \langle \sigma, v[fc/f] \rangle} \text{ OP_FRAC_CAP}$$

$$\frac{}{\langle \sigma, \mathbf{fix} \, (g, x : t, e : t') \, v \rangle \rightarrow \langle \sigma, e[x/v][g/\mathbf{fix} \, (g, x : t, e : t')] \rangle} \text{ OP_APP_FIX}$$

$$\frac{}{\langle \sigma, (\mathbf{fun} \, x : t \rightarrow e) \, v \rangle \rightarrow \langle \sigma, e[x/v] \rangle} \text{ OP_APP_LAMBDA}$$

$$\frac{\langle \sigma, e \rangle \rightarrow \langle \sigma', e' \rangle}{\langle \sigma, C[e] \rangle \rightarrow \langle \sigma, C[e'] \rangle} \text{ OP_CONTEXT}$$

$$\begin{array}{c}
\frac{\langle \sigma, e \rangle \rightarrow \mathbf{err}}{\langle \sigma, C[e] \rangle \rightarrow \mathbf{err}} \quad \text{OP_CONTEXT_ERR} \\
\\
\frac{0 \leq k_1, k_2 \quad l \text{ fresh}}{\langle \sigma, \mathbf{matrix} \ k_1 \ k_2 \rangle \rightarrow \langle \sigma + \{l \mapsto_1 M_{k_1, k_2}\}, l \cdot 1 \rangle} \quad \text{OP_MATRIX} \\
\\
\frac{}{\langle \sigma + \{l \mapsto_1 m_{k_1, k_2}\}, \mathbf{free} \ l \cdot 1 \rangle \rightarrow \langle \sigma, () \rangle} \quad \text{OP_FREE} \\
\\
\frac{}{\langle \sigma + \{l \mapsto_f m_{k_1, k_2}\}, \mathbf{share} \ l \cdot f \rangle \rightarrow \langle \sigma + \{l \mapsto_{\frac{1}{2}f} m_{k_1, k_2}\} + \{l \mapsto_{\frac{1}{2}f} m_{k_1, k_2}\}, (l \cdot \frac{1}{2}f, l \cdot \frac{1}{2}f) \rangle} \quad \text{OP_SHARE} \\
\\
\frac{f \leq 1 \quad v \equiv l \cdot \frac{1}{2}f}{\langle \sigma + \{l \mapsto_{\frac{1}{2}f} m_{k_1, k_2}\} + \{l \mapsto_{\frac{1}{2}f} m_{k_1, k_2}\}, \mathbf{unshare} \ v \ v \rangle \rightarrow \langle \sigma + \{l \mapsto_f m_{k_1, k_2}\}, l \cdot f \rangle} \quad \text{OP_UNSHARE_EQ} \\
\\
\frac{l \neq l'}{\langle \sigma + \{l \mapsto_{\frac{1}{2}f} m_{k_1, k_2}\} + \{l' \mapsto_{\frac{1}{2}f} m'_{k_1, k_2}\}, \mathbf{unshare} \ (l \cdot \frac{1}{2}f) \ (l' \cdot \frac{1}{2}f') \rangle \rightarrow \mathbf{err}} \quad \text{OP_UNSHARE_NEQ} \\
\\
\frac{\begin{array}{l} \sigma' \equiv \sigma + \{l_1 \mapsto_{fc_1} m_{1k_1, k_2}\} + \{l_2 \mapsto_{fc_2} m_{2k_2, k_3}\} \quad v_1 \equiv l_1 \cdot f_1 \quad v_2 \equiv l_2 \cdot f_2 \\ v_3 \equiv l_3 \cdot 1 \end{array}}{\langle \sigma' + \{l_3 \mapsto_1 m_{3k_1, k_3}\}, \mathbf{gemm} \ v_1 \ v_2 \ v_3 \rangle \rightarrow \langle \sigma' + \{l_3 \mapsto_1 (m_1 \ m_2 + m_3)_{k_1, k_3}\}, ((v_1, v_2), v_3) \rangle} \quad \text{OP_GEMM_MATCH} \\
\\
\frac{\begin{array}{l} k_2 \neq k'_2 \\ \sigma' \equiv \sigma + \{l_1 \mapsto_{fc_1} m_{1k_1, k_2}\} + \{l_2 \mapsto_{fc_2} m_{2k'_2, k_3}\} \\ v_1 \equiv l_1 \cdot f_1 \quad v_2 \equiv l_2 \cdot f_2 \quad v_3 \equiv l_3 \cdot 1 \end{array}}{\langle \sigma' + \{l_3 \mapsto_1 m_{1k_1, k_3}\}, \mathbf{gemm} \ v_1 \ v_2 \ v_3 \rangle \rightarrow \mathbf{err}} \quad \text{OP_GEMM_MISMATCH}
\end{array}$$

3 Interpretation

3.1 Definitions

Operationally, $Heap \sqsubseteq Loc \times Permission \times Matrix$ (a multiset), denoted with a σ .

Define its *interpretation* to be $Loc \rightarrow Permission \times Matrix$ with $\star : Heap \times Heap \rightarrow Heap$ as follows:

$$(\varsigma_1 \star \varsigma_2)(l) \equiv \begin{cases} \varsigma_1(l) & \text{if } l \in \text{dom}(\varsigma_1) \wedge l \notin \text{dom}(\varsigma_2) \\ \varsigma_2(l) & \text{if } l \in \text{dom}(\varsigma_2) \wedge l \notin \text{dom}(\varsigma_1) \\ (f_1 + f_2, m) & \text{if } (f_1, m) = \varsigma_1(l) \wedge (f_2, m) = \varsigma_2(l) \wedge f_1 + f_2 \leq 1 \\ \text{undefined} & \text{otherwise} \end{cases}$$

Commutativity and associativity of \star follows from that of $+$.

$\varsigma_1 \star \varsigma_2$ is *defined* if it is for all $l \in \text{dom}(\varsigma_1) \cup \text{dom}(\varsigma_2)$.

Implicitly denote $\varsigma \equiv \mathcal{H}[\sigma] \equiv \star_{(l, f, m) \in \sigma} [l \mapsto_f m]$.

The n -fold iteration for the *StepsTo* (functional) relation, is also a (functional) relation:

$$\forall n. \mathbf{err} \rightarrow^n \mathbf{err} \quad \langle \sigma, v \rangle \rightarrow^n \langle \sigma, v \rangle \quad \langle \sigma, e \rangle \rightarrow^0 \langle \sigma, e \rangle \quad \langle \sigma, e \rangle \rightarrow^{n+1} ((\langle \sigma, e \rangle \rightarrow) \rightarrow^n)$$

Hence, all bounded iterations end in either an **err**, a heap-and-expression or a heap-and-value.

3.2 Interpretation

$$\mathcal{V}_k[\mathbf{unit}] = \{(\emptyset, *)\}$$

$$\mathcal{V}_k[\mathbf{bool}] = \{(\emptyset, true), (\emptyset, false)\}$$

$$\mathcal{V}_k[\mathbf{int}] = \{(\emptyset, n) \mid 2^{-63} \leq n \leq 2^{63} - 1\}$$

$$\mathcal{V}_k[\mathbf{elt}] = \{(\emptyset, f) \mid f \text{ a IEEE Float64 } \}$$

$$\mathcal{V}_k[f \mathbf{mat}] = \{(\{l \mapsto_{2^{-f}} -\}, l)\}$$

$$\mathcal{V}_k[!t] = \{(\emptyset, \mathbf{Many} \ v) \mid (\emptyset, v) \in \mathcal{V}_k[t]\}$$

$$\mathcal{V}_k[\forall fc. t] = \{(\varsigma, \mathbf{fun} \ fc \rightarrow v) \mid \forall f. (\varsigma, (\mathbf{fun} \ fc \rightarrow v)[f]) \in \mathcal{V}_k[t[fc/f]]\}$$

$$\mathcal{V}_k[t_1 \otimes t_2] = \{(\varsigma_1 \star \varsigma_2, (v_1, v_2)) \mid (\varsigma_1, v_1) \in \mathcal{V}_k[t_1] \wedge (\varsigma_2, v_2) \in \mathcal{V}_k[t_2]\}$$

$$\begin{aligned} \mathcal{V}_k[t' \multimap t] = \{(\varsigma_v, v) \mid (v \equiv \mathbf{fun} \ x : t \rightarrow e \vee v \equiv \mathbf{fix}(g, x : t', e : t)) \wedge \\ \forall j \leq k, (\varsigma_{v'}, v') \in \mathcal{V}_j[t']. \varsigma_v \star \varsigma_{v'} \text{ defined} \Rightarrow (\varsigma_v \star \varsigma_{v'}, v \ v') \in \mathcal{C}_j[t]\} \end{aligned}$$

$$\begin{aligned} \mathcal{C}_k[t] = \{(\varsigma_s, e_s) \mid \forall j < k, \sigma_r. \varsigma_s \star \varsigma_r \text{ defined} \Rightarrow \langle \sigma_s + \sigma_r, e_s \rangle \rightarrow^j \mathbf{err} \vee \exists \sigma_f, e_f. \\ \langle \sigma_s + \sigma_r, e_s \rangle \rightarrow^j \langle \sigma_f + \sigma_r, e_f \rangle \wedge (e_f \text{ is a value} \Rightarrow (\varsigma_f \star \varsigma_r, e_f) \in \mathcal{V}_{k-j}[t])\} \end{aligned}$$

$$\mathcal{I}_k[\cdot]\theta = \{\emptyset\}$$

$$\mathcal{I}_k[\Delta, x : t]\theta = \{\delta[x \mapsto v_x] \mid \delta \in \mathcal{I}_k[\Delta]\theta \wedge (\emptyset, v_x) \in \mathcal{V}_k[\theta(t)]\}$$

$$\mathcal{L}_k[\cdot]\theta = \{(\emptyset, [])\}$$

$$\mathcal{L}_k[\Gamma, x : t]\theta = \{(\varsigma \star \varsigma_x, \gamma[x \mapsto v_x]) \mid (\varsigma, \gamma) \in \mathcal{L}_k[\Gamma]\theta \wedge (\varsigma_x, v_x) \in \mathcal{V}_k[\theta(t)]\}$$

$$\varsigma \equiv \mathcal{H}[\sigma] \equiv \star_{(l, f, m) \in \sigma} [l \mapsto_f m]$$

$$\begin{aligned} {}_k[\Theta; \Delta; \Gamma \vdash e : t] = \forall \theta, \delta, \gamma, \sigma. \text{dom}(\Theta) = \text{dom}(\theta) \wedge (\varsigma, \gamma) \in \mathcal{L}_k[\Gamma]\theta \wedge \delta \in \mathcal{I}_k[\Delta]\theta \Rightarrow \\ (\varsigma, \gamma(\delta(e))) \in \mathcal{C}_k[\theta(t)] \end{aligned}$$

4 Proofs

4.1 Lemmas

4.1.1 $\forall \sigma_s, \sigma_r, e. \varsigma_s \star \varsigma_r \text{ defined} \Rightarrow \forall n. \langle \sigma_s, e \rangle \rightarrow^n \langle \sigma_s + \sigma_r, e \rangle \rightarrow^n$

SUFFICES: By induction on n , consider only the cases $\langle \sigma_s, e \rangle \rightarrow \langle \sigma_f, e_f \rangle$ where $\sigma_s \neq \sigma_f$.

PROOF SKETCH: Only `OP_FREE`, `MATRIX`, `SHARE`, `UNSHARE_EQ`, `GEMM_MATCH` change the heap: the rest are either parametric in the heap or step to an **err**.

PROVE: $\langle \sigma_s + \sigma_r, e \rangle \rightarrow \langle \sigma_f + \sigma_r, e_f \rangle$.

$\langle 1 \rangle 1$. CASE: `OP_FREE`, $\sigma_s \equiv \sigma' + \{l \mapsto_1 m\}$, $\sigma_f = \sigma'$.

PROOF: Instantiate `OP_FREE` with $(\sigma' + \sigma_r) + \{l \mapsto_1 m\}$,
valid because $l \notin \text{dom}(\varsigma_r)$ by $\varsigma' \star [l \mapsto_1 m] \star \varsigma_r$ defined (assumption).

$\langle 1 \rangle 2$. CASE: `OP_MATRIX`

PROOF: Rule has no requirements on σ_s so will also work with $\sigma_s + \sigma_r$.

$\langle 1 \rangle 3$. CASE: `OP_SHARE`, $\sigma_s \equiv \sigma' + \{l \mapsto_f m\}$, $\sigma_f = \sigma' + \{l \mapsto_{\frac{1}{2}.f} m\} + \{l \mapsto_{\frac{1}{2}.f} m\}$.

PROOF: Union-ing σ_r does not remove $l \mapsto_f m$, so that can be split out of $\sigma_s + \sigma_r$ as before.

$\langle 1 \rangle 4$. CASE: `OP_UNSHARE_EQ`, $\sigma_s \equiv \sigma' + \{l \mapsto_{\frac{1}{2}.f} m\} + \{l \mapsto_{\frac{1}{2}.f} m\}$, $\sigma_f = \sigma' + \{l \mapsto_f m\}$.

$\langle 2 \rangle 1$. Union-ing σ_r does not remove $l \mapsto_{\frac{1}{2}.f} m$, so that can still be split out of $\sigma_s + \sigma_r$.

$\langle 2 \rangle 2$. There may also be other valid splits introduced by σ_r .

$\langle 2 \rangle 3$. However, by assumption of $\varsigma_s \star \varsigma_r$ defined, any splitting of $\sigma_s + \sigma_r$ will satisfy $f \leq 1$.

$\langle 1 \rangle 5$. CASE: `OP_GEMM_MATCH`

$\langle 2 \rangle 1$. By assumption of $\varsigma_s \star \varsigma_r$ defined, either l_1 (or l_2 , or both) are not in σ_r , or they are and the matrix values they point to are the same.

$\langle 2 \rangle 2$. The permissions (of l_1 and/or l_2) may differ, but `OP_GEMM_MATCH` universally quantifies over them and leaves them unchanged, so they are irrelevant.

$\langle 2 \rangle 3$. Only the pointed to matrix value at l_3 changes.

$\langle 2 \rangle 4$. SUFFICES: $l_3 \notin \pi_1[\sigma_r]$.

$\langle 2 \rangle 5$. By assumption of $\varsigma_s \star \varsigma_r$ defined, $l_3 \notin \text{dom}(\varsigma_r)$.

$\langle 2 \rangle 6$. Hence $l_3 \notin \pi_1[\sigma_r]$.

4.1.2 $\forall k, t. \mathcal{V}_k[t] \subseteq \mathcal{C}_k[t]$

Follows from definition of $\mathcal{C}_k[t]$, \rightarrow^j ($\forall n. \langle \sigma, v \rangle \rightarrow^n \langle \sigma, v \rangle$) for arbitrary $j \leq k$ and 4.1.1.

4.1.3 $\forall \delta, \gamma, v. \delta(\gamma(v))$ is a value.

By construction, δ and γ only map variables to values, and values are closed under substitution.

4.1.4 $\forall k, \sigma, \sigma', e, e', t. (\varsigma', e') \in \mathcal{C}_k[t] \wedge \langle \sigma, e \rangle \rightarrow \langle \sigma', e' \rangle \Rightarrow (\varsigma, e) \in \mathcal{C}_{k+1}[t]$

ASSUME: arbitrary $j < k + 1$, and σ_r such that $\varsigma \star \varsigma_r$ defined.

$\langle 1 \rangle 1$. CASE: $j = 0$. Clearly $\sigma_f = \sigma_s + \sigma_r$ and $e' = e$.

Remains to show that if e is a value then $(\varsigma_s \star \varsigma_r, e) \in \mathcal{V}_k[t]$.

This is true vacuously, because by assumption, e is not a value.

$\langle 1 \rangle 2$. CASE: $j \geq 1$. We have $\langle \sigma, e \rangle \rightarrow^j = \langle \sigma', e' \rangle \rightarrow^{j-1}$.

Instantiate $(\varsigma', e') \in \mathcal{C}_k[t]$, with $j - 1 < k$ and σ_r to conclude the required conditions.

4.1.5 $j \leq k \Rightarrow _k[\cdot] \subseteq _j[\cdot]$

Lemma 4.1.4 is the inductive step for this lemma for the $\mathcal{C}[\cdot]$ case.

Need to prove for $\mathcal{V}[\cdot]$, by induction on t and then index.

SUFFICES: Consider only $t \multimap t'$ case, rest use k directly on structure of type.

ASSUME: Arbitrary $j \leq k$ and $(\varsigma_{v'}, v') \in \mathcal{V}_k[t \multimap t']$.

PROVE: $(\varsigma_{v'}, v') \in \mathcal{V}_j[t \multimap t']$.

$\langle 1 \rangle 1$. v' is of the correct syntactic form (lambda or fixpoint) by assumption.

$\langle 1 \rangle 2$. ASSUME: arbitrary $j' \leq j$ and $(\varsigma_v, v) \in \mathcal{V}_{j'}[t]$ such that $\varsigma_{v'} \star \varsigma_v$ is defined.

$\langle 1 \rangle 3$. SUFFICES: to show $(\varsigma_{v'} \star \varsigma_v, v'v) \in \mathcal{C}_{j'}[t']$.

$\langle 1 \rangle 4$. This is true by instantiating $(\varsigma_{v'}, v') \in \mathcal{V}_k[t \multimap t']$ with $j' \leq k$ and $(\varsigma_v, v) \in \mathcal{V}_{j'}[t]$.

4.1.6 $\forall \Delta, \Gamma, t, k, \theta, \delta, \gamma. \delta \in \mathcal{I}_k[\Delta]\theta \wedge \gamma \in \pi_2[\mathcal{L}_k[\Gamma]\theta] \Rightarrow \text{dom}(\Delta) = \text{dom}(\delta) \text{ and } \text{dom}(\Gamma) = \text{dom}(\gamma)$

PROOF: By induction on Δ and Γ .

4.1.7 $\forall k, \Gamma, \Gamma', \theta, \sigma_+, \gamma_+. (\varsigma_+, \gamma) \in \mathcal{L}_k[\Gamma, \Gamma']\theta \wedge \Gamma, \Gamma' \text{ disjoint} \Rightarrow$
 $\exists \sigma, \gamma, \sigma', \gamma'. \sigma_+ = \sigma + \sigma' \wedge \gamma, \gamma' \text{ disjoint} \wedge \gamma_+ = \gamma \cup \gamma'$
 $\wedge (\varsigma, \gamma) \in \mathcal{L}_k[\Gamma] \wedge (\varsigma', \gamma') \in \mathcal{L}_k[\Gamma']$

PROOF: By induction on Γ' .

4.2 Soundness

$$\forall \Theta, \Delta, \Gamma, e, t. \Theta; \Delta; \Gamma \vdash e : t \Rightarrow \forall k. {}_k\llbracket \Theta; \Delta; \Gamma \vdash e : t \rrbracket$$

PROOF SKETCH: Induction over the typing judgements.

ASSUME: 1. Arbitrary $\Theta, \Delta, \Gamma, e, t$ such that $\Theta; \Delta; \Gamma \vdash e : t$.

2. Arbitrary $k, \theta, \delta, \gamma, \sigma$ such that:

- a. $\text{dom}(\Theta) = \text{dom}(\theta)$
- b. $(\varsigma, \gamma) \in \mathcal{L}_k\llbracket \Gamma \rrbracket \theta$
- c. $\delta \in \mathcal{I}_k\llbracket \Delta \rrbracket \theta$.

3. W.l.o.g., all variables are distinct,

hence $\text{dom}(\Delta)$ and $\text{dom}(\Gamma)$ are disjoint

so $\gamma \circ \delta = \delta \circ \gamma$ (as substitutions defined recursively over expressions).

PROVE: $(\varsigma, \gamma(\delta(e))) \in \mathcal{C}_k\llbracket \theta(t) \rrbracket$.

ASSUME: Arbitrary $j < k$ and σ_r , such that $\varsigma \star \varsigma_r$ defined.

SUFFICES: $\langle \sigma + \sigma_r, e \rangle \rightarrow^j \mathbf{err} \vee \exists \sigma_f, e_f. \langle \sigma + \sigma_r, e \rangle \rightarrow^j \langle \sigma_f + \sigma_r, e_f \rangle$
 $\wedge (e_f \text{ is a value} \Rightarrow (\varsigma_f \star \varsigma_r, e_f) \in \mathcal{V}_{k-j}\llbracket t \rrbracket)$.

SUFFICES: By 4.1.1, to show $\langle \sigma, e \rangle \rightarrow^j \mathbf{err} \vee \exists \sigma_f, e_f. \langle \sigma, e \rangle \rightarrow^j \langle \sigma_f, e_f \rangle$
 $\wedge (e_f \text{ is a value} \Rightarrow (\varsigma_f, e_f) \in \mathcal{V}_{k-j}\llbracket t \rrbracket)$

$\langle 1 \rangle 1$. CASE: **TY-LET**.

$\langle 2 \rangle 1$. By induction,

- 1. $\forall k. {}_k\llbracket \Theta; \Delta; \Gamma \vdash e : t \rrbracket$
- 2. $\forall k. {}_k\llbracket \Theta; \Delta; \Gamma', x : t \vdash e' : t' \rrbracket$.

$\langle 2 \rangle 2$. By 2b, 3 and 4.1.7, we know there exists the following (for all k):

- 1. $(\varsigma_e, \gamma_e) \in \mathcal{L}_k\llbracket \Gamma \rrbracket$
- 2. $\gamma = \gamma_e \cup \gamma_{e'}$
- 3. $\sigma = \sigma_e + \sigma_{e'}$.

$\langle 2 \rangle 3$. So, using $k, \theta, \delta, \gamma_e, \sigma_e$, we have $(\varsigma_e, \gamma_e(\delta(e))) \in \mathcal{C}_k\llbracket \theta(t) \rrbracket$.

$\langle 2 \rangle 4$. By $\langle 2 \rangle 2$ ($\gamma = \gamma_e \cup \gamma_{e'}$), have $(\varsigma_e, \gamma(\delta(e))) \in \mathcal{C}_k\llbracket \theta(t) \rrbracket$.

$\langle 2 \rangle 5$. By definition of $\mathcal{C}_k\llbracket \cdot \rrbracket$ and $\langle 2 \rangle 2$, we instantiate with j and $\sigma_r = \sigma_{e'}$ to conclude that $\langle \varsigma, \gamma(\delta(e)) \rangle$ either takes j steps to **err** or another heap-and-expression $\langle \sigma_f, \gamma(\delta(e_f)) \rangle$.

$\langle 2 \rangle 6$. CASE: j steps to **err**

By **OP-CONTEXT-ERR**, the whole expression reduces to **err** in $j < k$ steps.

$\langle 2 \rangle 7$. CASE: j steps to another heap-and-expression.

If it is not a value, then **OP-CONTEXT** runs j times and we are done.

$\langle 2 \rangle 8$. If it is, then $\exists i \leq j. (\varsigma_f, v_1) \in \mathcal{V}_{k-i}\llbracket \theta(t_1) \rrbracket \subseteq \mathcal{V}_{k-j}\llbracket \theta(t_1) \rrbracket$ by 4.1.3 and 4.1.5.

So, **OP-CONTEXT** runs i times, and then we have the following.

SUFFICES: $(\varsigma_f \star \varsigma_{e'}, \mathbf{let } x = v \mathbf{ in } \gamma(\delta(e'))) \in \mathcal{C}_{k-i}\llbracket \theta(t') \rrbracket$ by 4.1.4 i times.

SUFFICES: $(\varsigma_f \star \varsigma_{e'}, \gamma(\delta(e'))[x/v]) \in \mathcal{C}_{k-i-1}\llbracket \theta(t') \rrbracket$ by 4.1.4.

$\langle 2 \rangle 9$. By 4.1.5, $(\varsigma_{e'}, \gamma_{e'}[x \mapsto v]) \in \mathcal{L}_k\llbracket \Gamma', x : t \rrbracket \theta \subseteq \mathcal{L}_{k-i-1}\llbracket \Gamma', x : t \rrbracket \theta$.

$\langle 2 \rangle 10$. Instantiate 2 of step $\langle 2 \rangle 1$ with $k - i - 1, \theta, \delta, \gamma_{e'}[x \mapsto v], \sigma_{e'}$ to conclude

$$(\varsigma_{e'}, \gamma_{e'}[x \mapsto v](\delta(e'))) \in \mathcal{C}_{k-i-1}[\![\theta(t')]\!].$$

$\langle 2 \rangle 11$. By 3, we have $\gamma(\delta(e'))[x/v] = \gamma_{e'}[x \mapsto v](\delta(e'))$ and by 4.1.1 we conclude $(\varsigma_f \star \varsigma_{e'}, \gamma(\delta(e'))[x/v]) \in \mathcal{C}_{k-i-1}[\![\theta(t')]\!]$

$\langle 1 \rangle 2$. CASE: `TY_PAIR_ELIM`.

PROOF SKETCH: Similar to `TY_LET`, but with the following key differences.

$\langle 2 \rangle 1$. When $(\varsigma_f, v) \in \mathcal{V}_{k-i}[\![\theta(t_1) \otimes \theta(t_2)]\!]$, we have $v = (v_1, v_2)$.

$\langle 2 \rangle 2$. SUFFICES: $(\varsigma_{e'}, \gamma(\delta(e'))) \in \mathcal{C}_{k-i-1}[\![\theta(t')]\!]$ by 4.1.4 $i + 1$ times.

$\langle 2 \rangle 3$. By 4.1.5, $(\varsigma_{e'}, \gamma_{e'}[a \mapsto v_1, b \mapsto v_2]) \in \mathcal{L}_k[\![\Gamma', a : t_1, b : t_2]\!]\theta \subseteq \mathcal{L}_{k-i-1}[\![\Gamma', a : t_1, b : t_2]\!]\theta$.

$\langle 2 \rangle 4$. Instantiate $_{k-i-1}[\![\Theta; \Delta; \Gamma', a : t_1, b : t_2 \vdash e' : t']\!]$ with $\theta, \delta, \gamma_{e'}[a \mapsto v_1, b \mapsto v_2], \sigma_{e'}$.

$\langle 2 \rangle 5$. By 3 (for $\gamma = \gamma_e \cup \gamma_{e'}$ and a, b), conclude $(\varsigma_{e'}, \gamma(\delta(e'[a/v_1][b/v_2]))) \in \mathcal{C}_{k-i-1}[\![\theta(t')]\!]$.

$\langle 1 \rangle 3$. CASE: `TY_BANG_ELIM`.

PROOF SKETCH: Similar to `TY_LET`, but with the following key differences.

$\langle 2 \rangle 1$. When $(\varsigma_f, v) \in \mathcal{V}_{k-i}[\![\theta(!t)]\!]$, since $\mathcal{V}_{k-i}[\![\theta(!t)]\!] = \mathcal{V}_{k-i}[\![\theta(t)]\!]$, we have $\varsigma_f = \emptyset$ and $v = \mathbf{Many} \ v'$ for some $(\emptyset, v') \in \mathcal{V}_{k-i}[\![\theta(t)]\!]$.

$\langle 2 \rangle 2$. SUFFICES: $(\varsigma_{e'}, \mathbf{let} \ \mathbf{Many} \ x = \mathbf{Many} \ v' \ \mathbf{in} \ \gamma(\delta(e'))) \in \mathcal{C}_{k-i}[\![\theta(t)]\!]$.

$\langle 2 \rangle 3$. SUFFICES: $(\varsigma_{e'}, \gamma(\delta(e'))[x/v]) \in \mathcal{C}_{k-i-1}[\![\theta(t)]\!]$ by 4.1.4 $i + 1$ times.

$\langle 2 \rangle 4$. Instantiate $_{k-i-1}[\![\Theta; \Delta, x : t, \Gamma' \vdash e' : t']\!]$ with $\theta, \delta_{e'} = \delta[x \mapsto v'], \gamma_{e'}, \sigma_{e'}$.

$\langle 2 \rangle 5$. By 3, $(\varsigma_{e'}, \gamma(\delta(e'))[x/v]) \in \mathcal{C}_{k-i-1}[\![\theta(t)]\!]$.

$\langle 1 \rangle 4$. CASE: `TY_UNIT_ELIM`.

PROOF SKETCH: Similar to `TY_LET`, but with the following key differences.

$\langle 2 \rangle 1$. When $(\varsigma_f, v) \in \mathcal{V}_{k-i}[\![\mathbf{unit}]\!]$, we have $\varsigma_f = \emptyset$ and $v = ()$.

$\langle 2 \rangle 2$. SUFFICES: $(\varsigma_{e'}, \gamma(\delta(e'))) \in \mathcal{C}_{k-i-1}[\![\theta(t')]\!]$ by 4.1.4 $i + 1$ times.

$\langle 2 \rangle 3$. By 4.1.5, $(\varsigma_{e'}, \gamma_{e'}) \in \mathcal{L}_k[\![\Gamma']\!]\theta \subseteq \mathcal{L}_{k-i-1}[\![\Gamma']\!]\theta$.

$\langle 2 \rangle 4$. Instantiate $_{k-i-1}[\![\Theta; \Delta; \Gamma' \vdash e' : t']\!]$ with $\theta, \delta, \gamma_{e'}, \sigma_{e'}$.

$\langle 2 \rangle 5$. By 3 $(\varsigma_{e'}, \gamma(\delta(e'))) \in \mathcal{C}_{k-i-1}[\![\theta(t')]\!]$.

$\langle 1 \rangle 5$. CASE: `TY_BOOL_ELIM`.

PROOF SKETCH: Similar to `TY_UNIT_ELIM` but with `OP_IF_{-}\{\mathbf{TRUE}, \mathbf{FALSE}\}`, $\varsigma_f = \emptyset$ and $v = \mathbf{Many} \ \mathbf{true}$ or $v = \mathbf{Many} \ \mathbf{false}$.

$\langle 1 \rangle 6$. CASE: `TY_BANG_INTRO`.

$\langle 2 \rangle 1$. We have, $e = v$ for some value $v \neq l \cdot f$, $\Gamma = \emptyset$ and so $\forall k. \ _k[\![\Theta; \Delta; \cdot \vdash v : t]\!]$ by induction.

$\langle 2 \rangle 2$. SUFFICES: $(\emptyset, \mathbf{Many} \ \delta(v)) \in \mathcal{C}_k[\![\theta(t)]\!]$ by 2b ($\varsigma = \emptyset, \gamma = []$).

$\langle 2 \rangle 3$. Instantiate $_k[\![\Theta; \Delta; \cdot \vdash v : t]\!]$ with $\theta, \delta, \gamma = [], \sigma = \emptyset$ to obtain $(\emptyset, \delta(v)) \in \mathcal{C}_k[\![\theta(t)]\!]$.

- ⟨2⟩4. Instantiate $(\emptyset, \delta(v)) \in \mathcal{C}_k[\![\theta(t)]\!]$ with $j = 0$, $\sigma_r = \emptyset$ and 4.1.3 ($\delta(v)$ is a value), to conclude $(\emptyset, \delta(v)) \in \mathcal{V}_k[\![\theta(t)]\!]$.
- ⟨2⟩5. By definition of $\mathcal{V}_k[\![\theta(t)]\!]$, 4.1.3 and 4.1.2 we have $(\emptyset, \mathbf{Many} \delta(v)) \in \mathcal{C}_k[\![\theta(t)]\!]$.
- ⟨1⟩7. CASE: `TY_PAIR_INTRO`.
- ⟨2⟩1. By 2b, 3 and 4.1.7, we know there exists the following (for all k):
1. $(\varsigma_1, \gamma_1) \in \mathcal{L}_k[\![\Gamma_1]\!]$
 2. $(\varsigma_2, \gamma_2) \in \mathcal{L}_k[\![\Gamma_2]\!]$
 3. $\gamma = \gamma_1 \cup \gamma_2$
 4. $\sigma = \sigma_1 + \sigma_2$.
- ⟨2⟩2. By induction,
1. $\forall k. {}_k[\![\Theta; \Delta; \Gamma_1 \vdash e_1 : t_1]\!]$
 2. $\forall k. {}_k[\![\Theta; \Delta; \Gamma_2 \vdash e_2 : t_2]\!]$.
- ⟨2⟩3. Instantiate the first with $k, \theta, \delta, \gamma_1, \sigma_1$.
- ⟨2⟩4. By that and ⟨2⟩1, $(\varsigma_1, \gamma_1(\delta(e_1))) = (\varsigma_1, \gamma(\delta(e_1))) \in \mathcal{C}_k[\![\theta(t)]\!]$.
- ⟨2⟩5. So, $\langle \sigma_1 + \sigma_2, \gamma_1(\delta(e_1)) \rangle$ either takes j steps to **err** or a heap-and-expression $\langle \sigma_{1f}, e_{1f} \rangle$.
- ⟨2⟩6. CASE: j steps to **err**
By `OP_CONTEXT_ERR`, the whole expression reduces to **err** in $j < k$ steps.
- ⟨2⟩7. CASE: j steps to another heap-and-expression.
If it is not a value, then `OP_CONTEXT` runs j times and we are done.
- ⟨2⟩8. If it is, then $\exists i_1 \leq j. (\varsigma_{1f}, v_1) \in \mathcal{V}_{k-i_1}[\![\theta(t_1)]\!] \subseteq \mathcal{V}_{k-j}[\![\theta(t_1)]\!]$ by 4.1.3 and 4.1.5.
So, `OP_CONTEXT` runs i_1 times, and then we have the following.
SUFFICES: By 4.1.4, $(\varsigma_{1f} \star \varsigma_2, (v_1, e_2)) \in \mathcal{C}_{k-i_1}[\![\theta(t_1 \otimes t_2)]\!]$.
- ⟨2⟩9. Instantiate the second IH with $k, \theta, \delta, \gamma_2, \sigma_2$.
- ⟨2⟩10. So, $\langle \sigma_{1f} \star \sigma_2, \gamma_2(\delta(e_2)) \rangle$ either takes j steps to **err** or a heap-and-expression $\langle \sigma_{2f}, e_{2f} \rangle$.
- ⟨2⟩11. CASE: j steps to **err**
By `OP_CONTEXT_ERR`, the whole expression reduces to **err** in $j < k$ steps.
- ⟨2⟩12. CASE: j steps to another heap-and-expression.
If it is not a value, then `OP_CONTEXT` runs j times and we are done.
- ⟨2⟩13. If it is, then $\exists i_2 \leq j. (\varsigma_{2f}, v_2) \in \mathcal{V}_{k-i_2}[\![\theta(t_2)]\!] \subseteq \mathcal{V}_{k-j}[\![\theta(t_2)]\!]$ by 4.1.3 and 4.1.5.
So, `OP_CONTEXT` runs i_2 times, and then we have the following.
SUFFICES: By 4.1.4, $(\varsigma_{1f} \star \varsigma_{2f}, (v_1, v_2)) \in \mathcal{V}_{k-i_1-i_2}[\![\theta(t_1) \otimes \theta(t_2)]\!]$.
- ⟨2⟩14. By 4.1.5 and $k - i_1 - i_2 \leq k - i_1, k - i_2$, have
 $(\varsigma_{1f}, v_1) \in \mathcal{V}_{k-i_1}[\![\theta(t_1)]\!] \subseteq \mathcal{V}_{k-i_1-i_2}[\![\theta(t_1)]\!]$ and
 $(\varsigma_{2f}, v_2) \in \mathcal{V}_{k-i_2}[\![\theta(t_2)]\!] \subseteq \mathcal{V}_{k-i_1-i_2}[\![\theta(t_2)]\!]$ as needed.
- ⟨1⟩8. CASE: `TY_LAMBDA`.
SUFFICES: By 4.1.2, to show $(\varsigma, \gamma(\delta(\mathbf{fun} x : t \rightarrow e))) \in \mathcal{V}_k[\![\theta(t \multimap t')]\!]$.
ASSUME: Arbitrary $j \leq k$, $(\varsigma_v, v) \in \mathcal{V}_j[\![\theta(t)]\!]$ such that $\varsigma \star \varsigma_v$ is defined.
SUFFICES: $(\varsigma \star \varsigma_v, \gamma(\delta(\mathbf{fun} x : t \rightarrow e)) v) \in \mathcal{C}_j[\![\theta(t')]\!]$.
SUFFICES: $(\varsigma \star \varsigma_v, \gamma(\delta(e))[x/v]) \in \mathcal{C}_{j-1}[\![\theta(t')]\!]$ by 4.1.4.

- ⟨2⟩1. By induction, $\forall k. {}_k\llbracket \Theta; \Delta; \Gamma, x : t \vdash e \rrbracket$.
- ⟨2⟩2. Instantiate it $j - 1, \theta, \gamma[x \mapsto v], \sigma + \sigma_v$.
- ⟨2⟩3. Hence, $(\varsigma \star \varsigma_v, \gamma[x \mapsto v](\delta(e))) \in \mathcal{C}_{j-1}\llbracket \theta(t) \rrbracket$.
- ⟨2⟩4. By 3, $\gamma[x \mapsto v](\delta(e)) = \gamma(\delta(e))[x/v]$, we are done.
- ⟨1⟩9. CASE: TY_APP.
- ⟨2⟩1. By 2b, 3 and 4.1.7, we know there exists the following (for all k):
1. $(\varsigma_e, \gamma_e) \in \mathcal{L}_k\llbracket \Gamma_e \rrbracket$
 2. $(\varsigma_{e'}, \gamma_{e'}) \in \mathcal{L}_k\llbracket \Gamma_{e'} \rrbracket$
 3. $\gamma = \gamma_e \cup \gamma_{e'}$
 4. $\sigma = \sigma_e + \sigma_{e'}$.
- ⟨2⟩2. By induction,
1. $\forall k. {}_k\llbracket \Theta; \Delta; \Gamma \vdash e : t' \multimap t \rrbracket$
 2. $\forall k. {}_k\llbracket \Theta; \Delta; \Gamma' \vdash e' : t' \rrbracket$.
- ⟨2⟩3. Instantiate the first with $\theta, k, \delta, \gamma_e, \sigma_e$ to conclude $(\varsigma_e, \gamma_e(\delta(e))) \in \mathcal{C}_k\llbracket \theta(t') \multimap \theta(t) \rrbracket$.
- ⟨2⟩4. Instantiate *this* with j and $\sigma_{e'}$ and use ⟨2⟩1 to conclude $\langle \sigma_e + \sigma_{e'}, \gamma(\delta(e)) \rangle$ either takes j steps to **err** or a heap-and-expression $\langle \sigma_f + \sigma_{e'}, e_f \rangle$.
- ⟨2⟩5. CASE: j steps to **err**
By OP_CONTEXT_ERR, the whole expression reduces to **err** in $j < k$ steps.
- ⟨2⟩6. CASE: j steps to another heap-and-expression.
If it is not a value, then OP_CONTEXT runs j times and we are done.
- ⟨2⟩7. If it is, then $\exists i_e \leq j. (\varsigma_f, v_e) \in \mathcal{V}_{k-i_e}\llbracket \theta(t') \multimap \theta(t) \rrbracket \subseteq \mathcal{V}_{k-j}\llbracket \dots \rrbracket$ by 4.1.3 and 4.1.5.
So, OP_CONTEXT runs i_e times, and then we have the following.
SUFFICES: By 4.1.4 i_e times, $(\varsigma_f \star \varsigma_{e'}, v_e e') \in \mathcal{C}_{k-i_e}\llbracket \theta(t') \rrbracket$.
- ⟨2⟩8. By 4.1.5, $(\varsigma_{e'}, \gamma_{e'} \in \mathcal{L}_k\llbracket \Gamma' \rrbracket \theta \subseteq \mathcal{L}_{k-i_e}\llbracket \Gamma' \rrbracket \theta$.
- ⟨2⟩9. So, instantiate the second IH with $k - i_e, \theta, \delta, \gamma_{e'}, \sigma_{e'}$ to conclude $(\varsigma_{e'}, \gamma_{e'}(\delta(e'))) \in \mathcal{C}_{k-i_e}\llbracket \theta(t') \rrbracket$.
- ⟨2⟩10. Instantiate *this* with $j - i_e$ and σ_f to conclude $\langle \sigma_f + \sigma_{e'}, \gamma_{e'}(\delta(e')) \rangle$ either takes $j - i_e$ steps to **err** or $\langle \sigma_f + \sigma_{f'}, e_{f'} \rangle$.
- ⟨2⟩11. CASE: $j - i_e$ steps to **err**
By OP_CONTEXT_ERR, the whole expression reduces to **err** in $j - i_e < k - i_e$ steps.
- ⟨2⟩12. CASE: $j - i_e$ steps to another heap-and-expression.
If it is not a value, then OP_CONTEXT runs $j - i_e$ times and we are done.
- ⟨2⟩13. If it is, then $\exists i_{e'} \leq j - i_e. (\varsigma_{f'}, v_{e'}) \in \mathcal{V}_{k-i_e-i_{e'}}\llbracket \theta(t') \rrbracket$ by 4.1.3.
So, OP_CONTEXT runs $i_{e'}$ times, and then we have the following.
SUFFICES: By 4.1.4 $i_{e'}$ times, $(\varsigma_f \star \varsigma_{f'}, v_e v_{e'}) \in \mathcal{C}_{k-i_e-i_{e'}}\llbracket \theta(t') \rrbracket$.
- ⟨2⟩14. Instantiate $(\varsigma_f, v_e) \in \mathcal{V}_{k-i_e}\llbracket \theta(t') \multimap \theta(t) \rrbracket$ with $k - i_e - i_{e'} \leq k - i_e$ and $(\varsigma_{v'}, v_{e'}) \in \mathcal{V}_{k-i_e-i_{e'}}\llbracket \theta(t') \rrbracket$, to conclude $(\varsigma_f \star \varsigma_{f'}, v_e v_{e'}) \in \mathcal{C}_{k-i_e-i_{e'}}\llbracket \theta(t) \rrbracket$ as needed.

⟨1⟩10. CASE: TY_GEN.

PROVE: $(\sigma, \gamma(\delta(\mathbf{fun} \, fc \rightarrow e))) \in \mathcal{C}_k[\![\theta(\forall \, fc. t)]\!]$.

⟨1⟩11. CASE: TY_SPC.

PROVE: $(\sigma, \gamma(\delta(e[f]))) \in \mathcal{C}_k[\![\theta(t[f c/f])]\!]$.

⟨1⟩12. CASE: TY_FIX.

PROVE: $(\sigma, \gamma(\delta(\mathbf{fix}(g, x : t, e : t')))) \in \mathcal{C}_k[\![\theta(! (t \multimap t'))]\!]$.

SUFFICES: to show $\dots \in \mathcal{V}_k[\![!(\theta(t) \multimap \theta(t'))]\!]$, by 4.1.2.

⟨2⟩1. ASSUME: Arbitrary $j < k$ and $(\sigma, v) \in \mathcal{V}_j[\![\theta(t)]\!]$.

⟨2⟩2. SUFFICES: $(\sigma, \mathit{letManyG} \, g \, v) \in \mathcal{C}_j[\![\theta(t')]\!]$.

⟨2⟩3. LET: $e_1 = e[g/\mathbf{fun} \, x : t \rightarrow \mathit{letManyG} \, g \, x]$.

⟨2⟩4. SUFFICES: by 4.1.4, $(\sigma, (\mathbf{fun} \, x : t \rightarrow e_1) \, v) \in \mathcal{C}_{j-1}[\![\theta(t')]\!]$.

⟨2⟩5. SUFFICES: by 4.1.4, $(\sigma, e_1[x/v]) \in \mathcal{C}_{j-2}[\![\theta(t')]\!]$.

⟨2⟩6. By induction, we have $\llbracket \Theta; \Delta, g : t \multimap t'; x : t \vdash e : t' \rrbracket$.

⟨2⟩7. Instantiate this with $\theta, j-2, \delta[g \mapsto \mathbf{fun} \, x : t \rightarrow e_1], \gamma = [x \mapsto v], \sigma$ (???).

PROVE: $(\sigma, \mathbf{fun} \, x : t \rightarrow e_1) \in \mathcal{V}_{j-2}[\![\theta(t) \multimap \theta(t')]\!]$.

⟨3⟩1. SUFFICES: by 4.1.4, $(\sigma', e_1[x/v']) \in \mathcal{C}_{j-2}[\![\theta(t')]\!]$ for arbitrary $(\sigma', v') \in \mathcal{V}_{j-2}[\![\theta(t)]\!]$.

⟨3⟩2. We can again use the induction hypothesis $\llbracket \Theta; \Delta, g : t \multimap t'; x : t \vdash e : t' \rrbracket$.

⟨3⟩3. But since it's true for $\mathcal{C}_0[\![\cdot]\!]$ (base case), it's true by induction ???

⟨2⟩8. Lastly, we show $\delta(\gamma(e)) = e_1[x/v]$, which follows by their definitions, to conclude $(\sigma, e_1[x/v]) \in \mathcal{C}_{j-2}[\![\theta(t')]\!]$.

⟨1⟩13. CASE: TY_VAR_LIN.

PROVE: $(\sigma, \gamma(\delta(x))) \in \mathcal{C}_k[\![\theta(t)]\!]$.

⟨2⟩1. $\Gamma = \{x : t\}$ by assumption of TY_VAR_LIN.

⟨2⟩2. SUFFICES: $(\sigma, \gamma(x)) \in \mathcal{C}_k[\![\theta(t)]\!]$ by 3.

⟨2⟩3. By 2b, there exist $(\sigma_x, v_x) \in \mathcal{V}_k[\![\theta(t)]\!]$, such that $\sigma = \sigma_x$ and $\gamma = [x \mapsto v_x]$.

⟨2⟩4. Hence, $(\sigma_x, v_x) \in \mathcal{C}_k[\![\theta(t)]\!]$, by 4.1.2.

⟨1⟩14. CASE: TY_VAR.

PROVE: $(\sigma, \gamma(\delta(x))) \in \mathcal{C}_k[\![\theta(t)]\!]$.

⟨2⟩1. $x : t \in \Delta$ and $\Gamma = \emptyset$ by assumption of TY_VAR.

⟨2⟩2. SUFFICES: $(\emptyset, \delta(x)) \in \mathcal{C}_k[\![\theta(t)]\!]$ by 3 and 2b.

⟨2⟩3. By 2c, there exists v_x such that $(\emptyset, v_x) \in \mathcal{V}_k[\![\theta(t)]\!]$.

$\langle 2 \rangle 4$. Hence, $(\emptyset, v_x) \in \mathcal{C}_k[\![\theta(t)]\!]$, by 4.1.2.

$\langle 1 \rangle 15$. CASE: `TY_UNIT_INTRO`.

PROVE: $(\sigma, \gamma(\delta(\ () \))) \in \mathcal{C}_k[\![\theta(\mathbf{unit})]\!]$.

$\langle 1 \rangle 16$. CASE: `TY_BOOL_TRUE`, `TY_BOOL_FALSE`, `TY_INT_INTRO`, `TY_ELT_INTRO`.

Similar to `TY_UNIT_INTRO`.

5 Grammar Definition

m	$::=$		matrix expressions
		M	matrix variables
		$m + m'$	matrix addition
		$m \ m'$	matrix multiplication
		(m)	S
f	$::=$		fractional capability
		fc	variable
		1	whole capability
		$\frac{1}{2}f$	
t	$::=$		linear type
		unit	unit
		bool	boolean (true/false)
		int	63-bit integers
		elt	array element
		$f \text{ arr}$	arrays
		$f \text{ mat}$	matrices
		$!t$	multiple-use type
		$\forall fc.t$	bind fc in t frac. cap. generalisation
		$t \otimes t'$	pair
		$t \multimap t'$	linear function
		(t)	S parentheses
p	$::=$		primitive
		not	boolean negation
		$(+)$	integer addition
		$(-)$	integer subtraction
		$(*)$	integer multiplication
		$(/)$	integer division
		$(=)$	integer equality
		$(<)$	integer less-than
		$(+.)$	element addition
		$(-.)$	element subtraction
		$(*.)$	element multiplication
		$(/.)$	element division
		$(=.)$	element equality
		$(<.)$	element less-than
		set	array index assignment
		get	array indexing
		share	share array
		unshare	unshare array
		free	free array
		array	Owl: make array
		copy	Owl: copy array
		sin	Owl: map sine over array
		hypot	Owl: $x_i := \sqrt{x_i^2 + y_i^2}$
		asum	BLAS: $\sum_i x_i $

	axpy		BLAS: $x := \alpha x + y$
	dot		BLAS: $x \cdot y$
	rotmg		BLAS: see its docs
	scal		BLAS: $x := \alpha x$
	amax		BLAS: $\text{argmax } i : x_i$
	setM		matrix index assignment
	getM		matrix indexing
	shareM		share matrix
	unshareM		unshare matrix
	freeM		free matrix
	matrix		Owl: make matrix
	copyM		Owl: copy matrix
	copyM_to		Owl: copy matrix onto another
	sizeM		dimension of matrix
	trnsp		transpose matrix
	gemm		BLAS: $C := \alpha A^{T?} B^{T?} + \beta C$
	symm		BLAS: $C := \alpha AB + \beta C$
	posv		BLAS: Cholesky decomp. and solve
	potrs		BLAS: solve with given Cholesky
	syrk		BLAS: $C := \alpha A^{T?} A^{T?} + \beta C$
v	::=		values
	p		primitives
	x		variable
	$()$		unit introduction
	true		true
	false		false
	k		integer
	$l \cdot f$		heap location
	el		array element
	Many v		!-introduction
	fun $fc \rightarrow v$		frac. cap. abstraction
	$v[f]$		frac. cap. specialisation
	(v, v')		pair introduction
	fun $x : t \rightarrow e$	bind x in e	abstraction
	fix $(g, x : t, e : t')$	bind $g \cup x$ in e	fixpoint
	(v)	S	parentheses
e	::=		expression
	p		primitives
	x		variable
	let $x = e$ in e'	bind x in e'	let binding
	$()$		unit introduction
	let $() = e$ in e'		unit elimination
	true		true
	false		false
	if e then e_1 else e_2		if
	k		integer

		$l \cdot f$		heap location
		el		array element
		Many e		!-introduction
		let Many $x = e$ in e'		!-elimination
		fun $fc \rightarrow e$		frac. cap. abstraction
		$e[f]$		frac. cap. specialisation
		(e, e')		pair introduction
		let $(a, b) = e$ in e'	bind $a \cup b$ in e'	pair elimination
		fun $x : t \rightarrow e$	bind x in e	abstraction
		$e e'$		application
		fix $(g, x : t, e : t')$	bind $g \cup x$ in e	fixpoint
		(e)	S	parentheses
C	$::=$			evaluation contexts
		let $x = [-]$ in e	bind x in e	let binding
		let $() = [-]$ in e		unit elimination
		if $[-]$ then e_1 else e_2		if
		Many $[-]$!-introduction
		let Many $x = [-]$ in e		!-elimination
		fun $fc \rightarrow [-]$		frac. cap. abstraction
		$[-][f]$		frac. cap. specialisation
		$([-], e)$		pair introduction
		$(v, [-])$		pair introduction
		let $(a, b) = [-]$ in e	bind $a \cup b$ in e	pair elimination
		$[-]e$		application
		$v[-]$		application
Θ	$::=$			fractional capability environment
		\cdot		
		Θ, fc		
Γ	$::=$			linear types environment
		\cdot		
		$\Gamma, x : t$		
		Γ, Γ'		
Δ	$::=$			intuitionistic types environment
		\cdot		
		$\Delta, x : t$		
σ	$::=$			heap (multiset of triples)
		$\{\}$		empty heap
		$\sigma + \{l \mapsto_f m_{k_1, k_2}\}$		location l points to matrix m
$StepsTo$	$::=$			result of small step
		$\langle \sigma, e \rangle$		heap and expression
		err		error