



ABBOTSLEIGH

Student's Name:

Student Number:

--	--	--	--	--	--	--	--	--	--

Teacher's Name:

2025

TRIAL HIGHER SCHOOL CERTIFICATE EXAMINATION

Mathematics Extension 1

General Instructions

- Reading time – 10 minutes
- Working time – 2 hours
- Write using black pen
- Calculators approved by NESA may be used
- A reference sheet is provided separately
- Diagrams are not to scale unless indicated
- Ensure your HSC candidate number is on each booklet
- For questions in Section I, answer on the sheet provided
- For questions in Section II,
 - show all necessary working
 - start a new booklet for each question

Total marks: **Section I – 10 marks** (pages 3-8)
70

- Attempt Questions 1–10
- Allow about 15 minutes for this section

Section II – 60 marks (pages 10-17)

- Attempt Questions 11–15
- Allow about 1 hour and 45 minutes for this section

Year 11 Mathematics Extension 1 outcomes

ME11-1 uses algebraic and graphical concepts in the modelling and solving of problems involving functions and their inverses

ME11-2 manipulates algebraic expressions and graphical functions to solve problems

ME11-3 applies concepts and techniques of inverse trigonometric functions and simplifying expressions involving compound angles in the solution of problems

ME11-4 applies understanding of the concept of a derivative in the solution of problems, including rates of change, exponential growth and decay and related rates of change

ME11-5 uses concepts of permutations and combinations to solve problems involving counting or ordering

ME11-7 communicates making comprehensive use of mathematical language, notation, diagrams and graphs

Year 12 Mathematics Extension 1 outcomes

ME12-1 applies techniques involving proof or calculus to model and solve problems

ME12-2 applies concepts and techniques involving vectors and projectiles to solve problems

ME12-3 applies advanced concepts and techniques in simplifying expressions involving compound angles and solving trigonometric equations

ME12-4 uses calculus in the solution of applied problems, including differential equations and volumes of solids of revolution

ME12-5 applies appropriate statistical processes to present, analyse and interpret data

ME12-7 evaluates and justifies conclusions, communicating a position clearly in appropriate mathematical forms

Section I

10 Marks

Attempt Questions 1–10

Allow about 15 minutes for this section

Use the multiple-choice answer sheet for Questions 1-10.

1 Given that $\overrightarrow{AD} = \begin{pmatrix} -2 \\ 3 \end{pmatrix}$ and $\overrightarrow{DO} = \begin{pmatrix} 6 \\ -4 \end{pmatrix}$, what is $\overrightarrow{OA}$?

A. $\begin{pmatrix} -4 \\ 1 \end{pmatrix}$

B. $\begin{pmatrix} -4 \\ -1 \end{pmatrix}$

C. $\begin{pmatrix} 4 \\ 1 \end{pmatrix}$

D. $\begin{pmatrix} 4 \\ -1 \end{pmatrix}$

2 Consider the polynomial $P(x) = 4x^2 + 5x - 10$.

When $(2x - a) \times P(x)$ is divided by $(x - 1)$ the remainder is 3.

What is the value of a ?

A. -1

B. 1

C. 2

D. 5

Section I continues on page 4

3 Which of the following is an anti-derivative of $\frac{1}{\sqrt{9-4x^2}}$?

A. $\frac{\sin^{-1}\left(\frac{2x}{3}\right)}{2} + c$

B. $\frac{\sin^{-1}\left(\frac{3x}{2}\right)}{3} + c$

C. $\frac{\sin^{-1}(2x)}{3} + c$

D. $\frac{\sin^{-1}\left(\frac{2x}{3}\right)}{6} + c$

4 A factory produces light switches. Each switch has a 6% chance of being faulty. A quality control manager selects a random sample of 80 switches. Let X represent the number of faulty switches in the sample, where X is a binomial variable. What is the standard deviation of X , correct to 2 decimal places?

A. 2.12


B. 2.13

C. 4.51

D. 4.8

Section I continues on page 5

- 5 The direction field for a differential equation is shown below.



Which of the differential equations below could give this direction field?

- A. $\frac{dy}{dx} = \frac{y}{2y + x}$
- B. $\frac{dy}{dx} = -\frac{y}{2y + x}$
- C. $\frac{dy}{dx} = \frac{y}{-2y + x}$
- D. $\frac{dy}{dx} = \frac{y}{2y - x}$

Section I continues of page 6

- 6 The vectors $\underline{d}$, $\underline{u}$, $\underline{b}$, $\underline{q}$ are each chosen to be either $\begin{pmatrix} 2 \\ 1 \end{pmatrix}$ or $\begin{pmatrix} 3 \\ -1 \end{pmatrix}$.
If each possible selection is equally likely, what is the probability that

$$\underline{d} + \underline{u} + \underline{b} + \underline{q} = \begin{pmatrix} 10 \\ 0 \end{pmatrix} ?$$

- A. $\frac{1}{24}$
- B. $\frac{3}{16}$
- C. $\frac{1}{8}$
- D. $\frac{3}{8}$
- 7 For the vectors $\underline{a} = 3\underline{i} + 2\underline{j}$ and $\underline{b} = 9\underline{i} + 3\underline{j}$, which of the following represents $|\underline{a} + \underline{b}|$?
- A. $\sqrt{37}$
- B. $\sqrt{103}$
- C. 12
- D. 13

Section I continues on page 7

- 8 The function $f(x)$ exists for all real numbers x .

The function $g(x)$ is defined such that $g(x) = f(x) + \frac{1}{f(x)}$.

If $f(2) = -1$ and $f'(2) > 0$, which of the following is true about $g(x)$?

- A. $g(2) = -2$ and $g'(2) > 0$
- B. $g(2) = 1$ and $g'(2) = 0$
- C. $g(2) = -2$ and $g'(2) = 0$
- D. $g(2) = 1$ and $g'(2) < 0$

- 9 A first-order differential equation is given by

$$\frac{dy}{dx} = 2y \ln y$$

Which of the following is a solution to the differential equation?

- A. $y = e^{2e^x}$
- B. $y = e^{e^{2x}}$
- C. $y = e^{e^{x^2}}$
- D. $y = 2e^{e^x}$

Section I continues on page 8

- 10** For a particular function $f(x)$, it is known that $f(0) = 0$ and $x \times f(x) > 0$ for all non-zero values of x .

It is also known that $\int_{-2}^2 f(x) \, dx = 4$ and $\int_{-2}^2 |f(x)| \, dx = 8$.

Which of the following gives the value of $\int_{-2}^0 f(|x|) \, dx$?

- A. -2
- B. 2
- C. 4
- D. 6

End of Section I

-BLANK PAGE-

Section II

60 Marks

Attempt Questions 11–15

Allow about 1 hour and 45 minutes for this section

Answer each question in a separate writing booklet. Use extra booklets if needed.

For questions in Section II, your responses should include relevant mathematical reasoning and/or calculations.

Question 11 (12 marks) Use a SEPARATE writing booklet

Marks

(a) For the vectors $\underline{a} = 10\underline{i} + 2\underline{j}$ and $\underline{b} = \underline{i} - \underline{j}$, evaluate each of the following.

(i) $3\underline{a} - \underline{b}$ 1

(ii) $(\underline{a} + 2\underline{b}) \cdot (\underline{a} - 2\underline{b})$ 2

(b) Using the substitution $u = x + 4$, find the exact value of $\int_{-4}^{-3} x(x+4)^{10} dx$. 3

(c) In how many different ways can all the letters of the word **TENNANT** be arranged in a line? 2

(d) Find the constant term in the expansion of $\left(2x - \frac{1}{x}\right)^{10}$. 2

(e) Solve $\sqrt{2} \cos\left(x + \frac{\pi}{4}\right) + \sqrt{2} \cos\left(x - \frac{\pi}{4}\right) = 1$ for $-\pi \leq x \leq \pi$. 2

End of Question 11

Question 12 (12 marks) Use a SEPARATE writing booklet


Marks

(a) (i) Show that $\sec^2\left(\frac{\pi}{4} - \frac{x}{2}\right) = \frac{2}{1 + \sin x}$. 2

(ii) Hence, or otherwise, evaluate $\int_0^\pi \frac{2}{1 + \sin x} dx$. 2

(b) The volume of a cylinder with a fixed height h cm, and radius r cm is given by $V = \pi r^2 h$. If the volume is increasing at a rate of $20 \text{ cm}^3 \text{ s}^{-1}$, show that $\frac{dr}{dt} = \frac{10}{\pi r h} \text{ cm s}^{-1}$. 2

(c) The region, **D**, is bounded by the y -axis, the x -axis, the line with equation $x = \ln 4$ and the curve with equation $y = e^x + 2e^{-x}$, as shown in the diagram. 3



Find the volume of the solid of revolution formed when the region **D** is rotated about the x -axis. Leave your answer in exact form.

(d) The polynomial $P(x) = 8x^4 - 38x^3 + 9x^2 + ax + b$ has a double root at $x = 3$. 3

Find the values of a and b .

End of Question 12

Question 13 (12 marks) Use a SEPARATE writing booklet

Marks

- (a) Use mathematical induction to show that for all integers $n \geq 1$, **3**
 $4^{2n} - 1$ is divisible by 3.

- (b) Consider the function given by $f(x) = \tan^{-1} \left(\frac{\cos x}{\sin x + 1} \right)$ for $-\frac{\pi}{2} < x < \frac{\pi}{2}$.

- (i) Find $f'(x)$. **2**

- (ii) Hence sketch the graph of $y = f(x)$. **1**

- (c) Solve the differential equation $\frac{dy}{dx} = x\sqrt{(1-x^2)(1-y^2)}$ given that it **3**
passes through the point $(1, 0)$. Give your answer with y as the subject.

- (d) Prove that **3**

$$\frac{1}{{}_{2025}C_{26}} = \frac{26}{25} \left(\frac{1}{{}_{2024}C_{25}} - \frac{1}{{}_{2025}C_{25}} \right)$$

End of Question 13

- (a) A team of ecologists is re-introducing a group of 3000 Komodo dragons to an island in Indonesia. From observations on other islands, the ecologists know that populations of more than 3000 are unsustainable.

They propose the following logistic model of population growth

$$\frac{dK}{dt} = rK(3000 - K)$$

where K is the number of Komodo dragons on the island, t is the time in years and r is a constant.

- (i) Ecologists know that when the maximum growth rate occurs, all the Komodo dragons find a mate. Half of these pairs successfully raise one juvenile Komodo dragon. **2**

Show that $r = \frac{1}{6000}$.

- (ii) The ecologists initially release 500 Komodo dragons on the island. **3**

Use the fact that $\frac{1}{K(3000 - K)} = \frac{1}{3000} \left(\frac{1}{K} + \frac{1}{3000 - K} \right)$ to show

that the population of Komodo dragons can be given by

$$K = \frac{3000}{5e^{-0.5t} + 1}.$$


- (iii) How long will it take for the Komodo population to reach 95% of the maximum sustainable population? Give your answer correct to 2 decimal places. **1**

Question 14 continues on page 14

- (b) Particle A is projected from a vertical tower that is H metres above the ground with an initial speed of $v \text{ m s}^{-1}$ at an acute angle of α with the horizontal plane.

At the same time, from another point D metres horizontally away from the base of the vertical tower, particle B is projected with an initial speed of $u \text{ m s}^{-1}$ at an acute angle of β with the horizontal plane.

This is shown in the diagram below.



Taking the position of the base of the tower as the origin, the positions of the projectiles at time t after they are launched are as follows.

$$\vec{r}_A = \begin{pmatrix} vt \cos \alpha \\ vt \sin \alpha - \frac{g}{2}t^2 + H \end{pmatrix} \quad \text{Particle } A$$

$$\vec{r}_B = \begin{pmatrix} ut \cos \beta + D \\ ut \sin \beta - \frac{g}{2}t^2 \end{pmatrix} \quad \text{Particle } B \quad (\text{DO NOT prove these.})$$

Question 14 continues on page 15

- (i) It is known that $u \sin \beta > v \sin \alpha$. (DO NOT prove this.)

3

Show that for the two particles to collide before either reaches the ground, then:

$$2u \sin \beta (u \sin \beta - v \sin \alpha) > gH .$$

- (ii) Show, by expanding the right-hand side, that:

1

$$\sin(2\beta) + \cos(2\beta) = \sqrt{2} \sin\left(2\beta + \frac{\pi}{4}\right).$$

- (iii) If $u = v$, $H = \frac{u^2}{g}$ and $\alpha = \frac{\pi}{2} - \beta$, show that $\frac{3\pi}{8} < \beta < \frac{\pi}{2}$

3

if the two particles collide before hitting the ground.

End of Question 14

- (a) *You may use the information on page 19 to answer this question.*

At Hogwarts School, the probability of a troll being in the dungeon on any given day is 5%.

Students are becoming disgruntled with the number of trolls in the dungeon and are concerned one may appear during their examination block. They demand a sample size of n is taken and the sample proportion $\hat{p}$ is calculated.

It is assumed that $\hat{p}$ is normally distributed with $E(\hat{p}) = p$ and

$$\text{Var}(\hat{p}) = \frac{p(1-p)}{n}.$$

The students will report Hogwarts School to the Ministry of Magic if a troll is in the dungeon during their 10-day examination block. That is, if $\hat{p} \geq 0.1$.

To protect the Hogwarts school's reputation, the Ministry of Magic suggests they choose a sample size so that the chance of students reporting Hogwarts is less than 0.62%.

- (i) Show that the mean and standard deviation of the sample proportion **1**

are given by 0.05 and $\sqrt{\frac{19}{400n}}$ respectively.

- (ii) Hence, or otherwise, find the approximate sample size required. **3**


Question 15 continues on page 17

- (b) Three different points O , V , and U form an isosceles triangle with unit length sides such that $|\overrightarrow{OV}| = |\overrightarrow{OU}| = 1$.

The points C and D are chosen such that $\overrightarrow{VC} = \overrightarrow{CD} = \overrightarrow{DU} = \frac{1}{3}\overrightarrow{VU}$

(that is, the points C and D split the line into thirds.)

Let $\underline{v} = \overrightarrow{OV}$ and $\underline{u} = \overrightarrow{OU}$, as shown in the diagram.



- (i) Show that $\overrightarrow{VD} = \frac{2}{3}\underline{u} - \frac{2}{3}\underline{v}$, and that $\overrightarrow{UC} = \frac{2}{3}\underline{v} - \frac{2}{3}\underline{u}$. 1
- (ii) The point A lies on the line OV such that $\overrightarrow{OA} = \lambda\underline{v}$ and $0 < \lambda < 1$. 2
 Show that $\overrightarrow{AD} = \frac{2}{3}\underline{u} + \left(\frac{1}{3} - \lambda\right)\underline{v}$.
- (iii) The point B similarly divides the line OU in the same ratio such that $\overrightarrow{BU} = \lambda\underline{u}$. 4

The lines AD and BC are perpendicular, as shown in the diagram.


By first finding an expression for $\overrightarrow{BC}$ in terms of $\underline{u}$, $\underline{v}$ and λ , show

that $\frac{8}{17} \leq \underline{u} \cdot \underline{v} \leq 1$.

End of Paper

-BLANK PAGE-

Table of values $P(Z \leq z)$ for the normal distribution $N(0,1)$



Z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
0.0	0.5000	0.5040	0.5080	0.5120	0.5160	0.5199	0.5239	0.5279	0.5319	0.5359
0.1	0.5398	0.5438	0.5478	0.5517	0.5557	0.5596	0.5636	0.5675	0.5714	0.5753
0.2	0.5793	0.5832	0.5871	0.5910	0.5948	0.5987	0.6026	0.6064	0.6103	0.6141
0.3	0.6179	0.6217	0.6255	0.6293	0.6331	0.6368	0.6406	0.6443	0.6480	0.6517
0.4	0.6554	0.6591	0.6628	0.6664	0.6700	0.6736	0.6772	0.6808	0.6844	0.6879
0.5	0.6915	0.6950	0.6985	0.7019	0.7054	0.7088	0.7123	0.7157	0.7190	0.7224
0.6	0.7257	0.7291	0.7324	0.7357	0.7389	0.7422	0.7454	0.7486	0.7517	0.7549
0.7	0.7580	0.7611	0.7642	0.7673	0.7704	0.7734	0.7764	0.7794	0.7823	0.7852
0.8	0.7881	0.7910	0.7939	0.7967	0.7995	0.8023	0.8051	0.8078	0.8106	0.8133
0.9	0.8159	0.8186	0.8212	0.8238	0.8264	0.8289	0.8315	0.8340	0.8365	0.8389
1.0	0.8413	0.8438	0.8461	0.8485	0.8508	0.8531	0.8554	0.8577	0.8599	0.8621
1.1	0.8643	0.8665	0.8686	0.8708	0.8729	0.8749	0.8770	0.8790	0.8810	0.8830
1.2	0.8849	0.8869	0.8888	0.8907	0.8925	0.8944	0.8962	0.8980	0.8997	0.9015
1.3	0.9032	0.9049	0.9066	0.9082	0.9099	0.9115	0.9131	0.9147	0.9162	0.9177
1.4	0.9192	0.9207	0.9222	0.9236	0.9251	0.9265	0.9279	0.9292	0.9306	0.9319
1.5	0.9332	0.9345	0.9357	0.9370	0.9382	0.9394	0.9406	0.9418	0.9429	0.9441
1.6	0.9452	0.9463	0.9474	0.9484	0.9495	0.9505	0.9515	0.9525	0.9535	0.9545
1.7	0.9554	0.9564	0.9573	0.9582	0.9591	0.9599	0.9608	0.9616	0.9625	0.9633
1.8	0.9641	0.9649	0.9656	0.9664	0.9671	0.9678	0.9686	0.9693	0.9699	0.9706
1.9	0.9713	0.9719	0.9726	0.9732	0.9738	0.9744	0.9750	0.9756	0.9761	0.9767
2.0	0.9772	0.9778	0.9783	0.9788	0.9793	0.9798	0.9803	0.9808	0.9812	0.9817
2.1	0.9821	0.9826	0.9830	0.9834	0.9838	0.9842	0.9846	0.9850	0.9854	0.9857
2.2	0.9861	0.9864	0.9868	0.9871	0.9875	0.9878	0.9881	0.9884	0.9887	0.9890
2.3	0.9893	0.9896	0.9898	0.9901	0.9904	0.9906	0.9909	0.9911	0.9913	0.9916
2.4	0.9918	0.9920	0.9922	0.9925	0.9927	0.9929	0.9931	0.9932	0.9934	0.9936
2.5	0.9938	0.9940	0.9941	0.9943	0.9945	0.9946	0.9948	0.9949	0.9951	0.9952
2.6	0.9953	0.9955	0.9956	0.9957	0.9959	0.9960	0.9961	0.9962	0.9963	0.9964
2.7	0.9965	0.9966	0.9967	0.9968	0.9969	0.9970	0.9971	0.9972	0.9973	0.9974
2.8	0.9974	0.9975	0.9976	0.9977	0.9977	0.9978	0.9979	0.9979	0.9980	0.9981
2.9	0.9981	0.9982	0.9982	0.9983	0.9984	0.9984	0.9985	0.9985	0.9986	0.9986
3.0	0.9987	0.9987	0.9987	0.9988	0.9988	0.9989	0.9989	0.9989	0.9990	0.9990

-BLANK PAGE-