# A STUDY OF T-AVOIDING ELEMENTS OF COXETER GROUPS

## By Taryn Laird

## A Thesis

Submitted in Partial Fulfillment

of the Requirements for the Degree of

Master of Science

in Mathematics

Northern Arizona University

May 2016

Approved:

Dana Ernst, Ph.D., Chair

Michael Falk, Ph.D.

Nándor Sieben, Ph.D.

## ABSTRACT

## A STUDY OF T-AVOIDING ELEMENTS OF COXETER GROUPS

## Taryn Laird

Coming soon. We are finishing writing everything else first so we know the appropriate roadmap to be in this part.

# Contents

List of 1	Figures	iv	
Chapte	er 1 Preliminaries	1	
1.1	Introduction	1	
1.2	Coxeter Systems	1	
1.3	Fully Commutative Elements	7	
1.4	Heaps	Ö	
Chapte	er 2 Star Reductions and Property T	14	
2.1	Star Reductions	14	
2.2	Property T	17	
2.3	Non-Cancellable Elements	19	
Chapte	er 3 T-Avoiding Elements in Types $\widetilde{A}_n, A_n, D_n, F_n$ , and $I_2(m)$	23	
3.1	Types $\widetilde{A}_n$ and $A_n$	23	
3.2	Type $D_n$	24	
3.3	Type $F_n$	25	
3.4	Type $I_2(m)$	28	
Chapter 4 T-Avoiding Elements in Type $B_n$		29	
4.1	Tools for the Classification	29	
4.2	Classification of T-Avoiding Elements in Type $B_n$	31	
Chapte	$\mathbf{r}$ 5 T-Avoiding Elements in Type $\widetilde{C}_n$	39	
5.1	Classification of T-Avoiding Elements in Type $\widetilde{C}_n$	39	
5.2	Future Work	53	
Bibliog	Bibliography		

# List of Figures

1.1	A few labeled Coxeter graphs	3
1.2	Irreducible finite Coxeter systems	5
1.3	Irreducible affine Coxeter systems	6
1.4	Irreducible FC-finite Coxeter systems	9
1.5	Labeled hasse diagram for the heap of an element in $FC(C_6)$	10
1.6	A lattice point representation for the heap of an FC element in $W(\widetilde{C}_6)$ .	12
1.7	Two heaps of a non-FC element in $W(\widetilde{C}_4)$	12
1.8	Impermissible configurations for heaps of $FC(\widetilde{C}_n)$	13
2.1	A visual representation of an element that is left star reducible	15
2.2	Visualization of Example 2.1.1	16
2.3	Visualization of Example 2.1.3	16
2.4	Heap of an element with Property T	18
2.5	Heap of a non-trivially T-avoiding element in $W(C_4)$	19
2.6	Heap of a weak star reducible element of $FC(B_4)$	20
2.7	Heap of a non-cancellable element of $FC(B_4)$	21
2.8	Heap of a single sandwich stack in $W(\widetilde{C}_n)$	21
2.9	Heap of a stack of sandwich stacks in $W(\widetilde{C}_n)$	21
3.1	Visual representation of non-trivial T-avoiding elements in $W(D_5)$ .	25
3.2	Heap of a single bowtie in $W(F_5)$	26
3.3	Heap of a stack of bowties in $W(F_5)$	27
3.4	Heap of a non-trivial T-avoiding element in $W(F_6)$	28
5.1	pushed-down representation of a heap	40

## Chapter 1

# **Preliminaries**

#### 1.1 Introduction

This thesis is organized as follows. After necessary background information is presented in Section 1.2, we then introduce the class of fully commutative elements in Section 1.3. Then, in Sectio ?? we discuss a visual representation for elements of Coxeter groups, called heaps. In Section 2.1, we introduce the concept of a star reducible Coxeter group and in Section 2.2 we introduce the notion of a T-avoiding element which is exactly an element that does not begin or end with a product of noncommuting generators. In Section 2.3 we state the classification of elements that are non-cancellable in Coxeter systems of type  $\widetilde{C}_n$ . We then state classifications and conjectures regarding T-avoiding elements in several Coxeter systems in Chapter 3. Following this we introduce the main results for this thesis, the classification of T-avoiding elements in Coxeter systems of type  $B_n$  and  $\widetilde{C}_n$ . In Section 4.1 we introduce the necessary lemmas and definitions for the classification in Section 4.2 in which we show there are no non-trivial T-avoiding elements in Coxeter systems of type  $B_n$ . In Section 5.1 we classify the non-trivial T-avoiding elements in Coxeter systems of type  $\widetilde{C}_n$ . We conclude with some open questions in Section 5.2.

## 1.2 Coxeter Systems

A Coxeter system is a pair (W, S) consisting of a finite set S of generating involutions and a group W, called a Coxeter group, with presentation

$$W = \langle S \mid (st)^{m(s,t)} = e \text{ for } m(s,t) < \infty \rangle,$$

where e is the identity, m(s,t) = 1 if and only if s = t, and m(s,t) = m(t,s). It turns out that the elements of S are distinct as group elements and that m(s,t) is the order

of st [10]. We call m(s,t) the bond strength of s and t.

Since s and t are elements of order 2, the relation  $(st)^{m(s,t)} = e$  can be rewritten as

$$\underbrace{sts\cdots}_{m(s,t)} = \underbrace{tst\cdots}_{m(s,t)} \tag{1.1}$$

with  $m(s,t) \geq 2$  factors. If m(s,t) = 2, then st = ts is called a commutation relation. Otherwise, if  $m(s,t) \geq 3$ , then the relation in (1.1) is called a braid relation. Replacing  $\underbrace{sts\cdots}_{m(s,t)}$  with  $\underbrace{tst\cdots}_{m(s,t)}$  will be referred to as a commutation if m(s,t) = 2 and a braid move if  $m(s,t) \geq 3$ .

We can represent a Coxeter system (W, S) with a unique Coxeter graph  $\Gamma$  having

- (1) vertex set S and
- (2) edges  $\{s,t\}$  for each  $m(s,t) \geq 3$  labeled by its corresponding bond strength.

Since m(s,t)=3 occurs frequently, it is customary to omit this label. Note that s and t are not connected by a single edge in the graph if and only if m(s,t)=2. There is a one-to-one correspondence between Coxeter systems and Coxeter graphs. That is, given a Coxeter graph  $\Gamma$ , we can uniquely reconstruct the corresponding Coxeter system. If (W,S) is a Coxeter system with corresponding Coxeter graph  $\Gamma$ , we may denote the Coxeter group as  $W(\Gamma)$  and the generating set as  $S(\Gamma)$  for clarity. Also, the Coxeter system (W,S) is said to be *irreducible* if and only if  $\Gamma$  is connected. If the graph  $\Gamma$  is disconnected, the connected components correspond to factors in a direct product of the corresponding Coxeter groups [10]. The Coxeter graphs given in Figure 1.1 correspond to the Coxeter systems that will be primarily addressed in this thesis. Notice that the vertices are labeled with the corresponding generators to provide context when talking about the different generating sets  $S(\Gamma)$ .




Figure 1.1: A few labeled Coxeter graphs.

#### Example 1.2.1.

- (a) The Coxeter system of type  $A_n$  is given by the graph in Figure 1.1(a). We can construct the corresponding Coxeter group  $W(A_n)$  with generating set  $S(A_n) = \{s_1, s_2, \ldots, s_n\}$  and defining relations
  - (1)  $s_i^2 = e$  for all i;
  - (2)  $s_i s_j = s_j s_i$  when |i j| > 1;
  - (3)  $s_i s_j s_i = s_j s_i s_j$  when |i j| = 1.

The Coxeter group  $W(A_n)$  is isomorphic to the symmetric group  $\operatorname{Sym}_{n+1}$  under the correspondence  $s_i \mapsto (i, i+1)$ , where (i, i+1) is the adjacent transposition that swaps i and i+1.

- (b) The Coxeter system of type  $B_n$  is given by the graph in Figure 1.1(c). We can construct the corresponding Coxeter group  $W(B_n)$  with generating set  $S(B_n) = \{s_0, s_1, \ldots, s_{n-1}\}$  and defining relations
  - (1)  $s_i^2 = e$  for all i;
  - (2)  $s_i s_j = s_j s_i$  when |i j| > 1;
  - (3)  $s_i s_j s_i = s_j s_i s_j$  when |i j| = 1 for  $i, j \in \{1, 2, \dots, n 1\}$ ;
  - $(4) \ s_0 s_1 s_0 s_1 = s_1 s_0 s_1 s_0.$

The Coxeter group  $W(B_n)$  is isomorphic to  $\operatorname{Sym}_n^B$ , where  $\operatorname{Sym}_n^B$  is the group of signed permutations on the set  $\{1, 2, \ldots, n\}$ .

- (c) The Coxeter system of type  $\widetilde{C}_n$  is seen in Figure 1.1(d). We can construct the corresponding Coxeter group  $W(\widetilde{C}_n)$  with generating set  $S(\widetilde{C}_n) = \{s_0, s_1, \ldots, s_n\}$  and defining relations
  - (1)  $s_i^2 = e$  for all i;
  - (2)  $s_i s_j = s_i s_i$  when |i j| > 1 for  $i \in \{0, 2, \dots, n\}$ ;
  - (3)  $s_i s_j s_i = s_j s_i s_j$  when |i j| = 1 for  $i \in \{1, 2, \dots, n 1\}$ ;
  - (4)  $s_0 s_1 s_0 s_1 = s_1 s_0 s_1 s_0$ ;
  - (5)  $s_n s_{n-1} s_n s_{n-1} = s_{n-1} s_n s_{n-1} s_n$ .

Note that  $W(\widetilde{C}_n)$  has n+1 generators.

The Coxeter graphs given in Figure 1.2 correspond to the collection of irreducible finite Coxeter systems, whose corresponding Coxeter groups are finite, while the Coxeter graphs given in Figure 1.3 are the so-called irreducible affine Coxeter systems, which are infinite [10]. Note that  $W(B_n)$  is one of the irreducible finite Coxeter groups so it is finite while  $W(\widetilde{C}_n)$  is one of the affine groups making it infinite. The irreducible affine Coxeter systems are unique in that if a vertex is removed along with the corresponding edges from the Coxeter graph, the newly created graph will result in a Coxeter system with a finite Coxeter group.

Given a Coxeter system (W, S), a word  $s_{x_1}s_{x_2}\cdots s_{x_m}$  in the free monoid  $S^*$  on S is called an *expression* for  $w \in W$  if it is equal to w when considered as a group element. If m is minimal among all expressions for w, the corresponding word is called a *reduced expression* for w. In this case, we define the *length* of w to be l(w) := m. Each element  $w \in W$  may have multiple reduced expressions that represent it. If we wish to emphasize a specific, possibly reduced, expression for  $w \in W$  we will represent it as  $\mathbf{w} = s_{x_1} s_{x_2} \cdots s_{x_m}$  (using sans serif font). If  $u, v \in W(\Gamma)$ , we say that the product




Figure 1.2: Irreducible finite Coxeter systems.

of group elements uv is reduced if l(uv) = l(u) + l(v). Matsumoto's Theorem, which follows, tells us more about how reduced expressions for a given group element are related.

**Proposition 1.2.2** (Matsumoto, [7]). Let (W, S) be a Coxeter system. If  $w \in W$ , then given a reduced expression for w we can obtain every other reduced expression for w by a sequence of braid moves and commutations of the form

$$\underbrace{sts\cdots}_{m(s,t)} \to \underbrace{tst\cdots}_{m(s,t)}$$

where  $s, t \in S$  and  $m(s, t) \geq 2$ .

It follows from Matsumoto's Theorem that if a generator s appears in a reduced expression for  $w \in W$ , then s appears in all reduced expressions for w. Let  $w \in W$  and define the *support* of w, denoted supp(w), to be the set of all generators that




Figure 1.3: Irreducible affine Coxeter systems.

appear in any reduced expression for w. If supp(w) = S, we say that w has full support.

Given  $w \in W$  and a fixed reduced expression w for w, any subsequence of w is called a *subexpression* of w. We will refer to a subexpression consisting of a consecutive subsequence of w as a *subword* of w.

**Example 1.2.3.** Let  $w \in W(A_7)$  and let  $\mathbf{w} = s_7 s_2 s_4 s_5 s_3 s_2 s_3 s_6$  be a fixed expression

for w. Then we have

```
s_7 s_2 s_4 s_5 s_3 s_2 s_3 s_6 = s_7 s_4 s_2 s_5 s_3 s_2 s_3 s_6
= s_7 s_4 s_5 s_2 s_3 s_2 s_3 s_6
= s_7 s_4 s_5 s_3 s_2 s_3 s_3 s_6
= s_7 s_4 s_5 s_3 s_2 s_6,
```

where the purple highlighted text corresponds to a commutation, the teal highlighted text corresponds to a braid move, and the red highlighted text corresponds to cancellation. This shows that the original expression w is not reduced. However, it turns out that  $s_7s_4s_5s_3s_2s_6$  is reduced. Thus l(w) = 6 and  $\sup(w) = \{s_2, s_3, s_4, s_5, s_6, s_7\}$ .

Let (W, S) be a Coxeter system of type  $\Gamma$  and let  $w \in W(\Gamma)$ . We define the *left descent set* and *right descent set* of w as follows:

$$\mathcal{L}(w) := \{ s \in S \mid l(sw) < l(w) \}$$

and

$$\mathcal{R}(w) := \{ s \in S \mid l(ws) < l(w) \}.$$

In [2] it is shown that  $s \in \mathcal{L}(w)$  (respectively,  $\mathcal{R}(w)$ ) if and only if there is a reduced expression for w that begins (respectively, ends) with s.

**Example 1.2.4.** The following list consists of all reduced expressions some  $w \in W(B_4)$ :

```
s_0s_1s_2s_1s_3 s_0s_2s_1s_2s_3

s_0s_1s_2s_3s_1 s_2s_0s_1s_2s_3
```

We see that l(w) = 5 and w has full support. Also, we see that  $\mathcal{L}(w) = \{s_0, s_2\}$  while  $\mathcal{R}(w) = \{s_1, s_3\}$ .

## 1.3 Fully Commutative Elements

Let (W, S) be a Coxeter system of type  $\Gamma$  and let  $w \in W(\Gamma)$ . Following [13], we define a relation  $\sim$  on the set of reduced expressions for w. Let  $\mathsf{w}_1$  and  $\mathsf{w}_2$  be two reduced expressions for w. We define  $\mathsf{w}_1 \sim \mathsf{w}_2$  if we can obtain  $\mathsf{w}_2$  from  $\mathsf{w}_1$  by applying a single commutation move of the form  $st \mapsto ts$  where m(s,t) = 2. Now, define the equivalence relation  $\approx$  by taking the reflexive transitive closure of  $\sim$ . Each equivalence class under  $\approx$  is called a *commutation class*. If w has a single commutation class, then we say that w is fully commutative (FC).

The set of FC elements of  $W(\Gamma)$  is denoted by  $FC(\Gamma)$ . Given some  $w \in FC(\Gamma)$ , and a starting reduced expression for w, observe that the definition of FC states that

one only needs to perform commutations to obtain all reduced expressions for w, but the following result due to Stembridge [13] states that when w is FC, performing commutations is the only possible way to obtain another reduced expression for w.

**Proposition 1.3.1** (Stembridge, [13]). Let (W, S) be a Coxeter system. An element  $w \in W$  is FC if and only if no reduced expression for w contains  $\underbrace{sts\cdots}_{w \in W}$  as a subword

for all  $m(s,t) \geq 3$ .

In other words, w is FC if and only if no reduced expression provides the opportunity to apply a braid move. For example, for a Coxeter system of type  $B_n$  an element is FC if no reduced expression contains the subwords  $s_0s_1s_0s_1$ ,  $s_1s_0s_1s_0$ ,  $s_ks_{k+1}s_k$ , and  $s_{k+1}s_ks_{k+1}$  where 0 < k < n-2. In a Coxeter system of type  $C_n$ , an element is FC if no reduced expression for the element contains the subwords seen above and does not contain the subwords  $s_{n-1}s_ns_{n-1}s_n$  and  $s_ns_{n-1}s_ns_{n-1}$ .

**Example 1.3.2.** Let  $w \in W(\widetilde{C}_4)$  and let  $w = s_0 s_1 s_2 s_0 s_3 s_1$  be a reduced expression for w. Although it is not immediately obvious, there is no possible way to perform a braid move in any reduced expression for w. Hence w is FC.

**Example 1.3.3.** Let  $w_1 = s_1 s_0 s_4 s_1 s_3 s_5 s_2 s_4 s_6$  be a reduced expression for  $w \in W(\widetilde{C}_6)$ . Applying the commutation  $s_4 s_2 \mapsto s_2 s_4$ , we can obtain another reduced expression for w, namely  $w_2 = s_1 s_0 s_4 s_1 s_3 s_5 s_4 s_2 s_6$ , which is in the same commutation class as w. However, applying the braid move  $s_2 s_3 s_2 \mapsto s_3 s_2 s_3$ , we obtain another reduced expression  $w = s_1 s_3 s_2 s_3 s_4 s_0$ . Note that since  $w_3$  was obtained by applying a braid move,  $w_3$  is in a different commutation class than  $w_1$  and  $w_2$ . Since w has at least two commutation classes, one containing  $w_1$  and  $w_2$  and another containing  $w_3$ , w is not FC by Proposition 1.3.1.

Stembridge classified the Coxeter systems whose groups contain a finite number of FC elements, the so-called FC-finite Coxeter groups. Both  $W(A_n)$  and  $W(B_n)$  are finite Coxeter groups, and thus are FC-finite. On the other hand,  $W(\widetilde{C}_n)$  is infinite and happens to also contain infinitely many FC elements. However, there exist infinite Coxeter groups that contain finitely many FC elements. For example,  $W(E_n)$  for  $n \geq 9$  (see Figure 1.4) is infinite, but contains only finitely many FC elements.

**Proposition 1.3.4** (Stembridge, [13]). The irreducible FC-finite Coxeter systems are of type  $A_n$  with  $n \ge 1$ ,  $B_n$  with  $n \ge 2$ ,  $D_n$  with  $n \ge 4$ ,  $E_n$  with  $n \ge 6$ ,  $E_n$  with  $n \ge 4$ ,  $E_n$  with  $E_$ 

The irreducible FC-finite Coxeter graphs are given in Figure 1.4. Note that the irreducible finite Coxeter systems given in Figure 1.2 certainly have only a finite

number of FC elements. We have not yet encountered the Coxeter groups determined by graphs in Figures 1.4(d) for  $n \geq 9$ , 1.4(e) for  $n \geq 5$ , 1.4(f) for  $n \geq 5$ . All of these Coxeter systems are infinite for sufficiently large n, yet contain only finitely many FC elements.




Figure 1.4: Irreducible FC-finite Coxeter systems.

## 1.4 Heaps

We now discuss a visual representation of Coxeter group elements. Each reduced expression can be associated with a labeled partially ordered set (poset) called a heap. Heaps provide a visual representation of a reduced expression while preserving the relations among the generators. We follow the development of heaps for straight line Coxeter groups found in [1], [3], and [13].

Let (W, S) be a Coxeter system of type  $\Gamma$ . Suppose  $\mathbf{w} = s_{x_1} s_{x_2} \cdots s_{x_r}$  is a fixed reduced expression for  $w \in W(\Gamma)$ . As in [13], we define a partial ordering on the indices  $\{1, 2, \ldots, r\}$  by the transitive closure of the relation < defined via j < i if i < j and  $s_{x_i}$  and  $s_{x_j}$  do not commute. In particular, since  $\mathbf{w}$  is reduced, j < i if  $s_{x_i} = s_{x_j}$  by transitivity. This partial order is referred to as the *heap* of  $\mathbf{w}$ , where i is labeled by  $s_{x_i}$ . Note that for simplicity we are omitting the labels of the underlying poset yet

retaining the labels of the corresponding generators.

It follows from [13] that heaps are well-defined up to commutation class. That is, given two reduced expressions  $\mathbf{w}_1$  and  $\mathbf{w}_2$  for  $w \in W$  that are in the same commutation class, the heaps for  $\mathbf{w}_1$  and  $\mathbf{w}_2$  will be equal. In particular, if  $w \in FC(\Gamma)$ , then w has one commutation class, and thus w has a unique heap. Conversely, if  $\mathbf{w}_1$  and  $\mathbf{w}_2$  are in different commutation classes, then the heap for  $\mathbf{w}_1$  will be distinct from the heap for  $\mathbf{w}_2$ .

**Example 1.4.1.** Let  $\mathbf{w} = s_6 s_4 s_2 s_5 s_3 s_1 s_4 s_0 s_1$  be a reduced expression for  $w \in FC(\widetilde{C}_6)$ . We see that  $\mathbf{w}$  is indexed by  $\{1, 2, 3, 4, 5, 6, 7, 8, 9\}$ . As an example,  $9 \leq 8$  since  $8 \leq 9$  and  $s_0$  and  $s_1$  do not commute. The labeled Hasse diagram for the heap poset is seen in Figure 1.5.




Figure 1.5: Labeled hasse diagram for the heap of an element in  $FC(\widetilde{C}_6)$ .

Let w be a reduced expression for an element  $w \in W(\widetilde{C}_n)$ . As in [1] and [3] we can represent a heap for w as a set of lattice points embedded in  $\{0, 1, 2, \ldots, n\} \times \mathbb{N}$ . To do so, we assign coordinates (not unique)  $(x, y) \in \{0, 1, 2, \ldots, n\} \times \mathbb{N}$  to each entry of the labeled Hasse diagram for the heap of w in such a way that:

- (1) An entry with coordinates (x, y) is labeled  $s_i$  (or i) in the heap if and only if x = i;
- (2) If an entry with coordinates (x, y) is greater than an entry with coordinates (x', y') in the heap then y > y'.

Although the above is specific to  $W(\widetilde{C}_n)$ , the same construction works for any straight line Coxeter graph with the appropriate adjustments made to the label set and assignment of coordinates. Specifically, for type  $A_n$  our label set is  $\{1, 2, \ldots, n\}$  and for type  $B_n$  our label set is  $\{0, 1, \ldots, n-1\}$ .

In the case of any straight line Coxeter graph it follows from the definition that (x,y) covers (x',y') in the heap if and only if  $x=x'\pm 1,\ y'< y$ , and there are no entries (x'',y'') such that  $x''\in\{x,x'\}$  and y'< y''< y. This implies that we can completely reconstruct the edges of the Hasse diagram and the corresponding heap poset from a lattice point representation. The lattice point representation can help us visualize arguments that are potentially complex. Note that in our heaps the entries

fully exposed to the top (respectively, bottom) correspond to the generators occurring in the left (respectively, right) descent set of the corresponding reduced expression. Since there are multiple heap representations when  $w \in W(\Gamma)$  is not FC, it is possible that a block that is fully exposed in one heap may not be fully exposed in a different heap representing w.

Let w be a reduced expression for  $w \in W(\widetilde{C}_n)$ . We denote the lattice representation of the heap poset in  $\{0, 1, 2, \dots n\} \times \mathbb{N}$  described in the preceding paragraphs via  $H(\mathbf{w})$ . If w is FC, then the choice of reduced expression for w is irrelevant and we will often write H(w) and we refer to H(w) as the heap of w. Note that we will use the same notation for heaps in Coxeter groups of all types with straightline Coxeter graphs.

Let  $\mathbf{w} = s_{x_1} s_{x_2} \cdots s_{x_r}$  be a reduced expression for  $w \in W(\widetilde{C}_n)$ . If  $s_{x_i}$  and  $s_{x_j}$  are adjacent generators in the Coxeter graph with i < j, then we must place the point labeled by  $s_{x_i}$  at a level that is above the level of the point labeled by  $s_{x_j}$ . Because generators in a Coxeter graph that are not adjacent do commute, points whose x-coordinates differ by more than one can slide past each other or land in the same level. To emphasize the covering relations of the lattice point representation we will enclose each entry in the heap in a square with rounded corners (called a block) in such a way that if one entry covers another the blocks overlap halfway. In addition, we will also label each square for  $s_i$  with i.

There are potentially many ways to illustrate a heap of an arbitrary reduced expression, each differing by the vertical placement of the blocks. For example, we can place blocks in vertical positions as high as possible, as low as possible, or some combination of low/high. In this thesis, we choose what we view to be the best representation of the heap for each example and when illustrating the heaps of arbitrary reduced expressions we will discuss the relative position of the entries but never the absolute coordinates.

**Example 1.4.2.** Let  $\mathbf{w} = s_6 s_4 s_2 s_5 s_3 s_1 s_4 s_0 s_1$  be a reduced expression for  $w \in \mathrm{FC}(\widetilde{C}_6)$  as seen in Example 1.4.1. Figure 1.6 shows a possible lattice point representation for H(w). Since w is FC this is the unique heap representation for w. Because H(w) is the unique heap we can obtain  $\mathcal{L}(w)$  (respectively,  $\mathcal{R}(w)$ ) from the blocks that are fully exposed to the top (respectively, bottom) of the heap. We see that  $\mathcal{L}(w) = \{s_2, s_4, s_6\}$  and  $\mathcal{R}(w) = \{s_1, s_4\}$ .

**Example 1.4.3.** Let  $w_1 = s_0 s_2 s_4 s_3 s_2 s_1$  be a reduced expression for  $w \in W(\widetilde{C}_4)$ . Applying the commutation move  $s_2 s_4 \mapsto s_4 s_2$ , we can obtain another reduced expression for w, namely  $w_2 = s_0 s_4 s_2 s_3 s_2 s_1$ , which is in the same commutation class as  $w_1$ , and hence has the same heap. However, applying the braid move  $s_2 s_3 s_2 \mapsto s_3 s_2 s_3$ , we obtain another reduced expression  $w_3 = s_0 s_4 s_3 s_2 s_3 s_1$ . Note that since  $w_3$  was obtained by applying a braid move,  $w_3$  is in a different commutation class than  $w_1$  and  $w_2$ .




Figure 1.6: A lattice point representation for the heap of an FC element in  $W(\widetilde{C}_6)$ .

Representations of  $H(w_1)$ ,  $H(w_2)$ , and  $H(w_3)$  are seen in Figure 1.7, where the braid relation is colored in teal. Notice that from the heaps we see that  $\mathcal{L}(w) = \{0, 2, 4\}$  and  $\mathcal{R} = \{s_1, s_3\}$ . However, if we only had one heap or the other, we would miss some elements in the left and right descent sets as  $s_3$  is not fully exposed to the bottom of the heap in Figure ?? and  $s_2$  is not fully exposed to the top of the heap in 1.7(b).




Figure 1.7: Two heaps of a non-FC element in  $W(\widetilde{C}_4)$ .

It will be extremely useful for us to be able to quickly determine whether a heap corresponds to an element in  $FC(B_n)$  or  $FC(\widetilde{C}_n)$ . The next proposition is a special case of [13, Proposition 3.3] and follows quickly when one considers the consecutive subwords that are impermissible in reduced expressions for elements in  $FC(B_n)$  and  $FC(\widetilde{C}_n)$  as discussed in Section 1.3.

**Proposition 1.4.4.** If  $w \in FC(\widetilde{C}_n)$ , then H(w) cannot contain any of the configurations seen in Figure 1.8, where 0 < k < n-1 and we use a square with a dotted boundary to emphasize that no element of the heap may occupy the corresponding position.

Since  $W(B_n)$  is a parabolic subgroup of  $W(\widetilde{C}_n)$ , we can use Figure 1.8 to classify the impermissible configurations for elements of  $FC(B_n)$ . In particular, the impermissible configurations for elements of  $FC(B_n)$  are those seen in Figures 1.8(a), 1.8(b) 1.8(c), and 1.8(d).




Figure 1.8: Impermissible configurations for heaps of  $FC(\widetilde{C}_n)$ .

## Chapter 2

# Star Reductions and Property T

#### 2.1 Star Reductions

The notion of a star operation was originally introduced by Kazhdan and Lusztig in [11] for simply-laced Coxeter systems (i.e.,  $m(s,t) \leq 3$  for all  $s,t \in S$ ), and was later generalized to all Coxeter systems in [12]. If  $I = \{s,t\}$  is a pair of non-commuting generators of a Coxeter group W, then I induces four partially defined maps from W to itself, known as star operations. A star operation, when it is defined, increases or decreases the length of an element to which it is applied by 1. For our purposes it is enough to only define the star operations that decrease the length of an element by 1, and as a result we will not develop the notion in full generality.

Let (W, S) be a Coxeter system of type  $\Gamma$  and let  $I = \{s, t\} \subseteq S$  be a pair of generators with  $m(s, t) \geq 3$ . Let  $w \in W(\Gamma)$  such that  $s \in \mathcal{L}(w)$ . We define w to be left star reducible by s with respect to t if there exists  $t \in \mathcal{L}(sw)$ . We analogously define w to be right star reducible by s with respect to t. Observe that w is left (respectively, right) star reducible if and only if w = stu (respectively, w = uts), where the product on the right hand side of the equation is reduced and  $m(s, t) \geq 3$ . We say that w is star reducible if it is either left or right star reducible.

**Example 2.1.1.** Let  $\mathbf{w} = s_0 s_1 s_0 s_2$  be a reduced expression for  $w \in W(B_4)$ . We see that w is left star reducible by  $s_0$  with respect to  $s_1$  to  $s_1 s_0 s_2$  since  $m(s_0, s_1) = 4$  and  $s_0 \in \mathcal{L}(w)$  while  $s_1 \in \mathcal{L}(s_0 w)$ . Notice that w is FC and  $\mathcal{R}(w) = \{s_2, s_0\}$ . We see that  $ws_2 = s_0 s_1 s_0$  and  $ws_0 = s_0 s_1 s_2$ . Note that in both instances  $s_1 \notin \mathcal{R}(ws_2) = \{s_0\}$  and  $s_1 \notin \mathcal{L}(ws_0) = \{s_2\}$ . Because of this w is not right star reducible.

It may be helpful to visualize star reductions in terms of heaps. Let (W, S) be a Coxeter system with straightline Coxeter graph  $\Gamma$  and let  $I = \{s, t\} \subseteq S$  be a pair of generators with  $m(s, t) \geq 3$ . Suppose w is left star reducible by s with respect to t. Then there exists a heap for w where the block for s is fully exposed to the top

such that removing the block for s off of the top allows for t to now be fully exposed to the top of the heap. Similarly if w is right star reducible by s with respect to t, then there exists a heap for w where the block for s is fully exposed to the bottom of the heap such that removing the block for s off of the bottom allows for t to now be fully exposed to the bottom. Conversely, if a heap for  $w \in W(\Gamma)$  has this property, then w is star reducible. In Figure 2.1 we see the top portion of two possible heap representations of an element that is left star reducible by s with respect to t, where the dotted square represents no block may occupy this space. Notice that flipping the heap upside down in Figure 2.1 will result in a heap that is right star reducible. It is important to note that if the group element we are evaluating for star reducibility is not FC, then we must consider all heap representations for the element before concluding that an element is not star reducible.



Figure 2.1: A visual representation of an element that is left star reducible.

The following example utilizes heaps to show that an element is star reducible.

**Example 2.1.2.** Let  $w = s_0 s_1 s_0 s_2$  be a reduced expression for  $w \in W(B_4)$ . Note that w is FC. By Example 2.1.1 we know that w is left star reducible by  $s_0$  with respect to  $s_1$ . In Figure 2.2(a), we see the heap for w. Notice that the block for  $s_0$  is fully exposed to the top of the heap. Removing the block for  $s_0$  gives the heap in Figure 2.2(b). Notice that the block for  $s_1$  is now fully exposed to the top of the heap. However, notice that the blocks for  $s_0$  and  $s_2$  are fully exposed to the bottom. In removing either of these we are unable to fully expose  $s_1$  to the bottom. Thus we can see that w is not right star reducible.

Notice that if w is not FC, then we are not be able to say that w is not star reducible when viewing a single heap as there could be a different heap for w in which we are able to fully expose an element that was previously blocked.

**Example 2.1.3.** Let  $w_1 = s_3 s_1 s_2 s_1 s_0 s_1 s_3 s_0 s_2 s_4$  be a reduced expression for  $w \in W(\widetilde{C}_3)$ . The heap for w is given in Figure 2.3(a), where we have highlighted a braid in teal. Notice that this heap appears to not be star reducible since if we were to remove the block for  $s_1$  or  $s_3$  we would still not fully expose  $s_2$  to the top of the heap. The same goes for fully exposing blocks in the bottom of the heap. However, when we



Figure 2.2: Visualization of Example 2.1.1.

perform the braid move resulting in the heap seen in Figure 2.3(b) it is now obvious that the element is star reducible. Thus when considering a non-FC element for star reducibility via the heap, it is very important to consider all heaps for that element.




Figure 2.3: Visualization of Example 2.1.3

Using the notion of star reduction we are now able to introduce the concept of a star reducible Coxeter group. Let (W, S) be a Coxeter group of type  $\Gamma$ . We say that  $W(\Gamma)$  is star reducible if every element of  $FC(\Gamma)$  is star reducible to a product of commuting generators. That is,  $W(\Gamma)$  is star reducible if when we apply star reductions on the left or right repeatedly to  $w \in FC(\Gamma)$ , eventually we obtain a product of commuting generators. Notice that we are restricting to just the FC elements in  $W(\Gamma)$ . Visually a star reducible Coxeter group can be thought of in the following way. Given a heap in  $FC(\Gamma)$ , we are able to systematically remove fully exposed blocks from the top or bottom of the heap and have a block that was

previously not fully exposed become fully exposed until we are left with a heap that can be drawn as a single row.

In [9], Green classified all star reducible Coxeter groups.

**Proposition 2.1.4** (Green, [9]). Let (W, S) be a Coxeter system of type  $\Gamma$ . Then (W, S) is star reducible if and only if each component of  $\Gamma$  is either a complete graph with labels  $m(s,t) \geq 3$ , or is one of the following types: type  $A_n$   $(n \geq 1)$ , type  $B_n$   $(n \geq 2)$ , type  $D_n$   $(n \geq 4)$ , type  $F_n$   $(n \geq 4)$ , type  $H_n$   $(n \geq 2)$ , type  $I_2(m)$   $(m \geq 3)$ , type  $\widetilde{A}_n$   $(n \geq 3)$  and n even ), type  $\widetilde{C}_n$   $(n \geq 3)$  and n odd ), type  $\widetilde{E}_6$ , or type  $\widetilde{F}_5$ .  $\square$ 

### 2.2 Property T

In [9], Green utilizes the following theorem to help classify the star reducible Coxeter groups.

**Proposition 2.2.1** (Green, [9]). Let (W, S) be a star reducible Coxeter system of type  $\Gamma$ , and let  $w \in W$ . Then one of the following possibilities occurs for some Coxeter generators s, t, u with  $m(s, t) \neq 2$ ,  $m(t, v) \neq 2$ , and m(s, u) = 2:

- (1) w is a product of commuting generators;
- (2) w has a reduced product w = stu;
- (3) w has a reduced product w = uts;
- (4) w has a reduced product w = svtu.

Notice that Items (2) and (3) indicate an element that is left or right star reducible, respectively. Also notice that an element w that has the form of Item (1) does not meet the conditions of Items (2) and (3). In particular, w is not star reducible if Condition (1). Lastly, notice that if an element w is of the form of Item (4) and not of the form of Items (2)–(3), then w is not star reducible. Notice Items (2), (3), and (4) are not mutually exclusive if w is .

Motivated by Items (1) and (4) above, we define the notions of Property T and T-avoiding. Let (W, S) be a Coxeter system of type  $\Gamma$  and let  $w \in W$ . We say that w has Property T if and only if there exists a reduced product for w such that w = stu or w = uts where  $m(s,t) \geq 3$ . That is, w has Property T if there exists a reduced expression for w that begins or ends with a product of non-commuting generators. An element  $w \in W(\Gamma)$  is called T-avoiding if w does not have Property T. This implies that a T-avoiding element is not star reducible.

Since elements that are star reducible also have Property T we already know how to visualize Property T in terms of heaps. It is clear that a product of commuting generators is T-avoiding, which we state as a proposition.

**Proposition 2.2.2.** Let (W, S) be a Coxeter system of type  $\Gamma$ . If  $w \in W(\Gamma)$  such that w is a product of commuting generators, then w is T-avoiding.

Visually a product of commuting generators is a one row heap, it is clear a one row heap will not portray the characteristic of Property T as seen in Figure 2.1.

We will call an element that is a product of commuting generators trivially T-avoiding. If w is T-avoiding and not a product of commuting generators, we will say that w is non-trivially T-avoiding. It is not clear that such elements exist. Referring back to Green's classification (Proposition 2.2.1) of what elements in star reducible Coxeter groups look like, we see that Item (1) corresponds to an element w being trivially T-avoiding, Items (2) and (3) refer to the element w having Property T on the left and right, respectively and Item (4) refers to an element being non-trivially T-avoiding provided no reduced expression for the element exhibits Items (2) and (3). In star reducible Coxeter systems, every FC element is star reducible to a product of commuting generators, which implies that no FC element can be non-trivially T-avoiding in such groups. For example, as will be seen in the following sections, the Coxeter systems of type  $A_n$  and  $B_n$  have no non-trivial T-avoiding elements, while the Coxeter systems of type  $D_n$  do have non-trivially T-avoiding elements.

**Example 2.2.3.** Let  $w \in W(A_5)$  with reduced expression  $w = s_1 s_3 s_5$ . It turns out that since w is a product of commuting generators, by Proposition 2.2.2 we know that w is trivially T-avoiding.

**Example 2.2.4.** Let  $w \in W(A_5)$  with reduced expression  $\mathbf{w}_1 = s_5 s_3 s_2 s_4 s_1$ . At first glance it may appear that w does not have Property T since both  $s_1$  and  $s_4$  commute as well as  $s_3$  and  $s_5$ . However, note that applying the commutation move  $s_4 s_2 \mapsto s_2 s_4$  results in  $\mathbf{w}_2 = s_1 s_2 s_4 s_3 s_5$ . Hence w has Property T since  $m(s_1, s_2) = 3$  and there is a reduced expression for w that begins with  $s_1 s_2$ . In Figure 2.4 we see the heap for w. Note that we can see Property T in the bottom of the heap highlighted in orange.



Figure 2.4: Heap of an element with Property T.

**Example 2.2.5.** Let  $w \in W(\widetilde{C}_4)$  with reduced expression  $w = s_0 s_2 s_4 s_1 s_3 s_0 s_2 s_4$ . It turns out that w is FC and non-trivially T-avoiding. The heap for w is seen in Figure 2.5. Notice that no matter which block we remove that is fully exposed to




Figure 2.5: Heap of a non-trivially T-avoiding element in  $W(\widetilde{C}_4)$ .

the top of the heap no new element becomes fully exposed. The same applies to the bottom of the heap. Thus, w is non-trivially T-avoiding.

One thing to notice here is that all Coxeter groups have trivially T-avoiding elements as they all contain products of commuting generators. The more interesting non-trivially T-avoiding elements do not appear in all Coxeter groups. In Chapter 3 we will summarize what is known about the T-avoiding elements in Coxeter systems of types  $\widetilde{A}_n$ ,  $A_n$ ,  $D_n$ ,  $F_n$ , and  $I_2(m)$ , and in Chapters 4 and 5 we classify the T-avoiding elements in Coxeter systems of types  $B_n$  and  $\widetilde{C}_n$ .

#### 2.3 Non-Cancellable Elements

We now introduce the concept of weak star reducible, which is related to the notion of cancellable in [5]. Let (W, S) be a Coxeter system of type  $\Gamma$  and let  $I = \{s, t\} \subseteq S$  be a pair of noncommuting generators. If  $w \in FC(\Gamma)$ , then w is left weak star reducible by s with respect to t to sw if

- (1) w is left star reducible by s with respect to t, and
- (2)  $tw \notin FC(\Gamma)$ .

Notice that Condition (2) implies that l(tw) > l(w). Also note that we are restricting out definition of weak star reducible to the set of FC elements of  $W(\Gamma)$ . We analogously define right weak star reducible by s with respect to t to ws. We say that w is weak star reducible if w is either left or right weak star reducible. Otherwise, we say that w is non-cancellable or weak star irreducible. Notice that from this we know that weak star reducible implies star reducible. However, star reducible does not imply weak star reducible.

**Example 2.3.1.** Let  $w = s_0 s_1 s_0 s_2$  be a reduced expression for  $w \in W(B_4)$  as in Example 2.1.1. From Example 2.1.1 we know that w is left star reducible. However,  $tw = s_1 s_0 s_1 s_0 s_2$ , which is not in  $FC(B_4)$ . Thus, we see that w is left weak star

reducible by  $s_0$  with respect to  $s_1$  to  $s_1s_0s_2$ . In addition, Example 2.1.1 showed that w is not right star reducible and hence w is not right weak star reducible. However, since w is left weak star reducible we know that w is not non-cancellable.

Again it might be useful to visualize the concept of weak star reducible in terms of heaps. Recall that in Section 2.1 we described what a star reduction looks like in terms of heap. Since the definition of weak star reducible includes that a heap is star reducible we again need to have those properties. In addition, for a heap to be weak star reducible, adding the block that becomes fully exposed when a block is removed from the heap must create a braid in the heap forcing the new larger heap to not be FC. That is, one of the impermissible configurations seen in Section 1.4 will appear at the top or bottom of the heap.

**Example 2.3.2.** Let  $w = s_0 s_1 s_0 s_2$  be a reduced expression for  $w \in W(B_4)$  as in Example 2.3.1. Recall in Figure 2.2 the heap for w was seen along with what it star reduced to. In Figure 2.6 we see that adding  $s_1$  to the top of the heap creates a braid which is highlighted in orange. Therefore, w is left star reducible by  $s_0$  with respect to  $s_1$ , to  $w = s_1 s_0 s_2$ .




Figure 2.6: Heap of a weak star reducible element of  $FC(B_4)$ .

**Example 2.3.3.** Let  $w \in FC(B_4)$  and let  $w = s_0s_1$  be a reduced expression for w. Note that w is left (respectively, right) star reducible by  $s_0$  with respect to  $s_1$  (respectively, by  $s_1$  with respect to  $s_0$ ). However,  $s_1s_0s_1 \in FC(B_4)$  (respectively,  $s_0s_1s_0 \in FC(B_4)$ ). Thus w is non-cancellable. Visually the heap appears in Figure 2.7. Clearly when  $s_0$  is added to the bottom of the heap, the new heap is still in  $FC(B_4)$  and the same can be said when  $s_1$  is added to the top of the heap.

In [3], Ernst classified the non-cancellable elements in Coxeter groups of type  $W(B_n)$  and  $W(\widetilde{C}_n)$ . We will state part of the classification here as it is important to the development of the non-trivial T-avoiding elements in  $W(\widetilde{C}_n)$ . To see the full classification see [3, Sections 4.2 and 5].



Figure 2.7: Heap of a non-cancellable element of  $FC(B_4)$ .




Figure 2.8: Heap of a single sandwich stack in  $W(\widetilde{C}_n)$ .

Before we state the classification we first define a specific group element in  $W(\widetilde{C}_n)$  which we will refer to as a single  $sandwich\ stack$ , which is seen in Figure 2.8.

We can stack single sandwich stacks together and get a sandwich stack. This is done by removing the top most layer of the heap and adding a new single sandwich stack to the stack. A sandwich stack is seen in Figure 2.9. These heaps are referenced in the classification of non-trivial T-avoiding elements in  $W(\widetilde{C}_n)$ .




Figure 2.9: Heap of a stack of sandwich stacks in  $W(\widetilde{C}_n)$ .

In [3], Ernst completely classified the non-cancellable elements in Coxeter systems of types  $B_n$  and  $\widetilde{C}_n$ . In the type  $\widetilde{C}_n$  case, the sandwich stacks are the only T-avoiding elements with full support. There are two other types of non-cancellable elements

that were classified in [3]. The first does not have full support which is important to our later classification. The second clearly has Property-T on the top.

## Chapter 3

# T-Avoiding Elements in Types $\widetilde{A}_n, A_n, D_n, F_n$ , and $I_2(m)$

In this Chapter we classify the non-trivial T-avoiding elements of Coxeter systems in types  $\widetilde{A}_n$ ,  $A_n$ ,  $D_n$ ,  $F_n$  and  $I_2(m)$ .

# 3.1 Types $\widetilde{A}_n$ and $A_n$

We start with T-avoiding elements in Coxeter systems of type  $\widetilde{A}_n$  and  $A_n$ . We first focus on non-trivial T-avoiding elements in  $W(\widetilde{A}_n)$ .

**Proposition 3.1.1.** If  $n \geq 2$  and n is odd, then there are no non-trivial T-avoiding elements in  $W(\widetilde{A}_n)$ . Otherwise, if  $n \geq 2$  and n is even then  $W(\widetilde{A}_n)$  contains non-trivial T-avoiding elements.

*Proof.* This is [6, Proposition 3.1.2] after a translation of terminology.

The classification seen in [6] did not specifically classify the non-trivial T-avoiding elements for type  $\widetilde{A}_n$  for n odd. Since  $W(\widetilde{A}_n)$  for n odd is not star reducible, the non-trivial T-avoiding elements could be FC. The following is our conjecture regarding what the non-trivial T-avoiding elements are in  $W(\widetilde{A}_n)$  for n odd.

**Conjecture 3.1.2.** The only non-trivial T-avoiding elements in  $W(\widetilde{A}_n)$  for n odd are of the form  $w = (s_0 s_2 \cdots s_{n-2} s_n s_1 s_3 \cdots s_{n-3} s_{n-1})^k$  for  $k \in \mathbb{Z}^+$ .

Recall that  $W(\widetilde{A}_n)$ , for n even, is not a star reducible Coxeter group. Hence it makes sense that the T-avoiding elements in  $W(\widetilde{A}_n)$ , for n even, can be FC. Further, as  $W(A_n)$  is a parabolic subgroup of  $W(\widetilde{A}_n)$  and  $W(A_n)$  is a star reducible Coxeter group, any FC non-trivial T-avoiding elements must have full support. First notice, that  $w = (s_0 s_2 \cdots s_{n-2} s_n s_1 s_3 \cdots s_{n-3} s_{n-1})^k$  is reduced, FC, and has full support. In

addition, w is in fact T-avoiding. As stated in the conjecture we believe that these are the only non-trivial T-avoiding elements. However, it is not immediately obvious that there are not any non-FC non-trivial T-avoiding elements. Classifying these non-trivial T-avoiding elements remains an open problem. We now proceed with the classification of T-avoiding elements in Coxeter groups of type  $A_n$ .

Corollary 3.1.3. There are no non-trivially T-avoiding elements in  $W(A_n)$ .

*Proof.* Notice that the Coxeter graph of type  $A_n$  can be obtained from the Coxeter graph of type  $\widetilde{A}_k$ , for k > n. This is done by removing the appropriate number of vertices and edges from the Coxeter graph of type  $\widetilde{A}_k$ . Since  $W(\widetilde{A}_k)$  for k even has no non-trivial T-avoiding elements this forces  $W(A_n)$  to not have non-trivial T-avoiding elements.  $\square$ 

## 3.2 Type $D_n$

In this section summarize the classification of the T-avoiding elements in Coxeter systems of type  $D_n$ . Recall that  $W(D_n)$  is a star reducible Coxeter group and as a result any potential non-trivial T-avoiding elements are not FC.

**Proposition 3.2.1.** The are non-trivial T-avoiding elements in  $W(D_n)$  for  $n \geq 4$ .

We now will classify these elements as seen in [8]. Before we do so we state interval notation useful to the classification from [8, Definition 2.3.1]. For  $2 \le i \le j$  denote the element  $s_i s_{i+1} \cdots s_{j-1} s_j$  by [i,j]. For  $i \ge 3$ , denote  $s_1 s_3 s_4 \cdots s_i$  by [1,i] and for  $j \ge 2$  denote  $s_1 s_2 s_3 \cdots s_j$  by [0,j]. If  $0 \le j < i$  and  $i \ge 2$  define  $[j,i] = [i,j]^{-1}$ . Finally, for  $i \le -3$  and  $j \ge 3$  denote  $s_1 s_{i-1} s_{i-2} \cdots s_4 s_3 s_2 s_3 s_4 \cdots s_j$  by [-i,j]. The following determines the classification for T-avoiding elements in  $W(D_n)$ .

**Proposition 3.2.2.** Let  $w \in W(B_n)$  be non-trivially T-avoiding. Then  $w = w_n u$  reduced for some  $m \leq n$ , where u is a product of commuting generators such that  $\sup(u) \subseteq s_{m+2}, s_{m+3}, s_{m+4}, \ldots s_n$  and

$$w_n = \begin{cases} [2,0][4,0] \cdots [n-2,0][n,0][n-k,n-2k] \cdots [n-1,n-2][n,n] \\ [2,0][4,0] \cdots [m-2,0][m,0][m-k,m-2k] \cdots [m-1,m-2][m,m] \end{cases}$$

where the first case above corresponds to n even, the second case corresponds to n odd, m = n - 1 and

$$k = \begin{cases} \frac{n}{2} - 2 & \text{if, } n \text{ is even} \\ \frac{n-1}{2} - 2 & \text{if, } n \text{ is odd.} \end{cases}$$

*Proof.* This is [8, Lemmas 2.2.18 and 2.3.4]. Although it is not immediately obvious,  $w_n$  is reduced and not FC.

In Figure 3.1, we see two different elements that are T-avoiding in  $W(D_5)$ . Notice that the blocks that are highlighted in red alternate, this prevents the teal highlighted braid from forcing its way to the top or the bottom of the heap. Due to the fork in the graph we must make slight alterations to heaps for  $W(D_n)$ . Specifically we allow  $s_0$  and  $s_1$  to occupy the same horizontal placement.




Figure 3.1: Visual representation of non-trivial T-avoiding elements in  $W(D_5)$ .

## 3.3 Type $F_n$

In this section we classify what is known regarding the T-avoiding elements in the Coxeter systems of type  $F_n$  for  $n \geq 4$ . Note that all of the following results are unpublished.

We start with the Coxeter system of type  $F_5$ . Recall that  $W(F_5)$  is a star reducible Coxeter group so any non-trivial T-avoiding elements will not be FC. Before we begin the classification we introduce the notion of a specific element in  $W(F_5)$  called a bowtie, which is given by the heap in Figure 3.2. Note that in Figure 3.2(a), the orange blocks correspond to the elements that have bond strength 4. It turns out that the expression determined by this heap is in fact reduced. Looking at the heap in Figure 3.2(b), we have highlighted a braid in teal. We can obtain a "stack of bowties"

by removing the top most layer of the given heap for the bowtie and adding a new single bowtie to the stack. Doing this repeatedly results in the heap seen in Figure 3.3. Similar to a single bowtie, the expression that corresponds to a stack of bowties is reduced and not FC. These heaps are referenced in the following unpublished theorem by Cross, Ernst, Hills-Kimball, and Quaranta in 2012, which classifies the T-avoiding elements in the Coxeter systems of type  $F_5$ .




Figure 3.2: Heap of a single bowtie in  $W(F_5)$ .

**Proposition 3.3.1.** The only non-trivial T-avoiding elements in  $W(F_5)$  are stacks of bowties.

As a result of the classification in type  $F_5$ , Cross et al. were also able to classify the T-avoiding elements in  $W(F_4)$ .

Corollary 3.3.2. There are no non-trivial T-avoiding elements in the Coxeter system of type  $F_4$ .

*Proof.* Since there are no non-trivial T-avoiding elements in  $W(F_5)$  that do not have full support, we know that there are not any non-trivial T-avoiding elements in  $W(F_4)$ . Because if there were non-trivial T-avoiding elements they would also be non-trivially T-avoiding in  $W(F_5)$ .

Cross et al. conjectured that in Coxeter systems of type  $F_n$  for  $n \geq 5$ , an element is non-trivially T-avoiding if and only if it is a stack of bowties multiplied by a product of commuting generators. In 2013, Gilbertson and Ernst worked with this conjecture and quickly found it to be false. The heap seen in Figure 3.4 corresponds to a non-trivial T-avoiding element in Type  $F_6$  that is not a bowtie. It turns out that like the bowties discussed above these elements can also be stacked to create an infinite




Figure 3.3: Heap of a stack of bowties in  $W(F_5)$ .

number of non-trivial T-avoiding elements. In addition, as n gets large there are a number of modifications that can be made that result in additional non-trivially T-avoiding elements. From this we conjecture that the classification of T-avoiding elements in Coxeter systems of type  $F_n$  for  $n \geq 6$  gets complicated very quickly. Classifying T-avoiding elements in  $W(F_n)$  for  $n \geq 6$  remains an open problem.




Figure 3.4: Heap of a non-trivial T-avoiding element in  $W(F_6)$ 

## **3.4 Type** $I_2(m)$

We next classify the T-avoiding elements in Coxeter systems of type  $I_2(m)$ . Note that in Coxeter systems of type  $I_2(m)$ , the only products of commuting generators have length 1. Although the following is a quick result, we believe that it does not already appear in the literature.

**Theorem 3.4.1.** There are no non-trivial T-avoiding elements in Coxeter systems of type  $I_2(m)$ .

Proof. The graph for the Coxeter system of  $I_2(m)$  appears in Figure 1.2(c). Note that the graph consists of two vertices, namely,  $s_1$  and  $s_2$ , and a single edge with weight m. Also, recall that  $W(I_2(m))$  is a star reducible Coxeter group. This implies that any non-trivial T-avoiding elements in  $W(I_2(m))$  must not be FC, as all of the FC elements have Property T or are trivially T-avoiding. The only non-FC element in  $W(I_2(m))$  is the element of length m that has exactly two reduced expressions consisting of alternating products of  $s_1$  and  $s_2$ . Cleary, this element begins and ends with a product of noncommuting generators. Thus, this element has Property T. Hence  $W(I_2(m))$  has no non-trivial T-avoiding elements.

## Chapter 4

# T-Avoiding Elements in Type $B_n$

In this section we classify the T-Avoiding elements in Coxeter systems of type  $B_n$ . We start by introducing some combinatorial tools for type  $B_n$  and then finish with a proof of the classification in type  $B_n$ . Note that the proof for Coxeter systems of type  $B_n$  closely follows the classification of T-avoiding elements of type  $D_n$  seen in [8].

#### 4.1 Tools for the Classification

Recall from Example 1.2.1 that  $W(B_n) \cong \operatorname{Sym}_n^B$ . This implies that we can represent each element  $w \in W(B_n)$  as a signed permutation. That is we can write  $w \in W(B_n)$  using one-line notation

$$w = [w(1), w(2), \dots, w(n-1), w(n)],$$

where we write a bar underneath a number in place of a negative sign in order to simplify notation. In particular, for each  $s_i \in S(B_n)$ , i = 1, 2, ..., n-1, we have

$$s_i = [1, 2, \dots, i-1, i+1, i, i+2, \dots, n-1, n]$$

and we identify  $s_0 \in S(B_n)$  via

$$s_0 = [\underline{1}, 2 \dots, n].$$

Further w(-i) = -w(i) for  $|i| \in \{1, 2, ..., n\}$ . The following propositions provide insight into what happens to a given on-line notation when we multiply by  $s_i$  on the right or the left.

**Proposition 4.1.1.** Let  $w \in W(B_n)$  with corresponding signed permutation

$$w = [w(1), w(2), \dots w(n)].$$

Suppose  $s_i \in S(B_n)$ . If  $i \ge 1$ , then multiplying w on the right by  $s_i$  has the effect of interchanging w(i) and w(i+1) in the one-line notation. If i = 0, then multiplying w on the right by  $s_i$  has the effect of switching the sign of w(1).

*Proof.* This follows from [2, Section 8.1 and A3.1].  $\square$ 

**Proposition 4.1.2.** Let  $w \in W(B_n)$  with corresponding signed permutation

$$w = [w(1), w(2), \dots w(n)].$$

Suppose  $s_i \in S(B_n)$ . If  $i \geq 1$ , then multiplying on the left by  $s_i$  has the effect of interchanging the entries whose absolute values are i and i+1 in the one-line notation. If i = 0, then multiplying w on the left by  $s_i$  has the effect of switching the sign of the entry whose absolute value is 1.

*Proof.* This follows from [2, Section 8.1 and A3.1].  $\square$ 

Suppose  $w \in W(B_n)$  has reduced expression  $w = s_{x_1} s_{x_2} \cdots s_{x_n}$ . We may construct the signed permutation of w from left to right as it is the easier way to multiply based upon the above propositions. We provide an example of this construction below.

**Example 4.1.3.** Let  $w \in W(B_6)$  with a given reduced expression  $\mathbf{w} = s_0 s_1 s_3 s_4 s_5 s_2$ . Then we iteratively build the signed permutation as follows. First,  $s_0 = [-1, 2, 3, 4, 5, 6]$  by definition. Next  $s_0 s_1 = [2, -1, 3, 4, 5, 6]$  since multiplying by  $s_1$  on the right hand side switches the values in position 1 and position 2. Repeating this we get  $s_0 s_1 s_3 = [2, -1, 4, 3, 5, 6]$  and ultimately we end with w = [2, 4, -1, 5, 6, 3].

Notice that if we were to construct the signed permutation for w from right to left, we would start with  $s_2 = [1, 3, 2, 4, 5, 6]$ . Next we would have  $s_5 s_2 = [1, 3, 2, 4, 6, 5]$ . However,  $s_4 s_5 s_2 = [1, 3, 2, 5, 6, 4]$ . Notice this time we were not able to just switch w(i) and w(i + 1) instead we found 4 and 5 and switched their relative positions, which is more difficult than constructing the one-line notation notation left to right, which is why we choose to construct left to right.

Given the one-line notation for an element  $w \in W(B_n)$  we can easily calculate the left and right descent sets of w. The following proposition explains how.

**Proposition 4.1.4.** Let  $w \in W(B_n)$ . Then

$$\mathcal{R}(w) = \{ s_i \in S \mid w(i) > w(i+1) \}$$

where w(0) = 0 by definition.

*Proof.* This is [2, Proposition 8.1.2].

We now will introduce the concept of signed pattern avoidance, which will help with the classification of the T-avoiding elements in Coxeter systems of type  $B_n$ . Our approach mimics the one found in [8]. Let  $w \in W(B_n)$ . We say that w avoids the consecutive pattern abc if there is no  $i \in \{1, 2, ..., n-2\}$  such that (w(i), w(i+1), w(i+2)) is in the same relative order as (a, b, c), where typically one takes  $\{a, b, c\} = \{1, 2, 3\}$ . We say that w avoids the signed consecutive pattern abc if there is no  $i \in \{1, 2, ..., n-2\}$  such that (|w(i)|, |w(i+1)|, |w(i+2)|) is in the same consecutive order as (|a|, |b|, |c|) and such that  $\operatorname{sgn}(w(i)) = \operatorname{sgn}(a)$ ,  $\operatorname{sgn}(w(i+1)) = \operatorname{sgn}(b)$ , and  $\operatorname{sgn}(w(i+2)) = \operatorname{sgn}(c)$ .

**Example 4.1.5.** Let  $w \in W(B_4)$  with signed permutation

$$w = [2, 4, 1, 3].$$

We see that w has the signed consecutive pattern  $\underline{231}$ , since (|w(1)|, |w(2)|, |w(3)|) are in the same relative order as (|-2|, |3|, |-1|), and  $\operatorname{sgn}(w(1)) = \operatorname{sgn}(-2)$ ,  $\operatorname{sgn}(w(2)) = \operatorname{sgn}(3)$ , and  $\operatorname{sgn}(w(3)) = \operatorname{sgn}(-1)$ . However, w avoids the signed consecutive pattern  $1\underline{2}3$ .

Occasionally, we will need to factor a reduced expression  $\mathbf{w}$  for  $w \in W(B_n)$  in a specific manner. Let  $I = \{s, t\}$  for  $s, t \in S(B_n)$ . Define  $w^I$  as the set of all  $w \in W(B_n)$  such that  $\mathcal{L}(w) \cap I = \emptyset$  and define  $W_I = \langle s, t \rangle$ . In [10], it is shown that any element  $w \in W(B_n)$  can be written as  $w = w_I w^I$  reduced where  $w_I \in W_I$  and  $w^I \in W^I$ .

## 4.2 Classification of T-Avoiding Elements in Type $B_n$

Time permitting we will streamline the corollaries into the lemmas themselves or as a remark at the end of the lemmas

In this section we will classify the T-avoiding elements in Coxeter groups of type  $B_n$ . Our main result in this section is Theorem 4.2.17. First we need some preparatory lemmas.

**Lemma 4.2.1.** Let  $s, t \in S(B_n)$  such that m(s, t) = 3. Then w has a reduced expression ending in sts if and only if w has the consecutive pattern 321.

Proof. Let  $i \geq 1$ , let  $I = \{s_i, s_{i+1}\}$  and write  $w = w^I w_I$ . Note that since  $m(s_i, s_{i+1}) = 3$ ,  $s_0 \notin I$ . Observe that if w has a reduced expression ending in the product of two non-commuting generators  $s_i s_{i+1}$  or  $s_{i+1} s_i$ , then we have  $w_I \in \{s_i s_{i+1}, s_{i+1} s_i\}$ .

Suppose w has the consecutive pattern 321. Then there is some i such that w(i) > w(i+1) > w(i+1). By Proposition 4.1.4,  $s_i, s_{i+1} \in \mathcal{R}(w)$ . Since  $W(B_n)$  is a simply laced Coxeter group, w ends in  $s_i s_{i+1} s_{i+1}$  or  $s_{i+1} s_i s_{i+1}$ .

Conversely, suppose w ends in  $s_i s_{i+1} s_i$ . This implies that either  $w_I = s_i s_{i+1}$  or  $w_I = s_{i+1} s_i$  which implies that  $s_i, s_{i+1} \in \mathcal{R}(w)$ . Since  $s_i, s_{i+1} \in \mathcal{R}(w)$ , we see that w(i) > w(i+1) > w(i+2) by Proposition 4.1.4. Thus w has the consecutive pattern 321. Therefore, w has a reduced expression ending in sts if and only if w has the consecutive pattern 321.

Corollary 4.2.2. Let  $s, t \in S(B_n)$  such that m(s, t) = 3. Then w has a reduced expression beginning with sts if and only if  $w^{-1}$  has the consecutive pattern 321.

Proof. Let  $s, t \in S(B_n)$  such that m(s, t) = 3, and  $s_0 \notin \{s, t\}$ . We know that w has no reduced expressions beginning with sts if and only if  $w^{-1}$  has no reduced expression ending with sts which by Theorem 4.2.2 happens only if  $w^{-1}$  avoids the consecutive pattern 321.

**Lemma 4.2.3.** If  $i \neq 0$ , then w has a reduced expression ending in  $s_i s_{i+1}$  if and only if w has the consecutive pattern 231.

Proof. Suppose that w has the consecutive pattern 231. Then there is some i such that w(i+1) > w(i) > w(i+2). By Proposition 4.1.4,  $s_{i+1} \in \mathcal{R}(w)$ . Now multiplying on the right by  $s_{i+1}$  we see that  $ws_{i+1}(i+1) = w(i+2)$  and  $ws_{i+1}(i) = w(i)$ . We know that w(i+2) < w(i), which implies that  $s_i \in \mathcal{R}(ws_{i+1})$ , and hence w has a reduced expression that ends in  $s_is_{i+1}$ .

Conversely, suppose that w has a reduced expression ending in  $s_i s_{i+1}$ . Then w(i+2) < w(i+1) and w(i) < w(i+1). Since  $s_i \in \mathcal{R}(w s_{i+1})$  we have  $w(i+2) = w s_{i+1}(i+1) < w s_{i+1}(i) = w(i)$ . Thus we have that w(i+1) > w(i) > w(i+2). Hence w has the consecutive pattern 231.

Therefore, w has a reduced expression ending in st if and only if w has the consecutive pattern 231.

Corollary 4.2.4. If  $i \neq 0$ , then w has a reduced expression beginning with  $s_i s_{i+1}$  if and only if  $w^{-1}$  has the consecutive pattern 231.

*Proof.* Let  $s, t \in S(B_n)$  such that m(s, t) = 3, and  $s_0 \notin \{s, t\}$ . We know that w has no reduced expressions beginning with st if and only if  $w^{-1}$  has no reduced expression ending with st which by Theorem 4.2.2 happens only if  $w^{-1}$  avoids the consecutive pattern 231.

**Lemma 4.2.5.** If  $i \neq 0$ , then w has a reduced expression ending in  $s_{i+1}s_i$  if and only if w has the consecutive pattern 312.

*Proof.* Suppose that w has the consecutive pattern 312. Then there is some i such that w(i) > w(i+2) > w(i+1). By Proposition 4.1.4 we see that  $s_i \in \mathcal{R}(w)$ . Multiplying on the right by  $s_i$  we get  $ws_i(i+1) = w(i)$  and  $ws_i(i+2) = w(i+2)$ .

By above w(i) > w(i+2), and by Proposition 4.1.4  $s_{i+1} \in \mathcal{R}(ws_i)$ . This implies that w has a reduced expression ending in  $s_{i+1}s_i$ .

Conversely suppose w ends in a reduced expression with  $s_{i+1}s_i$ . Then  $w_I = s_{i+1}s_i$ . We see that w(i) > w(i+1) and w(i+2) > w(i+1). Since  $s_{i+1} \in \mathcal{R}(ws_i)$ , we have  $w(i+2) = ws_i(i+2) < ws_i(i+1) = w(i)$ . From this we have w(i) > w(i+2), so w(i) > w(i+2) > w(i+1). Hence, w has the consecutive pattern 312.

Therefore, w has a reduced expression ending in ts if and only if w has the consecutive pattern 312.

Corollary 4.2.6. If  $i \neq 0$ , then w has a reduced expression beginning with  $s_{i+1}s_i$  if and only if  $w^{-1}$  has the consecutive pattern 312.

Proof. Let  $s, t \in S(B_n)$  such that m(s,t) = 3, and  $s_0 \notin \{s,t\}$ . We know that w has no reduced expressions beginning with ts if and only if  $w^{-1}$  has no reduced expression ending with ts which by Theorem 4.2.2 happens only if  $w^{-1}$  avoids the consecutive pattern 312.

**Lemma 4.2.7.** Let  $w \in W(B_n)$ . Then w has a reduced expression ending in  $s_1s_0$  if and only if w(0) > w(1) and -w(1) > w(2).

Proof. Suppose  $w \in W(B_n)$  such that w ends with  $s_1s_0$ . Then  $s_0 \in \mathcal{R}(w)$  and  $s_1 \in \mathcal{R}(ws_0)$ . This implies that  $ws_0(1) > ws_0(2)$  by Proposition 4.1.4. We see that  $ws_0(1) = w(-1) = -w(1)$  and  $ws_0(2) = 2$ . Hence  $-w(1) = ws_0(1) > ws_0(2) = w(2)$ . Further, since  $s_0 \in \mathcal{R}(w)$ , we see that w(0) > w(1).

Conversely, suppose  $w \in W(B_n)$  such that w(0) > w(1) and -w(1) > w(2). Since w(0) > w(1) so  $s_0 \in \mathcal{R}(w)$ . Multiplying on the right by  $s_0$  we see that  $ws_0(1) = -w(1)$  and  $ws_0(2) = w(2)$ . Note that since  $ws_0(1) = -w(1) > w(2) = ws_0(2)$ ,  $s_1 \in \mathcal{R}(ws_0)$ . Thus w ends with  $s_1s_0$ .

Therefore, w has a reduced expression ending in  $s_1s_0$  if and only if w(0) > w(1) and -w(1) > w(2).

Corollary 4.2.8. Let  $w \in W(B_n)$ . Then w has a reduced expression beginning in  $s_0s_1$  if and only if  $w^{-1}(0) > w^{-1}(1)$  and  $-w^{-1}(1) > w^{-1}(2)$ .

*Proof.* Let  $w \in W(B_n)$ . We know that w has no reduced expressions beginning in  $s_0s_1$  if and only if  $w^{-1}$  has no reduced expressions ending in  $s_0s_1$ . By Lemma 4.2.7 we know that this occurs if and only if  $w^{-1}(0) > w^{-1}(1)$  and  $-w^{-1}(1) > w^{-1}(2)$ .  $\square$ 

**Lemma 4.2.9.** Let  $w \in W(B_n)$ . Then w has a reduced expression ending in  $s_0s_1$  if and only if w(0) > w(2) and w(1) > w(2).

Proof. Suppose  $w \in W(B_n)$  such that w ends with  $s_0s_1$ . Then  $s_1 \in \mathcal{R}(w)$  and  $s_0 \in \mathcal{R}(ws_1)$ . Then  $ws_1(0) > ws_1(1)$ . We see that  $ws_1(0) = 0$  and  $ws_1(1) = w(2)$ . This implies that  $0 = ws_1(0) > ws_1(1) = 2$ . Further, since  $s_1 \in \mathcal{R}(w)$  this implies that w(1) > w(2). Thus if w ends with  $s_0s_1$ , then w(1) > w(2) and w(0) > w(2).

Conversely, suppose  $w \in W(B_n)$  such that w(1) > w(2) and w(0) > w(2). This implies that  $s_1 \in \mathcal{R}(W)$ . Multiplying w on the right by  $s_1$  we see that  $ws_1(0) = w(0)$  and  $ws_1(1) = w(2)$ . Note that since  $ws_1(0) = w(0) > w(2) = ws_1(1)$ ,  $s_0 \in \mathcal{R}(ws_1)$ . Thus w ends with  $s_0s_1$ .

Therefore, w has a reduced expression ending in  $s_0s_1$  if and only if w(1) > w(2) and w(0) > w(2).

Corollary 4.2.10. Let  $w \in W(B_n)$ . Then w has a reduced expression beginning in  $s_1s_0$  if and only if  $w^{-1}(0) > w^{-1}(2)$  and  $w^{-1}(1) > w^{-1}(2)$ .

*Proof.* Let  $w \in W(B_n)$ . We know that w has no reduced expressions beginning in  $s_1s_0$  if and only if  $w^{-1}$  has no reduced expressions ending in  $s_1s_0$ . By Lemma 4.2.7 we know that this occurs if and only if  $w^{-1}(0) > w^{-1}(2)$  and  $w^{-1}(1) > w^{-1}(2)$ .

**Lemma 4.2.11.** Let  $w \in W(B_n)$  such that each entry for w in the one-line notation is positive and both w and  $w^{-1}$  avoid the consecutive patterns 321, 231, and 312, then w is a product of commuting generators.

*Proof.* This follows from an appropriate translation of [8, Lemma 2.2.9].

**Lemma 4.2.12.** Let  $w \in W(B_n)$  be trivially T-avoiding and let  $i \in \{1, 2, ..., n\}$ . Then w satisfies all the following conditions:

- (1)  $w(j) > \min\{w(i-1), w(i)\}\$  for all j > i;
- (2)  $w(k) < \max\{w(i-1), w(i)\}\$  for all k < i-1;
- (3) If w(i), w(i+1) > 0, then w(j) > 0 for all  $j \ge i$ ;
- (4) If w(i), w(i+1) < 0, then w(j) < 0 for all  $j \le i+1$ .

Proof. Suppose there is some least j > i such that  $w(j) \le \min\{w(i-1), w(i)\}$ . Note that j > i so  $j \ne i$ , and  $j \ne i-1$  so  $w(j) < \min\{w(i-1), w(i)\}$ . Note that j is the least so  $w(j-2) \ge \min\{w(i+1), w(i)\} > w(j)$ . This implies that either w(j-1) > w(j-2) > w(j) or w(j-2) > w(j-1) > w(j), which implies w has the consecutive pattern 231 or 321, which is a contradiction to w being a non-trivial T-avoiding element by Lemmas 4.2.1 and 4.2.5. Thus proving (1).

Suppose there exists a maximal k < i - 1 such that  $w \ge \max\{w(i-1), w(i)\}$ . Note that k < i - 1 so  $k \ne i$  and  $k \ne i - 1$ . Then  $w(k) > \max\{w(i-1), w(i)\}$ . Since

k is maximal  $w(k+1) \leq \max\{w(i-1), w(i)\}$  and  $w(k+2) \leq \max\{w(i-1), w(i)\}$ . This implies that either w(k+2) < w(k+1) < w(k) or w(k+1) < w(k+2) < w(k), which implies w has the consecutive pattern 321 or 312, which is a contradiction to w being a non-trivial T-avoiding element by Lemmas 4.2.1 and 4.2.3. Thus proving (2).

It is easy to see that assertion (1) implies (3) and assertion (2) implies (4).

**Lemma 4.2.13.** Let  $w \in W(B_n)$  such that w has the consecutive pattern  $\underline{2}31$ . Then w has Property T.

*Proof.* Let  $w \in W(B_n)$  such that w has the consecutive pattern <u>2</u>31.

Case (1): Suppose w has the one-line notation  $w = [\underline{2}, 3, 1]$ . This implies that  $w = s_1 s_0 s_2$ . Clearly, w begins with a product of non-commuting generators. Thus w has Property T.

Case (2): Suppose that w has the one-line notation  $w = [\underline{a}, b, c, *, ..., *]$  where  $\underline{a}bc$  corresponds to the signed consecutive pattern  $\underline{2}31$ , and \* indicates unknown values for w(i) for i = 4, 5, ..., n. We now consider the possible signed consecutive pattern bc\*. The following are the possibilities:  $312, 31\underline{2}, 321, 32\underline{1}, 213$ , or  $21\underline{3}$ . We know that b and c must be positive since they are positive in w and we also know that b > c by the original signed consecutive pattern. Note that by Lemmas 4.2.1, 4.2.3, and 4.2.7 all of these patterns imply that w ends or begins with a product of noncommuting generators. Thus w has Property T.

Case (3): Suppose that w has the one-line notation  $w = [*, ..., *, \underline{a}, b, c]$  where  $\underline{abc}$  corresponds to the signed consecutive pattern  $\underline{2}31$ , and \* indicates unknown values for w(i) for i = 1, 2, ..., n - 3. We now consider the possible signed consecutive pattern  $*\underline{ab}$ . The following are the possibilities:  $1\underline{2}3$ ,  $\underline{1}23$ ,  $\underline{2}13$ ,  $\underline{2}13$ ,  $\underline{3}12$ , or  $\underline{3}12$ . Note that by Lemmas 4.2.3, 4.2.7, and 4.2.9 all of these patterns implies that w ends or begins with a product of noncommuting generators. Thus w has Property T.

Case (4): Suppose that w has the one-line notation  $w = [*, ..., *, \underline{a}, b, c, *, ..., *]$  where  $\underline{a}bc$  corresponds to the signed consecutive pattern  $\underline{2}31$ , and \* indicates unknown values for w(i) for  $|w(i)| \neq a, b, c$ . In this case we can apply either Case (2) or Case (3) and we can conclude that w ends or begins with a product of noncommuting generators. Thus w has Property T.

Therefore, if  $w \in W(B_n)$  contains the consecutive pattern 231, then w has Property T.

**Lemma 4.2.14.** Let  $w \in W(B_n)$  such that w has the consecutive pattern  $\underline{231}$ . Then w has Property T.

*Proof.* Let  $w \in W(B_n)$  such that w has the consecutive pattern  $\underline{231}$ .

Case (1): Suppose w has the one-line notation  $w = [\underline{2}, 3, \underline{1}]$ . This implies that  $w = s_0 s_1 s_0 s_2$ . Clearly, w begins with a product of non-commuting generators. Thus w has Property T.

- Case (2): Suppose that w has the one-line notation  $w = [\underline{a}, b, \underline{c}, *, \dots, *]$  where  $\underline{abc}$  corresponds to the signed consecutive pattern  $\underline{231}$ , and \* indicates unknown values for w(i) for  $i = 4, 5, \dots n$ . We now consider the possible signed consecutive pattern  $\underline{bc}$ . The following are the possibilities:  $\underline{312}$ ,  $\underline{312}$ ,  $\underline{321}$ ,  $\underline{321}$ ,  $\underline{213}$ , or  $\underline{213}$ . We know that b must be positive since it is positive in w, c must be negative since it is negative in w, and we also know that |b| > |c| by the original signed consecutive pattern. Note that by Lemmas 4.2.1, 4.2.3, and 4.2.7 all of these patterns imply that w ends or begins with a product of noncommuting generators. Thus w has Property T.
- Case (3): Suppose that w has the one-line notation  $w = [*, \ldots, *, \underline{a}, b, \underline{c}]$  where  $\underline{abc}$  corresponds to the signed consecutive pattern  $\underline{231}$ , and \* indicates unknown values for w(i) for  $i = 1, 2, \ldots, n-3$ . We now consider the possible signed consecutive pattern \*ab. The following are the possibilities:  $1\underline{23}$ ,  $1\underline{23}$ ,  $1\underline{23}$ ,  $1\underline{23}$ ,  $1\underline{23}$ ,  $1\underline{23}$ , or  $1\underline{23}$ . We know that  $1\underline{23}$  must be negative,  $1\underline{23}$  must be positive and  $1\underline{23}$  by the original signed permutation. Note that by Lemmas 4.2.3, 4.2.7, and 4.2.9 all of these patterns implies that  $1\underline{23}$  or  $1\underline{23}$  by the original signed permutation. Note that by Lemmas 4.2.3, 4.2.7, and 4.2.9 all of these patterns implies that  $1\underline{23}$  or  $1\underline{23}$  by the original signed permutation.
- Case (4): Suppose that w has the one-line notation  $w = [*, ..., *, \underline{a}, b, \underline{c}, *, ..., *]$  where  $\underline{a}b\underline{c}$  corresponds to the signed consecutive pattern  $\underline{2}3\underline{1}$ , and \* indicates unknown values for w(i) for  $|w(i)| \neq a, b, c$ . In this case we can apply either Case (2) or Case (3) and we can conclude that w ends or begins with a product of noncommuting generators. Thus w has Property T.

Therefore, if  $w \in W(B_n)$  contains the consecutive pattern  $\underline{231}$ , then w has Property T.

**Lemma 4.2.15.** Let  $w \in W(B_n)$  such that w has the consecutive pattern  $\underline{1}23$ . Then either w has Property T or is a trivial T-avoiding element.

*Proof.* Let  $w \in W(B_n)$  such that w has the consecutive pattern <u>1</u>23.

- Case (1): Suppose w has the one-line notation  $w = [\underline{1}23]$ . This implies that  $w = s_0$ . Clearly, w is a trivial T-avoiding element as it is a single generator.
- Case (2): Suppose that w has the one-line notation  $w = [\underline{a}, b, c, *, ..., *]$  where  $\underline{abc}$  corresponds to the signed consecutive pattern  $\underline{1}23$ , and \* indicates unknown values for w(i) for i = 4, 5, ..., n. We now consider the possible signed consecutive patterns bc\*. The following are the possibilities: 231, 23 $\underline{1}$ , 132, 13 $\underline{2}$ , 123, 12 $\underline{3}$ . We know that b and c are positive, and we also know that |b| < |c| by the original signed consecutive pattern. Note that by Lemmas 4.2.1, 4.2.3, and 4.2.7 all of these patterns imply that w ends or begins with a product of noncommuting generators. Thus w has Property T.
- Case (3): Suppose that w has the one-line notation  $w = [*, ..., *, \underline{a}, b, c]$  where  $\underline{abc}$  corresponds to the signed consecutive pattern  $\underline{1}23$ , and \* indicates unknown values for w(i) for i = 1, 2, ..., n 3. We now consider the possible signed consecutive patterns  $*\underline{ab}$ . The following are the possibilities:  $3\underline{1}2$ ,  $3\underline{1}2$ ,  $2\underline{1}3$ ,  $2\underline{1}3$ ,  $1\underline{2}3$ , or  $1, 2\underline{3}3$ .

We know that a must be negative, b must be positive and |a| < |b| by the original signed permutation. Note that by Lemmas 4.2.3, 4.2.7, and 4.2.9 all of these patterns implies that w ends or begins with a product of noncommuting generators. Thus w has Property T.

Case (4): Suppose that w has the one-line notation  $w = [*, ..., *, \underline{a}, b, c, *, ..., *]$  where  $\underline{abc}$  corresponds to the signed consecutive pattern  $\underline{1}23$ , and \* indicates unknown values for w(i) for  $|w(i)| \neq a, b, c$ . In this case we can apply either Case (2) or Case (3) and we can conclude that w ends or begins with a product of noncommuting generators. Thus w has Property T.

Therefore, if  $w \in W(B_n)$  contains the consecutive pattern 123, then w has Property T or is a trivial T-avoiding element.

**Lemma 4.2.16.** Let  $w \in W(B_n)$  such that w has the consecutive pattern  $\underline{1}32$ . Then either w has Property T or is a trivial T-avoiding element.

*Proof.* Let  $w \in W(B_n)$  such that w has the consecutive pattern 132.

Case (1): Suppose w has the one-line notation  $w = [\underline{1}32]$ . This implies that  $w = s_0 s_2$ . Clearly, w is a trivial T-avoiding element as it is a product of commuting generators.

Case (2): Suppose that w has the one-line notation  $w = [\underline{a}, b, c, *, ..., *]$  where  $\underline{a}bc$  corresponds to the signed consecutive pattern  $\underline{1}32$ , and \* indicates unknown values for w(i) for i = 4, 5, ..., n. We now consider the possible signed consecutive pattern bc\*. The following are the possibilities:  $231, 23\underline{1}, 132, 13\underline{2}, 123, \text{ or } 12\underline{3}$ . We know that b and c are positive, and we also know that |b| < |c| by the original signed consecutive pattern. Note that by Lemmas 4.2.1, 4.2.3, and 4.2.7 all of these patterns imply that w ends or begins with a product of noncommuting generators. Thus w has Property T.

Case (3): Suppose that w has the one-line notation  $w = [*, ..., *, \underline{a}, b, c]$  where  $\underline{a}bc$  corresponds to the signed consecutive pattern  $\underline{1}32$ , and \* indicates unknown values for w(i) for i = 1, 2, ..., n - 3. We now consider the possible signed consecutive pattern  $*\underline{a}b$ . The following are the possibilities:  $3\underline{1}2$ ,  $3\underline{1}2$ ,  $2\underline{1}3$ ,  $3\underline{2}1$ , or  $3\underline{2}1$ . We know that a must be negative, b must be positive and |a| < |b| by the original signed permutation. Note that by Lemmas 4.2.3, 4.2.7, and 4.2.9 all of these patterns implies that w ends or begins with a product of noncommuting generators. Thus w has Property T.

Case (4): Suppose that w has the one-line notation  $w = [*, ..., *, \underline{a}, b, c, *, ..., *]$  where  $\underline{abc}$  corresponds to the signed consecutive pattern  $\underline{1}32$ , and \* indicates unknown values for w(i) for  $w(i) \neq a, b, c$ . In this case we can apply either Case (2) or Case (3) and we can conclude that w ends or begins with a product of noncommuting generators. Thus w has Property T.

Therefore, if  $w \in W(B_n)$  contains the consecutive pattern 132, then w has Property T or is a trivial T-avoiding element.

We now are ready to tackle one of the main results of this thesis.

**Theorem 4.2.17.** There are no non-trivial T-avoiding elements in  $W(B_n)$ .

*Proof.* Suppose that  $w \in W(B_n)$  is a non-trivial T-avoiding element. There are  $2^3 \cdot 3! = 48$  possible choices of signed consecutive patterns for w(1)w(2)w(3) where w = [w(1), w(2), w(3), \*, ..., \*].

123	<u>1</u> 23	1 <u>2</u> 3	12 <u>3</u>	<u>12</u> 3	<u>123</u>	1 <u>23</u>	<u>123</u>
132	<u>1</u> 32	1 <u>3</u> 2	132	<u>13</u> 2	<u>132</u>	1 <u>32</u>	<u>132</u>
213	<u>2</u> 13	2 <u>1</u> 3	21 <u>3</u>	<u>21</u> 3	<u>213</u>	2 <u>13</u>	<u>213</u>
231	<u>2</u> 31	2 <u>3</u> 1	23 <u>1</u>	<u>23</u> 1	<u>231</u>	2 <u>31</u>	<u>231</u>
312	<u>3</u> 12	3 <u>1</u> 2	31 <u>2</u>	<u>31</u> 2	<u>312</u>	3 <u>12</u>	<u>312</u>
321	<u>3</u> 21	3 <u>2</u> 1	32 <u>1</u>	<u>32</u> 1	<u>321</u>	3 <u>21</u>	<u>321</u>

We can use Lemma 4.2.1 and Corollary 4.2.2 to eliminate the signed consecutive patterns highlighted in turquoise. We can use Lemma 4.2.5 and Corollary 4.2.4 to eliminate the signed consecutive patterns highlighted in red. We can use Lemma 4.2.3 and Corollary 4.2.6 to eliminate the consecutive patterns highlighted in green. We can use Lemma 4.2.7 and Corollary 4.2.8 to eliminate the signed consecutive patterns highlighted in yellow. We can use Lemma 4.2.9 and Corollary 4.2.10 to eliminate signed consecutive patterns highlighted in brown. We can use Lemmas 4.2.11 to eliminate the signed consecutive patterns highlighted in blue. We can use Lemmas 4.2.13 and 4.2.14 to eliminate signed consecutive patterns highlighted in purple. Finally we can use Lemmas 4.2.15 and 4.2.16 to eliminate signed consecutive patterns highlighted in orange. Since all of the above patterns are eliminated as possibilities for w(1)w(2)w(3) and there are no other signed consecutive patterns that are possible for these positions, w is not be a non-trivial T-avoiding element in the Coxeter group of type  $B_n$ .

The upshot of Theorem 4.2.17 is that the only T-avoiding elements in Coxeter systems of type  $B_n$  are products of commuting generators.

## Chapter 5

# T-Avoiding Elements in Type $\widetilde{C}_n$

# 5.1 Classification of T-Avoiding Elements in Type $\widetilde{C}_n$

In this section we will classify the T-avoiding elements in Coxeter systems of type  $\widetilde{C}_n$ . Since  $W(A_n)$  and  $W(B_n)$  are parabolic subgroups of  $W(\widetilde{C}_n)$  and these groups have no non-trivial T-avoiding elements, any non-trivial T-avoiding elements of  $W(\widetilde{C}_n)$  must have full support. We will first show that there are no non-trivial T-avoiding elements that are not FC in  $W(\widetilde{C}_n)$ .

Before we begin the proof we must first define the notion of a pushed-down representation of a heap. First recall that there are potentially many ways to draw the lattice point representation of a heap, each differing by the amount of vertical space in between blocks. We wish to fix one such representation. Let w be a reduced expression for  $w \in W(\widetilde{C}_n)$ . We construct the pushed-down representation of a heap of H(w) by placing all blocks in the lowest possible vertical position of the heap. Loosely speaking, the heap has been constructed by first giving all blocks fully exposed to the bottom the same vertical position, and then all blocks are as low as possible in the heap. Notice that we can now label the rows in the heap from bottom to top where the bottom-most row is row 1 and proceeding naturally upward from there.

We now define the height of a braid. Given the presence of braid in the heap for an element w, we say that the *height of the braid* is the row number in which the uppermost block involved in the braid is located in the pushed-down representation. It is important to note that in the pushed-down representation a braid may not appear in consecutive rows. That is, some of the blocks may be lower in the heap and the braid may not be immediately apparent.

**Example 5.1.1.** Let  $\mathbf{w} = s_0 s_1 s_3 s_2 s_1 s_0 s_1 s_3$  be a reduced expression for  $w \in W(C_3)$ . The pushed-down representation of a heap for  $\mathbf{w}$  is given in Figure 5.1. The height of the braid  $s_1 s_2 s_1$  which is highlighted in teal in Figure 5.1 is 5 since the upper block for  $s_1$  is located in the fifth row of the pushed-down representation heap. Notice that

the block for  $s_3$  can slide up higher in the heap. If we were to slide the block for  $s_3$  up until it hit the block for  $s_2$  we would obtain the braid  $s_3s_2s_3$  in the heap. In this case, the height for the braid  $s_3s_2s_3$  is also 5.





Figure 5.1: pushed-down representation of a heap


**Theorem 5.1.2.** There are no non-trivial T-avoiding elements in  $W(\widetilde{C}_n)\backslash FC(\widetilde{C}_n)$ .

Proof. We proceed by contradiction. Let  $w \in W(\widetilde{C}_n)\backslash FC(\widetilde{C}_n)$  such that w has full support and w is T-avoiding. Consider all possible pushed-down representation for heaps of w. Choose a representation that has a minimal height braid among all braids appearing in all heaps for w and let k represent the height. That is, choose a heap with the braid with lowest possible braid. There may be a tie, in which case choose your favorite. Without loss of generality we will call the generators involved in the braid  $s_i, s_{i+1}$  where the bond strength will be given in the specific case we are working in. In the following cases whenever we refer to a block being in a specific row, we are considering the pushed-down representation of the heap. However, in order to consider the braids that we are looking for we need to allow some flexibility when referring to the absolute vertical position of a given block.

Case (1): Suppose the lowest braid occurs in the bottom-most row where k=3 (respectively, k=4) if  $m(s_i,s_{i+1})=3$  (respectively,  $m(s_i,s_{i+1})=4$ ). In this case, we are assuming that the braid is located in consecutive rows with the upper-most block in the above specified row and the lowest block in the heap located in the bottom-most row of the heap. Without loss of generality assume  $s_{i+1}$  is in the bottom-most row of the heap. Clearly, the block for  $s_{i-1}$  must be in the bottom-most row of the heap as well, otherwise w has Propety T which is a contradiction to the original choice of w. From this restricting our view to the subheap of w that contains the braid we are considering we see that the heap of w has the following form




where the striped heap block represents the fourth block in the braid if  $m(s_i, s_{i+1}) = 4$ . Applying the braid move we get the subheap seen here




which clearly has Property T since  $s_{i-1}$  is now in the 1st row of the pushed-down representation. This is a contradiction to the way in which we chose w. T For the rest of the cases we will assume that  $k \geq 4$ .

Case (2): Suppose the braid has height k and assume the braid does not contain  $s_0, s_1, s_{n-1}$  or  $s_n$ . Without loss of generality assume  $s_i$  is in the (k)th row of the heap and if necessary we have brought the blocks for  $s_{i-1}$  and  $s_i$  up next to  $s_i$  in row k. We now consider what can be in the (k-3)th row of the heap in two cases.

Subcase (a): Assume that the block for  $s_{i-1}$  is in the heap in (k-3)th and we allow for the block for  $s_{i+1}$  to be in the same row as well, but it does not necessarily have to be. In the following pictures the block for  $s_{i+1}$  will be represented in a purple striped block to indicate that it could be present but it does not have to be. The following is the subheap that we are considering




where we have highlighted the braid in teal. Notice that the block for  $s_{i-2}$  is present in the row (k-2)th. If it was not there, w would have had the braid  $s_{i-1}s_is_{i-1}$  since  $m(s_{i-1}, s_i) = 3$  and we would have had a heap with a lower braid to choose. Applying the braid move to the heap we get the following subheap




which has a new braid in it. This braid which we have highlighted in red for emphasis has height k-1. In applying the braid move we have obtained a subheap which has a braid that has a lower height than our original choice. This is a contradiction to the way in which we chose our heap.

Subcase (b): Assume that the block for  $s_{i+1}$  is in the (k-3)th row of the heap and the block for  $s_{i-1}$  does not appear in the (k-3)th and  $s_{k-2}$  does not appear in the (k-2)th. The following is the subheap we are considering




where the dotted square represents that no block may occupy this space and the braid is highlighted in teal. Applying the braid move in the subheap we obtain the following heap




where the braid now occurs in the (k-1)th. This contradicts the way in which we chose the heap for w. From this we gather that the braid must contain  $s_0, s_1, s_{n-1}$  or  $s_n$ .

Case 3: Suppose the braid is located in the kth row and assume the braid contains  $s_2$  or  $s_{n-2}$ . Without loss of generality we assume that the braid contains  $s_2$  as the other argument is symmetric to the one presented here, and assume  $s_2$  is in the kth row. Notice that if the braid contains  $s_2s_3s_2$ , we are in Case (2), as a result we assume our braid is not of the form  $s_2s_3s_2$ . Assume that if necessary the blocks  $s_1$  and  $s_2$  have been brought up next to  $s_2$  in row  $s_2$ . We now consider what can be in the  $s_2$ -th and  $s_2$ -th rows of the heap in two cases.

Subcase (a): Assume the block for  $s_1$  is in row k-3, and  $s_0$  is in row k-4 but  $s_3$  is not in the (k-3)th row and  $s_2$  is not in the (k-4)th row. Then the subheap we are considering is as follows




where the braid we are considering in the heap is highlighted in teal. Notice that the block for  $s_0$  is in the (k-2)th row. It must be here, as if it was not w would not be reduced. Applying the braid move we get the following heap




where a new braid has appeared highlighted in red. Notice that the height of this new braid is k-1. This braid is lower in the heap than our original braid. This is a

contradiction to the original choice of heap.


Subcase (b): Assume the block for  $s_0$  is in the (k-2)th row, and  $s_1$  and  $s_3$  are in the (k-3)th row. Then the subheap we are considering is




where the braid is highlighted in teal. Applying the braid move we get the following subheap




where no new braids have appeared, and in fact the original braid is higher now with height k+1. Notice however, that the (k-3)th row will not be the bottom row of our heap because then w would contain Property T a contradiction to our assumption. With this in mind we consider the (k-4)th row. Notice that  $s_0$  will not appear in the (k-4)th row as we would have a lower braid. This implies that the (k-4)th row contains at least one of  $s_2$  or  $s_4$ . With this in mind we represent this with the following subheap




where we have highlighted the additions in red and striped blocks. Here these striped blocks represent that at least one of the blocks appear but possibly both. Notice that if this new row is row 1, then w would have Property T. Again, this implies that this will not be the bottom row of the heap. Repeating this process again we see that  $s_1$  will not be in row k-5 since this would create a lower braid, thus we must have at least one of  $s_3$  or  $s_5$  in the (k-5)th row. We represent this with the following subheap




where again we have highlighted the additions in red and striped blocks. Again, if the (k-5)th row is the first row, then w would have Property T. This implies that the (k-5)th row is not the bottom-most row in our heap. Iterating this process we obtain the subheap as follows



where again we see that if the row containing the blocks for  $s_{n-2}$  and  $s_n$  corresponds to row 1, then w would have Property T. From this we obtain the following subheap




where the striped red blocks correspond to a portion of an FC element. Originally we said that the white inner triangle of this heap must be empty, however this contradicts [4, Lemma 3.3] which says that elements that have the triangle of white space must be filled completely in order to maintain that the heap is FC. That is, in the blank space between the purple blocks  $s_0$ ,  $s_1$  and the striped red blocks  $s_1$ ,  $s_3$  in the 2nd row every block will actually be present. This leads to the heap seen here




where we see that this has led to  $s_0$  being located in the (k-4)th row. Notice that a new braid has now appeared in the heap. We have highlighted this in orange. This new braid has height k-1. This is a contradiction as the braid appears lower than the original braid we chose.

Case (4): Suppose the braid is located in the kth row and assume the braid contains  $s_1$  or  $s_{n-1}$ . Without loss of generality we assume that the braid contains  $s_1$ , as the the other argument is symmetric to the one presented here and assume  $s_1$  is in the kth row. Assume that, if necessary, the blocks that complete the braid have been brought up next to  $s_1$  in the kth row. We now consider what can happen in the (k-3)th row and (k-4)th row in two cases. Notice that if the braid is  $s_1s_2s_1$ , then we are in Case 1, so assume the braid consists of  $s_0$  and  $s_1$ .

Subcase (a): Assume the braid involves  $s_1$  and  $s_0$  and the block for  $s_2$  is located in the (k-3)th row. Then the subheap we are considering follows here




where the braid we are considering is highlighted in orange. Applying the braid move we get the following subheap



in which we see that the original braid is now located higher in the heap with height k+1. By Case (1), we know that the original heap we started with implies that  $s_0$  and  $s_2$  are located above row 1. This implies that the subheap has more rows underneath to fill in.


Subcase (i): We first consider if the block for  $s_1$  is located in row k-4 and  $s_3$  is allowed but not required to be there. This leads to the following heap




where a new braid appears which we have highlighted in red. This new braid has height k-2 which is lower than the height of the original braid that we chose. This

is a contradiction to the way in which we chose w.


Subcase (ii): Now we consider if the block for  $s_3$  is in the (k-4)th row and  $s_1$  is not. This leads to the subheap seen here



where again there are no new braids present. However, the bottom row of the subheap is not the bottom row of the heap for w since otherwise w would have Property T. Using this notion we extend our heap to look like




where again we see no new braids. Again, we know that the bottom row of the subheap is not the bottom row of the heap for w since otherwise w would have Property T. Iterating this process we obtain a heap that looks like




where the zig-zag pattern continues. That is, the red row of blocks continues without blocks above or below these blocks. Notice that if the zig-zag was to end before reaching  $s_0$ , the heap would have Property T, which is a contradiction to the way in which we chose w. Suppose that the zig-zag continues on after reaching  $s_0$ . Then reverting back to the original configuration of the subheap we are able to drop the block for  $s_0$  down and create a lower braid. This is a contradiction to the way in which we chose the heap.

Case (5): Suppose the braid is located in the kth row and assume the braid contains  $s_0$  or  $s_n$ . Without loss of generality we assume that the braid contains  $s_0$  as the other argument is symmetric to the one presented here, and assume  $s_0$  is in row k. We now consider what is in row k-4. Notice that  $s_0$  can not be in the (k-4)th row as w would not be reduced. This implies that  $s_2$  is in the (k-4)th row and we get the subheap below.



where we have highlighted the braid in orange. Applying the braid move we get the following subheap



which does not have any new braids. However, notice that the height of the braid is now k-1. This is a contradiction to our original assumption that the heap we started with contains the lowest braid.

Therefore, it follow that  $W(\widetilde{C}_n)$  does not contain any not FC non-trivial T-avoiding elements.

We will now consider the elements in  $W(\widetilde{C}_n)$  that are FC where we first will classify non-trivial T-avoiding elements in  $W(\widetilde{C}_n)$  for n odd and then proceed to the classification for n even.

**Theorem 5.1.3.** If n is odd, then there are no non-trivial T-avoiding elements in the Coxeter system of type  $\widetilde{C}_n$ .

*Proof.* Consider the Coxeter system of type  $\widetilde{C}_n$ . By Theorem 5.1.2 we know that  $W(\widetilde{C}_n)$  contains no non-trivial T-avoiding elements that are not FC. Recall  $W(\widetilde{C}_n)$  is a star reducible Coxeter group, which implies that  $W(\widetilde{C}_n)$  contains no non-trivial T-avoiding elements that are FC. Thus as  $W(\widetilde{C}_n)$  has no non-trivial T-avoiding elements that are FC or not FC,  $W(\widetilde{C}_n)$  has no non-trivial T-avoiding elements.

We next will classify the non-trivial T-avoiding elements in the Coxeter system of type  $\widetilde{C}_n$  for n even. Recall that  $W(\widetilde{C}_n)$  for n even is not a star reducible Coxeter group. In Theorem 5.1.2 we showed that  $W(\widetilde{C}_n)$  does not have non-trivial T-avoiding elements that are not FC. This leaves us with only the FC elements to check.

**Theorem 5.1.4.** If n is even, then the only non-trivial T-avoiding elements in  $W(\widetilde{C}_n)$  are sandwich stacks.

Proof. Let  $w \in W(\widetilde{C}_n)$ . By Theorem 5.1.2, we know that w is an FC element. Further, we can restrict our search down to non-trivial non-cancellable elements as they are not star reducible, and thus do not contain Property T. In Section 2.3 we stated the classification of the only non-cancellable element with full support. Recall this to be sandwich stacks. Thus the only non-trivial T-avoiding elements in  $W(\widetilde{C}_n)$  for n odd are sandwich stacks whose top and bottom rows in the heap consist of  $s_0, s_2, \ldots s_n$ .

#### 5.2 Future Work

In Sections 3.1–3.4, we relayed the known results involving T-avoiding elements in types  $\widetilde{A}_n$ ,  $A_n$ ,  $D_n$ ,  $F_4$ ,  $F_5$ , and proved results involving T-avoiding elements in type  $I_2(m)$ . It remains to be shown that the conjecture in Section 3.1 regarding the classification of the non-trivial T-avoiding elements in type  $\widetilde{A}_n$  holds. The classification of non-trivial T-avoiding elements in Coxeter systems of type  $F_n$  for  $n \geq 6$  still remains open.

We also mentioned several other Coxeter systems in Figures 1.2 and 1.3. The classification of non-trivial T-avoiding elements in the Coxeter systems of type  $E_n$  remains an open problem. However, we do know that these groups have non-trivial T-avoiding elements as  $W(D_n)$  (which has non-trivial T-avoiding elements) is a parabolic subgroup of  $W(E_n)$ . The classification of non-trivial T-avoiding elements in the Coxeter systems of type  $H_n$  is also an open problem.

A majority of the irreducible affine Coxeter systems currently do not have a classification of the non-trivial T-avoiding elements. Specifically, Coxeter systems of type  $\widetilde{B}_n$ ,  $\widetilde{D}_n$ ,  $\widetilde{E}_6$ ,  $\widetilde{E}_7$ ,  $\widetilde{E}_8$ , and  $\widetilde{G}_4$  do not have a classification. Future work could include classifying the non-trivial T-avoiding elements of the Coxeter systems mentioned above.

## **Bibliography**

- [1] S.C. Billey and B.C. Jones. Embedded factor patterns for Deodhar elements in Kazhdan-Lusztig theory. *Ann. Comb.*, 11(3-4):285-333, 2007.
- [2] A Björner and F Brenti. Combinatorics of Coxeter groups. 2005.
- [3] D.C. Ernst. Non-cancellable elements in type affine C Coxeter groups. Int. Electron. J. Algebr., 8, 2010.
- [4] D.C. Ernst. Diagram calculus for a type affine C Temperley–Lieb algebra, II. arXiv1101.4215v1, 2012.
- [5] C.K. Fan. Structure of a Hecke algebra quotient. J. Amer. Math. Soc., 10:139– 167, 1997.
- [6] C.K. Fan and R.M. Green. On the affine Temperley–Lieb algebras. *Jour. L.M.S.*, 60:366–380, 1999.
- [7] M. Geck and G. Pfeiffer. Characters of finite Coxeter groups and Iwahori–Hecke algebras. 2000.
- [8] T. Gern. Leading Coefficients of Kazhdan-Lusztig Polynomials in Type D. PhD, University of Colorado, 2013.
- [9] R.M. Green. Star reducible Coxeter groups. Glas. Math. J., 48:583-609, 2006.
- [10] J.E. Humphreys. Reflection Groups and Coxeter Groups. 1990.
- [11] D. Kazhdan and G. Lusztig. Representations of Coxeter groups and Hecke algebras. *Inven. Math.*, 53:165–184, 1979.
- [12] G. Lusztig. Cells in affine Weyl groups, I. In *Collection*, pages 255–287. 1985.
- [13] J.R. Stembridge. On the fully commutative elements of Coxeter groups. *J. Algebr. Comb.*, 5:353–385, 1996.