

# THE MEANING OF LIFE

By Tony Stark

A Thesis

Submitted in Partial Fulfillment  
of the Requirements for the Degree of  
Master of Science  
in Mathematics

Northern Arizona University

July 2013

Approved:

Évariste Galois, Ph.D., Chair

John von Neumann, Ph.D.

Srinivasa Ramanujan, Ph.D.

ABSTRACT

THE MEANING OF LIFE

Tony Stark

Everything you always wanted to know will be discussed.

# Contents

List of Figures . . . . .	iv
<b>Chapter 1 Preliminaries</b>	<b>1</b>
1.1 Coxeter Systems . . . . .	1
1.2 Fully Commutative Elements . . . . .	4
<b>Chapter 2 Heaps</b>	<b>9</b>
2.1 Heaps . . . . .	9
<b>Chapter 3 Star Operations and NonCancellable Elements</b>	<b>14</b>
3.1 Star Operations . . . . .	14
3.2 NonCancellable Elements . . . . .	16
<b>Chapter 4 Property T and T-Avoiding</b>	<b>19</b>
4.1 Property T and T-Avoiding Elements . . . . .	19
4.2 T-Avoiding Elements in Types $\tilde{A}_n$ and $A_n$ . . . . .	22
4.3 T-Avoiding Elements in Type $D_n$ . . . . .	23
4.4 T-Avoiding Elements in Type $F_n$ . . . . .	24
4.5 T-Avoiding Elements in Type $I_2(m)$ . . . . .	25
<b>Chapter 5 T-Avoiding Elements in Type <math>B_n</math></b>	<b>26</b>
5.1 Tools for the classification . . . . .	26
5.2 Classification of T-Avoiding Elements in Type $B_n$ . . . . .	26
<b>Bibliography</b>	<b>28</b>

## List of Figures

1.1	Coxeter graphs corresponding to the irreducible finite Coxeter systems.	5
1.2	Coxeter graphs corresponding to the irreducible affine Coxeter systems.	6
1.3	Coxeter graphs corresponding to the irreducible FC-finite Coxeter systems. . . . .	8
2.1	Labeled hasse diagram for the heap of an element in $\text{FC}(\tilde{C}_6)$ . . . . .	10
2.2	A lattice point representation for the heap of an FC element in $W(\tilde{C}_6)$ .	11
2.3	Two heaps of a non-FC element in $W(\tilde{C}_4)$ . . . . .	12
2.4	Impermissible subheaps for elements in $\text{FC}(\tilde{C}_n)$ . . . . .	13
3.1	Visualization of Example 3.1.1. . . . .	15
3.2	Visualization of a noncancellable element of $\text{FC}(B_4)$ . . . . .	18
4.1	Heaps of an element with Property T and a T-Avoiding element . . .	20
4.2	Heap of a non-trivially T-Avoiding element in $W(\tilde{C}_4)$ . . . . .	21

## Chapter 1

# Preliminaries

### 1.1 Coxeter Systems

A *Coxeter system* is a pair  $(W, S)$  consisting of a finite set  $S$  of generating involutions and a group  $W$ , called a *Coxeter group*, with presentation

$$W = \langle S \mid (st)^{m(s,t)} = e \text{ for } m(s,t) < \infty \rangle,$$

where  $e$  is the identity,  $m(s,t) = 1$  if and only if  $s = t$ , and  $m(s,t) = m(t,s)$ . It turns out that the elements of  $S$  are distinct as group elements and that  $m(s,t)$  is the order of  $st$  [10]. We call  $m(s,t)$  the *bond strength* of  $s$  and  $t$ .

Since  $s$  and  $t$  are elements of order 2, the relation  $(st)^{m(s,t)} = e$  can be rewritten as

$$\underbrace{sts \cdots}_{m(s,t)} = \underbrace{tst \cdots}_{m(s,t)} \quad (1.1)$$

with  $m(s,t) \geq 2$  factors. If  $m(s,t) = 2$ , then  $st = ts$  is called a *commutation relation*. Otherwise, if  $m(s,t) \geq 3$ , then the relation in (1.1) is called a *braid relation*. Replacing  $\underbrace{sts \cdots}_{m(s,t)}$  with  $\underbrace{tst \cdots}_{m(s,t)}$  will be referred to as a *commutation* if  $m(s,t) = 2$  and a *braid move* if  $m(s,t) \geq 3$ .

We can represent a Coxeter system  $(W, S)$  with a unique *Coxeter graph*  $\Gamma$  having

- (1) vertex set  $S$  and
- (2) edges  $\{s, t\}$  for each  $m(s,t) \geq 3$  labeled by its corresponding bond strength  $m(s,t)$ .

Since  $m(s, t) = 3$  occurs frequently, it is customary to omit this label. Note that  $s$  and  $t$  are not connected by a single edge in the graph if and only if  $m(s, t) = 2$ . There is a one-to-one correspondence between Coxeter systems and Coxeter graphs. That is, given a Coxeter graph  $\Gamma$ , we can uniquely reconstruct the corresponding Coxeter system. If  $(W, S)$  is a Coxeter system with corresponding Coxeter graph  $\Gamma$ , we may denote the Coxeter group as  $W(\Gamma)$  and the generating set as  $S(\Gamma)$  for clarity. Also, the Coxeter system  $(W, S)$  is said to be *irreducible* if and only if  $\Gamma$  is connected. Further, if the graph  $\Gamma$  is disconnected, the connected components correspond to factors in a direct product of the corresponding Coxeter groups [10].

**Example 1.1.1.**

- (a) The Coxeter system of type  $A_n$  is given by the graph in Figure 1.1(a). We can construct the corresponding Coxeter group  $W(A_n)$  with generating set  $S(A_n) = \{s_1, s_2, \dots, s_n\}$  and defining relations

- (1)  $s_i^2 = e$  for all  $i$ ;
- (2)  $s_i s_j = s_j s_i$  when  $|i - j| > 1$ ;
- (3)  $s_i s_j s_i = s_j s_i s_j$  when  $|i - j| = 1$ .

The Coxeter group  $W(A_n)$  is isomorphic to the symmetric group  $\text{Sym}_{n+1}$  under the correspondence  $s_i \mapsto (i, i + 1)$ , where  $(i, i + 1)$  is the adjacent transposition that swaps  $i$  and  $i + 1$ .

- (b) The Coxeter system of type  $B_n$  is given by the graph in Figure 1.1(b). We can construct the corresponding Coxeter group  $W(B_n)$  with generating set  $S(B_n) = \{s_0, s_1, \dots, s_{n-1}\}$  and defining relations

- (1)  $s_i^2 = e$  for all  $i$ ;
- (2)  $s_i s_j = s_j s_i$  when  $|i - j| > 1$ ;
- (3)  $s_i s_j s_i = s_j s_i s_j$  when  $|i - j| = 1$  for  $i, j \in \{1, 2, \dots, n - 1\}$ ;
- (4)  $s_0 s_1 s_0 s_1 = s_1 s_0 s_1 s_0$ .

The Coxeter group  $W(B_n)$  is isomorphic to  $\text{Sym}_n^B$ , where  $\text{Sym}_n^B$  is the group of signed permutations on the set  $\{1, 2, \dots, n\}$ .

- (c) The Coxeter system of type  $\tilde{C}_n$  is seen in Figure 1.2(d). We can construct the corresponding Coxeter group  $W(\tilde{C}_n)$  with generating set  $S(\tilde{C}_n) = \{s_0, s_1, \dots, s_n\}$  and defining relations

- (1)  $s_i^2 = e$  for all  $i$ ;

- (2)  $s_i s_j = s_j s_i$  when  $|i - j| > 1$  for  $i \in \{1, 2, \dots, n - 1\}$ ;
- (3)  $s_i s_j s_i = s_j s_i s_j$  when  $|i - j| = 1$  for  $i \in \{1, 2, \dots, n - 1\}$ ;
- (4)  $s_0 s_1 s_0 s_1 = s_1 s_0 s_1 s_0$ ;
- (5)  $s_n s_{n-1} s_n s_{n-1} = s_{n-1} s_n s_{n-1} s_n$ .

Note that  $W(\tilde{C}_n)$  has  $n + 1$  generators.

The Coxeter graphs given in Figure 1.1 correspond to the collection of irreducible finite Coxeter groups, while the Coxeter graphs given in Figure 1.2 are the so called irreducible *affine Coxeter groups*, which are infinite [10]. Note that  $W(\tilde{C}_n)$  is one of the affine groups making it infinite. The irreducible affine Coxeter systems are unique in that if a vertex is removed along with the corresponding edges from the Coxeter graph, the newly created graph will result in a Coxeter system with a finite Coxeter group. This thesis will focus on Coxeter groups of type  $B_n$  and  $\tilde{C}_n$ .

Given a Coxeter system  $(W, S)$ , a word  $s_{x_1} s_{x_2} \cdots s_{x_m}$  in the free monoid  $S^*$  on  $S$  is called an *expression* for  $w \in W$  if it is equal to  $w$  when considered as a group element. If  $m$  is minimal among all expressions for  $w$ , the corresponding word is called a *reduced expression* for  $w$ . In this case, we define the *length* of  $w$  to be  $l(w) := m$ . Each element  $w \in W$  may have multiple reduced expressions that represent it. If we wish to emphasize a specific, possibly reduced, expression for  $w \in W$  we will represent it as  $\mathbf{w} = s_{x_1} s_{x_2} \cdots s_{x_m}$ . The following theorem tells us more about how reduced expressions for a given group element are related.

**Theorem 1.1.2** (Matsumoto, [6]). *Let  $(W, S)$  be a Coxeter system. If  $w \in W$ , then given a reduced expression for  $w$  we can obtain every other reduced expression for  $w$  by a sequence of braid moves and commutations of the form*

$$\underbrace{sts \cdots}_{m(s,t)} \rightarrow \underbrace{tst \cdots}_{m(s,t)}$$

where  $s, t \in S$  and  $m(s, t) \geq 2$ . □

It follows from Matsumoto's Theorem that if a generator  $s$  appears in a reduced expression for  $w \in W$ , then  $s$  appears in all reduced expressions for  $w$ . Let  $w \in W$  and define the *support* of  $w$ , denoted  $\text{supp}(w)$ , to be the set of all generators that appear in any reduced expression for  $w$ . If  $\text{supp}(w) = S$ , we say that  $w$  has *full support*.

Given  $w \in W$  and a fixed reduced expression  $\mathbf{w}$  for  $w$ , any subsequence of  $\mathbf{w}$  is called a *subexpression* of  $\mathbf{w}$ . We will refer to a subexpression consisting of a consecutive subsequence of  $\mathbf{w}$  as a *subword* of  $\mathbf{w}$ . If  $u, v \in W(\Gamma)$ , we say that the product of group elements  $uv$  is *reduced* if  $l(uv) = l(u) + l(v)$ .

**Example 1.1.3.** Let  $w \in W(A_7)$  and let  $\mathbf{w} = s_7 s_2 s_4 s_5 s_3 s_2 s_3 s_6$  be a fixed expression for  $w$ . Then we have

$$\begin{aligned}
 s_7 s_2 s_4 s_5 s_3 s_2 s_3 s_6 &= s_7 s_4 s_2 s_5 s_3 s_2 s_3 s_6 \\
 &= s_7 s_4 s_5 s_2 s_3 s_2 s_3 s_6 \\
 &= s_7 s_4 s_5 s_3 s_2 s_3 s_3 s_6 \\
 &= s_7 s_4 s_5 s_3 s_2 s_6,
 \end{aligned}$$

where the **blue** highlighted text corresponds to a commutation, the **teal** highlighted text corresponds to a braid move, and the **red** highlighted text corresponds to cancellation. This shows that the original expression  $\mathbf{w}$  is not reduced. However, it turns out that  $s_7 s_4 s_5 s_3 s_2 s_6$  is reduced. Thus  $l(w) = 6$  and  $\text{supp}(w) = \{s_2, s_3, s_4, s_5, s_6, s_7\}$ .

Let  $(W, S)$  be a Coxeter system of type  $\Gamma$  and let  $w \in W(\Gamma)$ . We define the *left descent set* and *right descent set* of  $w$  as follows:

$$\mathcal{L}(w) := \{s \in S \mid l(sw) < l(w)\}$$

and

$$\mathcal{R}(w) := \{s \in S \mid l(ws) < l(w)\}.$$

In [2] it is shown that  $s \in \mathcal{L}(w)$  (respectively,  $\mathcal{R}(w)$ ) if and only if there is a reduced expression for  $w$  that begins (respectively, ends) with  $s$ .

**Example 1.1.4.** The following list consists of all reduced expressions some  $w \in W(B_4)$ :

$$\begin{array}{cc}
 s_0 s_1 s_2 s_1 s_3 & s_0 s_2 s_1 s_2 s_3 \\
 s_0 s_1 s_2 s_3 s_1 & s_2 s_0 s_1 s_2 s_3
 \end{array}$$

We see that  $l(w) = 5$  and  $w$  has full support. Also, we see that  $\mathcal{L}(w) = \{s_0, s_2\}$  while  $\mathcal{R}(w) = \{s_1, s_3\}$ .

## 1.2 Fully Commutative Elements

Let  $(W, S)$  be a Coxeter system of type  $\Gamma$  and let  $w \in W(\Gamma)$ . Following [13], we define a relation  $\sim$  on the set of reduced expressions for  $w$ . Let  $\mathbf{w}_1$  and  $\mathbf{w}_2$  be two reduced expressions for  $w$ . We define  $\mathbf{w}_1 \sim \mathbf{w}_2$  if we can obtain  $\mathbf{w}_2$  from  $\mathbf{w}_1$  by applying a single commutation move of the form  $st \mapsto ts$  where  $m(s, t) = 2$ . Now, define the equivalence relation  $\approx$  by taking the reflexive transitive closure of  $\sim$ . Each equivalence class under  $\approx$  is called a *commutation class*. If  $w$  has a single commutation class, then we say that  $w$  is *fully commutative* (FC).






Figure 1.1: Coxeter graphs corresponding to the irreducible finite Coxeter systems.

The set of FC elements of  $W(\Gamma)$  is denoted by  $\text{FC}(\Gamma)$ . Given some  $w \in \text{FC}(\Gamma)$ , and a starting reduced expression for  $w$ , observe the definition of FC states that one only needs to perform commutations to obtain all reduced expressions for  $w$ , but the following result due to Stembridge [13] states that when  $w$  is FC, performing commutations is the only possible way to obtain another reduced expression for  $w$ .

**Theorem 1.2.1** (Stembridge, [13]). *Let  $(W, S)$  be a Coxeter system. An element  $w \in W$  is FC if and only if no reduced expression for  $w$  contains  $\underbrace{sts \cdots}_{m(s,t)}$  as a subword for all  $m(s, t) \geq 3$ .  $\square$*

In other words,  $w$  is FC if and only if we never have the opportunity to apply a braid move. In particular, for a Coxeter group of type  $B_n$  an element is FC if it does not contain the subwords  $s_0 s_1 s_0 s_1$ ,  $s_1 s_0 s_1 s_0$ ,  $s_k s_{k+1} s_k$ , and  $s_{k+1} s_k s_{k+1}$  where  $0 < k < n + 1$ . In a Coxeter group of type  $\tilde{C}_n$ , an element is FC if it does not contain the subwords seen above and does not contain the subwords  $s_{n-1} s_n s_{n-1} s_n$




Figure 1.2: Coxeter graphs corresponding to the irreducible affine Coxeter systems.

and  $s_n s_{n-1} s_n s_{n-1}$ .

**Example 1.2.2.** Let  $w \in W(\tilde{C}_4)$  and let  $\mathbf{w} = s_0 s_1 s_2 s_0 s_3 s_1$  be a reduced expression for  $w$ . We see that

$$s_0 s_1 \textcolor{violet}{s_2 s_0} s_3 s_1 = s_0 s_1 s_0 s_2 \textcolor{violet}{s_3 s_1} = s_0 s_1 s_0 s_2 s_1 s_3,$$

where the **purple** highlighted text indicates applying a commutation. Although it is not immediately obvious, these are all of the reduced expressions for the given  $w$ . Note that there is no possible way to perform a braid move. Hence  $w$  is FC.

**Example 1.2.3.** Let  $\mathbf{w} = s_1 s_0 s_4 s_1 s_3 s_5 s_2 s_4 s_6$  be a reduced expression for  $w \in \text{FC}(\tilde{C}_6)$ . Applying the commutation  $s_4 s_2 = s_2 s_4$ , we can obtain another reduced expression

for  $w$ , namely  $\mathbf{w}_2 = s_1 s_0 s_4 s_1 s_3 s_5 s_4 s_2 s_6$ , which is in the same commutation class as  $\mathbf{w}$ . However, applying the braid move  $s_2 s_3 s_2 = s_3 s_2 s_3$ , we obtain another reduced expression  $\mathbf{w} = s_1 s_3 s_2 s_3 s_4 s_0$ . Note that since  $\mathbf{w}_3$  was obtained by applying a braid move,  $\mathbf{w}_3$  is in a different commutation class than  $\mathbf{w}_1$  and  $\mathbf{w}_2$ . It turns out that  $w$  has exactly two commutation classes, one consisting of  $\mathbf{w}_1$  and  $\mathbf{w}_2$  and another consisting of  $\mathbf{w}_3$ . So  $w$  is not FC.

**Example 1.2.4.** Let  $w \in W(\tilde{C}_4)$  and let  $\mathbf{w} = s_0 s_1 s_2 s_0 s_1 s_2$  be a reduced expression for  $w$ . We see that

$$s_0 s_1 \textcolor{purple}{s_3} s_0 s_1 s_2 = s_0 s_1 s_0 \textcolor{purple}{s_3} \textcolor{purple}{s_1} s_2 = \textcolor{orange}{s_0} \textcolor{orange}{s_1} s_0 \textcolor{orange}{s_1} s_3 s_2,$$

where the **purple** highlighted text indicates applying a commutation and the **orange** highlighted text indicates applying a braid move. Thus  $w$  is not FC by Theorem 1.2.1.

Stembridge classified the Coxeter Systems that contain a finite number of FC elements, the so-called *FC-finite Coxeter groups*. Both  $W(A_n)$  and  $W(B_n)$  are finite Coxeter groups, and thus are FC-finite. On the other hand,  $W(\tilde{C}_n)$  is infinite and happens to also contain infinitely many FC elements. However, there exist some infinite Coxeter groups that contain finitely many FC elements. For example,  $W(E_n)$  for  $n \geq 9$  (see Figure 1.3) is infinite, but contains only finitely many FC elements.

**Theorem 1.2.5** (Stembridge, [13]). *The FC-finite irreducible Coxeter systems are of type  $A_n$  with  $n \geq 1$ ,  $B_n$  with  $n \geq 2$ ,  $D_n$  with  $n \geq 4$ ,  $E_n$  with  $n \geq 6$ ,  $F_n$  with  $n \geq 4$ ,  $H_n$  with  $n \geq 3$ , and  $I_2(m)$  with  $5 \leq m < \infty$ .  $\square$*

The irreducible FC-finite Coxeter graphs are given in Figure 1.3. Note that we have already encountered some of the FC-finite Coxeter groups in Figure 1.1. Since these are finite Coxeter groups it is clear that they will have a finite number of FC elements. However, we haven't yet encountered the Coxeter groups determined by graphs in Figures 1.3(d), 1.3(e), 1.3(f). All of these Coxeter systems are infinite for large  $n$ , yet contain only finitely many FC elements.




Figure 1.3: Coxeter graphs corresponding to the irreducible FC-finite Coxeter systems.

## Chapter 2

# Heaps

### 2.1 Heaps

We now discuss a visual representation of Coxeter group elements. Each reduced expression can be associated with a labeled partially ordered set (poset) called a heap. Heaps provide a visual representation of a reduced expression while preserving the relations among the generators. We follow the development of heaps for straight line Coxeter groups found in [1], [3], and [13].

Let  $(W, S)$  be a Coxeter system of type  $\Gamma$ . Suppose  $\mathbf{w} = s_{x_1}s_{x_2}\cdots s_{x_r}$  is a fixed reduced expression for  $w \in W(\Gamma)$ . As in [13], we define a partial ordering on the indices  $\{1, 2, \dots, r\}$  by the transitive closure of the relation  $\triangleleft$  defined via  $j \triangleleft i$  if  $i < j$  and  $s_{x_i}$  and  $s_{x_j}$  do not commute. In particular, since  $\mathbf{w}$  is reduced,  $j \triangleleft i$  if  $s_{x_i} = s_{x_j}$  by transitivity. This partial order is referred to as the *heap* of  $\mathbf{w}$ , where  $i$  is labeled by  $s_{x_i}$ . Note that for simplicity we are omitting the labels of the underlying poset but retaining the labels of the corresponding generators.

It follows from [13] that heaps are well-defined up to commutation class. That is, given two reduced expressions  $\mathbf{w}_1$  and  $\mathbf{w}_2$  for  $w \in W$  that are in the same commutation class, then the heaps for  $\mathbf{w}_1$  and  $\mathbf{w}_2$  will be equal. In particular, if  $w \in \text{FC}(\Gamma)$ , then  $w$  has one commutation class, and thus  $w$  has a unique heap. Conversely, if  $\mathbf{w}_1$  and  $\mathbf{w}_2$  are in different commutation classes, then there will not be a unique heap representation for  $w$ .

**Example 2.1.1.** Let  $\mathbf{w} = s_6s_4s_2s_5s_3s_1s_4s_0s_1$  be a reduced expression for  $w \in \text{FC}(\tilde{C}_6)$ . We see that  $\mathbf{w}$  is indexed by  $\{1, 2, 3, 4, 5, 6, 7, 8, 9\}$ . As an example,  $8 \triangleleft 9$  since  $0 < 1$  and  $s_0$  and  $s_1$  do not commute. The labeled Hasse diagram for the heap poset is seen in Figure 2.1.

Let  $\mathbf{w}$  be a reduced expression for an element in  $w \in W(\tilde{C}_n)$ . As in [1] and [3] we can represent a heap for  $\mathbf{w}$  as a set of lattice points embedded in  $\{0, 1, 2, \dots, n\} \times \mathbb{N}$ .




Figure 2.1: Labeled hasse diagram for the heap of an element in  $\text{FC}(\tilde{C}_6)$ .

To do so, we assign coordinates (not unique)  $(x, y) \in \{0, 1, 2, \dots, n\} \times \mathbb{N}$  to each entry of the labeled Hasse diagram for the heap of  $\mathbf{w}$  in such a way that:

- (1) An entry with coordinates  $(x, y)$  is labeled  $s_i$  (or  $i$ ) in the heap if and only if  $x = i$ ;
- (2) If an entry with coordinates  $(x, y)$  is greater than an entry with coordinates  $(x', y')$  in the heap then  $y > y'$ .

Although the above is specific to  $W(\tilde{C}_n)$ , the same construction works for any straight line Coxeter graph with the appropriate adjustments made to the labeling set and assignment of coordinates. Specifically, for type  $A_n$  our label set is  $\{1, 2, \dots, n\}$  and for type  $B_n$  our label set is  $\{0, 1, \dots, n-1\}$ .

In the case of any straight line Coxeter graph it follows from the definition that  $(x, y)$  covers  $(x', y')$  in the heap if and only if  $x = x' \pm 1$ ,  $y > y'$ , and there are no entries  $(x'', y'')$  such that  $x'' \in \{x, x'\}$  and  $y' < y'' < y$ . This implies that we can completely reconstruct the edges of the Hasse diagram and the corresponding heap poset from a lattice point representation. The lattice point representation can help us visualize arguments that are potentially complex. Note that in our heaps the entries in the top correspond to the generators occurring in the right descent set of the corresponding reduced expression.

Let  $\mathbf{w}$  be a reduced expression for  $w \in W(\tilde{C}_n)$ . We denote the lattice representation of the heap poset in  $\{0, 1, 2, \dots, n\} \times \mathbb{N}$  described in the preceding paragraphs via  $H(\mathbf{w})$ . If  $w$  is FC, then the choice of reduced expression for  $w$  is irrelevant and we will often write  $H(w)$  and we refer to  $H(w)$  as the heap of  $w$ . Note that we will use the same notation for heaps in Coxeter groups of all types with straightline Coxeter graphs.

There are potentially many ways to illustrate a heap of an arbitrary reduced expression, each differing by the vertical placement of the blocks. For example, we can place blocks in vertical positions as high as possible, as low as possible, or some combination of low/high. In this thesis, we choose what we view to be the best representation of the heap for each example and when illustrating the heaps of arbitrary reduced expressions we will discuss the relative position of the entries but never the absolute coordinates.

Let  $\mathbf{w} = s_{x_1}s_{x_2}\cdots s_{x_r}$  be a reduced expression for  $w \in W(\tilde{C}_n)$ . If  $s_{x_i}$  and  $s_{x_j}$  are adjacent generators in the Coxeter graph with  $i < j$ , then we must place the point labeled by  $s_{x_j}$  at a level that is *above* the level of the point labeled by  $s_{x_i}$ . Because generators in a Coxeter graph that are not adjacent do commute, points whose  $x$ -coordinates differ by more than one can slide past each other or land in the same level. To emphasize the covering relations of the lattice point representation we will enclose each entry in the heap in a square with rounded corners in such a way that if one entry covers another the squares overlap halfway. In addition, we will also label each square for  $s_i$  with  $i$ .

**Example 2.1.2.** Let  $\mathbf{w} = s_6s_4s_2s_5s_3s_1s_4s_0s_1$  be a reduced expression for  $w \in \text{FC}(\tilde{C}_6)$  as seen in Example 1.4.1. Figure 2.2 shows a possible lattice point representation for  $H(w)$ .




Figure 2.2: A lattice point representation for the heap of an FC element in  $W(\tilde{C}_6)$ .

**Example 2.1.3.** Let  $\mathbf{w}_1 = s_0s_2s_4s_3s_2s_1$  be a reduced expression for  $w \in W(\tilde{C}_4)$ . Applying the commutation  $s_4s_2 = s_2s_4$ , we can obtain another reduced expression for  $w$ , namely  $\mathbf{w}_2$ , which is in the same commutation class as  $\mathbf{w}_1$  and hence has the same heap. However, applying the braid move  $s_2s_3s_2 = s_3s_2s_3$ , we obtain another reduced expression  $\mathbf{w}_3 = s_0s_4s_3s_2s_3s_1$ . Note that since  $\mathbf{w}_3$  was obtained by applying a braid move,  $\mathbf{w}_3$  is in a different commutation class than  $\mathbf{w}_1$  and  $\mathbf{w}_2$ . Representations of  $H(\mathbf{w}_1)$ ,  $H(\mathbf{w}_2)$ , and  $H(\mathbf{w}_3)$  are seen in Figure 2.3 where the braid relation is colored in orange.

It will be extremely useful for us to be able to quickly determine whether a heap corresponds to an element in  $\text{FC}(B_n)$ , and  $\text{FC}(\tilde{C}_n)$ . The next proposition is a special case of [13, Proposition 3.3] and follows quickly when one considers the consecutive subwords that are impermissible in reduced expressions for elements in  $\text{FC}(B_n)$  and  $\text{FC}(\tilde{C}_n)$  seen in Section 1.2.

**Theorem 2.1.4.** *If  $w \in \text{FC}(\tilde{C}_n)$ , then  $H(w)$  cannot contain any of the subheaps seen in Figure 2.4, where  $0 < k < n + 1$  and we use a square with a dotted boundary to emphasize that no element of the heap occupies the corresponding position.*  $\square$




Figure 2.3: Two heaps of a non-FC element in  $W(\tilde{C}_4)$

Since  $W(B_n)$  is parabolic subgroups of  $W(\tilde{C}_n)$  we can use Figure 2.4 to classify the impermissible subheaps for elements of  $FC(B_n)$ . The impermissible subheaps for elements of  $FC(B_n)$  are those seen in Figures 2.4(a), 2.4(b) 2.4(c), and 2.4(d).






Figure 2.4: Impermissible subheaps for elements in  $\text{FC}(\tilde{C}_n)$ .

## Chapter 3

# Star Operations and NonCancellable Elements

### 3.1 Star Operations

The notion of star operations was originally introduced by Kazhdan and Lusztig in [11] for simply laced Coxeter systems (i.e.,  $m(s, t) \leq 3$  for all  $s, t \in S$ ), and was later generalized to all Coxeter systems in [12]. If  $I = \{s, t\}$  is a pair of non-commuting generators of a Coxeter group  $W$ , then  $I$  induces four partially defined maps from  $W$  to itself, known as *star operations*. A star operation, when it is defined, increases or decreases the length of an element to which it is applied by 1. For our purposes it is enough to only define the star operations that decrease the length of an element by 1, and as a result we will not develop the notion in full generality.

Let  $(W, S)$  be a Coxeter system of type  $\Gamma$  and let  $I = \{s, t\} \subseteq S$  be a pair of noncommuting generators whose product has order  $m$ . Let  $w \in W(\Gamma)$  such that  $s \in \mathcal{L}(w)$ . We define  $w$  to be *left star reducible by  $s$  with respect to  $t$*  if there exists  $t \in \mathcal{L}(sw)$  and  $m(s, t) \geq 3$ . We analogously define  $w$  to be *right star reducible by  $s$  with respect to  $t$* . Observe that if  $m(s, t) \geq 3$ , then  $w$  is left (respectively, right) star reducible if and only if there is a reduced group element for  $w$  such that  $w = stu$  (respectively,  $w = uts$ ). We say that  $w$  is *star reducible* if it is either left or right star reducible.

**Example 3.1.1.** Let  $w \in W(B_4)$  and let  $\mathbf{w} = s_0s_1s_0s_2s_3$  be a reduced expression for  $w$ . We see that  $w$  is left star reducible by  $s_0$  with respect to  $s_1$  to  $s_1s_0s_2s_3$ , since  $m(s_0, s_1) = 4$  and  $s_0 \in \mathcal{L}(w)$  while  $s_1 \in \mathcal{L}(s_0w)$ . Also  $w$  is right star reducible by  $s_3$  with respect to  $s_2$  to  $s_0s_1s_0s_2$ , since  $m(s_2, s_3) = 3$  and  $s_3 \in \mathcal{R}(w)$  and  $s_2 \in \mathcal{R}(ws_3)$ .

It may be helpful to visualize star reductions in terms of heaps. Figure 3.1(a) represents  $H(\mathbf{w})$ . Note that we can see  $s_0$  is in the left descent set of  $w$  since  $s_0$  is

in the top row of the heap. Furthermore, multiplying on the left by  $s_0$  we get the heap in Figure 3.1(b). Again, since  $s_1$  is in the top row of the heap,  $s_1 \in \mathcal{L}(s_1 w)$ . In Figure 3.1(a) we also see that  $s_3$  is in the right descent set of  $w$  since  $s_3$  is in the bottom row of the heap. Multiplying on the left by  $s_3$  we can see that  $s_2$  would be in the bottom level of the heap so  $s_2 \in \mathcal{R}(ws_3)$ . From this we can interpret visually an element  $w \in W(\Gamma)$  is right star reducible (respectively, left star reducible) if there exists a heap that we can pull a block off the bottom row of the heap (respectively, top of the heap) and a new block that wasn't previously in the bottom row (respectively, top row) is now in the bottom row (respectively, top row) of the heap. That is, we can systematically dismantle the heap for a given element by pulling blocks off of the bottom row (respectively, top row) of the heap and have the heap decrease in height, meaning there are fewer rows of the heap than there were in the original, or a block that was previously trapped in the second (respectively, second to last) row of the heap is now free to be in either the first or second (respectively, second to last or last) row. If the heap has the same number of rows as the original and if now blocks that were in the second or second to last row of the heap can now be in the first or last row of the heap when we attempt pulling a single block off the top and a single block off the bottom and we try all possible combination of single blocks in the top row and bottom row, then the heap is not star reducible.



Figure 3.1: Visualization of Example 3.1.1.

Using the notion of star reduction we are now able to introduce the concept of a star reducible Coxeter group. We say that a Coxeter group  $W(\Gamma)$ , or its Coxeter graph  $\Gamma$ , is *star reducible* if every element of  $\text{FC}(\Gamma)$  is star reducible to a product of commuting generators. That is,  $W(\Gamma)$  is star reducible if when we apply star operations repeatedly to  $w \in \text{FC}(\Gamma)$ , eventually we obtain a product of commuting generators. Using heaps to visualize a star reducible Coxeter group given a heap in  $\text{FC}(\Gamma)$ , we are able to pull the top or bottom most block off of the heap and have the heap decrease in height (the number of rows that it consists of) or have a new

block come into the top or bottom most rows. We can perform this pulling of blocks off the top and bottom of the heap systematically until the heap consists of one row, corresponding to an element that is a product of commuting generators. For example in Figure 3.1(a), we were able to apply a star reduction to remove the topmost block corresponding to the generator  $s_0$  and obtain the new heap seen in Figure 3.1(b). We see that the heap in Figure 3.1(b) has one less row than Figure 3.1(a). We could do the same with the block on bottom of Figure 3.1(b) corresponding to  $s_3$  and have the heap again decrease in the number of rows it has. Performing one more star reduction in pulling the brick off the top of Figure 3.1(b) which corresponds to  $s_1$  in the reduced expression would leave us with a heap that is only 1 row which is a product of commuting generators. Thus Figure 3.1(a) can be star reduced to a product of commuting generators. In [8], Green classified all star reducible Coxeter groups.

**Theorem 3.1.2** (Green, [8]). *Let  $W(\Gamma)$  be a Coxeter group with (finite) generating set  $S$ . Then  $W(\Gamma)$  is star reducible if and only if each component of  $\Gamma$  is either a complete graph with labels  $m(s, t) \geq 3$ , or is one of the following types: type  $A_n$  ( $n \geq 1$ ), type  $B_n$  ( $n \geq 2$ ), type  $D_n$  ( $n \geq 4$ ), type  $F_n$  ( $n \geq 4$ ), type  $H_n$  ( $n \geq 2$ ), type  $I_2(m)$  ( $m \geq 3$ ), type  $\tilde{A}_{n-1}$  ( $n \geq 3$  and  $n$  odd), type  $\tilde{C}_{n-1}$  ( $n \geq 4$  and  $n$  even), type  $\tilde{E}_6$  or type  $\tilde{F}_5$ .*  $\square$

In [9], Green defined star reducible Coxeter groups to have *Property F*. We go a bit farther and say that an element of  $FC(\Gamma)$  that is star reducible to a product of commuting generators will have Property F. In the same paper Green defined a Coxeter group with *Property S* to be a Coxeter group in which each element  $w \in W(\Gamma) \setminus FC(\Gamma)$  is star reducible to a product of noncommuting generators either in  $\mathcal{L}(w)$  or in  $\mathcal{R}(w)$ . Again we extend the definition to a specific element in the  $W(\Gamma)$ . That is,  $w \in W(\Gamma) \setminus FC(\Gamma)$  has Property S if  $w$  is star reducible to a product of non-commuting generators in  $\mathcal{L}(w)$  or  $\mathcal{R}(w)$ .

## 3.2 NonCancellable Elements

We now introduce the concept of weak star reducible, which is related to the notion of cancellable in [4]. Let  $(W, S)$  be a Coxeter system of type  $\Gamma$  and let  $I = \{s, t\} \subseteq S$  be a pair of noncommuting generators of the Coxeter group  $W(\Gamma)$ . If  $w \in FC(\Gamma)$ , then  $w$  is *left weak star reducible by  $s$  with respect to  $t$  to  $sw$*  if

- (1)  $w$  is left star reducible by  $s$  with respect to  $t$  and;
- (2)  $tw \notin FC(W)$ .

Notice that (2) implies that  $l(tw) > l(w)$ . Also note that we are restricting our definition of weak star reducible to the set of FC elements of  $W(\Gamma)$ . We analogously define *right weak star reducible by  $s$  with respect to  $t$  to  $ws$* . We say that  $w$  is *weak star reducible* if  $w$  is either left or right weak star reducible. Otherwise, we say that  $w$  is *non-cancellable* or *weak star irreducible*.

**Example 3.2.1.** Let  $w \in \text{FC}(B_4)$  and let  $\mathbf{w} = s_0s_1s_0s_2s_3$  be a reduced expression for  $w$  as in Example 3.1.1. By Example 3.1.1 we know that  $w$  is left star reducible. Also,  $tw = s_1s_0s_1s_0s_2s_3$  which is not in  $\text{FC}(B_4)$ . Thus, we see that  $w$  is left weak star reducible by  $s_0$  with respect to  $s_1$  to  $s_1s_0s_2s_3$ . In addition, Example 3.1.1 showed that  $w$  is right star reducible. But,  $wt = s_0s_1s_0s_2s_3s_2$  which is not in  $\text{FC}(B_4)$ . Thus,  $w$  is right weak star reducible by  $s_3$  with respect to  $s_2$  to  $s_0s_1s_0s_2$ . This implies that  $w$  is not non-cancellable.

Again it might be useful to visualize the concept of weak star reducible in terms of heaps. Recall in Figure 3.1(a) we have a representation for  $w$  as described in Example 3.2.1. In Section ??, we described the systematic dismantling of the heap that could occur with star reductions being repeatedly applied to the heap in Figure 3.1(a). In Figure 3.2 we can see that when we multiply  $w$  by  $s_1$  on the right we end up with a braid, which is highlighted in orange. Since the heap in Figure 3.2 has the impermissible subheap seen in Figure 2.4(b),  $s_1w \notin \text{FC}(B_4)$ . When using heaps to identify whether an element  $w$  of  $\text{FC}(\Gamma)$  is noncancellable or not there are two key properties that must be observed. The first of the properties is that the  $H(w)$  must be able to be dismantled from the top or bottom so that the heap resulting from pulling the top block off or the bottom block off is one row shorter or allows for a new brick to be in the top most or bottom most row of the heap. The second property that must be observed is that given the block that appears in the top or bottom of the heap when the star operation is performed, adding another of that block to the top or bottom of the original heap respectively will result in an impermissible subheap appearing. If both of these properties occur, then the element that corresponds to the heap is not noncancellable.

**Example 3.2.2.** Let  $w \in \text{FC}(B_4)$  and let  $\mathbf{w} = s_0s_1$  be a reduced expression for  $w$ . Note that  $w$  is left (respectively, right) star reducible by  $s_0$  with respect to  $s_1$  (respectively, by  $s_1$  with respect to  $s_0$ ). However,  $s_1s_0s_1 \in \text{FC}(B_4)$  (respectively,  $s_0s_1s_0 \in \text{FC}(B_4)$ ). Thus  $w$  is non-cancellable.




Figure 3.2: Visualization of a noncancellable element of  $\text{FC}(B_4)$ .

## Chapter 4

# Property T and T-Avoiding

### 4.1 Property T and T-Avoiding Elements

As mentioned in Section 3.1 Green classified all star reducible Coxeter groups. In [8], Green utilizes the following theorem to help classify the star reducible Coxeter groups.

**Theorem 4.1.1** (Green, [8]). *Let  $W$  be a star reducible Coxeter group, and let  $w \in W$ . Then one of the following possibilities occurs for some Coxeter generators  $s, t, u$  with  $m(s, t) \neq 2$ ,  $m(t, u) \neq 2$ , and  $m(s, u) = 2$ :*

- (1)  *$w$  is a product of commuting generators;*
- (2)  *$w$  has a reduced expression beginning with  $st$ ;*
- (3)  *$w$  has a reduced expression ending in  $ts$ ;*
- (4)  *$w$  has a reduced expression beginning with  $sut$ .* □

In the following discussion we will give name to elements that exhibit the property in Item (1), as well as name elements that exhibit the properties seen in Items (2) and (3), and name elements that exhibit the property seen in Item (4). We will then focus on discussing the irreducible finite and affine Coxeter groups that have elements with properties seen in Items (1) and (4).

We first begin by defining the notion of Property T. Let  $(W, S)$  be a Coxeter system of type  $\Gamma$  and let  $w \in W$ . We say that  $w$  has *Property T* if and only if there exists a reduced product for  $w$  such that  $w = stu$  or  $w = uts$  where  $m(s, t) \geq 3$ . That is,  $w$  has Property T if there exists a reduced expression for  $w$  that begins or ends with a product of non-commuting generators. It should be noted that by the symmetry of the definition  $w$  has Property T if and only if  $w^{-1}$  has Property T.

**Example 4.1.2.** Let  $w \in W(A_5)$  with reduced expression  $\mathbf{w} = s_1 s_4 s_2 s_3 s_5$ . At first glance it may appear that  $w$  does not have Property T, since both  $s_1$  and  $s_4$  commute as well as  $s_3$  and  $s_5$ . However, note that applying a commutation to  $s_4 s_2$  results in  $\mathbf{w}_1 = s_1 s_2 s_4 s_3 s_5$ . Hence  $w$  has Property T, since  $m(s_1, s_2) = 3$  and there is a reduced expression for  $w$  that begins with  $s_1 s_2$ .

**Example 4.1.3.** Let  $w \in W(A_5)$  with reduced expression  $\mathbf{w} = s_1 s_3 s_5$ . It turns out that since  $w$  is a product of commuting generators there is no reduced expression for  $w$  that begins or ends with a pair of non-commuting generators. This implies that  $w$  does not have Property T.

As with star reducible elements it may be helpful to visualize Property T through heaps. Figure 4.1(a) provides a representation of an element in  $W(\Gamma)$  with Property T and Figure 4.1(b) provides a representation of an element without Property T. Notice that if we were to remove the block for  $s_1$  in the bottom row of Figure 4.1(a), the heap would become one less row in height and we would have a new bottom row in the heap. However, in Figure 4.1(b), we are not able to remove any bricks and have a new brick come to the top or bottom row as the heap is just one row. From this we can gather that when we are using heaps to visualize whether or not an element of  $W(\Gamma)$  has Property T we must observe either of the following things. The first of these is that the heap decreased in height. That is, there is one less row than the original heap. The second thing that we could observe is that when we remove a block from the heap a new element, that was originally trapped in the second (respectively, second to last) row is now able to move into the first (respectively, last) row of the heap. This implies that an element that does not have Property-T, has every element in the second and second to last row of the heap blocked by two blocks in the first and last rows of the heap, as this would imply that we could not remove a block and have a new element come to the first or last row as it would still be blocked by the other element that remained.




Figure 4.1: Heaps of an element with Property T and a T-Avoiding element

An element  $w \in W(\Gamma)$  is called *T-avoiding* if  $w$  and  $w^{-1}$  do not have Property T. We will call an element that is a product of commuting generators *trivially T-avoiding*.



The reason behind this comes in the following theorem. If  $w$  is T-avoiding and not a product of commuting generators, we will say that  $w$  is *non-trivially T-avoiding*.

**Theorem 4.1.4.** *Let  $(W, S)$  be a Coxeter system of type  $\Gamma$  and let  $w \in W(\Gamma)$  such that  $w$  is a product of commuting generators. Then  $w$  is T-avoiding.  $\square$*

Visually this is seen in Example 4.1(b) elements in  $W(\Gamma)$  that are products of commuting generators are always going to be one row in the heap. This implies that we are not able to remove generators and have elements come into rows that they were previously in as the heap is only one row and there can be no lateral movement when we remove bricks.

**Example 4.1.5.** Let  $w \in W(A_5)$  and let  $\mathbf{w} = s_1 s_3 s_5$ . Then by Example 4.1.3, we know that  $w$  is T-avoiding and since  $w$  is a product of commuting generators,  $w$  is trivially T-avoiding.

**Example 4.1.6.** Let  $w \in W(\tilde{C}_4)$  with reduced expression  $\mathbf{w} = s_0 s_2 s_4 s_1 s_3 s_0 s_2 s_4$ . It turns out that  $w$  is FC and non-trivially T-avoiding. The heap for  $w$  is seen in Figure 4.2. Notice that no matter which block we remove from the top row of  $w$  no new element can be pushed into the topmost row. The same applies to the bottom row. Hence there is not a single block that can be removed from the top that allows a new element to come into the top row and no single block that can be removed from the bottom row to allow a new element to come into the bottom row. Thus,  $w$  is non-trivially T-avoiding.




Figure 4.2: Heap of a non-trivially T-Avoiding element in  $W(\tilde{C}_4)$ .

Referring back to Green's classification of what elements in star reducible Coxeter groups look like, we see that Item (1) corresponds to the element  $w$  being trivially T-avoiding, Items (2) and (3) refer to the element  $w$  having Property T at the beginning and end respectively, and Item (4) corresponds to the element  $w$  being T-avoiding. Notice that this implies that some Coxeter groups will have elements that are non-trivially T-avoiding while others may not. For example, as will be seen in the following

sections, the Coxeter groups of type  $A_n$  and  $B_n$  have no non-trivial T-avoiding elements, while the Coxeter group of type  $D_n$  does have non-trivial T-avoiding elements. One interesting thing to note about star reducible Coxeter groups is that if they contain non-trivially T-avoiding elements, these elements will not be FC. This is because FC elements in star reducible Coxeter groups are star reducible and star reducible elements in the way in which we have defined them are analogous to the element having Property T.

We will now extend the definition of Property T to Coxeter groups similar to Properties F and S seen in Section 3.1. That is, we say a Coxeter group  $W(\Gamma)$  has Property T if all elements in  $W(\Gamma)$  that are not the products of commuting generators have Property T. It is clear that if a Coxeter group has Property T, then it also has Property S. It remains an open question whether or not if a Coxeter group has Property S, then it has Property T. If this is the case, then by a remark following the definition in [9], completes the classification of the T-avoiding elements of any connected, nonbranching Coxeter graph of finite or affine type, except the Coxeter group  $\tilde{F}_4$ .

One thing to notice here is that all Coxeter groups have trivial T-avoiding elements as they all contain products of commuting generators. The more interesting non-trivial T-avoiding elements do not appear in all Coxeter groups. The remainder of this thesis discusses what is currently known regarding T-avoiding elements in the irreducible finite Coxeter groups and the irreducible affine Coxeter groups, as well as it classifies the T-avoiding elements in Coxeter groups of types  $B_n$  and  $\tilde{C}_n$ . In the next few sections we will summarize what is known about the T-avoiding elements in Coxeter groups of types  $\tilde{A}_n$ ,  $A_n$ ,  $D_n$ ,  $F_n$  and  $I_2(m)$ .

## 4.2 T-Avoiding Elements in Types $\tilde{A}_n$ and $A_n$

We start by classifying the T-avoiding elements in Coxeter groups of type  $\tilde{A}_n$  and  $A_n$ . Specifically we will classify the T-avoiding elements in  $W(A_n)$  and then move to  $W(\tilde{A}_n)$ . Clearly  $W(A_n)$  contains trivial T-avoiding elements and  $W(A_n)$  contains products of commuting generators. Now we will classify the non-trivial T-avoiding elements in  $W(A_n)$ . Recall that  $W(A_n)$  is a star reducible Coxeter group which by our observations above implies that if  $W(A_n)$  has non-trivial T-Avoiding elements, these elements will be not FC. In [5], the non-trivial T-avoiding elements were classified. The classification is seen in the following theorem.

**Theorem 4.2.1.** *Then there are no non-trivially T-avoiding elements in  $W(A_n)$ .*

*Proof.* This is a consequence of [5, Proposition 3.1.2.]. □

We will now proceed into the classification of T-avoiding elements in  $W(\tilde{A}_n)$ . Similar to  $W(A_n)$ ,  $W(\tilde{A}_n)$  will also have trivial T-avoiding elements since  $W(\tilde{A}_n)$  contains elements that are products of commuting generators. We will now proceed to classifying non-trivial T-avoiding elements in  $W(\tilde{A}_n)$ . The following classification is a result from [5].

**Theorem 4.2.2.** *If  $n \geq 2$  and  $n$  is even, then there are no non-trivial T-avoiding elements in  $W(\tilde{A}_n)$ . Otherwise, if  $n \geq 2$  and  $n$  is odd then  $W(\tilde{A}_n)$  contains non-trivial T-avoiding elements.*

*Proof.* This is [5, Proposition 3.1.2.]. □

This implies that in the Coxeter group of type  $\tilde{A}_n$ , when  $n$  is even there are no non-trivial T-avoiding elements, while when  $n$  is odd, there are non-trivial T-avoiding elements. One interesting observation to go along with this is  $W(\tilde{A}_n)$  is star reducible for  $n$  even but not star reducible for  $n$  odd. The classification seen in [5] did not specifically classify the non-trivial T-avoiding elements for type  $\tilde{A}_n$  for  $n$  odd. Before we state our conjecture as to what the non-trivial T-Avoiding elements are in  $W(\tilde{A}_n)$  we will discuss shortly what the heaps look like in  $W(\tilde{A}_n)$ . Because the Coxeter graph for type  $\tilde{A}_n$  is not a straight line Coxeter graph, the heaps can no longer be 2 dimensional and now must take on a 3 dimensional representation. The nature of the Coxeter graph of type  $\tilde{A}_n$  leads us to represent heaps in the sense of a turret on a castle where the blocks are stacked in a circular manner. We now continue with our conjectured classification. Because type  $A_n$  does not contain any non-trivial T-avoiding elements we know that the non-trivial T-avoiding elements in  $W(\tilde{A}_n)$  for  $n$  odd must have full support. We conjecture that the only non-trivial T-avoiding elements are castle turrets that have no missing blocks in the walls. However, this remains an open problem.

### 4.3 T-Avoiding Elements in Type $D_n$

In this section we will classify the T-avoiding elements in the Coxeter group of type  $D_n$ . We begin with trivial T-avoiding elements. Note that like  $W(A_n)$ ,  $W(D_n)$  has elements that are products of commuting generators. Hence,  $W(D_n)$  contains trivial T-avoiding elements. We will now classify the non-trivial T-avoiding elements of Coxeter groups of type  $D_n$ . Recall that  $W(D_n)$  is a star reducible Coxeter group and as a result of this any non-trivial T-avoiding element will not be fully commutative.

**Theorem 4.3.1.** *Let  $W = W(D_n)$ . Then  $W$  contains non-trivial T-avoiding elements.*

*Proof.* This is a consequence of [7, Section 2.2].  $\square$

In addition, to showing that there are non-trivial T-avoiding elements in type  $D_n$  Gern also classified the non-trivial T-avoiding elements as well. The following is his classification translated into heaps. **Once we figure out the heap include it here.** For the full details regarding his classification see [7]. Note that in his classification Gern refers to non-trivially T-avoiding elements as "bad."

#### 4.4 T-Avoiding Elements in Type $F_n$

In this section we classify what is known regarding the non-trivial T-avoiding elements in the Coxeter groups of type  $F_n$  for  $n \geq 4$ . Note that all of the following results are unpublished. First as with all other types we have seen so far,  $F_n$  has trivial T-avoiding elements since the groups contain products of commuting generators. For the rest of this section we will focus on the non-trivial T-avoiding elements in  $F_n$ . Recall that  $F$  is a star reducible Coxeter group so any non-trivial T-avoiding element in  $F_n$  will not be FC.

We start with the Coxeter group of type  $F_5$ . In 2012, Cross, Ernst, Hills-Kimball, and Quaranta classified all non-trivial T-avoiding elements in the following theorem.

**Theorem 4.4.1.** *An element in  $F_5$  is non-trivially T-avoiding if and only if it is a stack of bowties.*  $\square$

The above theorem references a stack of bowties. This refers to what the heap of the given non-trivial T-avoiding element looks like. We first restrict our attention to a single bowtie, this is seen in Figures ???. Note that in this figure, the orange blocks correspond to the elements that have bond strength 4 (Figure ???). Since the element is not FC we also wished to highlight the braid which is seen in the center of the element as colored in orange in Figure ???. In stacking the single bowties together, we get the stack of bowties referenced in the theorem which is seen in Figure ???. As a result of the classification in  $F_5$ , Cormier et. al. were also able to classify the non-trivial T-avoiding elements in  $F_4$ . The following is their classification.

**Corollary 4.4.2.** *There are no non-trivial T-avoiding elements in  $F_4$ .*  $\square$

As a result of their work, Cross et. al. conjectured that in Coxeter groups of type  $F_n$  for  $n \geq 5$ , an element is non-trivially T-avoiding if and only if it is a stack of bowties multiplied by a product of commuting generators. In 2013, Gilbertson and Ernst worked with this conjecture and quickly found out that it was incorrect. The heap seen in Figure ??? corresponds to a non-trivial T-avoiding element in  $F_6$ . It turns out that like the bowties discussed above these elements can also be stacked to

create and infinite number of non-trivial T-avoiding elements as well. In addition, as  $n$  gets large there are a number of things that can be altered in the group element that result in the element being non-trivially T-avoiding. This leads us to believe that there are potentially even more non-trivially T-avoiding elements in  $F_n$  for  $n \geq 6$ . From this we can see that the classification of T-avoiding elements in  $F_n$  for  $n \geq 6$  gets complicated very quickly. Hence, classifying T-avoiding elements in  $F_n$  for  $n \geq 5$  remains an open problem.

## 4.5 T-Avoiding Elements in Type $I_2(m)$

We next will classify the T-avoiding elements in Coxeter groups of type  $I_2(m)$ . Differing from the preceding classifications,  $W(I_2(m))$  has exactly 2 trivial T-avoiding elements as  $W(I_2(m))$  contains only 2 generators and as a result the only products of commuting generators are single generators. We now classify the non-trivial T-avoiding elements in  $W(I_2(m))$ . Although the following is a quick result, we believe that the result does not already appear in literature.

**Theorem 4.5.1.** *The Coxeter group  $W(I_2(m))$  has no non-trivial T-avoiding elements.*

*Proof.* The graph for the Coxeter group of type  $I_2(m)$  appears in Figure 1.1(c). Note that the graph consists of two vertices,  $s_1, s_2$ , and an edge with weight  $m(s, t)$ . First recall, that  $W(I_2(m))$  is a star reducible Coxeter group. This implies that any non-trivial T-avoiding elements in  $W(I_2(m))$  must not be FC. As all of the FC elements have Property T. We now consider the case when  $w \in W(I_2(m))$  is not FC. Since  $s_1$  and  $s_2$  are involutions this implies that the elements in  $W$  must alternate between  $s_1$  and  $s_2$ . Since  $w$  is not FC this implies that a reduced expression for  $w$  is  $w = \underbrace{s_1 s_2 \cdots s_1}_{m(s_1, s_2)}$ . By definition  $m(s_1, s_2) > 3$  and hence all elements in  $W(I_2(m))$  that

are not products of commuting generators have Property T, as they clearly have a reduced expression that begins with a product of non-commuting generators. Hence  $W(I_2(m))$  has no non-trivial T-avoiding elements.  $\square$

## Bibliography

- [1] S.C. Billey and B.C. Jones. Embedded factor patterns for Deodhar elements in Kazhdan–Lusztig theory. *Ann. Comb.*, 11(3–4):285–333, 2007.
- [2] A Björner and F Brenti. *Combinatorics of Coxeter groups*. 2005.
- [3] D.C. Ernst. Non-cancellable elements in type affine  $C$  Coxeter groups. *Int. Electron. J. Algebr.*, 8, 2010.
- [4] C.K. Fan. Structure of a Hecke algebra quotient. *J. Amer. Math. Soc.*, 10:139–167, 1997.
- [5] C.K. Fan and R.M. Green. On the affine Temperley–Lieb algebras. *Jour. L.M.S.*, 60:366–380, 1999.
- [6] M. Geck and G. Pfeiffer. *Characters of finite Coxeter groups and Iwahori–Hecke algebras*. 2000.
- [7] Tyson Gern. *Leading Coefficients of Kazhdan–Lusztig Polynomials in Type D*. Phd, University of Colorado, 2013.
- [8] R.M. Green. Star reducible Coxeter groups. *Glas. Math. J.*, 48:583–609, 2006.
- [9] R.M. Green. On the Markov trace for Temperley–Lieb algebras of type  $E_n$ . *J. Knot Th. Ramif.*, 18, 2007.
- [10] J.E. Humphreys. *Reflection Groups and Coxeter Groups*. 1990.
- [11] D. Kazhdan and G. Lusztig. Representations of Coxeter groups and Hecke algebras. *Inven. Math.*, 53:165–184, 1979.
- [12] G. Lusztig. Cells in affine Weyl groups, I. In *Collection*, pages 255–287. 1985.
- [13] J.R. Stembridge. On the fully commutative elements of Coxeter groups. *J. Algebr. Comb.*, 5:353–385, 1996.