

# **A Study of T-Avoiding Elements in Coxeter Groups**

Taryn Laird

Northern Arizona University  
Department of Mathematics and Statistics

**NAU Thesis Defense**

April 29, 2016

## Definition

A **Coxeter system** consists of a group  $W$  (called a **Coxeter group**) generated by a set  $S$  of involutions with presentation

$$W = \langle S \mid s^2 = e, (st)^{m(s,t)} = e \rangle$$

where  $m(s, t) \geq 2$  for all  $s \neq t$ .

## Definition

A **Coxeter system** consists of a group  $W$  (called a **Coxeter group**) generated by a set  $S$  of involutions with presentation

$$W = \langle S \mid s^2 = e, (st)^{m(s,t)} = e \rangle$$

where  $m(s, t) \geq 2$  for all  $s \neq t$ .

## Comment

Since  $s$  and  $t$  are involutions, the relation  $(st)^{m(s,t)} = e$  can be rewritten as

$$m(s, t) = 2 \implies st = ts \quad \} \quad \text{commutations}$$

## Definition

A **Coxeter system** consists of a group  $W$  (called a **Coxeter group**) generated by a set  $S$  of involutions with presentation

$$W = \langle S \mid s^2 = e, (st)^{m(s,t)} = e \rangle$$

where  $m(s, t) \geq 2$  for all  $s \neq t$ .

## Comment

Since  $s$  and  $t$  are involutions, the relation  $(st)^{m(s,t)} = e$  can be rewritten as

$$m(s, t) = 2 \implies st = ts \quad \} \quad \text{commutations}$$

$$\left. \begin{array}{l} m(s, t) = 3 \implies sts = tst \\ m(s, t) = 4 \implies stst = tsts \\ \vdots \end{array} \right\} \quad \text{braid relations}$$

## Definition

We can encode  $(W, S)$  with a unique Coxeter graph  $\Gamma$  having:

- vertex set  $S$ ;
- edges  $\{s, t\}$  labeled  $m(s, t)$  whenever  $m(s, t) \geq 3$ ;

## Definition

We can encode  $(W, S)$  with a unique Coxeter graph  $\Gamma$  having:


- vertex set  $S$ ;
- edges  $\{s, t\}$  labeled  $m(s, t)$  whenever  $m(s, t) \geq 3$ ;

## Comments

- if  $m(s, t) = 3$ , we omit label.
- If  $s$  and  $t$  are not connected in  $\Gamma$ , then  $s$  and  $t$  commute.
- Given  $\Gamma$ , we can uniquely reconstruct the corresponding  $(W, S)$ .


# Coxeter groups of type $A$

Coxeter groups of type  $A_n$  ( $n \geq 1$ ) are defined by:



# Coxeter groups of type A

Coxeter groups of type  $A_n$  ( $n \geq 1$ ) are defined by:




Then  $W(A_n)$  is generated by  $\{s_1, s_2, \dots, s_n\}$  and is subject to defining relations

1.  $s_i^2 = 1$  for all  $i$ ,
2.  $s_i s_j = s_j s_i$  if  $|i - j| > 1$ ,
3.  $s_i s_j s_i = s_j s_i s_j$  if  $|i - j| = 1$ .



# Coxeter groups of type A

Coxeter groups of type  $A_n$  ( $n \geq 1$ ) are defined by:



Then  $W(A_n)$  is generated by  $\{s_1, s_2, \dots, s_n\}$  and is subject to defining relations

1.  $s_i^2 = 1$  for all  $i$ ,
2.  $s_i s_j = s_j s_i$  if  $|i - j| > 1$ ,
3.  $s_i s_j s_i = s_j s_i s_j$  if  $|i - j| = 1$ .


$W(A_n)$  is isomorphic to the symmetric group,  $Sym_{n+1}$ , under the correspondence

$$s_i \mapsto (i, i + 1),$$

where  $(i, i + 1)$  is the adjacent transposition exchanging  $i$  and  $i + 1$ .


## Coxeter groups of type $B$

Coxeter groups of type  $B_n$  ( $n \geq 2$ ) are defined by:



## Coxeter groups of type $B$

Coxeter groups of type  $B_n$  ( $n \geq 2$ ) are defined by:




Then  $W(B_n)$  is generated by  $\{s_1, s_2, \dots, s_{n-1}\}$  and is subject to defining relations

1.  $s_i^2 = 1$  for all  $i$ ,
2.  $s_i s_j = s_j s_i$  if  $|i - j| > 1$ ,
3.  $s_i s_j s_i = s_j s_i s_j$  if  $|i - j| = 1$  and  $1 < i, j \leq n$ ,
4.  $s_0 s_1 s_0 s_1 = s_1 s_0 s_1 s_0$ .

## Coxeter groups of type $B$

Coxeter groups of type  $B_n$  ( $n \geq 2$ ) are defined by:




Then  $W(B_n)$  is generated by  $\{s_1, s_2, \dots, s_{n-1}\}$  and is subject to defining relations

1.  $s_i^2 = 1$  for all  $i$ ,
2.  $s_i s_j = s_j s_i$  if  $|i - j| > 1$ ,
3.  $s_i s_j s_i = s_j s_i s_j$  if  $|i - j| = 1$  and  $1 < i, j \leq n$ ,
4.  $s_0 s_1 s_0 s_1 = s_1 s_0 s_1 s_0$ .

$W(B_n)$  is a finite group of order  $n!2^n$  (wreath product of  $\mathbb{Z}_2$  and the symmetric group).


## Coxeter groups of type affine $C$

Coxeter groups of type  $\tilde{C}_n$  ( $n \geq 2$ ) are defined by:



## Coxeter groups of type affine $C$

Coxeter groups of type  $\tilde{C}_n$  ( $n \geq 2$ ) are defined by:




Here, we see that  $W(\tilde{C}_n)$  is generated by  $\{s_0, \dots, s_n\}$  and is subject to defining relations

1.  $s_i^2 = 1$  for all  $i$ ,
2.  $s_i s_j = s_j s_i$  if  $|i - j| > 1$ ,
3.  $s_i s_j s_i = s_j s_i s_j$  if  $|i - j| = 1$  and  $1 < i, j < n + 1$ ,
4.  $s_i s_j s_i s_j = s_j s_i s_j s_i$  if  $\{i, j\} = \{0, 1\}$  or  $\{n - 1, n\}$ .

## Coxeter groups of type affine $C$

Coxeter groups of type  $\tilde{C}_n$  ( $n \geq 2$ ) are defined by:




Here, we see that  $W(\tilde{C}_n)$  is generated by  $\{s_0, \dots, s_n\}$  and is subject to defining relations

1.  $s_i^2 = 1$  for all  $i$ ,
2.  $s_i s_j = s_j s_i$  if  $|i - j| > 1$ ,
3.  $s_i s_j s_i = s_j s_i s_j$  if  $|i - j| = 1$  and  $1 < i, j < n + 1$ ,
4.  $s_i s_j s_i s_j = s_j s_i s_j s_i$  if  $\{i, j\} = \{0, 1\}$  or  $\{n - 1, n\}$ .

$W(\tilde{C}_n)$  is an infinite group.

## Coxeter groups of type affine $C$

Coxeter groups of type  $\tilde{C}_n$  ( $n \geq 2$ ) are defined by:



Here, we see that  $W(\tilde{C}_n)$  is generated by  $\{s_0, \dots, s_n\}$  and is subject to defining relations

1.  $s_i^2 = 1$  for all  $i$ ,
2.  $s_i s_j = s_j s_i$  if  $|i - j| > 1$ ,
3.  $s_i s_j s_i = s_j s_i s_j$  if  $|i - j| = 1$  and  $1 < i, j < n + 1$ ,
4.  $s_i s_j s_i s_j = s_j s_i s_j s_i$  if  $\{i, j\} = \{0, 1\}$  or  $\{n - 1, n\}$ .

$W(\tilde{C}_n)$  is an infinite group.

### Comment

We can obtain  $W(A_n)$  and  $W(B_n)$  from  $W(\tilde{C}_n)$  by removing the appropriate generators and corresponding relations. In fact, we can obtain  $W(B_n)$  in two ways.



## Definition

A word  $s_{x_1}s_{x_2}\cdots s_{x_m} \in S^*$  is called an **expression** for  $w \in W$  if it is equal to  $w$  when considered as a group element.

## Definition

A word  $s_{x_1}s_{x_2}\cdots s_{x_m} \in S^*$  is called an **expression** for  $w \in W$  if it is equal to  $w$  when considered as a group element.

If  $m$  is minimal, it is a **reduced expression**, and the **length** of  $w$  is  $\ell(w) := m$ .

## Definition

A word  $s_{x_1}s_{x_2}\cdots s_{x_m} \in S^*$  is called an **expression** for  $w \in W$  if it is equal to  $w$  when considered as a group element.

If  $m$  is minimal, it is a **reduced expression**, and the **length** of  $w$  is  $\ell(w) := m$ .

Given  $w \in W$ , if we wish to emphasize a fixed, possibly reduced, expression for  $w$ , we represent it as

$$\overline{w} = s_{x_1}s_{x_2}\cdots s_{x_m}.$$

# Matsumoto's Theorem and Support

## Theorem (Matsumoto)

Any two reduced expressions for  $w \in W$  differ by a sequence of commutations and braid moves.

# Matsumoto's Theorem and Support

## Theorem (Matsumoto)

Any two reduced expressions for  $w \in W$  differ by a sequence of commutations and braid moves.

## Definition

We define  $\text{supp}(w)$  to be the set of generators appearing in any reduced expression for  $w$ . This is well defined by Matsumoto's theorem.

# Matsumoto's Theorem and Support

## Theorem (Matsumoto)

Any two reduced expressions for  $w \in W$  differ by a sequence of commutations and braid moves.

## Definition

We define  $\text{supp}(w)$  to be the set of generators appearing in any reduced expression for  $w$ . This is well defined by Matsumoto's theorem.

## Example

Let  $\overline{w} = s_2 s_1 s_2 s_3 s_1$  be a fixed expression for  $w \in W(A_3)$ . We see that

$$s_2 s_1 s_2 s_3 s_1$$

# Matsumoto's Theorem and Support

## Theorem (Matsumoto)

Any two reduced expressions for  $w \in W$  differ by a sequence of commutations and braid moves.

## Definition

We define  $\text{supp}(w)$  to be the set of generators appearing in any reduced expression for  $w$ . This is well defined by Matsumoto's theorem.

## Example

Let  $\overline{w} = s_2 s_1 s_2 s_3 s_1$  be a fixed expression for  $w \in W(A_3)$ . We see that

$$s_2 s_1 s_2 s_3 s_1 = s_1 s_2 s_1 s_3 s_1$$

# Matsumoto's Theorem and Support

## Theorem (Matsumoto)

Any two reduced expressions for  $w \in W$  differ by a sequence of commutations and braid moves.

## Definition

We define  $\text{supp}(w)$  to be the set of generators appearing in any reduced expression for  $w$ . This is well defined by Matsumoto's theorem.

## Example

Let  $\overline{w} = s_2 s_1 s_2 s_3 s_1$  be a fixed expression for  $w \in W(A_3)$ . We see that

$$s_2 s_1 s_2 s_3 s_1 = s_1 s_2 s_1 s_3 s_1 = s_1 s_2 s_1 s_1 s_3$$



# Matsumoto's Theorem and Support

## Theorem (Matsumoto)

Any two reduced expressions for  $w \in W$  differ by a sequence of commutations and braid moves.

## Definition

We define  $\text{supp}(w)$  to be the set of generators appearing in any reduced expression for  $w$ . This is well defined by Matsumoto's theorem.

## Example

Let  $\overline{w} = s_2 s_1 s_2 s_3 s_1$  be a fixed expression for  $w \in W(A_3)$ . We see that

$$s_2 s_1 s_2 s_3 s_1 = s_1 s_2 s_1 s_3 s_1 = s_1 s_2 s_1 s_1 s_3 = s_1 s_2 s_3$$

This implies that  $\overline{w}$  was not reduced. However, it turns out that  $s_1 s_2 s_3$  is a reduced expression for  $w$ . Then  $\text{supp}(w) = \{s_1 s_2 s_3\}$  and  $\ell(w) = 3$ .

# Fully Commutative Elements

## Definition

Let  $(W, S)$  be a Coxeter system of type  $\Gamma$ . We say that  $w \in W(\Gamma)$  is **fully commutative** (FC) if any two reduced expressions for  $w$  can be transformed into each other via iterated commutations. The set of FC elements is denoted  $FC(\Gamma)$ .

## Theorem (Stembridge)

$w \in FC(\Gamma)$  if and only if no reduced expression for  $w$  contains a braid.

## Comment

It follows from Stembridge that  $W(\tilde{C}_n)$  contains an infinite number of FC elements, while  $W(A_n)$  and  $W(B_n)$  do not.

# Fully Commutative Elements

## Comment

The elements of  $\text{FC}(\tilde{C}_n)$  are precisely those whose reduced expressions avoid the consecutive subwords  $s_i s_j s_i$  for  $m(s_i, s_j) = 3$ ,  $s_0 s_1 s_0 s_1$ , and  $s_{n-1} s_n s_{n-1} s_n$ .

# Fully Commutative Elements

## Comment

The elements of  $\text{FC}(\tilde{C}_n)$  are precisely those whose reduced expressions avoid the consecutive subwords  $s_i s_j s_i$  for  $m(s_i, s_j) = 3$ ,  $s_0 s_1 s_0 s_1$ , and  $s_{n-1} s_n s_{n-1} s_n$ .

## Example

Let  $\overline{w} = s_0 s_2 s_4 s_3 s_2 s_1$  be a reduced expression for  $w \in W(\tilde{C}_4)$ . We see that

$$s_0 \textcolor{red}{s_2} \textcolor{red}{s_4} s_3 s_2 s_1 = s_0 s_4 \textcolor{red}{s_2} \textcolor{red}{s_3} \textcolor{red}{s_2} s_1.$$

Since  $w$  has one of the forbidden consecutive subwords,  $w$  is **not** FC.

We introduce **heaps** through an example.

We introduce **heaps** through an example.

### Example


Let  $\overline{w} = s_4 s_5 s_1 s_0 s_2 s_4 s_1$  be a reduced expression for  $w \in W(B_6)$ .



We introduce **heaps** through an example.

### Example


Let  $\overline{w} = s_4 s_5 s_1 s_0 s_2 s_4 s_1$  be a reduced expression for  $w \in W(B_6)$ .



We introduce **heaps** through an example.

### Example

Let  $\overline{w} = s_4 s_5 s_1 s_0 s_2 s_4 s_1$  be a reduced expression for  $w \in W(B_6)$ .






We introduce **heaps** through an example.

### Example


Let  $\overline{w} = s_4 s_5 s_1 s_0 s_2 s_4 s_1$  be a reduced expression for  $w \in W(B_6)$ .



We introduce **heaps** through an example.

### Example


Let  $\overline{w} = s_4 s_5 s_1 s_0 s_2 s_4 s_1$  be a reduced expression for  $w \in W(B_6)$ .



We introduce **heaps** through an example.

### Example


Let  $\overline{w} = s_4 s_5 s_1 s_0 s_2 s_4 s_1$  be a reduced expression for  $w \in W(B_6)$ .



We introduce **heaps** through an example.

### Example


Let  $\overline{w} = s_4 s_5 s_1 s_0 s_2 s_4 s_1$  be a reduced expression for  $w \in W(B_6)$ .



We introduce **heaps** through an example.

### Example

Let  $\overline{w} = s_4 s_5 s_1 s_0 s_2 s_4 s_1$  be a reduced expression for  $w \in W(B_6)$ .



We introduce **heaps** through an example.

### Example

Let  $\overline{w} = s_4 s_5 s_1 s_0 s_2 s_4 s_1$  be a reduced expression for  $w \in W(B_6)$ .

