

# Chapter 7

## Integration

Unlike with differentiation, we will need a number of auxiliary definitions for beginning integration.

**Definition 7.1.** A set of points  $P = \{t_0, t_1, \dots, t_n\}$  is a *partition* of the closed interval  $[a, b]$  if  $a = t_0 < t_1 < \dots < t_{n-1} < t_n = b$ . If  $t_i - t_{i-1} = \frac{b-a}{n}$  for all  $i$ , we say that the partition is a *regular partition* of  $[a, b]$ . In this case, we may use the notation  $\Delta t := t_i - t_{i-1}$ .

**Exercise 7.2.** Give some partitions, regular and not regular, of  $[0, 1]$ ,  $[2, 4]$ , and  $[-1, 0]$ .

**Definition 7.3.** We say that a function is *bounded* if it has bounded image set.

*Important!* For the next four definitions, we assume that  $f$  is a bounded function with domain the closed interval  $[a, b]$ .

**Definition 7.4.** Let  $f$  be a bounded function with domain  $[a, b]$  and let  $\{t_0, t_1, \dots, t_n\}$  be a partition of  $[a, b]$ . We say that any sum  $S$  of the form

$$S = \sum_{i=1}^n f(x_i)(t_i - t_{i-1}),$$

where  $x_i \in [t_{i-1}, t_i]$  is a *Riemann sum* for  $f$  on  $[a, b]$ .

**Definition 7.5.** Let  $f$  be a bounded function with domain  $[a, b]$  and let  $P = \{t_0, t_1, \dots, t_n\}$  be a partition of  $[a, b]$ . For each  $i \in \{1, 2, \dots, n\}$ , define  $M_i := \sup\{f(x) \mid x \in [t_{i-1}, t_i]\}$ . We say that the sum

$$U_P(f) := \sum_{i=1}^n M_i(t_i - t_{i-1}),$$

is the *upper Riemann sum* for  $f$  with partition  $P$ .

**Definition 7.6.** Let  $f$  be a bounded function with domain  $[a, b]$  and let  $P = \{t_0, t_1, \dots, t_n\}$  be a partition of  $[a, b]$ . For each  $i \in \{1, 2, \dots, n\}$ , define  $m_i := \inf\{f(x) \mid x \in [t_{i-1}, t_i]\}$ . We say that the sum

$$L_P(f) := \sum_{i=1}^n m_i(t_i - t_{i-1}),$$

is the *lower Riemann sum* for  $f$  with partition  $P$ .

**Exercise 7.7.** Draw pictures that capture the concepts of upper and lower Riemann sums.

Contrary to the name, upper and lower Riemann sums are not always Riemann sums.

**Problem 7.8.** Give an example of an interval  $[a, b]$ , partition  $P$ , and bounded function  $f$  such that  $U_P(f)$  is not a Riemann sum.

**Problem 7.9.** Define  $f : [0, 1] \rightarrow \mathbb{R}$  via

$$f(x) = \begin{cases} 0, & x \in (0, 1] \\ 1, & x = 0. \end{cases}$$

- (a) Show that  $U_P(f) > 0$  for all partitions of  $[0, 1]$ .
- (b) Show that for any positive number  $\epsilon$  there is a partition  $P_\epsilon$  such that  $U_{P_\epsilon}(f) < \epsilon$ .
- (c) Fully describe all lower sums of  $f$  on  $[0, 1]$ .

**Problem 7.10.** Define  $f : [0, 1] \rightarrow \mathbb{R}$  via  $f(x) = x$ . For each  $n \in \mathbb{N}$ , let  $P_n$  be the regular partition of  $[0, 1]$  given by  $\{0, \frac{1}{n}, \frac{2}{n}, \dots, \frac{n-1}{n}, 1\}$ .

- (a) Compute  $U_{P_5}(f)$ .
- (b) Give a formula for  $U_{P_n}(f)$ .<sup>1</sup>
- (c) Compute  $L_{P_5}(f)$ .
- (d) Give a formula for  $L_{P_n}(f)$ .

**Problem 7.11.** Suppose that  $f$  is a bounded function on  $[a, b]$  with lower bound  $m$  and upper bound  $M$ . Show that for any partition  $P$  of  $[a, b]$ ,  $U_P(f) \leq M(b - a)$  and  $L_P(f) \geq m(b - a)$ .

**Problem 7.12.** Suppose that  $f$  is a bounded function on  $[a, b]$  and  $P$  is a partition of  $[a, b]$ . Show that  $L_P(f) \leq U_P(f)$ .

Note that as a result of the previous problems that the set of all upper and lower sums of  $f$  is a bounded point set.

More coming soon...

---

<sup>1</sup>Recall that the sum  $\sum_{i=1}^k i = \frac{k(k+1)}{2}$ .