

**PROTEOMIC CHARACTERIZATION OF THE 26S PROTEASOME IN
ARABIDOPSIS REVEALS INSIGHTS INTO COMPOSITION AND ASSEMBLY**

by

David C. Gemperline

A dissertation submitted in partial fulfillment of
the requirements for the degree of

Doctor of Philosophy

(Genetics)

at the

UNIVERSITY OF WISCONSIN-MADISON

2016

Date of final oral examination: September 26th 2016

The dissertation is approved by the following members of the Final Oral Committee:

Richard D. Vierstra, Professor, Genetics

Richard Amasino, Professor, Biochemistry

Josh Coon, Professor, Chemistry

Donna Fernandez, Professor, Botany

Patrick Masson, Professor, Genetics

Ahna Skop, Professor, Genetics

© Copyright by David C. Gemperline 2016

All Rights Reserved

To Erin, my wife.

ACKNOWLEDGMENTS

Science doesn't purvey absolute truth. Science is a mechanism, a way of trying to improve your knowledge of nature. It's a system for testing your thoughts against the universe and seeing whether they match. This works not just for the ordinary aspects of science, but for all of life.

— ISAAC ASIMOV (1988)

To my family, especially my wife Erin: I am forever grateful for your love and support, without which I would not be here today. The many adventures that we have already had are just the start of many more yet to come. To my parents, for instilling a curiosity in me that cannot be quenched, and for their unconditional love and support. To my father, Robert: for encouraging me to pursue the computational side of science and to my mother, Terri: for always encouraging me to teach others. To my brother, Aaron, for our many discussions, on science, life, and playing games. To my twin, Stephen, for always reminding me to have fun. To my good friend, Lucas, for all of your guidance and support when I was completely lost in code.

My scientific colleagues and mentors, you have pushed me to become a better scientist and to always examine the world with a critical eye. Rick Vierstra, thank you for always pushing me, and helping me realize that we can always improve.

Patrick Masson, thank you for providing a welcoming home during a time of change.

I owe much of my mentorship to the Vierstra Lab. Richard Marshall, you are a great friend, and a great mentor. Without your guidance, I would have been lost. To Nick Gladman, for always listening to my crazy ideas, and for being a great friend. To Aaron Lomax, and Robb Stankey for your companionship while drinking coffee. I will always look back on my time in Madison fondly. To my friends, Richard, Kurt, Carol, and others: thank you for always being there.

TABLE OF CONTENTS

Table of Contents	iv
List of Tables	x
List of Figures	xii
List of Abbreviations and Acronyms	xvii
Abstract	xxi
Chapter 1: Introduction to the Ubiquitin 26S Proteasome System	1
1.1 The Ubiquitin 26S Proteasome System (UPS)	1
1.2 The 26S Proteasome	6
1.3 Proteasome Purification Strategies	12
1.4 26S Proteasome Substrate Processing	18
1.5 Proteasome Assembly	21
1.5.1 20S Core Protease Assembly	21
1.5.2 19S Regulatory Particle Assembly RP base	33
1.5.3 19S Regulatory Particle Assembly RP lid	37
1.5.4 RP-CP Assembly and Alternative Capping Particles	39

1.5.5	Proteasome Assembly in Plants	39
1.6	Proteasome Isotypes	41
1.6.1	Immunoproteasome	41
1.6.2	Thymoproteasome proteasome	42
1.6.3	Testes-specific proteasome	43
1.6.4	$\alpha 4\text{-}\alpha 4$ proteasome Isotype	44
1.6.5	Plant Proteasome Isotypes	46
1.7	Conclusions	47
1.8	Literature Cited	51

Chapter 2: Morpheus Spectral Counter: A Computational Tool for Label-Free Quantitative Mass Spectrometry using the Morpheus Search Engine **66**

2.1	Summary	66
2.2	Main Text	67
2.3	Methods	81
2.3.1	Sample Preparation	81
2.3.2	Liquid Chromatography and High-Resolution Mass Spectrometry	83
2.3.3	Data Processing	84

2.3.4 Isoform Incorporation Rates	86
2.3.5 Speed and Accuracy Comparisons of Morpheus/MSpC to the TPP/ABACUS Pipeline	86
2.4 Discussion on Requiring Two Unique Peptides to Quantify a Protein	88
2.5 Tutorial	90
2.6 Acknowledgements	94
2.7 Literature Cited	94
 Chapter 3: Proteomic Characterization of Affinity-Purified Arabidopsis Pro- teasome Particles Identifies Sub-Complex Specific Interactors	
3.1 Statement of Attribution	97
3.2 Abstract	98
3.3 Introduction	100
3.4 Experimental Procedures	106
3.4.1 Transgenic Plants and Growth Conditions	106
3.4.2 Sequence and Phylogenetic Analyses	110
3.4.3 Proteasome Affinity Purifications using PAG1-FLAG and FLAG- RPT4a/b	111
3.4.4 SDS-PAGE and Native-PAGE Analysis and Silver Staining . .	112
3.4.5 MS/MS Sample Preparation	113

3.4.6	MS/MS Analyses	115
3.4.7	MS/MS Data Processing	117
3.4.8	Yeast-two hybrid, bimolecular fluorescent complementation, and HA-PAP1 immunopurification	119
3.5	Results	121
3.5.1	<i>Arabidopsis</i> Proteasomes Can Be Effectively Affinity-Purified Using Epitope-Tagged Versions of Either Isoform of the Reg- ulatory Particle Subunit RPT4	121
3.5.2	MS/MS Analyses of RPT4a/b Affinity-Purified Proteins Iden- tifies a Similar Complement of Proteins as PAG1-FLAG Affin- ity Purifications	128
3.5.3	Statistical Analysis of Affinity Enrichments for FLAG-RPT4a Show Little to No Difference When Compared to FLAG-RPT4b	133
3.5.4	Statistical Analyses of LFQ-MS/MS Data Show Affinity Prepa- rations Targeting Either the CP or RP Specifically Enrich for Their Respective Subcomplexes	135
3.5.5	Statistical Analysis of Affinity Purifications Enriched for the RP or CP Identifies Proteasome-Associated Proteins and Pu- tative Assembly Chaperones Specific to Each Subcomplex .	139

3.5.6 A Novel Plant Proteasome-Associated Protein, PAP1, Interacts with a Putative CP Assembly Chaperone	142
3.5.7 Inhibition of the Proteasome Results in Formation of Distinct Proteasome Sub-Species Containing Putative CP Assembly Chaperones	147
3.6 Discussion	153
3.7 Literature Cited	168
Chapter 4: Future Directions and Closing Remarks	212
4.1 Introduction	212
4.2 Future Directions	215
4.2.1 Proteasome Purifications	215
4.2.2 CP Assembly Chaperones	217
4.2.3 PAP1	223
4.2.4 RP Assembly Chaperones	229
4.2.5 Identifying CP Assembly Intermediates	229
4.2.6 Isoform-Specific Proteasomes	231
4.3 Concluding Remarks	232
4.4 Literature Cited	234

LIST OF TABLES

2.1 Table of dNSAF values for each 20S proteasome subunit generated by analyzing the proteasome spike in experiments with the Morpheus and MSpC pipeline	79
2.2 Table of adj_NSAF values for each 20S proteasome subunit (equivalent to dNSAF) generated by analzying the proteasome spike in experiments with the TPP and ABACUS pipeline	80
3.S1 Oligonucleotide Primer Sequences Used in this Study	201
3.S2 Contaminants Removed from MaxQuant's ProteinGroups.txt file . . .	202
3.S3 InterProScan analysis of <i>Arabidopsis</i> protein sequences for PBAC1-4 . .	203
3.S4 Iterative PSI-BLAST Analysis using Human PAC1-4 to Identify PBAC1-4	204
3.S5 Proteasome associated proteins identified after PAG1-based affinity pu- rification under minus ATP Conditions	205
3.S6 Proteasome associated proteins identified after PAG1-based affinity pu- rification under plus ATP Conditions	206
3.S7 Proteasome associated proteins identified after RPT4a-based affinity purification under minus ATP conditions	207

3.S8 Proteasome associated proteins identified after RPT4a-based affinity purification under plus ATP conditions	208
3.S9 Proteasome associated proteins identified after RPT4b-based affinity purification under minus ATP conditions	209
3.S10 Proteasome associated proteins identified after RPT4b-based affinity purification under plus ATP conditions	210
3.S11 Fold Change dNSAF values comparing Slice 1 and Smear from Figure 3.6	211
4.1 Candidate alleles for PBAC1-4	221

LIST OF FIGURES

1.1 A schematic representation of the E1 activating, E2 conjugating, and E3 ligase enzymatic cascade for the covalent attachment of the small polypeptide Ubiquitin to target protein lysines	3
1.2 The 26S proteasome is made up of two functionally distinct sub-particles.	7
1.3 The 20S Core Protease (CP)	9
1.4 Cryo EM Structure of the 19S Regulatory Particle (RP)	12
1.5 Electron microscopy images of 20S and 26S proteasomes from mammals and plants	14
1.6 Affinity purification of 26S proteasomes from <i>PAG1::PAG1-FLAG pag1-1</i> plants	19
1.7 Schematic representation of CP Assembly	24
1.8 Pba1 and Pba2 binding sites via their HbYX motifs into adjacent α -subunit pairs	30
1.9 Structural Diagram of opposing Pba1/2 Pba3/4 binding	32
1.10 Schematic representation of RP Assembly	36
1.11 Representation of subunits replaced for both the immunoproteasome and the thyoproteasome	45

1.12 Solvent accessible tag locations on the 26S proteasome for both PAG1 and RPT4	50
2.1 MSpC Graphic User Interface (GUI) and software flow chart	69
2.2 Confirmation of MSpC accuracy by analysis of MS/MS datasets generated with the Universal Proteome Standard 2 (UPS2)	71
2.3 Re- Analysis of MS/MS datasets generated with the Universal Proteome Standard 2 (UPS2)	73
2.4 Confirmation of MSpC accuracy by analysis of MS/MS datasets generated with affinity purified <i>Arabidopsis</i> 20S proteasomes spiked into a total cell lysate from <i>E. coli</i> . Following MS/MS analysis, the dNSAF and uNSAF values for each subunit/isoform were determined by Morpheus and MSpC	76
2.5 MSpC combined with Morpheus works faster than TPP combined with ABACUS. Speed comparisons were performed for 12, 24, 48, and 96 raw MS/MS files generated with the 20S proteasome/ <i>E. coli</i> lysate samples analyzed in 2.4	82
3.1 <i>rpt4a-1</i> and <i>rpt4b-2</i> mutants rescued with flag tagged variants	123

3.2 Protein composition of affinity-purified proteasomes from FLAG-RPT4a-, FLAG-RPT4b-, and PAG1-FLAG-expressing plants and from mock wild-type controls in the presence (+) or absence (-) of 20mM ATP	127
3.3 Label-free quantitative MS/MS analysis of affinity-purified proteins in the presence (+) or absence (-) of ATP	130
3.3 Label-free quantitative MS/MS analysis of affinity-purified proteins in the presence (+) or absence (-) of ATP	131
3.3 Label-free quantitative MS/MS analysis of affinity-purified proteins in the presence (+) or absence (-) of ATP	132
3.4 Volcano Plots comparing proteasome affinity purifications under both plus and minus ATP conditions relative to WT controls, allow identification of statistically significant interacting partners for the proteasome .	136
3.4 Volcano Plots comparing proteasome affinity purifications under both plus and minus ATP conditions relative to WT controls, allow identification of statistically significant interacting partners for the proteasome .	137
3.5 PAP1 interacts with the proteasome and with putative assembly chaperone PBAC1	145
3.6 Protein composition of PAG1-FLAG affinity-purified proteasomes from tissues treated with and without MG132	149

3.6 Protein composition of PAG1-FLAG affinity-purified proteasomes from tissues treated with and without MG132	150
3.7 Putative Model of 20S Proteasome Assembly in <i>Arabidopsis</i>	166
3.S1 Sequence Alignments of RPT4a and RPT4b	177
3.S2 Volcano Plots Comparing Protein Profiles of Proteasome Affinity Purifications Based on RPT4a or RPT4b subunits in the presence (+) or absence (-) of ATP	178
3.S3 Kernel Density Estimates (KDE) of the CP and RP distributions from label-free quantitative mass spectrometry show enrichment for the CP when purified via PAG1-FLAG, and for the RP when purified via FLAG-RPT4a/b	179
3.S4 Multiple Sequence Alignment for UMP1	180
3.S4 Multiple Sequence Alignment for UMP1	182
3.S5 Multiple Sequence Alignment for PBAC1	183
3.S5 Multiple Sequence Alignment for PBAC1	185
3.S6 Multiple Sequence Alignment for PBAC2	186
3.S6 Multiple Sequence Alignment for PBAC2	188
3.S7 Multiple Sequence Alignment for PBAC3	189
3.S7 Multiple Sequence Alignment for PBAC3	191

3.S8 Multiple Sequence Alignment for PBAC4	192
3.S8 Multiple Sequence Alignment for PBAC4	194
3.S9 Multiple Sequence Alignment for PAP1	196
3.S10 Full BiFC Panel and Controls	197
3.S11 Phylogenetic Analysis of PAP1 and PBAC1-4	198
3.S11 Phylogenetic Analysis of PAP1 and PBAC1-4	200
4.1 Recombinant protein expression of PAP1 and PBAC1-3	220
4.2 Insertion alleles for <i>PAP1</i> and qRT-PCR analysis	225
4.3 Glycerol gradient centrifugation analysis of <i>pap1-3</i>	227

LIST OF ABBREVIATIONS AND ACRONYMS

ACN	acetonitrile
ATP	adenosine triphosphate
BLAST	Basic local alignment search tool
C#	C sharp, a programming language
CP	Core Protease
Da	Dalton, the atomic mass unit
DMSO	Dimethyl sulfoxide
dNSAF	distributed Normalized Spectral Abundance Factor
DTT	Dithiothreitol
DUB	De-ubiquitylating enzyme
E1	Ubiquitin Activating Enzyme
E2	Ubiquitin Conjugating Enzyme
E3	Ubiquitin Ligase
ESI	Electrospray ionization
E-value	Expectation value
FA	formic acid
FASTA	A format for storing protein sequences
FDR	False discovery rate

GB	Gigabytes
GUI	Graphical user interface
h	Hour
HbYX	Hydrophobic amino acid, followed by a Tyrosine, followed by any amino acid
HCD	Higher-energy collisional dissociation
HEPES	4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid
HPLC	High-performance liquid chromatography
JA	Jasmonic acid
kDa	Kilodalton, the atomic mass unit
KS	Kolmogorov-Smirnov
LC	Liquid chromatography
LFQ	Label-free quantification
<i>m/z</i>	mass to charge ratio
MES	2-(N-morpholino)ethanesulfonic acid
min	Minute
MS/MS	Tandem mass spectrometry
MS	Mass spectrometry
MS1	Survey mass analysis
MS2	Fragment mass spectrum

MSpC	Morpheus Spectral Counter
NCE	Normalized collision energy
nLC	Nanoflow liquid chromatography
NSAF	Normalized Spectral Abundance Factor
PAGE	Polyacrylamide gel electrophoresis
PCR	Polymerase chain reaction
PDB	Protein Data Bank
PGPH	Peptidylglutamyl-peptide hydrolyzing
pK _a	Acid dissociation constant
PMSF	Phenylmethane sulfonyl fluoride
ppm	Part per million
PRIDE	PRoteomics IDEntifications database
PSI-BLAST	Position-specific iterated basic local alignment search tool
PSM	Peptide-spectrum match
PTM	Post-translational modification
.raw	Thermo Raw File
RP	Regulatory Particle
RT-PCR	Reverse transcription polymerase chain reaction
S/N	Signal-to-noise ratio

s	Second
SDS	Sodium dodecyl sulfate
SILAC	Stable isotope labeling by amino acids in cell culture
SILAC	Stable isotope labeling by amino acids in cell culture
TAIR10	The Arabidopsis Information Resource v10
TEMED	N,N,N',N'-tetramethylethylenediamine
TFA	trifluoroacetic acid
TMT	Tandem mass tag
TPP	Trans-proteomic pipeline
UBA	ubiquitin associated
UBL	ubiquitin like
uNSAF	unique Normalized Spectral Abundance Factor
UPLC	ultra-high performance liquid chromatography
UPS	Ubiquitin 26S Proteasome System
UPS2	Universal Proteomics Standard 2
UTR	Untranslated region
Y2H	Yeast-two-hybrd
YFP	Yellow fluorescent protein

ABSTRACT

Abstract

The 26S proteasome is the central proteolytic effector in the ubiquitin system that is responsible for degrading numerous regulators following their selective ubiquitylation. The 26S proteasome is made up of two distinct subcomplexes, the Regulatory Particle (RP), responsible for recognizing and unfolding ubiquitylated substrates, and the Core Protease (CP), responsible for degrading the unfolded polypeptide. Herein, I developed an affinity purification of the 26S proteasome that targeted the RP via two isoforms of the RPT4 subunit. With this newly developed affinity purification, in conjunction with a previously developed CP-based affinity purification of 26S proteasome, I was able to selectively enrich for either the RP or CP subcomplexes. Mass spectrometric analyses of these affinity purified complexes identify a large collection of interacting proteins that associate with either the core protease (CP) or regulatory particle (RP). While much is known about the construction of the yeast and mammalian particles, little is known about the pathways used to assemble the plant particle. Sequence comparisons, Y2H and BiFC studies revealed that some proteasome associated proteins are likely assembly chaperones, with several CP factors harboring a signature C-terminal HbYX motif that typically enables association with the α -subunit ring. Several of the RP-specific factors ap-

pear to be orthologs of the chaperones Nas2, Nas6, Hsm3. Whereas yeast assembles only a single particle type, mammals can assemble alternate proteasomes isotypes by replacing individual subunits with distinct isoforms (e.g., immunoproteasomes). In plants, most 26S proteasome subunits are encoded by paralogous genes with sufficient divergence to suggest that plants also accumulate a collection of particles. However, proteomic analysis of proteasomes selectively enriched using paralog-specific tags strongly imply that although plants possess this genetic diversity, the incorporation of these paralogs appears random. Taken together, these proteomic studies provide the first insights into plant proteasome assembly and diversity, and identify factors that may build the CP and RP subcomplexes and finally the 26S holo-particle.

Chapter 1

INTRODUCTION TO THE UBIQUITIN 26S PROTEASOME SYSTEM

Adapted in part from: R.S. Marshall*, D.C. Gemperline*, and R.D. Vierstra (2015). Purification of 26S proteasomes and their sub-complexes from plants. *Methods Mol. Biol.* DOI: 10.1007/978-1-4939-6533-5

*These authors contributed equally to the *Methods Mol. Biol.* work.

This chapter has expanded sections on the proteasomes role in plant biology, proteasome assembly, and proteasome isotypes.

1.1. The Ubiquitin 26S Proteasome System (UPS)

Selective proteolysis in plants plays a critical role in both regulating growth and development, and maintaining cellular homeostasis (Nelson *et al.*, 2014; Smalle and Vierstra, 2004; Vierstra, 1993, 2009). One of the principle pathways for protein degradation in plants and other eukaryotes is the ubiquitin-26S proteasome system

(UPS), which involves the attachment of polyubiquitin chains to target proteins followed by their recognition and degradation by the 26S proteasome, an exquisitely designed proteolytic machine (Bhattacharyya *et al.*, 2014; Finley, 2009; Vierstra, 2009). The UPS is highly conserved across all eukaryotes; it was first elucidated by elegant work in rabbit reticulocyte lysates (Ciechanover *et al.*, 1980a,b; Etlinger and Goldberg, 1977; Hershko *et al.*, 1980; Wilkinson *et al.*, 1980), and was subsequently identified in other animals, yeast and plants (Ciechanover *et al.*, 1984; Finley *et al.*, 1984, 1987; Glotzer *et al.*, 1991; Hochstrasser *et al.*, 1991; Shanklin *et al.*, 1987). Ubiquitin conjugation to target proteins is accomplished through a highly polymorphic, ATP-dependent cascade involving the sequential action of three enzyme classes, termed the E1 ubiquitin-activating enzymes, E2 ubiquitin-conjugating enzymes, and E3 ubiquitin-protein ligases (Berndsen and Wolberger, 2014; Smalle and Vierstra, 2004; Vierstra, 2009) (Figure 1.1). Selectivity in ubiquitylation is typically driven by the E3 family, which has dramatically expanded during plant evolution to include well over a thousand variants in Arabidopsis and other plant species (Hua *et al.*, 2013; Hua and Vierstra, 2011). Indeed 6% of the Arabidopsis proteome is devoted to UPS (Smalle and Vierstra, 2004).

The UPS helps regulate plant growth and development by regulating several key proteins including those involved in morphogenesis, light sensing, and the




Figure 1.1: A schematic representation of the E1 activating, E2 conjugating, and E3 ligase enzymatic cascade for the covalent attachment of the small polypeptide Ubiquitin to target protein lysines. Once a protein becomes polyubiquitylated it is efficiently recognized by the 26S proteasome. The rate of addition of ubiquitins, and the rate of removal by de-ubiquitylating enzymes (DUBs) determines the final equilibrium of degradation.

circadian clock (reviewed in (Vierstra, 2009). Some examples include E3 enzymes such as BIG BROTHER that controls organ size (Disch *et al.*, 2006), F-BOX-LIKE 17 that helps regulate spermatogenesis (Kim *et al.*, 2008), and UNUSUAL FLORAL ORGANS which regulates floral development (Samach *et al.*, 1999). The UPS also helps regulate plants response to light by degrading phytochrome A (Clough and Vierstra, 1997; Shanklin *et al.*, 1987), a key photoreceptor in plants. An E3 ligase ZTL also aides in removing TOC1 involved in circadian clock regulation (Más *et al.*, 2003).

Additionally, the UPS also aides in the regulation of key plant hormonal responses including auxin, abscisic acid, jasmonic acid (JA), giberellins (GA), and ethylene. (reviewed in (Santner and Estelle, 2010)). Key hormone response factors are degraded by a variety of E3 complexes including TIR1 involved in auxin regulation (Dharmasiri *et al.*, 2005), COI1, involved JA sensing (Katsir *et al.*, 2008), and GID1 involved in GA responses (Murase *et al.*, 2008). An elegant mechanism requires the direct binding of these key hormones either to their respective E3 complexes, or key proteins associated with these E3's to promote the efficient ubiquitylation and subsequent degradation of their hormone response factors (Shabek and Zheng, 2014).

The UPS is both co-opted and inhibited by several plant pathogens that helps

aide in their virulence. The plant pathogen *Ralstonia solanacearum* encodes a suite of E3 enzymes that co-opt host machinery to evade plant defense responses (Angot *et al.*, 2006). Additionally, *Pseudomonas syringae*, a tomato pathogen, secretes a small molecule that irreversibly binds the host 26S proteasome to prevent plants from actively ubiquitylating and degrading key virulence factors (Schellenberg *et al.*, 2010).

Self-incompatibility is a key process in plants that encourages plants to outcross helping aide in plant evolution and diversity by preventing inbreeding (Zhang *et al.*, 2009). Upon incompatible pollination an E3 enzyme ARC1 helps to ubiquitylate compatibility factors in the pistil leading to self-pollen rejection (Stone *et al.*, 2003). In another example S-RNase, which acts to degrade the RNA from self-pollen tubes preventing their growth, is actively ubiquitylated via SLF and SSK1 (McClure, 2004; Zhao *et al.*, 2010). From these and other analyses it is clear that the UPS either directly, or indirectly, regulates many aspects of plant growth and development including hormone responses, pathogen interactions, and self-incompatibility. Therefore, a fundamental understanding of the UPS system in plants has important agricultural impacts.

1.2. The 26S Proteasome

The 26S proteasome is the central enzymatic effector of the UPS, and is a 2.5 MDa particle located in the cytosol and nucleus of eukaryotic cells. It is composed of two functionally distinct sub-complexes; the 20S core protease (CP) that houses the proteolytic active sites, and the 19S regulatory particle (RP) that recognizes appropriate substrates (Figure 1.2 A and B; (Bhattacharyya *et al.*, 2014; Finley, 2009; Lander *et al.*, 2012; Lasker *et al.*, 2012; Unverdorben *et al.*, 2014)).

The CP has a barrel shape generated by four stacked hetero-heptameric rings, which contain seven α -subunits or seven β -subunits (termed PAA-PAG and PBA-PBG, respectively, in *Arabidopsis*) in an α 1-7/ β 1-7/ β 1-7/ α 1-7 configuration (Figure 1.3 A (side view- stacked rings), C (top view- α -subunits) and D (top view- β -subunits).

Upon assembly, a central chamber is formed at the β -ring interface that houses six peptidase catalytic sites provided by the β 1 (PBA), β 2 (PBB), and β 5 (PBE) subunits (Arendt and Hochstrasser, 1997; Heinemeyer *et al.*, 1997) (Figure 1.3 B and E). The active sites involve a catalytic triad, one residue of which is an N-terminal threonine that becomes exposed during CP assembly. Collectively these peptidases can cleave a broad range of protein sequences with peptidylglutamyl-peptide hydrolyzing (PGPH) (β 1), trypsin-like (β 2), and chymotryptic-like (β 5)





Figure 1.2: The 26S proteasome is made up of two functionally distinct sub-particles.. (A) The Core Protease (CP) is shown in red, while the Regulatory Particle (RP) is shown in blue for the RPT subunits, and in gold for the RPN sub-units (adapted from (Lasker *et al.*, 2012)). (B) Schematic representation of CP and RP functions. The RP is responsible for recognizing and unfolding polyubiquitylated substrates into a central axial chamber upon which the unfolded polypeptide is cleaved by the CP. The RP is also responsible for releasing the ubiquitin so that it can be recycled for re-use through the de-ubiquitylating activity of the metalloprotease RP subunit RPN11.

[Figure 1.3 caption follows on next page]



activities (Arendt and Hochstrasser, 1997; Groll *et al.*, 1999) (Figure 1.3 E). The α -rings create two antechambers with narrow opposing axial pores that are gated by extensions at the N-terminus of several subunits (Figure 1.3 B) (Groll *et al.*, 2000; Ruschak *et al.*, 2010). Through this distinctive architecture, the CP acts as a self-compartmentalized protease that will only degrade polypeptides that are deliberately recognized, unfolded, and imported into the β -ring chamber.


The CP is capped at one or both ends by the RP, which sits on top of the axial pores. The RP provides activities for recognition of ubiquitylated proteins, substrate unfolding and import, and release of the ubiquitin moieties before substrate degradation (Figure 1.2). Its binding to the CP is stabilized by ATP, which is thus a

Figure 1.3 (preceding page): The 20S Core Protease (CP). (A) An x-ray crystallographic structure of the 20S CP shows its barrel-like structure. The structure is composed of four stacked hetero-heptameric rings in an α 1-7, β 1-7, β 1-7, α 1-7 configuration. (B) This stacked configuration generates an interior structure amenable to cleaving unfolded polypeptides that enter through the axial pore. N-terminal extensions present in α -subunits gate the entry to the antechamber. The antechamber helps maintain the polypeptide in an unfolded state, and the β -ring chamber houses the catalytic activities of the CP. (C) Top-down view of the CP showing the α -subunit configuration (PDB ID 5CZ4) visualized in PyMOL (Schrodinger, 2015). (D) Top-down view of the CP showing the β -subunit configuration (PDB ID 5CZ4) visualized in PyMOL. (E) Catalytic activities of the 20S proteasome are conferred by the β 1 (peptidylglutamyl-peptide hydrolyzing (PGPH)), β 2 (Trypsin-like), and β 5 (Chymotrypsin-like) subunits. A slice through the middle of a model of the 20S proteasome (adapted from (Kisselev *et al.*, 2012)) shows the location and type of the three catalytic activities of the 20S proteasome. While this is a half-structure, each barrel actually houses 6 catalytically active locations, with three different catalytic activities.


necessary ingredient for purifying intact 26S proteasomes (Smith *et al.*, 2005). The RP itself consists of two sub-complexes; the base, which contains a hexameric ring of AAA-ATPases (RPT1-6) plus two non-ATPase subunits, RPN1 and RPN2; and the lid, which is composed of an additional 11 non-ATPase subunits, RPN3, RPN5-13 and DSS1/SEM1 (Figure 1.2; (Bhattacharyya *et al.*, 2014; Book *et al.*, 2010; Finley, 2009; Glickman *et al.*, 1998b; Russell *et al.*, 2013)). This lid/base demarcation was first revealed by the absence of lid subunits in proteasomes isolated from a *rpn10* yeast deletion strain, and it was hence thought that RPN10 helps enforce binding of the lid to the base (Glickman *et al.*, 1998a). However, more recent structural studies have demonstrated that RPN10 has a more indirect stabilizing role via its interaction with RPN9 (Lander *et al.*, 2012). The ring of RPT subunits in the base promotes substrate unfolding through ATP hydrolysis, and gates the α -ring axial pores through repositioning of the CP α -subunit extensions (Köhler *et al.*, 2001; Rabl *et al.*, 2008; Smith *et al.*, 2005). The N-terminal regions from proximal RPT pairs intertwine to create three spokes onto which most RPN subunits are scaffolded (Figure 1.4 A; (Beck *et al.*, 2012)).

Several RPT subunits harbor C-terminal *Hydrophobic-Tyrosine-Any* amino acid (HbYX) motifs important for their binding to the α -subunits in the CP. These C-terminal HbYX and HbYX-like extensions typically intercalate between adjacent

[Figure 1.4 caption follows on next page]



B



→ HbYX /
HbYX-like
C-terminal Extensions

α -subunits (Figure 1.4 B). The RPN6 subunit acts as a molecular clamp to tether the RP onto the CP (Figure 1.4 A) (Pathare *et al.*, 2012).

1.3. Proteasome Purification Strategies

Even before the realization that the 26S proteasome is a protease, sub-particles of the complexes were described. The first reports of proteasomes used avian erythroblast preparations enriched by differential ultracentrifugation followed by fractionation through a sucrose gradient (Schmid *et al.*, 1984). These 20S fractions isolated in the absence of added ATP were found to inhibit mRNA translation in a cell-free system, leading to early proposals that the identified complex repressed gene expression through a cryptic ribonuclease activity. This lead to the particle initially being named the “prosome” (Kremp *et al.*, 1986; Schmid *et al.*, 1984). Subsequent analyses of these preparations by SDS-PAGE and electron microscopy revealed the signature ladder of α - and β -subunits at 20-35 kDa, as well as their barrel-like architecture

Figure 1.4 (preceding page): Cryo EM Structure of the 19S Regulatory Particle (RP). (A) The RP structure as solved by Cryo-EM is shown (modified from (Lander *et al.*, 2012)). In the center in blue are the RPT1-6 subunits, while the RPN subunits are individually labeled. A 90-degree rotation is shown so that the other side of the RP can be visualized. (B) A schematic representation of how the RP subunit C-termini containing HbYX and HbYX-like motifs intercalate between the α -subunits (adapted from (Sokolova *et al.*, 2015)). Gaps between each α -subunit are shown, and the HbYX motifs are shown as red strands.

(Figure 1.5 A and B; (Baumeister *et al.*, 1988; Kremp *et al.*, 1986; Schmid *et al.*, 1984)).

Purification of the 20S fraction from HeLa cells followed by SDS-PAGE also gave rise to this stereotypical protein banding pattern and shape (Schmid *et al.*, 1984), and this was followed shortly thereafter by the first description of plant prosomes, purified from tobacco leaf extracts using similar sedimentation protocols in ATP-free buffers (Kremp *et al.*, 1986). In these later cases, the purified preparations had strong peptidase activity but little to no RNase activity, thus leading to the conclusion that the CP is actually a protease. Once its true function in protein turnover was confirmed, the moniker for the particle was changed to “proteasome” (Arrigo *et al.*, 1988).

Subsequently, the 20S particle was purified from other plant tissues, including dry pea seeds, potato tubers, mung bean seedlings, and leaves from both spinach and wheat (Murray *et al.*, 1997; Ozaki *et al.*, 1992; Schliephacke *et al.*, 1991; Skoda and Malek, 1992). These purifications were typically performed using sequential anion exchange and size-exclusion chromatography steps in the absence of ATP, hence only the CP was isolated. Their remarkable similarity in protein composition and structure, as observed by SDS-PAGE and electron microscopy, respectively, coupled with the fact that several of the plant subunits cross-reacted with antibodies against their yeast, human, rat and *Xenopus* counterparts, strongly implied that the CP was




Figure 1.5: Electron microscopy images of 20S and 26S proteasomes from mammals and plants. (A) Images of 20S proteasomes purified from rat skeletal muscle. On the left is an electron micrograph of the 20S particles negatively stained with sodium phosphotungstate, while on the right is a close-up image with overlaid contour plots generated by correlation averaging of approximately 300 individual images negatively stained with ammonium molybdate. (B) Images of the first 20S proteasomes purified from different plant species. On the left are proteasomes isolated from tobacco leaves, while on the right are proteasomes from potato tubers, both negatively stained with uranyl acetate. The typical barrel-shaped structures are indicated with red circles. (C) Images of 26S proteasomes purified from rat liver. On the left is an electron micrograph of the 26S particles negatively stained with uranyl acetate, while on the right is a close-up image with overlaid contour plots generated by correlation averaging of 215 individual images. (D) Images of 26S proteasomes purified from spinach leaves. On the left is an electron micrograph of the 26S particles negatively stained with uranyl acetate, while on the right is a close-up image with overlaid contour plots generated by correlation averaging of 450 individual images. In all cases, scale bars represent 25 nm for the electron micrograph images and 5 nm for close-up images generated by averaging. The images were modified from references (Baumeister *et al.*, 1988; Fujinami *et al.*, 1994; Kremp *et al.*, 1986; Schliephacke *et al.*, 1991; Yoshimura *et al.*, 1993).

conserved and widely distributed among eukaryotes (Schliephacke *et al.*, 1991).

The complete 26S proteasome (i.e. the CP capped at one or both ends by the RP) was subsequently discovered by the purification of ubiquitin conjugate-degrading activity from rabbit reticulocytes (Hough *et al.*, 1986). While it had been well established that major catabolic processes in animal cells involved the ATP-dependent proteolysis of selective substrates (Etlinger and Goldberg, 1977), the enzyme(s) responsible for this activity had yet to be identified. Taking advantage of the new ability to synthesize ubiquitylated substrates such as ¹²⁵I-labelled ubiquitin-lysozyme conjugates (Hough and Rechsteiner, 1986), a protocol was developed to purify the responsible ATP-dependent protease. Through a series of anion exchange and size exclusion chromatography steps followed by glycerol gradient sedimentation, all of which were performed in ATP-containing buffers, the responsible activity was isolated (Hough *et al.*, 1986, 1987). The active enzyme turned out to be the 20S proteasome (i.e. the CP) along with a number of additional polypeptides, which together formed a 26S particle, thus providing the first direct link between ubiquitylation and a protease (Ganoth *et al.*, 1988; Hough *et al.*, 1987; Waxman *et al.*, 1987). SDS-PAGE analysis of these preparations identified a host of new polypeptides in the 35-100 kDa range in addition to the known CP subunits, which were later shown to comprise a second stable complex, the RP. Shortly thereafter, the RP

was demonstrated to have ATPase activities attributable to the RPT subunits, which help in substrate unfolding and maintaining CP-RP association (Armon *et al.*, 1990). Electron microscopic images of the full 26S particle then revealed its diagnostic quaternary structure in which the CP is capped by one or two RPs that sit over the axial pores for substrate entry (Figure 1.5 C; (Peters *et al.*, 1991; Yoshimura *et al.*, 1993)).

The existence of a similar 26S proteasome in plants was initially implied by the detection of an ATP-dependent activity in oat and wheat germ extracts capable of degrading ubiquitylated proteins (Hatfield and Vierstra, 1989; Vierstra and Sullivan, 1988). This was followed some years later by the first isolation of a complete plant 26S proteasome holocomplex from spinach leaves (Fujinami *et al.*, 1994). As with the mammalian forms, purification was achieved by anion exchange and size exclusion chromatography, followed by glycerol gradient centrifugation, all in the presence of ATP to stabilize the CP-RP association. These spinach preparations were, like their rabbit reticulocyte counterparts, able to rapidly degrade ubiquitylated substrates in an ATP-dependent manner, and further analysis by native-PAGE, SDS-PAGE and electron microscopy revealed the complete subunit composition and caterpillar-like structure of the plant particle (Figure 1.5 D; (Fujinami *et al.*, 1994)). Similar purifications were successful using rice suspension culture cells and garlic cloves (Malik

et al., 2004; Yanagawa *et al.*, 1999), which were accompanied by the first demonstrations that proteasome inhibitors designed for their mammalian counterparts were effective with the plant particles, suggesting very similar enzymatic mechanisms (Ozaki *et al.*, 1992; Woffenden *et al.*, 1998).

Despite its prevalence as a genetic model, purification of the 26S proteasome from the flowering plant *Arabidopsis thaliana* was not reported until several years after other plant species (Yang *et al.*, 2004). First protocols involved differential PEG precipitation followed by anion exchange and size exclusion chromatography, with the latter exploiting the large size of the holoprotease. More recently, an improved one-step affinity method was developed (Book *et al.*, 2010), based on the strategies that had been successfully employed in yeast (Leggett *et al.*, 2005). Here, epitope-tagged proteasomes were generated by genetically replacing the subunit PAG1 with a variant bearing a C-terminal FLAG tag; this tagged particle could then be purified with appropriate affinity matrices, and released in non-denaturing conditions with FLAG peptide. This approach enables rapid and robust purification of the whole 26S proteasome complex when performed in the presence of ATP, or enables purification of the CP sub-particle when performed in the absence of ATP (Book *et al.*, 2010). This affinity method has considerable advantages compared to previous, conventional chromatographic approaches (Yang *et al.*, 2004) as it is both faster,

more reliable and produces higher yields per gram of tissue (~6 µg/g). This milder more rapid technique also prevents breakdown of some subunits, in particular RPN10, which is sensitive to post-homogenization proteolysis (Yang *et al.*, 2004). One caveat is that the epitope tag, given its exposed position and flexible structure, might be sensitive to proteolytic cleavage following tissue homogenization. For the PAG1-FLAG protocol, chymostatin was found to effectively block the interfering protease (Book *et al.*, 2010). An example of such preparations analyzed by SDS-PAGE followed by immunoblotting with antibodies against several proteasome subunits, is shown in Figure 1.6.

1.4. 26S Proteasome Substrate Processing

A variety of proteins help the 26S proteasome process ubiquitylated substrates. Some include key constituents of the complex itself, such as RPN11, which is a metalloprotease that uses a zinc-coordinated active site to release the ubiquitin moieties isopeptide-linked to substrates (Verma *et al.*, 2002; Worden *et al.*, 2014). Through RPN11 and other loosely associated deubiquitylating enzymes such as UBP6/USP14 (Hanna *et al.*, 2006; Sakata *et al.*, 2011), bound ubiquitins are actively recycled. Substrate selection by the 26S proteasome is dictated by several ubiquitin receptors intrinsic to the RP lid, including RPN1, RPN10, RPN13, and




Figure 1.6: Affinity purification of 26S proteasomes from *PAG1::PAG1-FLAG* *pag1-1* plants. (Book *et al.*, 2010) (A) SDS-PAGE analysis of proteins obtained at each affinity purification step. Total protein extracts from 10 day old wild-type (WT) and *PAG1::PAG1-FLAG* *pag1-1* plants were incubated with anti-FLAG affinity resin, washed, and competitively eluted with the FLAG peptide. The procedure was performed in the presence or absence of ATP, and the input, unbound, washed and eluted fractions were subjected to SDS-PAGE and the gel stained for protein with silver. The black arrowhead indicates the *PAG1-FLAG* protein, while the open arrowhead identifies nitrilase, which is non-specifically enriched during the purification. (B) Immunoblot detection of various 26S proteasome subunits in the affinity-purified preparations shown in A. Subunits tested include the CP subunits PAG1 and PBA1, the RP subunits RPT2, RPN1, RPN5, RPN10 and RPN12a, and the alternate capping particle PA200. Other proteins tested include the Rubisco small subunit and nitrilase. This figure was modified from reference (Book *et al.*, 2010).

DSS1/SEM1 (Elsasser *et al.*, 2002; Fatimababy *et al.*, 2010; Finley, 2009; Lin *et al.*, 2011; Paraskevopoulos *et al.*, 2014; Sakata *et al.*, 2012; Shi *et al.*, 2016; Van Nocker *et al.*, 1996). RPN10 binds ubiquitin via defined ubiquitin-interacting motifs (UIMs), of which yeast, human and *Arabidopsis* RPN10 contain 1, 2 and 3 in tandem, respectively (Fatimababy *et al.*, 2010; Finley, 2009; Fu *et al.*, 1998; Lin *et al.*, 2011; Van Nocker *et al.*, 1996). By contrast, RPN13 binds ubiquitin via a pleckstrin-like receptor for ubiquitin (PRU) domain, which is structurally distinct from UIMs but binds to the same hydrophobic patch on ubiquitin (Husnjak *et al.*, 2008; Schreiner *et al.*, 2008). More recently, DSS1/SEM1 was also found to be a proteasomal ubiquitin receptor (Paraskevopoulos *et al.*, 2014). It had previously resisted identification due to both its small size, which prevented visualization by standard protein stains following sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE), and its paucity of lysine and arginine residues, which complicated detection by conventional mass spectrometric methods. Only with the use of top-down mass spectrometry of 26S proteasome complexes was DSS1/SEM1 first detected in intact 26S proteasomes from *Arabidopsis* (Russell *et al.*, 2013). In addition to these core ubiquitin receptors, there are several extra-proteasomal ubiquitin-binding proteins that shuttle ubiquitylated cargo to the RP. They work by virtue of ubiquitin-associated (UBA) domains that bind ubiquitin, combined with a ubiquitin-like

(UBL) domain that interacts with the intrinsic ubiquitin receptors such as RPN10. Important shuttle factors in plants include RAD23, DSK2, and DDI1 (Farmer *et al.*, 2010; Fatimababy *et al.*, 2010; Finley, 2009; Lin *et al.*, 2011), though many other ubiquitin-binding proteins are known in other species (Husnjak and Dikic, 2012). Numerous other factors also associate sub-stoichiometrically with the mature CP and RP sub-complexes, including deubiquitylating enzymes, several E3 ligases and protein kinases, and a collection of protein folding chaperones (Besche *et al.*, 2014; Book *et al.*, 2010; Leggett *et al.*, 2002; Xie and Varshavsky, 2000).

1.5. Proteasome Assembly

Not surprisingly given its intricate architecture, construction of the eukaryotic 26S proteasome requires a large collection of assembly factors that work in synchrony. The most well-studied eukaryotic models for assembly include both yeast and mammalian systems; however, very little if anything is known about the assembly of the plant complex.

1.5.1. 20S Core Protease Assembly

In yeast and mammals five of the seven β -subunits ($\beta 1$, $\beta 2$, $\beta 5$, $\beta 6$, and $\beta 7$) are synthesized as an immature propeptide form with an N-terminal extension. These

β -subunits undergo autolytic cleavage exposing a catalytically active N-terminal threonine for the β 1, β 2, β 5 subunits upon CP maturation. (Gu and Enenkel, 2014; Lee *et al.*, 1990). This propeptide prevents inactivation of the mature form's N-terminal threonine by N-terminal acetylation, and also prevents premature activation of peptidase activities (Arendt, 1999; Arendt and Hochstrasser, 1997). While this propeptide processing occurs in *Arabidopsis*, and likely other plants, its role in plant proteasome assembly remains to be investigated (Book *et al.*, 2010). In yeast, complexes containing these immature propeptide forms were first identified in 13-16S fractions (smaller than the 20S fraction) suggesting that distinct assembly steps may occur (Frentzel *et al.*, 1994). Indeed in yeast these propeptides were found to be important in the assembly of the 20S CP as deletion of the β 5 propeptide was shown to be lethal by preventing assembly of the CP, and deletion of the β 2 propeptide accumulated aberrant complexes (Jager *et al.*, 1999). Several of these propeptides were found to interact with the first dedicated assembly chaperone discovered, Ump1, which was identified in a screen for null mutants defective in ubiquitin-mediated proteolysis in yeast (Li *et al.*, 2007; Ramos *et al.*, 1998). Ump1 along with other dedicated assembly chaperones aide in the assembly of 15S complexes, or proteasome half barrels containing α -subunits 1-7, and β -subunits 1-6, noticeably lacking the β 7 subunit (Li *et al.*, 2007; Marques *et al.*, 2007). Similar investigations

of the mammalian β 5 propeptide enabled identification of 15S half-barrels that also contained Ump1 and a similar complement of α - and β -subunits also lacking the β 7 subunit (Witt *et al.*, 2000). Later studies would go on to show that in both mammals and yeast, β 7 is the last subunit to enter the β -ring, resulting in Ump1 being degraded as the first substrate of nascent CP (Hirano *et al.*, 2008; Ramos *et al.*, 1998). Intriguingly, the alternative capping particle, PA200 co-purified with these 15S half-barrels suggesting that PA200 may be involved in proteasome assembly (see CP assembly summary in Figure 3.7) (Li *et al.*, 2007; Marques *et al.*, 2007).

An additional suite of assembly chaperones aide in construction of the CP including the proteasome biogenesis associated Pba1-4 proteins in yeast, and the proteasome assembly chaperones PAC1-4 in mammals. The yeast assembly chaperones Pba1 and Pba2 were initially discovered in affinity preparations of the CP in a genetic background that induced the formation of aberrant 15S half-barrels (Li *et al.*, 2007). MS/MS analyses of these 15S complexes separated by native-PAGE identified Pba1, Pba2, and Ump1 contained in these complexes suggesting that Pba1 and Pba2 played a role in CP assembly (Li *et al.*, 2007). In a separate study, Pba1 and Pba2, along with two additional assembly chaperones Pba3 and Pba4 were identified in a mutant screen for suppressors of a conditional dominant lethal allele of Rad53 which induces DNA damage (Le Tallec *et al.*, 2007). Pba1, and Pba2 were




Figure 1.7: Schematic representation of CP Assembly. Adapted from (Hirano *et al.*, 2008). Pba1-4 (yeast) and PAC1-4 (human) nomenclature is used. Free α -subunits are bound to prevent aberrant α -ring assembly by Pba1-4. The β 2 subunit along with Ump1 bind this α -ring, and subsequently the Pba(PAC)3/4 heterodimer leaves the assembly complex. Then the β 3 and β 4 subunits are incorporated and the β 5 and β 6 subunits are added forming a 15S half-barrel, possibly with the help of PA200. Two half-barrels come together and the last β -subunit, β 7 is added. N-terminal propeptides for β -subunits are autocatalytically cleaved, Ump1 is degraded as the first substrate of the nascent proteasome, and a fully functional CP is generated.

found to co-fractionate with both the precursor propeptide form of the $\beta 2$ subunit, and the mature form; however, Pba3 and Pba4 were found to exclusively fractionate with the propeptide form of $\beta 2$ (Le Tallec *et al.*, 2007). These data suggested that Pba3 and Pba4 aided in earlier stages of assembly, while Pba1 and Pba2 seemed to act at both early and later stages. Pba1 and Pba2 were found to form a heterodimer, while Pba3 and Pba4 were found to form a separate heterodimer (Le Tallec *et al.*, 2007).

An orthologous system to the yeast Pba1-4 system was elucidated in mammals. PAC1 and PAC2 were first identified in affinity preparations of an immature propeptide form of the $\beta 1$ subunit, and like their yeast counterparts (Pba1 and 2) were found to form a stable heterodimer (Hirano *et al.*, 2005). Consistent with their role in assembly they were found to co-fractionate with immature β -subunits, and Ump1 (Hirano *et al.*, 2005). Overexpression of these PAC1 and PAC2 chaperones yielded increases in half-barrel precursors, while knockdown by siRNA impaired assembly of the CP complex (Hirano *et al.*, 2005). An additional chaperone involved in assembling the α -ring was identified in PAC1 immunopurifications in human cell lines, and was termed PAC3 (Hirano *et al.*, 2006). siRNA knockdown of PAC3 attenuated α -ring formation and gave rise to the accumulation of free α -subunits, suggesting that PAC3 played a role in stabilization of the CP α -ring (Hirano *et al.*,

2006). Mammalian PAC4 was identified as an ortholog to yeast Pba4 through PSI-BLAST analyses (Kusmierczyk *et al.*, 2008) and together with PAC4 forms a heterodimer, like its yeast Pba3/4 counterpart (Le Tallec *et al.*, 2007).

The role of the PAC1/2 and PAC3/4 heterodimers in assembly, along with the exact ordered incorporation of β -subunits was aided by siRNA knockdown of specific β -subunits, and studying the resulting assembly products (Hirano *et al.*, 2008). Through this analysis Hirano *et al.* clearly delineated PAC3/4's role in early assembly, and found that PAC3 and PAC4 must leave the assembling half-barrel before the addition of the β_3 subunit. While this was demonstrated in mammals, observations of the yeast Pba3/4 heterodimer suggest a similar model in which the Pba3/4 heterodimer also remains bound to the α -ring until the β_2 subunit binds (Kusmierczyk *et al.*, 2008; Yashiroda *et al.*, 2008). While Pba(PAC)3/4 act early in assembly, Pba(PAC)1/2 have the propensity to bind the mature proteasome, and stay bound even after the incorporation of β -subunits.

A model of yeast and mammalian assembly adapted from Hirano *et al.* is as follows (Figure 3.7). After the α -ring is formed, with the help of Pba(PAC)1/2 and Pba(PAC)3/4, the β_2 subunit is incorporated along with its C-terminal tail that helps to recruit the β_3 subunit (Hirano *et al.*, 2008). Pba(PAC)3/4 is released upon β_3 subunit binding (Hirano *et al.*, 2008). Ump1 stabilizes the incorporation of additional

β -subunits by binding their inherently disordered propeptide regions (Li *et al.*, 2007). $\beta 4$, $\beta 5$, $\beta 6$, and $\beta 1$ are then incorporated (Hirano *et al.*, 2008). At this stage PA200 may help stabilize this precursor; however, its exact role in assembly is poorly understood. Two 15S half proteasomes come together, and finally the $\beta 7$ subunit is incorporated (Hirano *et al.*, 2008). The β -subunit propeptides are then cleaved exposing the catalytically active N-terminal threonines and Ump1 is degraded becoming the first substrate of the nascent CP (Li *et al.*, 2007). Pba(PAC)1/2 are then released, or can bind mature proteasomes until activation by the RP (Hirano *et al.*, 2008). Taken together these data are consistent with a model in which the heterodimeric pair Pba(PAC)3/4 acts at early stages to assemble free α -subunits into an α -ring while preventing the formation of aberrant α -subunit complexes (Hirano *et al.*, 2008). In contrast, the heterodimeric pair Pba(PAC)1/2 acts at both early and later stages in proteasome assembly and has the ability to bind mature CP (Wani *et al.*, 2016).

While the CP assembly process is conserved in yeast and mammals, the orthologs involved share very little sequence homology. The Ump1 orthologs share ~20% identity, and Pba1-4 and PAC1-4 share ~10% identity respectively (Murata *et al.*, 2009). Despite this low sequence identity, structural analysis revealed that the Pba3/4 heterodimer shared significant structural features with the human PAC3

protein, suggesting that these assembly chaperones may have retained structural similarities (Yashiroda *et al.*, 2008). The Hochstrasser group also speculated that the Pba1/2 and PAC1/2 complexes may be structurally related (Kusmierczyk *et al.*, 2011). Besides these speculative structural implications, Pba1 and Pba2 share some protein sequence features, including HbYX motifs important for their interaction with the α -ring (Kusmierczyk *et al.*, 2011). These features are conserved in their mammalian orthologs with PAC1 containing a HbYX motif, while PAC2 shares a HbYX-like HbY/F motif that is conserved in other eukaryotes outside the fungal kingdom (Kusmierczyk *et al.*, 2011).

Structural analysis of these CP assembly chaperones have also provided insight into their function. Pba1, and Pba2 were able to bind mature 20S proteasomes, and a co-crystal structure has been solved by X-ray crystallography (Stadtmauer *et al.*, 2012). The HbYX motif found in Pba1 was shown to intercalate between the α 5 and α 6 subunits, while the HbYX motif found in Pba2 was shown to intercalate between the α 6 and α 7 subunits (Figure 1.8). Crystal structures of the Pba3, Pba4 complex were shown to be similar to the PAC3 suggesting similar role (Yashiroda *et al.*, 2008). A co-crystal structure binding the α 5 subunit helped define the binding surface of the Pba3/4 heterodimer (Yashiroda *et al.*, 2008). This binding surface for Pba3/4 occurs on the opposite side of the α -ring as compared to the Pba1/2

binding surface providing further structural evidence for the distinct and separate roles that the Pba1/2 and Pba3/4 play in proteasome assembly (see Figure 1.9) (Stadtmueller *et al.*, 2012; Yashiroda *et al.*, 2008).

Outside of eukaryotes, proteasome assembly has also been studied in *archeal* and *eubacterial* proteasome models (Kusmierczyk *et al.*, 2011). While the assembly of these proteasomes can occur without assembly chaperones two chaperones, PbaA and PbaB were identified that likely aide in efficient assembly of the particle (Kusmierczyk *et al.*, 2011). PbaA and PbaB are evolutionarily related to yeast Pba1 and Pba2 and both harbor conserved HbYX motifs showing that these chaperones provide an ancient link to proteasome assembly evolution (Kusmierczyk *et al.*, 2011).






Figure 1.8: Pba1 and Pba2 binding sites via their HbYX motifs into adjacent α -subunit pairs. α -ring is shown with adjacent Pba1 and Pba2 binding surfaces that intercalate between α -subunits. Pba1 intercalates between α_5 and α_6 , while Pba2 intercalates between α_6 and α_7 (Yashiroda *et al.*, 2008).

[Figure 1.9 caption follows on next page]

A




B




α -ring + Pba1/2

C



α -ring + Pba3/4

D



Pba3/4 occlude β 3 subunit binding

Figure 1.9 (preceding page): Structural Diagram of opposing Pba1/2 Pba3/4 binding. Structures were obtained from the Protein Data Bank with PDB ID 4G4S for Pba1 and Pba2 bound to yeast 20S proteasome (Stadtmueller *et al.*, 2012), and PDB ID 2Z5C for Pba3 and Pba4 bound to the α 5 subunit(Yashiroda *et al.*, 2008) and visualized using PyMOL (Schrodinger, 2015). The α 5 subunit was used to align the structures so that the opposing Pba1/2 and Pba3/4 interfaces could be visualized in reference to the 20S α -ring. **(A)** 20S α -ring with alone (left), Pba1/2 heterodimer, and Pba3/4 heterodimer alone (middle), and 20S α -ring with bound Pba1-4 shows the opposing binding surfaces, with Pba1/2 binding to the top of the α -ring, and Pba3/4 binding to the bottom of the α -ring where β -subunits normally bind in the full 20S proteasome. **(B)** α -ring with Pba1/2 binding. **(C)** α -ring with Pba3/4 binding. **(D)** Binding of Pba3/4 prevents premature binding of the β -ring (shown as a grey ribbon structure), specifically to occlude the β 3 subunit (Yashiroda *et al.*, 2008).

1.5.2. 19S Regulatory Particle Assembly RP base

Like the chaperones that facilitate CP assembly, there are proteins that aide in proper formation of the RP. The majority of identified RP assembly chaperones are involved in the assembly of the RPT base ring. The RP base assembly chaperone Hsm3 was identified in the same screen for suppressors of Rad53 that identified the yeast CP assembly chaperones Pba1-4 (Le Tallec *et al.*, 2007), although HSM3 was not characterized until two years later (Le Tallec *et al.*, 2009). Several months later additional RP assembly chaperones (Nas2, Nas6 and Rpn14) were identified and characterized as components of an RP base assembly system (Funakoshi *et al.*, 2009). The majority of these RP base chaperones function by binding the C-terminal domains of RPT subunits preventing them from forming pre-mature contacts with the CP (Funakoshi *et al.*, 2009). However, their evolutionary history is likely distinct as they are made up of distinct domains including ARM/HEAT repeats in Hsm3, coiled-coil PDZ domains in Nas2, ankyrin repeats in Nas6, and WD40 repeats in Rpn14 (Funakoshi *et al.*, 2009).

Immunoprecipitation of Hsm3 showed that it interacted with several RP base subunits; however no α -subunits or RP lid subunits were detected suggesting that Hsm3 may be specific to the RP base (Le Tallec *et al.*, 2009). Fractionation of crude protein lysates from yeast showed that Hsm3 associated with free RP

but not with the mature 26S proteasome, suggesting that it may be involved in RP assembly (Le Tallec *et al.*, 2009). Assembly intermediates containing Hsm3, Rpn1, and Rpt2 were identified when RP base assembly was impaired (Funakoshi *et al.*, 2009). A mammalian ortholog of Hsm3, HSM3 was found to form a similar complex in mammals (Kaneko *et al.*, 2009). Recently, a co-crystal structure involving Hsm3, Rpt1, and Rpt2 showed that Hsm3 forms contacts with the C-terminal domains of both Rpt1 and Rpt2, providing further support for the formation of this pre-assembly complex and that these assembly chaperones function to prevent premature binding of the C-terminus of RPT subunits to the CP (Barrault *et al.*, 2012).

As described above, the C-terminal domains of RPT subunits contain HbYX motifs, which are important for binding the α -ring. Mutating the HbYX motif of either Rpt4 or Rpt6 by removing a single amino acid was sufficient to induce the formation of two separate pre-assembly complexes with one containing Hsm3 Rpn1, and Rpt1, 2, and 5 and another containing Rpn14, Rpt4, and Rpt6 (Park *et al.*, 2009). The formation of these pre-assembly complexes was blocked in triple null mutants for Hsm3, Bas6 and Rpn14 (Park *et al.*, 2009). Interestingly single mutants in either Nas6, or Rpn14 did not exhibit strong assembly defects; however, yeast deficient in both Nas6, and Rpn14 exhibited a strong assembly defect suggesting

that Nas6, and Rpn14 play semi-redundant roles (Saeki *et al.*, 2009). Despite this redundancy MS/MS analyses of tagged assembly chaperones and yeast-two-hybrid analyses showed Nas6 binding predominantly to Rpt3, while Rpn14 was found to predominantly bind to Rpt6 (Saeki *et al.*, 2009). The yeast Nas2 assembly chaperone was found to bind to Rpt5, and its mammalian ortholog NAS2 was shown to form a complex with both RPT4 and RPT5. Recent structural analyses of Nas2 suggest that it binds the C-terminus of Rpt5 likely acting to prevent pre-mature binding with the CP (Satoh *et al.*, 2014).

A general model emerges in which a majority of the RP base assembly chaperones act to prevent pre-mature association of unassembled RP base by actively binding the C-termini of RPT subunits (Tomko and Hochstrasser, 2013). This model involves four separate modules including [Hsm3 with Rpn1, Rpt1, and Rpt2], [Nas2 with Rpt4, and Rpt5], [Nas6 with Rpt3], and [Rpn14 with Rpt6]. While the exact ordered steps of RP base assembly are still unclear, the process appears to be conserved between both yeast and mammals. A model of RP base assembly using the yeast nomenclature is presented in Figure 1.10 showing how the individual modules may come together to form the RP base.




Figure 1.10: Schematic representation of RP Assembly. While the exact order of RP-base assembly is unclear, it involves several modules including [Hsm3 with Rpn1, Rpt1, and Rpt2], [Nas2 with Rpt4, and Rpt5], [Nas6 with Rpt3], and [Rpt14 with Rpt6].

1.5.3. 19S Regulatory Particle Assembly RP lid

Several modules and chaperones are involved in the assembly of the RP base; however, construction of the RP lid is not well understood. RP lid assembly has primarily been investigated in yeast by mutant analyses of specific RP lid subunits including mutants in Rpn7, Rpn9 and Rpn12 (Fukunaga *et al.*, 2010). MS/MS analyses of affinity purified lid complexes in these differing mutant backgrounds identified two different subassemblies, the first containing Rpn5, 6, 8, 9, and 11, and the second containing Rpn3, Rpn7, and Sem1 (Fukunaga *et al.*, 2010). Intriguingly, Sem1 was found to act as a bridge between Rpn3, and Rpn7, as cleavage of an engineered version of Sem1 containing a Tobacco Etch Virus (TEV)-cleavage site showed defects in RP lid assembly (Tomko and Hochstrasser, 2014). While Sem1 may play a role in proteasome biogenesis, alternative roles for Sem1 have been proposed. In the mature particle, Sem1 has been found to aide in recognition of ubiquitin chains which are linked at lysine 48 or 63 (Paraskevopoulos *et al.*, 2014). Proteasomes affinity purified from Sem1 mutants had only mild defects in stability, suggesting that its main function may be that of ubiquitin recognition (Paraskevopoulos *et al.*, 2014).

Structural studies of the RP lid, have identified conserved C-terminal helices that form a helical bundle important in assembly of this sub-complex (Estrin *et al.*,

2013). The assembly of the lid was investigated by expressing the full complement of yeast RP lid subunits in which the C-terminal helices were removed in varying combinations (Estrin *et al.*, 2013). In the most extreme case, when the RPN6 helical bundle was omitted, a similar arrangement as described above was identified with four distinct sub-complexes identified that included [Rpn5, 8, 9, and 11], [Rpn3, and Rpn7], [Rpn6 alone], and [Rpn12 alone] (Estrin *et al.*, 2013). Estrin *et al.* speculate on the exact ordering of the assembly, but this has not been thoroughly investigated.

While the exact ordering involving the other RP lid subunits is unclear, RPN12 is likely the last subunit to be added to the RP lid. In Rpn9 mutant cells, Rpn12 was completely absent from the RP lid (Fukunaga *et al.*, 2010). However, mutants in Rpn12 exhibited the full complement of RP lid subunits suggesting that Rpn12 was likely the last subunit added to the RP lid (Tomko and Hochstrasser, 2011). The conserved C-terminus of RPN12 was found to contact both the RP base and lid suggesting that RPN12 might be a final checkpoint for RP lid and base assembly (Tomko and Hochstrasser, 2011). When the RP lid helical bundle described above is impaired, RPN12 cannot bind, suggesting a mechanism by which RPN12 can scan this bundle for proper assembly (Estrin *et al.*, 2013).

1.5.4. RP-CP Assembly and Alternative Capping Particles

While there are dedicated chaperones for the CP and RP, an additional chaperone mediates assembly of the final particle (Ramos *et al.*, 1998). ECM29 stabilizes the association of assembled CP and RP and provides a final quality control checkpoint for mature 26S proteasomes (Besche *et al.*, 2014; Lehmann *et al.*, 2010). Finally, in some situations, the RP is replaced entirely by alternate capping particles such as PA200 (also known as Blm10) or CDC48 (Barthelme and Sauer, 2012; Book *et al.*, 2010; Schmidt *et al.*, 2005). The functions of these caps are not yet clear, but recent proposals for PA200 have it participating in 26S proteasome assembly, helping shuttle proteasomes into the nucleus, and/or generating a ubiquitin-independent proteasome containing CP and PA200 only (Dange *et al.*, 2011; Sadre-Bazzaz *et al.*, 2010; Weberruss *et al.*, 2013). However, in plants, mutants in PA200 exhibit no gross morphological defects showing that PA200 is not essential to plant growth and development (Book *et al.*, 2010).

1.5.5. Proteasome Assembly in Plants

While much is known about proteasome assemble in both yeast and mammals, very little is known about how plants assemble this particle. There is some evidence that plants likely use a similar process as an ortholog of PAC2, PBAC2, has been

identified in affinity preparations for the plant CP complex (Book *et al.*, 2010). A slightly different nomenclature is used in plants as compared to Pba in yeast and PAC in mammals as both of these can refer to proteasome α - and β -subunits (example, $\alpha 1$ is PAC1, and $\beta 1$ is PBA1 in *Arabidopsis*). Furthermore, PSI-BLAST-based bioinformatic analyses, similar to the one used to find orthologs between yeast and humans was able to identify some putative plant orthologs (PBAC1 and PBAC2) to the mammalian PAC1 and PAC2 assembly chaperones by two separate research groups, although their annotations in The *Arabidopsis* Information Resource (TAIR) are incomplete (Kusmierczyk *et al.*, 2011; Le Tallec *et al.*, 2007). These identified orthologs share very little sequence identity (<10%), although this is unsurprising given the poor sequence identity between the yeast and mammalian orthologs (Kusmierczyk *et al.*, 2011; Le Tallec *et al.*, 2007; Murata *et al.*, 2009). Genomic searches based on BLAST alone, have failed to identify orthologs of the CP assembly chaperones; however putative orthologues for NAS2, NAS6, and HSM3 are readily identifiable in the genome (Chapter 3). Together these data suggested that our analysis of plant proteasome assembly has remained incomplete. One goal of this thesis, explored in Chapter 3, was to identify additional CP and RP interacting proteins, potentially including some of these assembly chaperones.

1.6. Proteasome Isotypes

In yeast, proteasome subunits are typically encoded as a single subunit, however in animals specific subunits are replaced to form different proteasome isotypes. This replacement typically occurs due to tissue specific expression of alternative CP subunits. Several examples of isotypes exist, including the Immunoproteasome, the thymoproteasome proteasome, and the testes-specific proteasome.

1.6.1. Immunoproteasome

While the proteasome was first discovered in the 1970's it wasn't until 20 years later that an alternative form of the proteasome, the immunoproteasome was discovered (Brown *et al.*, 1991; Glynne *et al.*, 1991; Ortiz-Navarrete *et al.*, 1991). The immunoproteasome was discovered while trying to determine how small antigen presenting peptides are produced for the major histocompatibility complex (MHC). The MHC is a set of cell surface proteins involved in the acquired immune response (reviewed in (Neefjes *et al.*, 2011)). Small peptides resulting from normally present cytosolic proteins (MHC class I antigens) are loaded into the MHC and presented on a cell's surface helping establish a normal profile of surface peptides recognized by the immune system. Atypical protein production, whether through viral or bacterial infection, or by cells that are proliferating abnormally, such as cancerous

cells, result in atypical surface peptides causing the immune system to respond and recruit CD8+ T-killer cells to remove the aberrant cell. Several of the identified genes involved in this process were termed low molecular weight proteins, or LMPs and their protein sequences were found to be related to the proteasome. The cytokine Interferon-gamma, a small signaling protein involved in the immune response was found to induce expression of three alternative β -subunits of the proteasome (β 1i, β 2i, and β 5i), which are then incorporated to form the immunoproteasome (Figure 1.11) (Aki *et al.*, 1994; Groettrup *et al.*, 1996).

An elegant study that compared the enzymatic activity of affinity purified 20S proteasomes with immunoproteasomes using model peptide substrates determined that cleavage after acidic residues via the β 1's PGPH activity is nearly eradicated by the replacement of the β 1 subunit with the immunoproteasome β 1i's subunit which favors chymotryptic activities (Dahlmann *et al.*, 2000). Together these altered enzymatic activities allow for the production of peptides more favorable for antigen presentation to the MHC.

1.6.2. Thymoproteasome proteasome

A sub-type of the immunoproteasome was discovered in the thymus. A search for proteasome related genes identified an uncharacterized ortholog of the β -subunit

that was preferentially expressed in thymus tissue, termed β 5t (Murata *et al.*, 2007). This thymoproteasome was similar in composition to the immunoproteasome in that it contained the altered β 1i, and β 2i subunits; however, the β 5i subunit was replaced with this novel β 5t subunit (shown in Figure 1.11) (Murata *et al.*, 2007). Expression of the β 5t was even further sub-localized to cells in the thymic cortex, and mice lacking this subunit could not generate CD8+ T cells, normally involved in aiding in the destruction of foreign invaders (Murata *et al.*, 2007).

1.6.3. Testes-specific proteasome

Like the discovery of the thymoproteasome proteasome, the discovery of the testes-specific isotype was initially driven by a bioinformatics search for proteasome related genes (Uechi *et al.*, 2014). A search revealed an uncharacterized ortholog of the α 4 subunit that shared 85% amino acid identity, termed here α 4s (Uechi *et al.*, 2014). Analysis of α 4s's expression by qRT-PCR analysis showed that it was expressed in all tissues; however only α 4s protein accumulated in testes (Uechi *et al.*, 2014). α 4s is preferentially incorporated over the α 4 subunit in testes, however α 4s specific proteasomes did not show altered enzymatic activities, and its biological function remains unknown (Uechi *et al.*, 2014).

1.6.4. α 4- α 4 proteasome Isotype

When defects are present in the Pba(PAC)3/4 heterodimer the α 3 subunit is replaced by the α 4 subunit, making an α 4- α 4 specific proteasome isotype (Kusmierczyk *et al.*, 2008; Padmanabhan *et al.*, 2016). Intriguingly, this α 4- α 4 isotype occurs in both mammals and yeast, conferring increased resistance to heavy metal ions, suggesting an evolutionarily conserved role for these proteasome isotypes (Kusmierczyk *et al.*, 2008; Padmanabhan *et al.*, 2016). Additionally, α 4 levels are modulated by several oncogenes including BRCA1 (Li *et al.*, 2015), and the formation of these alternative subtypes is highly dependent on the relative availability of α 3 vs α 4. Together these data suggest that α 4- α 4 proteasome isotypes may play a role in the progression of cancer (Padmanabhan *et al.*, 2016).




Figure 1.11: Representation of subunits replaced for both the immunoproteasome and the thyoproteasome. Adapted from (Groettrup *et al.*, 2009). The three catalytically active sites of the proteasome $\beta 1$, $\beta 2$, and $\beta 5$ are replaced with the immunoproteasome variants $\beta 1i$, $\beta 2i$, and $\beta 5i$. Furthermore, in the thyoproteasome the $\beta 5i$ subunit is replaced with another variant the $\beta 5t$ subunit.

1.6.5. Plant Proteasome Isotypes

Outside of the proteasome isotypes discussed above, animal and yeast genomes typically encode a single gene for most proteasome subunits. In rice the RPT family has been duplicated such that four of the six RPT subunits are encoded as paralogs (Shibahara *et al.*, 2004). While the function of these duplicated subunits is unclear, they do exhibit alternative tissue specific expression patterns as determined by MS/MS analyses using total protein extracts. Specifically, the expression patterns differed between bran and callus suggesting that there is tissue specific regulation of these sets of RPT subunits. In the most extreme case, most subunits in *Arabidopsis* are encoded as paralogs, likely due to a genome duplication (Book *et al.*, 2010; Fu *et al.*, 1998). MS/MS analyses of proteasomes purified from *Arabidopsis* showed that most of these subunit paralogs are incorporated into the proteasome. Therefore, *Arabidopsis* has the propensity to assemble its proteasomes into a diverse array of proteasome isotypes by incorporating specific subsets of isoforms. Some genetic evidence exists that suggests that these paralogs may have differing functions. RPN1a and RPN1b play differing roles in embryogenesis (Brukhin *et al.*, 2005), and overexpression of RPN5a induces an early senescent phenotype while overexpression of RPN5b does not (Book *et al.*, 2009). Despite these observations, it is unclear if plants assemble their proteasome subunit isoforms into compositionally and func-

tionally distinct isotypes. One of the objectives of this thesis was to determine if *Arabidopsis* can assemble proteasome isotypes by leveraging label-free quantitative mass spectrometric analysis of affinity preparations enriched for either the RPT4a or RPT4b subunit isoforms (explored in Chapter 3). If plants can assemble distinct isotypes, preparations for either isoform may show enrichment for other subunit isoforms; however, if plants assemble its subunit isoforms in a random fashion the preparations should be similar.

1.7. Conclusions

The *Arabidopsis* 26S proteasome exists *in planta* as a diverse array of complexes containing multiple subunit isoforms and interacting proteins (Book *et al.*, 2010; Fu *et al.*, 1999; Yang *et al.*, 2004). However, there are several gaps in our knowledge of the plant particle. While the yeast and mammalian CP assembly systems are well described, very little is known about proteasome assembly in plants. Initial MS/MS analyses of the plant complex identified a likely ortholog related to PAC2, named PBAC2, suggesting that the plant system may be similar; however, no additional assembly chaperones were described (Book *et al.*, 2010). Several putative *Arabidopsis* assembly orthologs were identified through PSI-BLAST analyses including orthologs for PAC1, and PAC2 by both the Hoschstrasser group, and others

(Kusmierczyk *et al.*, 2011; Le Tallec *et al.*, 2007); however, putative orthologs of PAC3, and PAC4 remained to be identified, until this thesis work. Taken together these data suggested that our analysis of the *Arabidopsis* CP interactome was incomplete and further analyses of CP associated proteins was necessary. Previous MS/MS analyses of the plant proteasome also failed to identify obvious orthologs of RP assembly chaperones. This suggested that our analyses of the RP interactome, like our previous analysis of the CP interactome, were also incomplete. One reason these previous analyses may have failed to identify RP assembly chaperones is because our affinity purifications targeted the CP specifically. To overcome this potential problem, I developed an affinity purification that targeted the RP through two separate isoforms of RPT4, RPT4a, and RPT4b. I exploited the fact that RPT4's N-terminus was likely solvent accessible (Figure 1.12) and used a similar strategy developed previously by Book *et al.* (Book *et al.*, 2010). Through quantitative MS/MS analyses I defined the plant RP interactome, as shown in Chapter 3. Additionally, I investigated the possibility of forming proteasome isotypes by the incorporation of specific subunit isoforms in combination with either RPT4a or RPT4b. In summary I identify CP- and RP-specific interactors, many of which may be putative orthologs of both the plant CP and RP assembly chaperones, and show that proteasomes enriched for either RPTa or RPT4b are largely similar suggesting that plants assemble

their proteasome subunit isoforms randomly.




Figure 1.12: Solvent accessible tag locations on the 26S proteasome for both PAG1 and RPT4. A structural model of the 26S proteasome from yeast at subatomic resolution modified from PDB ID 4CR2 (Beck *et al.*, 2012). The RP subunits, as well as the CP α and β -rings are shown. Highlighted in red, and indicated by black arrowheads, are the positions where FLAG affinity tags have been successfully used to enrich for *Arabidopsis* 26S proteasomes.

1.8. Literature Cited

- Aki, M., Shimbara, N., Takashina, M., Akiyama, K., Kagawa, S., Tamura, T., Tanahashi, N., Yoshimura, T., Tanaka, K., and Ichihara, A.** (1994). Interferon-gamma induces different subunit organizations and functional diversity of proteasomes. *J. Biochem.* **115**: 257–69.
- Angot, A., Peeters, N., Lechner, E., Vailleau, F., Baud, C., Gentzbittel, L., Sartorel, E., Genschik, P., Boucher, C., and Genin, S.** (2006). *Ralstonia solanacearum* requires F-box-like domain-containing type III effectors to promote disease on several host plants. *Proc. Natl. Acad. Sci. U.S.A.* **103**: 14620–5.
- Arendt, C.S.** (1999). Eukaryotic 20S proteasome catalytic subunit propeptides prevent active site inactivation by N-terminal acetylation and promote particle assembly. *The EMBO Journal* **18**: 3575–3585.
- Arendt, C.S. and Hochstrasser, M.** (1997). Identification of the yeast 20S proteasome catalytic centers and subunit interactions required for active site formation. *Proc. Natl. Acad. Sci. U.S.A.* **94**: 7156–7161.
- Armon, T., Ganoth, D., and Hershko, A.** (1990). Assembly of the 26S complex that degrades proteins ligated to ubiquitin is accompanied by the formation of ATPase activity. *J. Biol. Chem.* **265**: 20723–20726.
- Arrigo, A.P., Tanaka, K., Goldberg, A.L., and Welch, W.J.** (1988). Identity of the 19S prosome particle with the large multi-functional protease complex of mammalian cells (the proteasome). *Nature* **331**: 192–194.
- Barrault, M.B., Richet, N., Godard, C., Murciano, B., Le Tallec, B., Rousseau, E., Legrand, P., Charbonnier, J.B., Le Du, M.H., Guerois, R., Ochsenbein, F., and Peyroche, A.** (2012). Dual functions of the Hsm3 protein in chaperoning and scaffolding regulatory particle subunits during the proteasome assembly. *Proc. Natl. Acad. Sci. U.S.A.* **109**: E1001–10.
- Barthelme, D. and Sauer, R.T.** (2012). Identification of the Cdc48-20S proteasome as an ancient AAA proteolytic machine. *Science* **337**: 843–846.
- Baumeister, W., Dahlmann, B., Hegerl, R., Kopp, F., Kühn, L., and Pfeifer, G.** (1988). Electron microscopy and image analysis of the multi-catalytic proteinase. *FEBS Lett.* **241**: 239–245.

- Beck, F., Unverdorben, P., Bohn, S., Schweitzer, A., Pfeifer, G., Sakata, E., Nickell, S., Plitzko, J.M., Villa, E., Baumeister, W., and Förster, F.** (2012). Near-atomic resolution structural model of the yeast 26S proteasome. *Proc. Natl. Acad. Sci. U.S.A.* **109**: 14870–14875.
- Berndsen, C.E. and Wolberger, C.** (2014). New insights into ubiquitin E3 ligase mechanisms. *Nat. Struct. Mol. Biol.* **21**: 301–307.
- Besche, H.C., Sha, Z., Kukushkin, N.V., Peth, A., Hock, E.M., Kim, W., Gygi, S.P., Gutierrez, J.A., Liao, H., Dick, L.R., and Goldberg, A.L.** (2014). Auto-ubiquitination of the 26S proteasome on Rpn13 regulates breakdown of ubiquitin conjugates. *EMBO J.* **33**: 1159–1176.
- Bhattacharyya, S., Yu, H., Mim, C., and Matouschek, A.** (2014). Regulated protein turnover: snapshots of the proteasome in action. *Nat. Rev. Mol. Cell Biol.* **15**: 122–133.
- Book, A.J., Gladman, N.P., Lee, S.S., Scalf, M., Smith, L.M., and Vierstra, R.D.** (2010). Affinity purification of the *Arabidopsis* 26S proteasome reveals a diverse array of plant proteolytic complexes. *J. Biol. Chem.* **285**: 25554–25569.
- Book, A.J., Smalle, J., Lee, K.H., Yang, P., Walker, J.M., Casper, S., Holmes, J.H., Russo, L.A., Buzzinotti, Z.W., Jenik, P.D., and Vierstra, R.D.** (2009). The RPN5 subunit of the 26S proteasome is essential for gametogenesis, sporophyte development, and complex assembly in *Arabidopsis*. *Plant Cell* **21**: 460–78.
- Brown, M.G., Driscoll, J., and Monaco, J.J.** (1991). Structural and serological similarity of MHC-linked LMP and proteasome (multicatalytic proteinase) complexes. *Nature* **353**: 355–7.
- Brukhin, V., Gheyselinck, J., Gagliardini, V., Genschik, P., and Grossniklaus, U.** (2005). The RPN1 subunit of the 26S proteasome in *Arabidopsis* is essential for embryogenesis. *Plant Cell* **17**: 2723–37.
- Ciechanover, A., Elias, S., Heller, H., Ferber, S., and Hershko, A.** (1980a). Characterization of the heat-stable polypeptide of the ATP-dependent proteolytic system from reticulocytes. *J. Biol. Chem.* **255**: 7525–8.
- Ciechanover, A., Finley, D., and Varshavsky, A.** (1984). Ubiquitin dependence of selective protein degradation demonstrated in the mammalian cell cycle mutant ts85. *Cell* **37**: 57–66.

- Ciechanover, A., Heller, H., Elias, S., Haas, A.L., and Hershko, A.** (1980b). ATP-dependent conjugation of reticulocyte proteins with the polypeptide required for protein degradation. *Proc. Natl. Acad. Sci. U.S.A.* **77**: 1365–1368.
- Clough, R.C. and Vierstra, R.D.** (1997). Phytochrome degradation. *Plant, Cell and Environment* **20**: 713–721.
- Dahlmann, B., Ruppert, T., Kuehn, L., Merforth, S., and Kloetzel, P.M.** (2000). Different proteasome subtypes in a single tissue exhibit different enzymatic properties. *J. Mol. Biol.* **303**: 643–53.
- Dange, T., Smith, D.M., Noy, T., Rommel, P.C., Jurzitz, L., Cordero, R.J., Legendre, A., Finley, D., Goldberg, A.L., and Schmidt, M.** (2011). The Blm10 protein promotes proteasomal substrate turnover by an active gating mechanism. *J. Biol. Chem.* **286**: 42830–42839.
- Dharmasiri, N., Dharmasiri, S., and Estelle, M.** (2005). The F-box protein TIR1 is an auxin receptor. *Nature* **435**: 441–445.
- Disch, S., Anastasiou, E., Sharma, V.K., Laux, T., Fletcher, J.C., and Lenhard, M.** (2006). The E3 ubiquitin ligase BIG BROTHER controls arabidopsis organ size in a dosage-dependent manner. *Curr. Biol.* **16**: 272–9.
- Elsasser, S., Gali, R.R., Schwickart, M., Larsen, C.N., Leggett, D.S., Müller, B., Feng, M.T., Tübing, F., Dittmar, G.A., and Finley, D.** (2002). Proteasome subunit Rpn1 binds ubiquitin-like protein domains. *Nat. Cell Biol.* **4**: 725–730.
- Estrin, E., Lopez-Blanco, J.R., Chacon, P., and Martin, A.** (2013). Formation of an intricate helical bundle dictates the assembly of the 26S proteasome lid. *Structure* **21**: 1624–35.
- Etlinger, J.D. and Goldberg, A.L.** (1977). A soluble, ATP-dependent proteolytic system responsible for degradation of abnormal proteins in reticulocytes. *Proc. Natl. Acad. Sci. U.S.A.* **74**: 54–58.
- Farmer, L.M., Book, A.J., Lee, K.H., Lin, Y.L., Fu, H., and Vierstra, R.D.** (2010). The RAD23 family provides an essential connection between the 26S proteasome and ubiquitylated proteins in Arabidopsis. *Plant Cell* **22**: 124–142.
- Fatimababy, A.S., Lin, Y.L., Usharani, R., Radjacommare, R., Wang, H.T., Tsai, H.L., Lee, Y., and Fu, H.** (2010). Cross-species divergence of the major recognition pathways of ubiquitylated substrates for ubiquitin/26S proteasome-mediated proteolysis. *FEBS J.* **277**: 796–816.

- Finley, D.** (2009). Recognition and processing of ubiquitin-protein conjugates by the proteasome. *Annu. Rev. Biochem.* **78**: 477–513.
- Finley, D., Ciechanover, A., and Varshavsky, A.** (1984). Thermolability of ubiquitin-activating enzyme from the mammalian cell cycle mutant ts85. *Cell* **37**: 43–55.
- Finley, D., Ozkaynak, E., and Varshavsky, A.** (1987). The yeast polyubiquitin gene is essential for resistance to high temperatures, starvation, and other stresses. *Cell* **48**: 1035–1046.
- Frentzel, S., Pesold-Hurt, B., Seelig, A., and Kloetzel, P.M.** (1994). 20 S proteasomes are assembled via distinct precursor complexes. Processing of LMP2 and LMP7 proproteins takes place in 13-16 S preproteasome complexes. *J. Mol. Biol.* **236**: 975–81.
- Fu, H., Girod, P.A., Doelling, J.H., Van Nocker, S., Hochstrasser, M., Finley, D., and Vierstra, R.D.** (1999). Structural and functional analyses of the 26S proteasome subunits from plants. *Mol. Biol. Rep.* **26**: 137–146.
- Fu, H., Sadis, S., Rubin, D.M., Glickman, M.H., Van Nocker, S., Finley, D., and Vierstra, R.D.** (1998). Multi-ubiquitin chain binding and protein degradation are mediated by distinct domains within the 26S proteasome subunit MCB1. *J. Biol. Chem.* **273**: 1970–1981.
- Fujinami, K., Tanahashi, N., Tanaka, K., Ichihara, A., Cejka, Z., Baumeister, W., Miyawaki, M., Sato, T., and Nakagawa, H.** (1994). Purification and characterization of the 26S proteasome from spinach leaves. *J. Biol. Chem.* **269**: 25905–25910.
- Fukunaga, K., Kudo, T., Toh-e, A., Tanaka, K., and Saeki, Y.** (2010). Dissection of the assembly pathway of the proteasome lid in *Saccharomyces cerevisiae*. *Biochem. Biophys. Res. Commun.* **396**: 1048–53.
- Funakoshi, M., Tomko, R.J., Kobayashi, H., and Hochstrasser, M.** (2009). Multiple assembly chaperones govern biogenesis of the proteasome regulatory particle base. *Cell* **137**: 887–899.
- Ganoth, D., Leshinsky, E., Eytan, E., and Hershko, A.** (1988). A multi-component system that degrades proteins conjugated to ubiquitin: resolution of factors and evidence for ATP-dependent complex formation. *J. Biol. Chem.* **263**: 12412–12419.

- Glickman, M.H., Rubin, D.M., Coux, O., Wefes, I., Pfeifer, G., Cjeka, Z., Baumeister, W., Fried, V.A., and Finley, D.** (1998a). A subcomplex of the proteasome regulatory particle required for ubiquitin-conjugate degradation and related to the COP9-signalosome and eIF3. *Cell* **94**: 615–623.
- Glickman, M.H., Rubin, D.M., Fried, V.A., and Finley, D.** (1998b). The regulatory particle of the *Saccharomyces cerevisiae* proteasome. *Mol. Cell. Biol.* **18**: 3149–3162.
- Glotzer, M., Murray, A.W., and Kirschner, M.W.** (1991). Cyclin is degraded by the ubiquitin pathway. *Nature* **349**: 132–138.
- Glynne, R., Powis, S.H., Beck, S., Kelly, A., Kerr, L.A., and Trowsdale, J.** (1991). A proteasome-related gene between the two ABC transporter loci in the class II region of the human MHC. *Nature* **353**: 357–60.
- Groettrup, M., Kirk, C.J., and Basler, M.** (2009). Proteasomes in immune cells: more than peptide producers? *Nat. Rev. Immunol.* **10**: 73–78.
- Groettrup, M., Kraft, R., Kostka, S., Standera, S., Stohwasser, R., and Kloetzel, P.M.** (1996). A third interferon-gamma-induced subunit exchange in the 20S proteasome. *Eur. J. Immunol.* **26**: 863–9.
- Groll, M., Bajorek, M., Köhler, A., Moroder, L., Rubin, D.M., Huber, R., Glickman, M.H., and Finley, D.** (2000). A gated channel into the proteasome core particle. *Nat. Struct. Biol.* **7**: 1062–1067.
- Groll, M., Heinemeyer, W., Jäger, S., Ullrich, T., Bochtler, M., Wolf, D.H., and Huber, R.** (1999). The catalytic sites of 20S proteasomes and their role in subunit maturation: a mutational and crystallographic study. *Proc. Natl. Acad. Sci. U.S.A.* **96**: 10976–10983.
- Gu, Z.C. and Enenkel, C.** (2014). Proteasome assembly. *Cell. Mol. Life Sci.* **71**: 4729–4745.
- Hanna, J., Hathaway, N.A., Tone, Y., Crosas, B., Elsasser, S., Kirkpatrick, D.S., Leggett, D.S., Gygi, S.P., King, R.W., and Finley, D.** (2006). The deubiquitinating enzyme Ubp6 functions non-catalytically to delay proteasomal degradation. *Cell* **127**: 99–111.
- Hatfield, P.M. and Vierstra, R.D.** (1989). Ubiquitin-dependent proteolytic pathway in wheat germ: isolation of multiple forms of the E1 ubiquitin-activating enzyme. *Biochemistry (Mosc.)* **28**: 735–742.

- Heinemeyer, W., Fischer, M., Krimmer, T., Stachon, U., and Wolf, D.H.** (1997). The active sites of the eukaryotic 20S proteasome and their involvement in subunit precursor processing. *J. Biol. Chem.* **272**: 25200–25209.
- Hershko, A., Ciechanover, A., Heller, H., Haas, A.L., and Rose, I.A.** (1980). Proposed role of ATP in protein breakdown: conjugation of proteins with multiple chains of the polypeptide of ATP-dependent proteolysis. *Proc. Natl. Acad. Sci. U.S.A.* **77**: 1783–1786.
- Hirano, Y., Hayashi, H., Iemura, S.i., Hendil, K.B., Niwa, S.i., Kishimoto, T., Kasahara, M., Natsume, T., Tanaka, K., and Murata, S.** (2006). Cooperation of Multiple Chaperones Required for the Assembly of Mammalian 20S Proteasomes. *Mol. Cell* **24**: 977–984.
- Hirano, Y., Hendil, K.B., Yashiroda, H., Iemura, S.i., Nagane, R., Hioki, Y., Natsume, T., Tanaka, K., and Murata, S.** (2005). A heterodimeric complex that promotes the assembly of mammalian 20S proteasomes. *Nature* **437**: 1381–1385.
- Hirano, Y., Kaneko, T., Okamoto, K., Bai, M., Yashiroda, H., Furuyama, K., Kato, K., Tanaka, K., and Murata, S.** (2008). Dissecting beta-ring assembly pathway of the mammalian 20S proteasome. *EMBO J.* **27**: 2204–13.
- Hochstrasser, M., Ellison, M.J., Chau, V., and Varshavsky, A.** (1991). The short-lived MAT2 transcriptional regulator is ubiquitinated in vivo. *Proc. Natl. Acad. Sci. U.S.A.* **88**: 4606–4610.
- Hough, R., Pratt, G., and Rechsteiner, M.** (1986). Ubiquitin-lysozyme conjugates: identification and characterization of an ATP-dependent protease from rabbit reticulocyte lysates. *J. Biol. Chem.* **261**: 2400–2408.
- Hough, R., Pratt, G., and Rechsteiner, M.** (1987). Purification of two high molecular weight proteases from rabbit reticulocyte lysate. *J. Biol. Chem.* **262**: 8303–8313.
- Hough, R. and Rechsteiner, M.** (1986). Ubiquitin-lysozyme conjugates: purification and susceptibility to proteolysis. *J. Biol. Chem.* **261**: 2391–2399.
- Hua, Z., Pool, J.E., Schmitz, R.J., Schultz, M.D., Shiu, S.H., Ecker, J.R., and Vierstra, R.D.** (2013). Epigenomic programming contributes to the genomic drift evolution of the F-Box protein superfamily in *Arabidopsis*. *Proc. Natl. Acad. Sci. U.S.A.* **110**: 16927–16932.

- Hua, Z. and Vierstra, R.D.** (2011). The cullin-RING ubiquitin-protein ligases. *Annu. Rev. Plant Biol.* **62**: 299–334.
- Husnjak, K. and Dikic, I.** (2012). Ubiquitin-binding proteins: decoders of ubiquitin-mediated cellular functions. *Annu. Rev. Biochem.* **81**: 291–322.
- Husnjak, K., Elsasser, S., Zhang, N., Chen, X., Randles, L., Shi, Y., Hofmann, K., Walters, K.J., Finley, D., and Dikic, I.** (2008). Proteasome subunit Rpn13 is a novel ubiquitin receptor. *Nature* **453**: 481–488.
- Jager, S., Groll, M., Huber, R., Wolf, D.H., and Heinemeyer, W.** (1999). Proteasome beta-type subunits: unequal roles of propeptides in core particle maturation and a hierarchy of active site function. *J. Mol. Biol.* **291**: 997–1013.
- Kaneko, T., Hamazaki, J., Iemura, S., Sasaki, K., Furuyama, K., Natsume, T., Tanaka, K., and Murata, S.** (2009). Assembly pathway of the Mammalian proteasome base subcomplex is mediated by multiple specific chaperones. *Cell* **137**: 914–25.
- Katsir, L., Schilmiller, A.L., Staswick, P.E., He, S.Y., and Howe, G.A.** (2008). COI1 is a critical component of a receptor for jasmonate and the bacterial virulence factor coronatine. *Proc. Natl. Acad. Sci. U.S.A.* **105**: 7100–7105.
- Köhler, A., Cascio, P., Leggett, D.S., Woo, K.M., Goldberg, A.L., and Finley, D.** (2001). The axial channel of the proteasome core particle is gated by the Rpt2 ATPase and controls both substrate entry and product release. *Mol. Cell* **7**: 1143–1152.
- Kim, H.J., Oh, S.A., Brownfield, L., Hong, S.H., Ryu, H., Hwang, I., Twell, D., and Nam, H.G.** (2008). Control of plant germline proliferation by SCF(FBL17) degradation of cell cycle inhibitors. *Nature* **455**: 1134–7.
- Kisselev, A.F., van der Linden, W.A., and Overkleeft, H.S.** (2012). Proteasome inhibitors: an expanding army attacking a unique target. *Chem. Biol.* **19**: 99–115.
- Kremp, A., Schliephacke, M., Kull, U., and Schmid, H.P.** (1986). Prosomes exist in plant cells too. *Exp. Cell. Res.* **166**: 553–557.
- Kusmierczyk, A.R., Kunjappu, M.J., Funakoshi, M., and Hochstrasser, M.** (2008). A multimeric assembly factor controls the formation of alternative 20S proteasomes. *Nat. Struct. Mol. Biol.* **15**: 237–244.

- Kusmierczyk, A.R., Kunjappu, M.J., Kim, R.Y., and Hochstrasser, M.** (2011). A conserved 20S proteasome assembly factor requires a C-terminal HbYX motif for proteasomal precursor binding. *Nat. Struct. Mol. Biol.* **18**: 622–9.
- Lander, G.C., Estrin, E., Matyskiela, M.E., Bashore, C., Nogales, E., and Martin, A.** (2012). Complete subunit architecture of the proteasome regulatory particle. *Nature* **482**: 186–191.
- Lasker, K., Förster, F., Bohn, S., Walzthoeni, T., Villa, E., Unverdorben, P., Beck, F., Aebersold, R., Sali, A., and Baumeister, W.** (2012). Molecular architecture of the 26S proteasome holocomplex determined by an integrative approach. *Proc. Natl. Acad. Sci. U.S.A.* **109**: 1380–1387.
- Le Tallec, B., Barrault, M.B., Courbeyrette, R., Guérois, R., Marsolier-Kergoat, M.C., and Peyroche, A.** (2007). 20S proteasome assembly is orchestrated by two distinct pairs of chaperones in yeast and mammals. *Mol. Cell* **27**: 660–674.
- Le Tallec, B., Barrault, M.B., Guerois, R., Carre, T., and Peyroche, A.** (2009). Hsm3/S5b participates in the assembly pathway of the 19S regulatory particle of the proteasome. *Mol. Cell* **33**: 389–99.
- Lee, L.W., Moomaw, C.R., Orth, K., McGuire, M.J., DeMartino, G.N., and Slaughter, C.A.** (1990). Relationships among the subunits of the high molecular weight proteinase, macropain (proteasome). *Biochim. Biophys. Acta* **1037**: 178–85.
- Leggett, D.S., Glickman, M.H., and Finley, D.** (2005). Purification of proteasomes, proteasome sub-complexes, and proteasome-associated proteins from budding yeast. *Methods Mol. Biol.* **301**: 57–70.
- Leggett, D.S., Hanna, J., Borodovsky, A., Crosas, B., Schmidt, M., Baker, R.T., Walz, T., Ploegh, H.L., and Finley, D.** (2002). Multiple associated proteins regulate proteasome structure and function. *Mol. Cell* **10**: 495–507.
- Lehmann, A., Niewienda, A., Jechow, K., Janek, K., and Enenkel, C.** (2010). Ecm29 fulfills quality control functions in proteasome assembly. *Mol. Cell* **38**: 879–888.
- Li, D., Dong, Q., Tao, Q., Gu, J., Cui, Y., Jiang, X., Yuan, J., Li, W., Xu, R., Jin, Y., Li, P., Weaver, D.T., Ma, Q., Liu, X., and Cao, C.** (2015). c-Abl regulates proteasome abundance by controlling the ubiquitin-proteasomal degradation of PSMA7 subunit. *Cell Rep* **10**: 484–96.

- Li, X., Kusmierczyk, A.R., Wong, P., Emili, A., and Hochstrasser, M.** (2007). beta-Subunit appendages promote 20S proteasome assembly by overcoming an Ump1-dependent checkpoint. *EMBO J.* **26:** 2339–49.
- Lin, Y.L., Sung, S.C., Tsai, H.L., Yu, T.T., Radjacompare, R., Usharani, R., Fatimababy, A.S., Lin, H.Y., Wang, Y.Y., and Fu, H.** (2011). The defective proteasome, not substrate recognition function, is responsible for the null phenotypes of the *Arabidopsis* proteasome subunit RPN10. *Plant Cell* **23:** 2754–2773.
- Malik, M.N., Spivack, W.D., Sheikh, A.M., and Fenko, M.D.** (2004). The 26S proteasome in garlic (*Allium sativum*): purification and partial characterization. *J. Agric. Food. Chem.* **52:** 3350–3355.
- Marques, A.J., Glanemann, C., Ramos, P.C., and Dohmen, R.J.** (2007). The C-terminal extension of the beta7 subunit and activator complexes stabilize nascent 20 S proteasomes and promote their maturation. *J. Biol. Chem.* **282:** 34869–76.
- McClure, B.** (2004). S-RNase and SLF Determine S-Haplotype-Specific Pollen Recognition and Rejection. *The Plant Cell Online* **16:** 2840–2847.
- Más, P., Kim, W.Y., Somers, D.E., and Kay, S.A.** (2003). Targeted degradation of TOC1 by ZTL modulates circadian function in *Arabidopsis thaliana*. *Nature* **426:** 567–570.
- Murase, K., Hirano, Y., Sun, T.P., and Hakoshima, T.** (2008). Gibberellin-induced DELLA recognition by the gibberellin receptor GID1. *Nature* **456:** 459–63.
- Murata, S., Sasaki, K., Kishimoto, T., Niwa, S., Hayashi, H., Takahama, Y., and Tanaka, K.** (2007). Regulation of CD8+ T cell development by thymus-specific proteasomes. *Science* **316:** 1349–53.
- Murata, S., Yashiroda, H., and Tanaka, K.** (2009). Molecular mechanisms of proteasome assembly. *Nat. Rev. Mol. Cell Biol.* **10:** 104–115.
- Murray, P.F., Giordano, C.V., Passeron, S., and Barneix, A.J.** (1997). Purification and characterization of the 20S proteasome from wheat leaves. *Plant Sci.* **125:** 127–136.
- Neefjes, J., Jongsma, M.L., Paul, P., and Bakke, O.** (2011). Towards a systems understanding of MHC class I and MHC class II antigen presentation. *Nat. Rev. Immunol.* **11:** 823–36.

- Nelson, C.J., Li, L., and Millar, A.H.** (2014). Quantitative analysis of protein turnover in plants. *Proteomics* **14**: 579–592.
- Ortiz-Navarrete, V., Seelig, A., Gernold, M., Frentzel, S., Kloetzel, P.M., and Hammerling, G.J.** (1991). Subunit of the '20S' proteasome (multicatalytic proteinase) encoded by the major histocompatibility complex. *Nature* **353**: 662–4.
- Ozaki, M., Fujinami, K., Tanaka, K., Amemiya, Y., Sato, T., Ogura, N., and Nakagawa, H.** (1992). Purification and initial characterization of the proteasome from the higher plant Spinacia oleracea. *J. Biol. Chem.* **267**: 21678–21684.
- Padmanabhan, A., Vuong, S.A., and Hochstrasser, M.** (2016). Assembly of an Evolutionarily Conserved Alternative Proteasome Isoform in Human Cells. *Cell Rep* **14**: 2962–74.
- Paraskevopoulos, K., Kriegenburg, F., Tatham, M.H., Rösner, H.I., Medina, B., Larsen, I.B., Brandstrup, R., Hardwick, K.G., Hay, R.T., Kragelund, B.B., Hartmann-Petersen, R., and Gordon, C.** (2014). Dss1 is a 26S proteasome ubiquitin receptor. *Mol. Cell* **56**: 453–461.
- Park, S., Roelofs, J., Kim, W., Robert, J., Schmidt, M., Gygi, S.P., and Finley, D.** (2009). Hexameric assembly of the proteasomal ATPases is templated through their C termini. *Nature* **459**: 866–70.
- Pathare, G.R., Nagy, I., Bohn, S., Unverdorben, P., Hubert, A., Körner, R., Nickell, S., Lasker, K., Sali, A., Tamura, T., Nishioka, T., Förster, F., Baumeister, W., and Bracher, A.** (2012). The proteasomal subunit Rpn6 is a molecular clamp holding the core and regulatory sub-complexes together. *Proc. Natl. Acad. Sci. U.S.A.* **109**: 149–154.
- Peters, J.M., Harris, J.R., and Kleinschmidt, J.A.** (1991). Ultrastructure of the approximately 26S complex containing the approximately 20S cylinder particle (multi-catalytic proteinase/proteasome). *Eur. J. Cell Biol.* **56**: 422–432.
- Rabl, J., Smith, D.M., Yu, Y., Chang, S.C., Goldberg, A.L., and Cheng, Y.** (2008). Mechanism of gate opening in the 20S proteasome by the proteasomal ATPases. *Mol. Cell* **30**: 360–368.
- Ramos, P.C., Höckendorff, J., Johnson, E.S., Varshavsky, A., and Dohmen, R.J.** (1998). Ump1p is required for proper maturation of the 20S proteasome and becomes its substrate upon completion of the assembly. *Cell* **92**: 489–499.

- Ruschak, A.M., Religa, T.L., Breuer, S., Witt, S., and Kay, L.E.** (2010). The proteasome antechamber maintains substrates in an unfolded state. *Nature* **467**: 868–871.
- Russell, J.D., Scalf, M., Book, A.J., Ladror, D.T., Vierstra, R.D., Smith, L.M., and Coon, J.J.** (2013). Characterization and quantification of intact 26S proteasome proteins by real-time measurement of intrinsic fluorescence prior to top-down mass spectrometry. *PLoS One* **8**: e58157.
- Sadre-Bazzaz, K., Whitby, F.G., Robinson, H., Formosa, T., and Hill, C.P.** (2010). Structure of a Blm10 complex reveals common mechanisms for proteasome binding and gate opening. *Mol. Cell* **37**: 728–735.
- Saeki, Y., Toh, E.A., Kudo, T., Kawamura, H., and Tanaka, K.** (2009). Multiple proteasome-interacting proteins assist the assembly of the yeast 19S regulatory particle. *Cell* **137**: 900–13.
- Sakata, E., Bohn, S., Mihalache, O., Kiss, P., Beck, F., Nagy, I., Nickell, S., Tanaka, K., Saeki, Y., Förster, F., and Baumeister, W.** (2012). Localization of the proteasomal ubiquitin receptors Rpn10 and Rpn13 by electron cryo-microscopy. *Proc. Natl. Acad. Sci. U.S.A.* **109**: 1479–1484.
- Sakata, E., Stengel, F., Fukunaga, K., Zhou, M., Saeki, Y., Förster, F., Baumeister, W., Tanaka, K., and Robinson, C.V.** (2011). The catalytic activity of Ubp6 enhances maturation of the proteasomal regulatory particle. *Mol. Cell* **42**: 637–649.
- Samach, A., Klenz, J.E., Kohalmi, S.E., Risseeuw, E., Haughn, G.W., and Crosby, W.L.** (1999). The UNUSUAL FLORAL ORGANS gene of *Arabidopsis thaliana* is an F-box protein required for normal patterning and growth in the floral meristem. *Plant J.* **20**: 433–45.
- Santner, A. and Estelle, M.** (2010). The ubiquitin-proteasome system regulates plant hormone signaling. *Plant J.* **61**: 1029–40.
- Satoh, T., Saeki, Y., Hiromoto, T., Wang, Y.H., Uekusa, Y., Yagi, H., Yoshihara, H., Yagi-Utsumi, M., Mizushima, T., Tanaka, K., and Kato, K.** (2014). Structural basis for proteasome formation controlled by an assembly chaperone nas2. *Structure* **22**: 731–43.
- Schellenberg, B., Ramel, C., and Dudler, R.** (2010). *Pseudomonas syringae* virulence factor syringolin A counteracts stomatal immunity by proteasome inhibition. *Mol. Plant. Microbe Interact.* **23**: 1287–93.

- Schliephacke, M., Kremp, A., Schmid, H.P., Köhler, K., and Kull, U.** (1991). Pro-somes (proteasomes) of higher plants. *Eur. J. Cell Biol.* **55**: 114–121.
- Schmid, H.P., Akhayat, O., Martins, C., Puvion, F., Köhler, K., and Scherrer, K.** (1984). The prosome: a ubiquitous morphologically distinct RNP particle associated with repressed mRNPs and containing specific scRNA and a characteristic set of proteins. *EMBO J.* **3**: 29–34.
- Schmidt, M., Haas, W., Crosas, B., Santamaria, P.G., Gygi, S.P., Walz, T., and Finley, D.** (2005). The HEAT repeat protein Blm10 regulates the yeast proteasome by capping the core particle. *Nat. Struct. Mol. Biol.* **12**: 294–303.
- Schreiner, P., Chen, X., Husnjak, K., Randles, L., Zhang, N., Elsasser, S., Finley, D., Dikic, I., Walters, K.J., and Groll, M.** (2008). Ubiquitin docking at the proteasome through a novel pleckstrin-homology domain interaction. *Nature* **453**: 548–552.
- Schrodinger, L.** (2015). The PyMOL Molecular Graphics System, Version 1.8.
- Shabek, N. and Zheng, N.** (2014). Plant ubiquitin ligases as signaling hubs. *Nat. Struct. Mol. Biol.* **21**: 293–296.
- Shanklin, J., Jabben, M., and Vierstra, R.D.** (1987). Red light-induced formation of ubiquitin-phytochrome conjugates: identification of possible intermediates of phytochrome degradation. *Proc. Natl. Acad. Sci. U.S.A.* **84**: 359–363.
- Shi, Y., Chen, X., Elsasser, S., Stocks, B.B., Tian, G., Lee, B.H., Shi, Y., Zhang, N., de Poot, S.A., Tuebing, F., Sun, S., Vannoy, J., Tarasov, S.G., Engen, J.R., Finley, D., and Walters, K.J.** (2016). Rpn1 provides adjacent receptor sites for substrate binding and deubiquitination by the proteasome. *Science* **351**.
- Shibahara, T., Kawasaki, H., and Hirano, H.** (2004). Mass spectrometric analysis of expression of ATPase subunits encoded by duplicated genes in the 19S regulatory particle of rice 26S proteasome. *Arch. Biochem. Biophys.* **421**: 34–41.
- Skoda, B. and Malek, L.** (1992). Dry pea seed proteasome: purification and enzymatic activities. *Plant Physiol.* **99**: 1515–1519.
- Smalle, J.A. and Vierstra, R.D.** (2004). The ubiquitin-26S proteasome proteolytic pathway. *Annu. Rev. Plant Biol.* **55**: 555–590.

- Smith, D.M., Kafri, G., Cheng, Y., Ng, D., Walz, T., and Goldberg, A.L.** (2005). ATP binding to PAN or the 26S ATPases causes association with the 20S proteasome, gate opening, and translocation of unfolded proteins. *Mol. Cell* **20**: 687–698.
- Sokolova, V., Li, F., Polovin, G., and Park, S.** (2015). Proteasome Activation is Mediated via a Functional Switch of the Rpt6 C-terminal Tail Following Chaperone-dependent Assembly. *Sci. Rep.* **5**: 14909.
- Stadtmueller, B.M., Kish-Trier, E., Ferrell, K., Petersen, C.N., Robinson, H., Myszka, D.G., Eckert, D.M., Formosa, T., and Hill, C.P.** (2012). Structure of a proteasome Pba1-Pba2 complex: implications for proteasome assembly, activation, and biological function. *J. Biol. Chem.* **287**: 37371–82.
- Stone, S.L., Anderson, E.M., Mullen, R.T., and Goring, D.R.** (2003). ARC1 is an E3 ubiquitin ligase and promotes the ubiquitination of proteins during the rejection of self-incompatible Brassica pollen. *Plant Cell* **15**: 885–98.
- Tomko, R. J., J. and Hochstrasser, M.** (2011). Incorporation of the Rpn12 subunit couples completion of proteasome regulatory particle lid assembly to lid-base joining. *Mol. Cell* **44**: 907–17.
- Tomko, R. J., J. and Hochstrasser, M.** (2014). The intrinsically disordered Sem1 protein functions as a molecular tether during proteasome lid biogenesis. *Mol. Cell* **53**: 433–43.
- Tomko, R.J. and Hochstrasser, M.** (2013). Molecular architecture and assembly of the eukaryotic proteasome. *Annu. Rev. Biochem.* **82**: 415–445.
- Uechi, H., Hamazaki, J., and Murata, S.** (2014). Characterization of the testis-specific proteasome subunit alpha4s in mammals. *J. Biol. Chem.* **289**: 12365–74.
- Unverdorben, P., Beck, F., Ied, P., Schweitzer, A., Pfeifer, G., Plitzko, J.M., Baumeister, W., and Förster, F.** (2014). Deep classification of a large cryo-EM dataset defines the conformational landscape of the 26S proteasome. *Proc. Natl. Acad. Sci. U.S.A.* **111**: 5544–5549.
- Van Nocker, S., Devereaux, Q., Rechsteiner, M., and Vierstra, R.D.** (1996). The Arabidopsis MBP1 gene encodes a conserved ubiquitin recognition component of the 26S proteasome. *Proc. Natl. Acad. Sci. U.S.A.* **93**: 856–860.

- Verma, R., Aravind, L., Oania, R., McDonald, W.H., Yates, J.R., Koonin, E.V., and Deshaies, R.J.** (2002). Role of the Rpn11 metalloprotease in deubiquitination and degradation by the 26S proteasome. *Science* **298**: 611–615.
- Vierstra, R.D.** (1993). Protein degradation in plants. *Annu. Rev. Plant Phys.* **44**: 385–410.
- Vierstra, R.D.** (2009). The ubiquitin-26S proteasome system at the nexus of plant biology. *Nat. Rev. Mol. Cell Biol.* **10**: 385–397.
- Vierstra, R.D. and Sullivan, M.L.** (1988). Hemin inhibits ubiquitin-dependent proteolysis in both a higher plant and yeast. *Biochemistry (Mosc.)* **27**: 3290–3295.
- Wani, P.S., Suppahia, A., Capalla, X., Ondracek, A., and Roelofs, J.** (2016). Phosphorylation of the C-terminal tail of proteasome subunit alpha7 is required for binding of the proteasome quality control factor Ecm29. *Sci. Rep.* **6**: 27873.
- Waxman, L., Fagan, J.M., and Goldberg, A.L.** (1987). Demonstration of two distinct high molecular weight proteases in rabbit reticulocytes, one of which degrades ubiquitin conjugates. *J. Biol. Chem.* **262**: 2451–2457.
- Weber, M.H., Savulescu, A.F., Jando, J., Bissinger, T., Harel, A., Glickman, M.H., and Enenkel, C.** (2013). Blm10 facilitates nuclear import of proteasome core particles. *EMBO J.* **32**: 2697–2707.
- Wilkinson, K.D., Urban, M.K., and Haas, A.L.** (1980). Ubiquitin is the ATP-dependent proteolysis factor I of rabbit reticulocytes. *J. Biol. Chem.* **255**: 7529–7532.
- Witt, E., Zantopf, D., Schmidt, M., Kraft, R., Kloetzel, P.M., and Kruger, E.** (2000). Characterisation of the newly identified human Ump1 homologue POMP and analysis of LMP7(beta 5i) incorporation into 20 S proteasomes. *J. Mol. Biol.* **301**: 1–9.
- Woffenden, B.J., Freeman, T.B., and Beers, E.P.** (1998). Proteasome inhibitors prevent tracheary element differentiation in *Zinnia* mesophyll cell cultures. *Plant Physiol.* **118**: 419–429.
- Worden, E.J., Padovani, C., and Martin, A.** (2014). Structure of the Rpn11-Rpn8 dimer reveals mechanisms of substrate deubiquitination during proteasomal degradation. *Nat. Struct. Mol. Biol.* **21**: 220–227.

- Xie, Y. and Varshavsky, A.** (2000). Physical association of ubiquitin ligases and the 26S proteasome. *Proc. Natl. Acad. Sci. U.S.A.* **97**: 2497–2502.
- Yanagawa, Y., Ohhashi, A., Murakami, Y., Saeki, Y., Yokosawa, H., Tanaka, K., Hashimoto, J., Sato, T., and Nakagawa, H.** (1999). Purification and characterization of the 26S proteasome from cultured rice (*Oryza sativa*) cells. *Plant Sci.* **149**: 33–41.
- Yang, P., Fu, H., Walker, J.M., Papa, C.M., Smalle, J., Ju, Y.M., and Vierstra, R.D.** (2004). Purification of the *Arabidopsis* 26S proteasome: biochemical and molecular analyses revealed the presence of multiple isoforms. *J. Biol. Chem.* **279**: 6401–6413.
- Yashiroda, H., Mizushima, T., Okamoto, K., Kameyama, T., Hayashi, H., Kishimoto, T., Niwa, S., Kasahara, M., Kurimoto, E., Sakata, E., Takagi, K., Suzuki, A., Hirano, Y., Murata, S., Kato, K., Yamane, T., and Tanaka, K.** (2008). Crystal structure of a chaperone complex that contributes to the assembly of yeast 20S proteasomes. *Nat. Struct. Mol. Biol.* **15**: 228–36.
- Yoshimura, T., Kameyama, K., Takagi, T., Ikai, A., Tokunaga, F., Koide, T., Tanahashi, N., Tamura, T., Cejka, Z., Baumeister, W., Tanaka, K., and Ichihara, A.** (1993). Molecular characterization of the 26S proteasome complex from rat liver. *J. Struct. Biol.* **111**: 200–211.
- Zhang, Y., Zhao, Z., and Xue, Y.** (2009). Roles of Proteolysis in Plant Self-Incompatibility. *Annu. Rev. Plant Biol.* **60**: 21–42.
- Zhao, L., Huang, J., Zhao, Z., Li, Q., Sims, T.L., and Xue, Y.** (2010). The Skp1-like protein SSK1 is required for cross-pollen compatibility in S-RNase-based self-incompatibility. *Plant J.* **62**: 52–63.

Chapter 2

MORPHEUS SPECTRAL COUNTER: A COMPUTATIONAL TOOL

FOR LABEL-FREE QUANTITATIVE MASS SPECTROMETRY

USING THE MORPHEUS SEARCH ENGINE

2.1. Summary

Label-free quantitative MS based on the Normalized Spectral Abundance Factor (NSAF) has emerged as a straightforward and robust method to determine the relative abundance of individual proteins within complex mixtures. Here, we present Morpheus Spectral Counter (MSpC) as the first computational tool that directly calculates NSAF values from output obtained from Morpheus, a fast, open-source, peptide-MS/MS matching engine compatible with high-resolution accurate-mass instruments. NSAF has distinct advantages over other MS-based quantification methods, including a higher dynamic range as compared to isobaric tags, no requirement to align and re-extract MS1 peaks, and increased speed. MSpC features an easy to use graphic user interface that additionally calculates both distributed and unique NSAF values to permit analyses of both protein families and isoforms/proteoforms. MSpC determinations of protein concentration were linear over

several orders of magnitude based on the analysis of several high-mass accuracy datasets either obtained from PRIDE or generated with total cell extracts spiked with purified *Arabidopsis* 20S proteasomes. The MSpC software was developed in C# and is open sourced under a permissive license with the code made available at http://dcgempeline.github.io/Morpheus_SpC/.

2.2. Main Text

Quantification of individual polypeptides within complex mixtures by MS is an extremely useful tool to understand proteomic changes in organisms during growth and development, and after environmental perturbation (Wong and Cagney, 2010). While a number of MS/MS strategies have been developed to measure protein abundance, including Stable Isotope Labeling by Amino Acids in Cell Culture (SILAC), labeling with isobaric tags, and Absolute Quantification of proteins (AQUA) (Gerber *et al.*, 2003; Ong *et al.*, 2002; Ross *et al.*, 2004; Thompson *et al.*, 2003), label-free quantification (LFQ) have become increasingly popular given their simplicity and low cost (Wong and Cagney, 2010; Zhang *et al.*, 2006). One LFQ strategy infers abundance from the number of observed peptide spectra matches (PSMs). For these PSM-based approaches, changes in protein abundance can be generated artificially when total PSMs differ among samples and because longer proteins

tend to produce more raw counts. For these reasons normalizing for both protein length and total PSMs is paramount. While this adjustment can be made in a number of ways; one of the most straight forward methods is to use Normalized Spectral Abundance Factor (NSAF), a length- and count-normalized measure for each protein (Zybailov *et al.*, 2006). Further improvements to the NSAF algorithm have been made by accounting for shared peptides in distributed NSAF (dNSAF), which distributes common PSMs among a family of isoforms/proteoforms based on the number of distinct PSMs observed for each isoform/proteoform, and unique NSAF (uNSAF), which ignores shared PSMs and only assigns distinct PSMs to each specific isoform/proteoform (Zhang *et al.*, 2010).

The Morpheus MS search engine was recently designed for high-resolution, accurate-mass data obtained from Orbitrap-based instruments to provide faster matching of spectra to peptides (Wenger and Coon, 2013). Unfortunately, no downstream automated tools are available to facilitate LFQ analysis, which can be quite challenging, if not impossible, to complete manually when accounting for shared peptides. To overcome this bottleneck, we developed Morpheus Spectral Counter (MSpC) as the first LFQ computational tool that integrates directly with Morpheus to calculate NSAF, dNSAF, uNSAF, and corrected PSM (Fermin *et al.*, 2011) values in complex protein samples. MSpC is fully automated, and only requires a Morpheus




Figure 2.1: MSpC Graphic User Interface (GUI) and software flow chart. (A) Screenshot of the GUI. The input requires the user to select a Morpheus search summary file containing experiments to be analyzed. The user can optionally select a whitelist to filter output, and select an output directory. Additional options can set peptide and protein FDR cutoffs, and method of quantification for output, including Normalized Spectral Abundance Factor (NSAF), distributed NSAF (dNSAF), and unique NSAF (uNSAF). A progress bar highlights completion of the analysis. (B) Data analysis flow chart. Experiments and groups of experiments to be analyzed are imported through the Morpheus summary file. PSMs and protein groups are filtered at the specified FDR cutoff with a default of 1%.

search summary file (summary.tsv) as input. The user interface (see Figure 2.1 (A)) allows one to select the summary file and displays the raw MS/MS files that will be analyzed by MSpC. Due to shared peptides being attributed to only one instance of a protein group in Morpheus's PSM file, PSMs are re-matched to all possible protein groups. PSMs are then cataloged as shared or as unique (distinctly matching one protein group) to generate NSAF, dNSAF, and uNSAF outputs. Finally, the output can be filtered for proteins of interest by specifying a comma delimited file containing unique identifiers and descriptions. Some important features of MSpC are its ability to handle fractionation experiments as input, and the ability to whitelist proteins of interest in the output by specifying a csv file (see Tutorial 2.5). Options exist to specify global PSM and protein group FDR rates (thus avoiding increased FDRs when one analyzes many experiments at once), to output NSAF, dNSAF, and uNSAF values, to require a minimum number of unique peptides to quantify a protein, and to specify an output directory. A progress bar indicates completion of the analysis by MSpC. To validate the accuracy of MSpC, we analyzed two MS/MS datasets available in PRIDE that were previously generated by high-energy collision-induced dissociation using Thermo Q-Exactive Orbitrap instruments.

Here, *Xenopus* egg (see top, Figure 2.2) and embryo (bottom, Figure 2.2) extracts




Figure 2.2: Confirmation of MSpC accuracy by analysis of MS/MS datasets generated with the Universal Proteome Standard 2 (UPS2). The array of UPS2 standards were spiked into *Xenopus laevis* egg (**Top**) and embryo (**Bottom**) extracts at a range of concentrations. Following MS/MS analysis, dNSAF values for each protein were determined by Morpheus and MSpC. (**A**) A log-log plot of dNSAF versus concentration for each UPS2 protein detected across each fmol range. (**B**) A log-log plot of average dNSAF vs average concentration of each group of UPS2 proteins at each fmol range: (50, 500, 5000, and 50,000 fmol).

were spiked at a 4:1 ratio with the Universal Proteome Standard 2 (UPS2), a mix of 48 purified proteins at defined molar ratios of 0.5, 5, 50, 500, 5000, and 50,000, with each ratio containing a different set of 8 of the 48 proteins. As shown in Figure 2.2 A, when the Morpheus/MSpC pipeline was used to calculate the average dNSAF value for each UPS2 protein, requiring only a single unique peptide to quantify, strong linear correlations ($R^2 = 0.886$ and 0.823) were obtained across a 1,000 fold change in abundance (50 fmol to 50,000 fmol). In fact, the R^2 values were similar to those obtained by others with PSM-based LFQ methods (Cox *et al.*, 2014; Tu *et al.*, 2014). This linear correlation was further strengthened when the dNSAF values were averaged for all UPS2 proteins within each of the concentration groups, with R^2 values of 0.994 and 0.992 for the egg and embryo datasets, respectively (see Figure 2.2 B). Notably, the slope of the concentration series was significantly less than unity, showing that NSAF measurements are not appropriate for absolute quantification, which was expected given that NSAF is a relative value.

We also reprocessed the UPS2 dataset using the option of requiring a minimum of two unique peptides for quantification, which should improve stringency. This option provided only a minor improvement in overall linearity for the average UPS2 dNSAF values, but decreased linearity when each UPS2 protein was considered individually and removed some UPS2 proteins at low concentrations (compare





Figure 2.3: Re- Analysis of MS/MS datasets generated with the Universal Proteome Standard 2 (UPS2). The array of UPS2 standards were spiked into *Xenopus laevis* egg (**Top**) and embryo (**Bottom**) extracts at a range of concentrations. Following MS/MS analysis, dNSAF values for each protein were determined by Morpheus and MSpC with a change from Figure 2.2 in that two unique peptides were required to quantify a protein. **(A)** A log-log plot of dNSAF versus concentration for each UPS2 protein detected across each fmol range. **(B)** A log-log plot of average dNSAF vs average concentration of each group of UPS2 proteins at each fmol range: (50, 500, 5000, and 50,000 fmol).

Figure 2.3 A to Figure 2.2 A). Consequently, caution should be exercised when selecting this option even though it might provide a slight improvement in stringency (see Discussion on Requiring Two Peptide 2.4). To demonstrate the utility and accuracy of MSpC as applied to our work, we analyzed 20S proteasomes isolated from *Arabidopsis thaliana*. This particle contains multiple subunits assembled in stoichiometric amounts, with many subunits encoded by two paralogous genes of sufficient amino acid identity (typically >90% (Yang *et al.*, 2004)) such that discrimination between paralogs can be challenging using LFQ approaches (Book *et al.*, 2010). To simulate changes in 20S proteasome abundance, we added varying amounts of trypsinized proteasomes (0.05 µg to 3 µg) to a fixed amount of trypsinized *Escherichia coli* lysate (0.5 µg) to generate proteasome/lysate ratios of ~0.091, 0.167, 0.333, 0.500, 0.667, 0.750 0.800, 0.857. The digests were then subjected to MS/MS and the dNSAF value for each subunit along with the uNSAF value for individual isoforms were calculated by the Morpheus/MSpC pipeline (see Methods 2.3). The data from this experiment are deposited in PRIDE with ID PXD003002. As shown in Figure 2.4, MSpC provided an excellent determination for the overall abundance of 20S proteasomes within a complex mixture, along with a good reflection of the abundance of individual subunits and their isoforms.

When the dNSAF values for all subunits for the *Arabidopsis* 20S proteasome

[Figure 2.4 caption follows on next page]



including their isoforms (representing 14 distinct subunits, 10 of which exist as isoform pairs) were summed, a very close approximation of the dNSAF/actual abundance was obtained (slope=0.875) with a very strong linear correlation ($R^2 = 0.99$) over a ~10-fold range in protein abundance. When each 20S proteasome subunit was analyzed individually, a strong linear response was also obtained ($R^2 > 0.90$) for a majority of subunits (Figure 2.4 C and Table 2.1).

For example, reasonably accurate concentration plots were obtained for the

Figure 2.4 (preceding page): Confirmation of MSpC accuracy by analysis of MS/MS datasets generated with affinity purified *Arabidopsis* 20S proteasomes spiked into a total cell lysate from *E. coli*. Following MS/MS analysis, the dNSAF and uNSAF values for each subunit/isoform were determined by Morpheus and MSpC. (A) A silver-stained SDS-PAGE gel of 20S proteasome samples affinity purified from 10-d-old *Arabidopsis* seedlings. The crude seedling extract(CE), sample buffer (SB), and affinity-purified 20S proteasome samples (Prot) are shown. (B) Quantification of trypsinized 20S proteasomes when mixed at varying ratios with trypsinized total protein lysates from *E. coli*. The spiked samples were subjected to MS/MS followed by data analysis with the Morpheus and MSpC. dNSAF values for each proteasome subunit were averaged across three technical replicates, then summed to obtain an estimate of abundance for the 20S proteasome, and plotted against their known ratios. The total protein load is listed at each point in ug. (C and D) dNSAF and uNSAF values determined from the data in panel B for individual subunits (C) and their isoforms (D) for several subunits of the 20S proteasome. (E) Quantification accuracy of the Morpheus/MSpC pipeline for determining the amount of each α and β subunit of the 20S proteasome. Single subunit isoforms are in black, whereas subunits having two isoforms are shown in black and grey to reflect the contributions of isoforms 1 and 2 respectively. Each bar represents the average of eight technical replicates (\pm SE). The dashed line represents the expected value of one assuming an equal stoichiometry of each subunit within the particle.

PAF (α 6) and PBD (β 4) subunits that are encoded by the PAF1/2 and PBF1/2 gene pairs, and for the PAG (α 7) and PBF (β 6) subunits that are encoded by single PAG1 and PBF1 genes (R^2 from 0.94 to 0.99). Even when we calculated uNSAF values for individual isoforms added to the *E. coli* lysate, strong linear responses were obtained (e.g., the PAF1/PAF2 and PBD1/PBD2 pairs) with robust correlations (R^2 from 0.89 to 0.95) (Figure 2.4 D). Taken together, MSpC worked well for relative LFQ analysis of a multi-subunit complex and its individual subunits and isoforms within a complex proteomic mixture.

The Morpheus/MSpC pipeline also allowed us to calculate the respective incorporation of each paralog in the complex (see Isoform Incorporation Methods 2.3.4). As shown in Figure 2.4E, these estimated/expected occupancies were close to unity for most subunits within both the α and β rings of the 20S proteasome. The only strong deviation was for PBD1/2 (β 4), which had a greater dNSAF value relative to other β subunits across the experiments analyzed (see Table 2.1). The calculations for uNSAF values also estimated the relative proportion of each isoform within the complex for those subunits expressed from paralogous genes. The data obtained are similar to prior studies of the complex involving quantitative top-down proteomic analysis of purified proteasome samples using ultra violet-intrinsic fluorescence to quantify tyrosine-containing subunits (Russell *et al.*, 2013). However, our MSpC

analysis provided a more complete picture as several subunit isoforms were difficult to quantify by fluorescence either because they lacked tryosine, or because their fluorescence peaks overlapped with those of other subunits/isoforms. Notably , the protein isoform ratios measured here agree well with the expression ratios for the paralogous genes (Book *et al.*, 2010), suggesting that the protein isoform abundance generally reflects the relative transcriptional activity of the gene pair. We consistently estimated slightly more α ring subunits (PAA-PAG) versus β ring subunits (PBA-PBG) in the final MSpC calculations (Figure 2.4 E). This deviation could represent enhanced detection of α ring versus β ring subunits, or more likely that purification via the tagged α ring subunit PAG1 also isolated assembly intermediates comprised of only α ring subunits.

We compared the Morpheus and MSpC pipeline to the next most comparable open source, spectral-count-based LFQ pipeline, The Trans Proteomic Pipeline (TPP) (Deutsch *et al.*, 2010) and ABACUS (Fermin *et al.*, 2011) using our datasets generated with the 20S proteasome/*E. coli* lysate mixture (see Tables 2.1 and 2.2). Morpheus/MSpC slightly outperformed TPP/ABACUS by having a greater overall accuracy (average linearity of 0.88 compared to 0.84), and by having more subunits showing an R^2 linear correlation greater than 0.9 (14/23 subunits for MSpC versus and 11/23 for ABACUS).

Table 2.1: Table of dNSAF values for each 20S proteasome subunit generated by analyzing the proteasome spike in experiments with the Morpheus and MSpC pipeline. The top half of the table lists α 1-7 (PAA-PAG) where 1 or 2 represent different isoforms) subunits, while the bottom half of the table lists β 1-7 subunits (PBA-PBG where 1 or 2 represent different isoforms)

Ratio	0.091	0.167	0.333	0.500	0.667	0.750	0.800	0.857		Pearsons
PAA1	0.0102	0.0180	0.0275	0.0392	0.0505	0.0489	0.0467	0.0530		0.952
PAA2	0.0063	0.0072	0.0118	0.0121	0.0139	0.0218	0.0351	0.0195		0.652
PAB1	0.0091	0.0117	0.0209	0.0391	0.0377	0.0377	0.0268	0.0397		0.721
PAB2	0.0017	0.0062	0.0147	0.0223	0.0295	0.0277	0.0303	0.0250		0.894
PAC1	0.0076	0.0092	0.0210	0.0339	0.0373	0.0395	0.0546	0.0512		0.955
PAD1	0.0056	0.0070	0.0103	0.0151	0.0160	0.0147	0.0149	0.0147		0.826
PAD2	0.0039	0.0068	0.0141	0.0162	0.0200	0.0225	0.0255	0.0228		0.959
PAE1	0.0046	0.0095	0.0208	0.0352	0.0395	0.0356	0.0389	0.0490		0.928
PAE2	0.0063	0.0082	0.0184	0.0240	0.0269	0.0282	0.0266	0.0293		0.925
PAF1	0.0106	0.0173	0.0239	0.0303	0.0292	0.0379	0.0406	0.0492		0.920
PAF2	0.0032	0.0083	0.0128	0.0155	0.0182	0.0184	0.0152	0.0209		0.847
PAG1	0.0080	0.0100	0.0208	0.0291	0.0394	0.0431	0.0498	0.0423		0.966
Ratio	0.091	0.167	0.333	0.500	0.667	0.750	0.800	0.857		Pearsons
PBA1	0.0076	0.0104	0.0170	0.0207	0.0246	0.0288	0.0337	0.0449		0.898
PBB1	0.0066	0.0074	0.0191	0.0276	0.0223	0.0206	0.0172	0.0227		0.498
PBB2	0.0009	0.0016	0.0034	0.0035	0.0110	0.0109	0.0160	0.0161		0.891
PBC1	0.0000	0.0000	0.0162	0.0309	0.0373	0.0344	0.0295	0.0388		0.877
PBC2	0.0000	0.0000	0.0000	0.0096	0.0144	0.0147	0.0146	0.0190		0.931
PBD1	0.0053	0.0097	0.0150	0.0147	0.0221	0.0228	0.0231	0.0271		0.955
PBD2	0.0094	0.0162	0.0286	0.0318	0.0426	0.0457	0.0441	0.0538		0.969
PBE1	0.0070	0.0097	0.0156	0.0224	0.0323	0.0373	0.0334	0.0519		0.914
PBE2	0.0007	0.0003	0.0036	0.0069	0.0080	0.0105	0.0106	0.0133		0.973
PBF1	0.0093	0.0122	0.0205	0.0310	0.0368	0.0412	0.0473	0.0459		0.991
PBG1	0.0085	0.0165	0.0203	0.0323	0.0331	0.0345	0.0334	0.0442		0.912
									Average	0.885

In addition to this modest improvement, we note that the Morpheus/MSpC pipeline required significantly less intermediary steps, thus accelerating the data analysis. Some of the additional steps in TPP/ABACUS could be automated from the command-line, but it would likely be a challenge for the average user. Importantly, we found that the Morpheus/MSpC pipeline was faster. Timing tests using the proteasome/*E. coli* spike data generated here showed that the Morpheus/MSpC

Table 2.2: Table of adj_NSAF values for each 20S proteasome subunit (equivalent to dNSAF) generated by analyzing the proteasome spike in experiments with the TPP and ABACUS pipeline. The top half of the table lists α 1-7 (PAA-PAG) where 1 or 2 represent different isoforms) subunits, while the bottom half of the table lists β 1-7 subunits (PBA-PBG where 1 or 2 represent different isoforms)

Ratio	0.091	0.167	0.333	0.500	0.667	0.750	0.800	0.857		Pearsons
PAA1	125.16	166.81	245.61	355.60	491.76	494.18	504.75	554.97		0.988
PAA2	57.52	70.52	114.60	126.35	132.06	162.25	358.13	154.62		0.493
PAB1	96.13	134.98	207.90	328.16	362.21	416.46	303.60	397.39		0.865
PAB2	29.67	79.42	153.34	282.91	365.22	299.47	310.19	299.93		0.849
PAC1	95.27	112.45	214.24	376.66	389.05	361.39	512.73	488.62		0.925
PAD1	62.58	76.16	106.03	196.03	179.37	182.99	164.04	188.71		0.794
PAD2	39.88	68.33	145.34	178.43	190.84	246.20	243.28	240.01		0.954
PAE1	50.41	97.53	193.18	368.87	464.79	397.27	468.64	534.16		0.951
PAE2	71.38	80.07	184.85	255.98	299.41	290.16	304.77	348.32		0.955
PAF1	132.67	179.60	207.68	253.51	253.04	327.11	334.73	460.76		0.835
PAF2	0.00	0.00	59.62	70.93	111.83	80.29	63.08	65.86		0.606
PAG1	108.09	142.82	271.18	389.94	495.94	530.90	632.72	540.86		0.966
<hr/>										
Ratio	0.091	0.167	0.333	0.500	0.667	0.750	0.800	0.857		Pearsons
PBA1	85.99	113.01	208.38	233.44	273.50	327.23	417.12	560.62		0.851
PBB1	47.77	80.66	195.35	324.53	303.26	220.36	200.51	216.77		0.440
PBB2	17.11	4.54	0.00	0.00	21.05	102.51	144.30	187.86		0.618
PBC1	27.19	40.65	165.46	345.14	364.19	386.59	338.47	373.16		0.885
PBC2	10.64	40.65	0.00	55.58	145.43	158.90	159.74	228.05		0.845
PBD1	50.14	94.46	168.58	167.50	244.25	248.17	252.65	319.61		0.945
PBD2	81.45	149.28	256.88	319.89	390.60	407.02	380.30	490.17		0.950
PBE1	81.01	107.76	196.21	302.90	365.03	404.25	378.67	441.03		0.981
PBE2	0.00	1.52	5.59	8.05	40.06	56.56	59.39	58.56		0.887
PBF1	71.53	112.26	190.45	226.89	259.90	366.60	431.42	420.23		0.943
PBG1	104.93	174.55	232.34	392.37	425.10	397.85	396.42	463.13		0.906
									Average	0.845

pipeline was 1.9-fold faster than the TPP/ABACUS pipeline (Figure 2.5). Such an improvement was expected given that Morpheus completes its searches on average 1.3 to 4.6 times faster than most other search engines available (Wenger and Coon, 2013). Given its simplicity of use, speed, and open source nature, MSpC combined with Morpheus is clearly advantageous over other PSM-based LFQ approaches currently available. Moreover, by being open source, MSpC should allow others to

extend its utility and to serve as a platform for integrating additional open source LFQ approaches into the Morpheus pipeline.

2.3. Methods

2.3.1. Sample Preparation

All chemicals were purchased from Sigma-Aldrich unless otherwise stated. 20S proteasomes were obtained as described previously (Book *et al.*, 2010) from *Arabidopsis thaliana* Col-0 ecotype seedlings in which the 20S proteasome subunit PAG1 ($\alpha 7$) was genetically replaced with a FLAG-tagged variant, with the minor modification of switching to a more stable HEPES buffer during purification. The FLAG peptide used for elution was removed by filtering through an Amicon Ultra 4 10K filter with the elution buffer also exchanged into 8 M urea. Total protein was quantified by the bicinchoninic acid protein assay (Thermo Scientific) using bovine serum albumin as the standard. Approximately 70 μ g of proteasomes were digested overnight at 37°C using a 1:30 trypsin/sample ratio. Peptides were acidified to a final concentration of 1% TFA, desalted on a Waters C18 Sep-Pak containing 50 mg sorbent material, and lyophilized. Total *E. coli* lysates were obtained from Bio-Rad (Cat. 163-2110) with 200 μ g digested as above. Both proteasome and *E. coli* peptides were dissolved in 5% acetonitrile, 95% water, and 0.1 % formic acid. Each MS analysis, performed




Figure 2.5: MSpC combined with Morpheus works faster than TPP combined with ABACUS. Speed comparisons were performed for 12, 24, 48, and 96 raw MS/MS files generated with the 20S proteasome/*E. coli* lysate samples analyzed in 2.4. On average, Morpheus/MSpC finished the calculations 1.9 times faster than TPP/ABACUS over a ~10-fold range of dataset size.

in triplicae, used 5 μ L volumes prepared with 3, 2, 1.5, 1, 0.5, 0.25, 0.1, or 0.05 μ g of digested proteasomes mixed with 0.5 μ g of digested *E. coli* proteins. This mixtures reflected proteasome/*E. coli* ratios of ~0.091, 0.167, 0.333, 0.500, 0.667, 0.750 0.800, and 0.857, respectively.

2.3.2. Liquid Chromatography and High-Resolution Mass Spectrometry

Samples were analyzed by ultra-high performance liquid chromatography (UPLC) (nanoAcuity, Waters Corporation) connected online to an electrospray ionization LTQ Velos Orbitrap mass spectrometer (Thermo Scientific). Separation employed a 100 x 365 μ m fused silica capillary micro-column packed with 20 cm of 1.7 μ m-diameter, 130- \AA pore size, C18 beads (Waters BEH), with an emitter tip pulled to approximately 1 μ m using a laser puller (Sutter Instruments). Peptides were loaded at a flow-rate of 400 nL/min for 30 min and then eluted over 120 min at a flow-rate of 300 nL/min with a 2% to 30% acetonitrile gradient in 0.1% formic acid. Full-mass scans were performed in the FT Orbitrap with a mass range of 300-1500 *m/z* at a resolution of 60,000, followed by ten MS/MS high energy C-trap dissociation scans of the ten highest intensity parent ions at 42% normalized collision energy and 7,500 resolution, with a mass range starting at 100 *m/z*. Dynamic exclusion was enabled with a repeat count of two over the duration of 30 s and an exclusion

window of 120 s.

2.3.3. Data Processing

Protein identifications were determined using the Morpheus search engine (Wenger and Coon, 2013). Raw data was searched with the Thermo module of Morpheus revision 151 downloaded and compiled from source code available at <http://sourceforge.net/projects/morpheus-ms/> using Microsoft Visual Studio 2013 professional edition. The following parameters were used to search all databases: unknown precursor charge states - +2, +3, +4; maximum number of MS/MS peaks = 400; assign charge states - enabled; de-isotope - disabled; generate target decoy database on the fly; protease trypsin (no proline rule); maximum missed cleavages = 2; initiator methionine behavior - variable; fixed modification of carbamidomethylation of cysteine; variable modification of oxidation of methionine; maximum variable modification isoforms per peptide = 1024; precursor mass tolerance = ± 2.1 Da monoisotopic (recommended parameters to account for neutral loss); precursor monoisotopic peak correction - disabled; product mass tolerance = ± 0.025 Da monoisotopic; consider modified forms as unique peptides - false; maximum threads = 8; minimize memory usage - false. MSpC quantification of Universal Proteome Standard 2 (UPS2) protein sequences exploited the MS/MS analysis of

individual USP2 proteins (Sigma-Aldrich) mixed at various concentrations with egg or embryo extracts from *Xenopus laevis* available in the PRoteomics IDEntifications (PRIDE) repository (Vizcaino *et al.*, 2013) using identifier - PXD000902 and the available proteomics database - pita_v1.71.protein.name.fa. For our analysis, both the raw MS/MS data and resulting FASTA files for the USP2 and egg and embryo proteomes were obtained from PRIDE. The database used for searching the MS/MS data of *Arabidopsis* proteasomes spiked into total *E. coli* peptides was generated by combining Uniprot K12 *E. coli* reference proteome UP000000625 with a common contaminant database, and then mixing the merged dataset with FASTA sequences for all proteoforms of all known proteasome subunits and associated proteins (Book *et al.*, 2010) obtained from the TAIR10_pep_20101214 FASTA database available within The *Arabidopsis* Information Resource (TAIR) version 10. All FASTA files are available for download in the Supporting Information. The datasets were analyzed by MSpC with a 1% PSM false discovery rate (FDR) and a 1% protein group FDR to determine NSAF, dNSAF, and uNSAF values for each protein group. Two separate analyses were performed in which one unique, or alternatively two unique peptides were required to quantify a protein group. NSAF values were calculated according to formulas 1a, 2a, and 3a from Figure 1 of Zhang *et al.* (Zhang *et al.*, 2010).

2.3.4. Isoform Incorporation Rates

For the individual subunit analysis of the 20S proteasome the isoform incorporation rates were treated as follows. Given that each proteasome subunit should be incorporated at equal stoichiometry within the 20S particle, we then tested whether the Morpheus/MSpC pipeline could calculate the relative abundance of each subunit and the distribution of isoforms. Here, we divided the dNSAF values for each subunit/isoform by the total number of dNSAF values for the entire complex across all eight total proteasome/*E. coli* lysate ratios tested. This averaged value provided a concentration-independent ratio for the incorporation of each subunit/isoform. We then normalized these values based on a 1/14 stoichiometry of each subunit within the complex to calculate the estimated occupancy versus the expected occupancy of each subunit.

2.3.5. Speed and Accuracy Comparisons of Morpheus/MSpC to the TPP/ABA-CUS Pipeline

The speed and accuracy of MSpC combined with Morpheus was compared to the next most comparable open source software suite for calculating NSAF values; i.e., TPP (Deutsch *et al.*, 2010) combined with ABACUS (Fermin *et al.*, 2011), using the proteasome spike-in experiment files as input. The .raw files were converted to

.mzML files by TPP Build 201411201551-6764 and then searched using the multi-threaded X!Tandem MS/MS search engine (Craig and Beavis, 2004) with the search parameters adjusted as close as possible to that used for the Morpheus searches. A decoy database was generated using the TPP tool DecoyFASTA for use with X!Tandem. Relevant X!Tandem parameters are listed here: parent monoisotopic mass error = \pm 2.1 Da, fragment mass error = \pm 0.025 Da, fixed modifications of carbamidomethylation (57.021464) on cysteine, and variable modification of oxidation (15.994915) on methionine, fully tryptic cleavages, missed cleavage sites = 2 maximum, no refinement and 8 threads. The configuration file used and the test datasets can be found in the Supporting Information. The data were analyzed in the TPP using Peptide Prophet (Keller *et al.*, 2002). Relevant settings are listed: minimum probability = 0.05; minimum peptide length = 7; accurate mass, and nonparametric decoy database to pin down false discovery rate; ignore +1 charged spectra; and run Protein Prophet (Nesvizhskii *et al.*, 2003) after Peptide Prophet. Once completed, all pepxml data from Peptide Prophet contained in a single folder was combined using the command line version of Protein Prophet from the TPP binaries with the following command: ProteinProphet.exe *.pep.xml interact-COMBINED.prot.xml. This post-analysis aggregation was required for running the spectral counting program ABACUS. Here, we note that there are no graphic user interfaces to perform this

post analysis aggregation, which makes this portion of the data analysis more difficult for those unfamiliar with setting up and running programs from the command line. The combined data was analyzed by ABACUS with the following parameters: best peptide probability = 0.99; minimum peptide probability = 0.99; experimental peptide probability = 0; and combined file probability = 0.99 to most accurately match a 1% FDR stringency settings in MSpC. dNSAF values were compared in Microsoft Excel using the CORREL function and squaring the result. Additional timing tests were performed with a subset of the calibration curve data (ratios 0.091, 0.167, 0.333, and 0.500 in triplicate corresponding to 12 .raw files) by increasing file input to 24 (2x), 48 (4x), and 96 (8x) .raw files to determine the time dependence on input size between both pipelines tested (Morpheus/MSpC versus TPP/ABACUS). The timing tests and all data analyses were performed on a computer running Windows 7 Ultimate, with 16 GB of random access memory, and an Intel Core i7-2700k with hyper-threading turned on for eight logical cores.

2.4. Discussion on Requiring Two Unique Peptides to Quantify a Protein

Occasionally, some researchers may want to use a more stringent criterion for quantification such as requiring a protein to have more than one unique identifying peptide. To see how this might affect our data analysis, we re-analyzed our results

shown in Figure 2.3, this time requiring two unique peptides to quantify an individual protein. The results point to a very small increase in linearity observed in the average plots; however, there is a slight decrease in linearity for the egg sample (0.886 to 0.865) and a larger decrease in linearity for the individual UPS2 protein plot for the embryo sample (0.827 to 0.723). The decrease in linearity in the embryo sample is due to the analysis removing a low abundance UPS2 protein (O00762ups) identified with only one unique peptide. While some have suggested that requiring more than one unique peptide to identify a protein is an ideal approach, requiring two peptides for identifications in database searches reduces the number of protein identifications in the target database more than those in the decoy database and results in increased false discovery rates (Gupta and Pevzner, 2009). While we recognize that researchers may want to implement more stringent requirements than what is typically used in database searching to quantify a set of proteins, there are two cases where requiring two unique peptides may not be ideal in a quantitative analysis. Firstly, low abundance proteins that have few PSMs might be identified by only a single peptide and thus be erroneously thrown out of the analysis. Secondly, there may be only one unique peptide that can differentiate between families of homologous proteins. In this second case, requiring two unique peptides would remove these homologous proteins from the MSpC analysis, even

if they had a large number of PSMs. Because of these reasons and because of the decreased linearity observed when requiring proteins to have two unique peptides (Figure 2.3) as compared to one unique peptide (Figure 2.2), we suggest caution in requiring more than one unique peptide per protein.

2.5. Tutorial

Morpheus Spectral Counter (MSpC) Tutorial

Requirements

- 64 bit Windows Installation: For Example Microsoft Windows 7 64 bit, Windows 8 and 8.1 64bit, or Windows 10 64bit
 - This is due to requirements from vendor libraries to process Thermo .raw files directly in this tutorial
 - Download and install proteowizard (<http://proteowizard.sourceforge.net/downloads.shtml>)
 - Installation of proteowizard correctly installs the latest Thermo Vendor .dll's to read .raw files directly
 - Download pre-compiled binaries of revision 151 of the Morpheus Mass Spectrometry Search Engine (Thermo Version)
(https://github.com/dcgemperline/Morpheus_SpC/releases/download/v1.0/Morpheus.Binaries.revision.151.zip), referred from here on out as just Morpheus
 - Download and install MSpC
(https://github.com/dcgemperline/Morpheus_SpC/releases/download/v1.0/MSpC_v1.0.zip)
 - MSpC depends on the latest .NET Runtime that will be installed with MSpC installer if it is not already installed
 - Download the data files from the PRIDE proteomics data repository using the following ID – PXD003002 Data Files (<http://www.ebi.ac.uk/pride/archive/>)
-

Data Analysis Starting from .raw files

1. Once all the requisite software is installed and the Data Files are downloaded you are ready to begin.
2. Unzip **Morpheus Binaries revision 151.zip** into your desired location and start **Morpheus (Thermo).exe** from the Morpheus (Thermo) folder.
3. Once Morpheus starts, verify at the top you are using **Morpheus (Thermo) revision 151**.
4. Add the following .raw files to Morpheus from the Raw Files folder contained in the Data Files folder (multiple files can be selected at once).
 - 04_28_15_A_rep1_c1.raw
 - 04_28_15_A_rep2_c1.raw

- 04_28_15_A_rep3_c1.raw
 - 04_28_15_B_rep1_c1.raw
 - 04_28_15_B_rep2_c1.raw
 - 04_28_15_B_rep3_c1.raw
 - 04_28_15_C_rep1_c1.raw
 - 04_28_15_C_rep2_c1.raw
 - 04_28_15_C_rep3_c1.raw
 - 04_28_15_D_rep1_c1.raw
 - 04_28_15_D_rep2_c1.raw
 - 04_28_15_D_rep3_c1.raw
5. Side Note on Fractionation: If you would like Morpheus, and thus MSpC to output summaries for a set of data by adding up all of the spectra identified in that dataset(as the case may be for fractionation data), place the files you would like to be summarized in seperate folders, such as SampleSet1, Sampleset2, and Morphues will generate protein_groups.tsv and PSMs.tsv specifically for each sampleset. Later in MSpC after selecting the summary.tsv file these will show up as SampleSet1*, SampleSet2*, etc.
 6. Browse for the fasta file contained in the Data Files Folder
uniprot_k12_e_coli_contams_plus_proteasome_and_interactors.fasta.
 7. Verify that Create Target-Decoy Database On The Fly is checked.
 8. Browse for an Output Folder. In this tutorial select browse, leave the default Desktop location highlighted and press Make a New Folder and rename it Morpheus Analysis. Press OK. You should now have a folder on the desktop called Morpheus Analysis, and Morpheus should say you are outputting data to the following output folder, where **NAME** is your username on your machine.
 - C:\Users\NAME\Desktop\Morpheus Analysis
 9. Set the Maximum threads to 2, for a dual core processor, 4 for a quad core processor, 8 for a quad core processor with hyperthreading, and 8 for an 8 core processor. These are reasonable values that will give you decent performance with Morpheus.
 10. Press Search, and the progress bar will indicate search progress for each raw file. On an Intel Core i7 2700K with Morpheus set to use 8 threads, this takes approximately 4.5 minutes.
 11. Close Morpheus if so desired.
 12. On completion of the search open MSpC.
 13. The indicators in Summary Directory Contains Valid Input should be red.
 14. Press the Select the Summary File button and navigate to the summary.tsv file contained in the Morpheus Analysis folder on the Desktop.
 - C:\Users\NAME\Desktop\Morpheus Analysis\summary.tsv
 15. The indicators in Summary Directory Contains Valid Input should now read green indicating that the Morpheus Analysis folder contains all the necessary files from Morpheus to run. (The files required to be in the same directory as summary.tsv are protein_groups.tsv and PSMs.tsv. This will automatically occur if you output all of your Morpheus output into a single output folder.)

16. (Optionally) Select the provided whitelist file contained in the Data Files folder that you downloaded to simplify the output.
 17. Select your desired PSM FDR and Protein FDR (although the default options of 1% are good default options).
 18. Check or uncheck boxes of the desired calculations (NSAF, dNSAF, uNSAF)
 19. (Optionally) Select an Output Directory, otherwise MSpC defaults to the current directory of summary.tsv as an output directory.
 - Let's press Select Output Directory then Make a New Folder here called MSpC Output, press OK.
 20. Press Summarize Data, and the progress bar will indicate the progress of MSpC. This should take ~ 30 seconds or less
 21. The MSpC Output folder will then contain NSAF summaries for each individual .raw file, as well as a summary for all .raw files analyzed. These files are in the tab delimited output format .tsv
-

Data Analysis Starting from previously searched Morpheus Data

1. It is strongly suggested, that you start with the Data Analysis from .raw files tutorial first, and then refer back to the tutorial for previously searched Morpheus data if necessary
2. Open MSpC
3. Select a Morpheus summary.tsv file in a directory that contains the entire output from Morpheus.
4. If all output is there, the indicators in Summary Directory Contains Valid Input should now read green indicating that the folder contains all the necessary files from Morpheus to run MSpC.
5. (Optionally) Select the provided whitelist file contained in the Data Files folder that you downloaded to simplify the output.
6. Select your desired PSM FDR and Protein FDR (although the default options of 1% are good default options).
7. Check or uncheck boxes of the desired calculations (NSAF, dNSAF, uNSAF)
8. (Optionally) Select an Output Directory, otherwise MSpC defaults to the current directory of summary.tsv as an output directory.
 - Let's press Select Output Directory then Make a New Folder here called MSpC Output, press OK.
9. Press Summarize Data, and the progress bar will indicate the progress of MSpC. This should take ~ 30 seconds or less
10. The MSpC Output folder will then contain NSAF summaries for each individual .raw file, as well as a summary for all .raw files analyzed. These files are in the tab delimited output format .tsv

2.6. Acknowledgements

D.C.G. was supported by a grant from the U.S. Department of Energy Office of Science; Office of Basic Energy Sciences; Chemical Sciences, Geosciences, and Biosciences Division (DE-FG02-88ER13968) and a graduate training fellowship from the NIH (5 T32 GM 7133-37). M.S. and L.M.S were supported by a grant from the National Institute of Health/National Institute of General Medical Sciences (1P50HG004942). The authors thank Erin Gemperline, Richard S. Marshall, and Josh Coon for critical reading of the manuscript, and additionally thank Derek Bailey for a critical code review.

2.7. Literature Cited

- Book, A.J., Gladman, N.P., Lee, S.S., Scalf, M., Smith, L.M., and Vierstra, R.D.** (2010). Affinity purification of the Arabidopsis 26S proteasome reveals a diverse array of plant proteolytic complexes. *J. Biol. Chem.* **285**: 25554–25569.
- Cox, J., Hein, M.Y., Luber, C.A., Paron, I., Nagaraj, N., and Mann, M.** (2014). Accurate proteome-wide label-free quantification by delayed normalization and maximal peptide ratio extraction, termed MaxLFQ. *Mol. Cell. Proteomics* **13**: 2513–2526.
- Craig, R. and Beavis, R.C.** (2004). TANDEM: matching proteins with tandem mass spectra. *Bioinformatics* **20**: 1466–1467.
- Deutsch, E.W., Mendoza, L., Shteynberg, D., Farrah, T., Lam, H., Tasman, N., Sun, Z., Nilsson, E., Pratt, B., Prazen, B., Eng, J.K., Martin, D.B., Nesvizhskii, A.I., and Aebersold, R.** (2010). A guided tour of the Trans-Proteomic Pipeline. *Proteomics* **10**: 1150–1159.

- Fermin, D., Basrur, V., Yocum, A.K., and Nesvizhskii, A.I.** (2011). ABACUS: a computational tool for extracting and pre-processing spectral count data for label-free quantitative proteomic analysis. *Proteomics* **11**: 1340–1345.
- Gerber, S.A., Rush, J., Stemman, O., Kirschner, M.W., and Gygi, S.P.** (2003). Absolute quantification of proteins and phosphoproteins from cell lysates by tandem MS. *Proc. Natl. Acad. Sci. USA* **100**: 6940–6945.
- Gupta, N. and Pevzner, P.A.** (2009). False discovery rates of protein identifications: a strike against the two-peptide rule. *J. Proteome Res.* **8**: 4173–81.
- Keller, A., Nesvizhskii, A.I., Kolker, E., and Aebersold, R.** (2002). Empirical statistical model to estimate the accuracy of peptide identifications made by MS/MS and database search. *Anal. Chem.* **74**: 5383–5392.
- Nesvizhskii, A.I., Keller, A., Kolker, E., and Aebersold, R.** (2003). A statistical model for identifying proteins by tandem mass spectrometry. *Anal. Chem.* **75**: 4646–4658.
- Ong, S.E., Blagoev, B., Kratchmarova, I., Kristensen, D.B., Steen, H., Pandey, A., and Mann, M.** (2002). Stable isotope labeling by amino acids in cell culture, SILAC, as a simple and accurate approach to expression proteomics. *Mol. Cell. Proteomics* **1**: 376–386.
- Ross, P.L., Huang, Y.N., Marchese, J.N., Williamson, B., Parker, K., Hattan, S., Khainovski, N., Pillai, S., Dey, S., Daniels, S., Purkayastha, S., Juhasz, P., Martin, S., Bartlet-Jones, M., He, F., Jacobson, A., and Pappin, D.J.** (2004). Multiplexed protein quantitation in *Saccharomyces cerevisiae* using amine-reactive isobaric tagging reagents. *Mol. Cell. Proteomics* **3**: 1154–1169.
- Russell, J.D., Scalf, M., Book, A.J., Ladror, D.T., Vierstra, R.D., Smith, L.M., and Coon, J.J.** (2013). Characterization and quantification of intact 26S proteasome proteins by real-time measurement of intrinsic fluorescence prior to top-down mass spectrometry. *PLoS One* **8**: e58157.
- Thompson, A., Schafer, J., Kuhn, K., Kienle, S., Schwarz, J., Schmidt, G., Neumann, T., Johnstone, R., Mohammed, A.K., and Hamon, C.** (2003). Tandem mass tags: a novel quantification strategy for comparative analysis of complex protein mixtures by MS/MS. *Anal. Chem.* **75**: 1895–1904.

- Tu, C., Li, J., Sheng, Q., Zhang, M., and Qu, J.** (2014). Systematic assessment of survey scan and MS2-based abundance strategies for label-free quantitative proteomics using high-resolution MS data. *J. Proteome Res.* **13**: 2069–2079.
- Vizcaino, J.A., Cote, R.G., Csordas, A., Dianes, J.A., Fabregat, A., Foster, J.M., Griss, J., Alpi, E., Birim, M., Contell, J., O'Kelly, G., Schoenegger, A., Ovelleiro, D., Perez-Riverol, Y., Reisinger, F., Rios, D., Wang, R., and Hermjakob, H.** (2013). The PRoteomics IDEntifications (PRIDE) database and associated tools: status in 2013. *Nucleic Acids Res.* **41**: D1063–9.
- Wenger, C.D. and Coon, J.J.** (2013). A proteomics search algorithm specifically designed for high-resolution tandem mass spectra. *J. Proteome Res.* **12**: 1377–1386.
- Wong, J.W. and Cagney, G.** (2010). An overview of label-free quantitation methods in proteomics by mass spectrometry. *Methods Mol. Biol.* **604**: 273–283.
- Yang, P., Fu, H., Walker, J., Papa, C.M., Smalle, J., Ju, Y.M., and Vierstra, R.D.** (2004). Purification of the *Arabidopsis* 26S proteasome: biochemical and molecular analyses revealed the presence of multiple isoforms. *J. Biol. Chem.* **279**: 6401–6413.
- Zhang, B., VerBerkmoes, N.C., Langston, M.A., Uberbacher, E., Hettich, R.L., and Samatova, N.F.** (2006). Detecting differential and correlated protein expression in label-free shotgun proteomics. *J. Proteome Res.* **5**: 2909–2918.
- Zhang, Y., Wen, Z., Washburn, M.P., and Florens, L.** (2010). Refinements to label free proteome quantitation: how to deal with peptides shared by multiple proteins. *Anal. Chem.* **82**: 2272–2281.
- Zybailov, B., Mosley, A.L., Sardiu, M.E., Coleman, M.K., Florens, L., and Washburn, M.P.** (2006). Statistical analysis of membrane proteome expression changes in *Saccharomyces cerevisiae*. *J. Proteome Res.* **5**: 2339–2347.

Chapter 3

PROTEOMIC CHARACTERIZATION OF AFFINITY-PURIFIED ARABIDOPSIS PROTEASOME PARTICLES IDENTIFIES SUB-COMPLEX SPECIFIC INTERACTORS

3.1. Statement of Attribution

Initial mutant characterization of *rpt4* mutants was performed by Drs. Kwang-Hee Lee and Richard S. Marshall, postdoctoral fellows in the Vierstra Lab. Additionally, Dr. Lee generated the 2XFLAG-RPT4b transgenic line in the *rpt4b-2* background. I created the 2XFLAG-RPT4a *rpt4a-1* transgenic lines. I developed the affinity purification via the RP with minor modifications from (Book *et al.*, 2010) by changing the primary buffer to HEPES. All mass-spectrometry sample preparation and data analysis in this chapter was completed by myself. The actual running of the mass spectrometer was either performed by Dr. Mark Scalf in Dr. Lloyd M. Smith's lab (Department of Chemistry, University of Wisconsin - Madison), or by Dr. Fionn McLoughlin in the Vierstra lab at Washington University. I cloned the CP interactors PAP1, PBAC1, PBAC2, PBAC3, and PBAC4, and performed yeast-two-hybrid (Y2H) analysis on these. While the streaking and yeast transformations were ac-

complished by myself, Dr. Marshall performed the final spotting of the yeast-two hybrid. All BiFC data was generated by myself. The HA-PAP1 transgenic line was created by Dr. Adam Book. I developed the affinity purification for HA-PAP1 and performed the immunoblot analyses. Native PAGE was performed by Dr. Marshall.

We intend to submit this chapter to J. Biol. Chem. for publication as a research article in the near future. A suggested list of authors is as follows: David Gemperline, Richard S. Marshall, Kwang-Hee Lee, Fionn McLoughlin, Mark Scalf, Lloyd M. Smith, and Richard D. Vierstra.

3.2. Abstract

The 26S proteasome is an ATP-dependent protease complex responsible for the selective removal of numerous eukaryotic proteins following their modification with polyubiquitin chains. It consists of two sub-complexes, a self-compartmentalized core protease (CP) that degrades polypeptides within a central chamber, and a regulatory particle (RP) that caps one or both ends of the CP and is responsible for recognizing and unfolding ubiquitylated substrates, and transporting them into the CP lumen. Previous mass spectrometric (MS) analyses of *Arabidopsis* 26S proteasomes affinity-purified via the CP subunit PAG1 ($\alpha 7$), detected a complex

assortment of particles that are assembled using paired isoforms for many of the 33 core subunits, as well as a small assortment of associated factors that might help in assembly, regulation, substrate specificity, and ubiquitin recycling. Here, we extended these studies by affinity purifying *Arabidopsis* 26S proteasomes with either tagged isoform of the RP subunit RPT4 (named RPT4a and RPT4b). Label-free quantitative (LFQ) MS analyses of the preparations detected both isoforms for many of the other CP and RP subunits, suggesting that plants incorporate subunit isoforms randomly into the holo-protease. We also identified an array of interacting factors, including likely orthologs of the yeast assembly chaperones Ump1 and Pba1-4 for the CP, and Nas2, Nas6, and Hsm3 for the RP. We additionally identified a novel plant-specific protein bound to the CP that interacts with the putative *Arabidopsis* Pba1 ortholog, and possibly contributes to CP assembly. By coupling native PAGE and MS analysis with proteasome preparations isolated from proteasome inhibitor-treated seedlings, we identified several assembly intermediates also harboring assembly chaperones and the proteasome regulators PA200 and ECM29. Taken together these data suggest that *Arabidopsis*, like yeast, dynamically employ a suite of assembly factors to construct the mature 26S particle.

3.3. Introduction

The selective removal of proteins that are non-functional or have key regulatory roles is essential to normal plant growth and development. A major pathway for this selective removal is the ubiquitin 26S proteasome system, in which target proteins are covalently modified with the 76-amino-acid polypeptide ubiquitin and subsequently degraded by the central enzymatic effector in this process, the 26S proteasome (reviewed in (Finley, 2009; Livneh *et al.*, 2016; Vierstra, 2009)). The 26S proteasome consists of two major sub-complexes, the core protease (CP), which is responsible for degrading the substrate, and the regulatory particle (RP), which caps the CP at one or both ends and is responsible for recognizing and unfolding ubiquitylated substrates. Compositionally, the CP is made up of 28 subunits configured into four stacked hetero-heptameric rings made up of α and β -subunits in an α 1-7/ β 1-7/ β 1-7/ α 1-7 arrangement. The N-termini of the β 1, β 2, and β 5 subunits are enclosed in a central chamber by the surrounding α -subunits, and together these β -subunits provide peptidylglutamyl-peptide-hydrolyzing activity (β 1), trypsin-like (β 2), and chymotrypsin-like (β 5) activities. The α -subunits possess N-terminal extensions that gate entry to the chamber, forming an axial pore that is only opened when substrates are actively processed by an associated RP, or when activated by alternative capping particles such as PA200 (Dange *et al.*, 2011;

Sadre-Bazzaz *et al.*, 2010).

The RP consists of two major sub-classes of proteins including six related AAA-ATPase (RPT) subunits (RPT1-6) and at least 15 non-AAA-ATPase (RPN) subunits (RPN1-3, 5-13, and SEM1) (Finley, 2009; Paraskevopoulos *et al.*, 2014; Russell *et al.*, 2013). The RPT subunits form a six-membered RPT ring responsible for unfolding the polypeptide substrate in an ATP dependent manner, and channeling the substrate through the axial pore into the central CP chamber. The RPN subunits have several differing functions including binding ubiquitylated substrates (RPN1, RPN10, RPN13, and SEM1 (Elsasser *et al.*, 2004; Paraskevopoulos *et al.*, 2014; Schreiner *et al.*, 2008; Shi *et al.*, 2016)), binding ubiquitin shuttle proteins such as DSK2 and RAD23 (Elsasser *et al.*, 2002; Farmer *et al.*, 2010; Fatimababy *et al.*, 2010; Lin *et al.*, 2011), and releasing ubiquitin bound to substrates through the deubiquitylating activity of RPN11 (Verma *et al.*, 2002; Yao and Cohen, 2002). RPN5-7 and RPN9 form a glove-like structure around the RPT ring providing structural support for adjacent subunits (Lander *et al.*, 2012; Lasker *et al.*, 2012; Unverdorben *et al.*, 2014), with RPN6 acting as a molecular clamp that holds the RP and CP together (Pathare *et al.*, 2012).

Over the last several years, analyses of the plant 26S proteasome have also revealed insights into its composition. Searches of the *Arabidopsis* genome showed

that plants typically encode two paralogs for most proteasome subunits (Fu *et al.*, 1998). Almost all of these paralogs were identified in purified preparations of the *Arabidopsis* proteasome that used either chromatographic strategies (Yang *et al.*, 2004), or an affinity purification method that exploited a FLAG tagged CP subunit PAG1 (Book *et al.*, 2010). Like in *Arabidopsis*, duplication of RPT genes have also occurred in monocots, with rice having duplicated genes for RPT1, 2, 4 and 5, suggesting that there may be some evolutionary benefit to having multiple copies of the RPT family (Shibahara *et al.*, 2004). Taken together these data suggest that plants have the capacity to generate a highly diverse set of proteasome isotypes by preferentially combining specific subsets of isoforms with others into the holo particle. Such a precedent for proteasome isotypes exist in mammals, which assemble the immunoproteasome and thymoproteasome isotypes by selectively incorporating alternative isoforms of the catalytic β -subunits (Murata *et al.*, 2007; Nandi *et al.*, 1996). Similarly, the mammalian testes proteasome isotype incorporates an alternative $\alpha 4$ isoform (Belote *et al.*, 1998; Uechi *et al.*, 2014). In plants, genetic evidence suggests that some subunit paralogs may have distinct functions. For instance, RPN1a and RPN1b play differing roles in embryogenesis (Brukhin *et al.*, 2005). Furthermore, overexpression of RPN5a induces an early senescent phenotype while overexpression of RPN5b does not (Book *et al.*, 2009). Despite these observations, it

remains unclear if plants deliberately assemble their proteasome subunit isoforms into compositionally and functionally distinct proteasome isotypes.

Outside of the core subunits, other proteins are known to associate with the proteasome transiently or sub-stoichiometrically. These proteasome-associated proteins, or PAPs, perform a variety of functions including, substrate processing, ubiquitin recycling, and proteasome assembly. DSK2 and RAD23 contribute to the shuttling of ubiquitylated substrates to the proteasome (Farmer *et al.*, 2010; Fatimababy *et al.*, 2010; Lin *et al.*, 2011), RPN13 helps bind ubiquitylated targets (Schreiner *et al.*, 2008), UCH37 and other de-ubiquitylating enzymes (DUBs) (Van-derLinden *et al.*, 2015) help process substrates bound to the RP, and a suite of assembly chaperones assist in the construction of both the CP and RP. While the route(s) for plant proteasome assembly are largely unknown, a possible model has been established in yeast and mammals (see Figure 1.7, and Figure 1.10). In both systems, the Pba3 chaperone (PAC3 in mammals) interacts with Pba4 (PAC4), forming a Pba(PAC)3/4 heterodimer aiding in early stages of CP assembly by binding the α 5 subunit, and likely adjacent α -subunits (Kunjappu and Hochstrasser, 2014; Yashiroda *et al.*, 2008). Similarly, Pba1 (PAC1) interacts with Pba2 (PAC2) to form a Pba(PAC)1/2 heterodimer that aide in assembly of the α -ring, and subsequently together with Ump1, helps to form 15S half-barrels consisting of α 1-7/ β 1-6 notably

lacking the $\beta 7$ subunit (Kunjappu and Hochstrasser, 2014; Marques *et al.*, 2007).

While the Pba(PAC)1/2 heterodimer act throughout assembly and can even bind mature CP (Stadtmueller *et al.*, 2012), the Pba(PAC)3/4 heterodimer leaves upon incorporation of the $\beta 3$ subunit prior to the formation of 15S half-barrels, thus only acting at early stages of assembly (Hirano *et al.*, 2008). Finally, CP maturation occurs concomitant with autolytic cleavage of β -subunit propeptide-appendages exposing proteolytic N-terminal threonines and subsequent degradation of Ump1, the first substrate of the CP (Ramos *et al.*, 1998). In addition to these dedicated CP assembly chaperones, in humans and yeast a different suite of other chaperones also guides assembly of the RP, including Nas2, Nas6, Hsm3, and Rpn14 (see Figure 1.10). These RP assembly chaperones typically bind to the C-terminal domains of specific RPT subunits forming modules (Nas2 with Rpt4 and 5; Nas6 with Rpt3; Hsm3 with Rpt1, Rpt2, and Rpn1; and Rpn14 with Rpt6) that, importantly, prevent the assembling RPT ring from forming a pre-mature CP-RP interface (Park *et al.*, 2010). In plants, MS/MS analyses of *Arabidopsis* proteasomes identified only one putative CP assembly chaperone, PBAC2 (the nomenclature used here is different from yeast (Pba) and mammals (PAC) as both are standard nomenclature for the $\beta 1$ and $\alpha 1$ subunits in *Arabidopsis*) (Book *et al.*, 2010), which is an ortholog to PAC2 based on PSI-BLAST sequence analyses (Kusmierczyk *et al.*, 2011; Le Tallec *et al.*,

2007). Together these data suggest that other plant CP assembly chaperones remain to be identified (Book *et al.*, 2010). Additionally, these proteomic analyses failed to identify obvious plant RP assembly orthologs, such as NAS2, NAS6 and HSM3, despite the identification of these orthologs in the *Arabidopsis* genome (Book *et al.*, 2010). Taken together, the current data suggest that our understanding of plant PAPs in general, and CP and RP assembly chaperones in particular, remains incomplete.

While tools exist in other organisms to affinity-purify the 26S proteasome via the RP, including Pro-A tagged variants of RPN11, and RPT1 in yeast (Leggett *et al.*, 2005, 2002), and RAD23-based strategies that bind RPN1 in mammals (Besche *et al.*, 2009), such systems were not available in plants until this work. We also reasoned that our existing CP-based affinity purification might miss some RP-specific PAPSs. To overcome this challenge and to explore the possibility of plants forming proteasome isotypes, we sought to develop an RP affinity-purification strategy using two FLAG epitope-tagged RP subunit isoforms, RPT4a and RPT4b. RPT4 was specifically chosen for affinity tagging as prior structural analyses indicated that its N-terminus was likely solvent exposed (PDB 4CR2) (Beck *et al.*, 2012). We also exploited the ATP dependence of the interaction between the CP and RP to enrich for both of these sub-complexes specifically (Book *et al.*, 2010; Liu *et al.*, 2006). These approaches

together allowed us to discover novel CP and RP interacting partners, including several putative assembly chaperones. Follow-up interaction studies imply a role for several of these associated proteins in CP assembly. Finally, proteomic analyses of native-gel separated complexes from tissues treated with the potent and specific proteasome inhibitor MG132 identified novel sub-complexes that contain distinct subgroups of these putative assembly chaperones, which presumably represent assembly intermediates. From these results, we propose a model that assigns several of these novel associated proteins with distinct phases of proteasome assembly.

3.4. Experimental Procedures

3.4.1. Transgenic Plants and Growth Conditions

The *rpt4a-1*, *rpt4a-3*, *rpt4a-4*, and *rpt4b-1*, *rpt4b-2*, and *rpt4b-3* T(transfer)-DNA insertion mutants in the *Arabidopsis thaliana* Columbia-0 ecotype (Col-0) (SALK_052372, SALK_128087C, SALK_135246, SALK_108556, SALK_108557, and SALK_101982C, respectively) were obtained from the *Arabidopsis* Biological Resource Center (The Ohio State University, Columbus, OH). All mutants were backcrossed three times to the Col-0 parent and then made homozygous by self-fertilization. The *PAG1::PAG1-FLAG pag1-1* germplasm was as previously described (Book *et al.*, 2010). The *rpt4a* and *rpt4b* alleles were tracked by genomic PCR using the T-DNA-specific left bor-

der primer (Lba1) in combination with gene-specific primers (see Table 3.S1 for all primers used in this study). The exact T-DNA insertion positions were determined by sequencing the T-DNA-specific PCR products using BigDye Terminator Sequencing v3.1 (University of Wisconsin - Madison Biotechnology Center). For RT-PCR, total RNA was extracted from 7-day old liquid grown seedlings of the indicated genotypes using the RNAeasy plant mini kit (Qiagen), and then converted to cDNA using oligo-(dT)₂₀ primers and the SuperScript III first-strand synthesis system (Invitrogen) both as according to manufactures instructions. RT-PCR of the converted cDNA was then performed with primer pairs (P1 - P9).

Transgenic plants expressing FLAG-tagged variants of *RPT4a* and *RPT4b* were constructed as follows. The genomic region encompassing the full coding sequence of *RPT4a* (*AT5G43010*), the 600bp sequence upstream of the ATG start codon, which included the 5'-untranslated region (UTR), and the 3'-UTR, was PCR-amplified from Col-0 genomic DNA using primer pair (P10 and P11) and cloned into the pDONR221 (Invitrogen) vector via the Gateway BP Clonase II reaction (Invitrogen). Codons for tandem FLAG (DYKDDDDK) tags were inserted using two rounds of site directed mutagenesis (Edelheit *et al.*, 2009) with primer pairs P12, P13 and P14, P15. The sequence-confirmed 2xFLAG-*RPT4a* clone was recombined into the plant transformation vector pMDC123 (which encodes resistance to the herbicide Basta

(glufosinate)) via the Gateway LR Clonase II reaction (Invitrogen). The genomic region including the full coding sequence of *RPT4b* (*AT1G45000*) (from start to stop codon, lacking the 3'UTR), plus the 2kb sequence upstream of the ATG start codon (including 5' UTR) was PCR-amplified from *A. thaliana* Col-0 genomic DNA using the primer pairs P16, P17 (containing the 2XFLAG tag, *NcoI*, and *SmaI* cleavage sites) and P18, P19 (containing a *PstI* and *NcoI* sites) for the coding sequence and upstream region, respectively. The amplified regions and the recipient pCAMBIA3301 vector were digested with *NcoI*, *SmaI*, and *PstI* (New England Biolabs (NEB)), and ligated together using T4 DNA ligase (NEB). The recombinant plasmid was introduced into the *Escherichia coli* TOP10 strain (Invitrogen), and confirmed as correct by sequencing.

The *PAP1* coding sequence was PCR-amplified from Col-0 cDNA using primer pair P20, P21, and cloned into the pDONR221 vector via the Gateway BP Clonase II reaction. The *PAP1* coding sequence was then recombined in-frame into the plant transformation vector pMDC99 which harbored a 3x-HA tag upstream of the recombination site (resulting in a 3x-HA tag, followed by a short linker RSSRGVHHMITT-TLYTKVEMK, followed by the initiator methionine for PAP1) driven by a *UBQ10* promoter. This was accompanied by a hygromycin B resistance gene. All constructs were transformed into plants by the *Agrobacterium tumefaciens*-mediated floral-dip

method with the GV3101 agrobacterium strain (Clough and Bent, 1998; Gelvin, 2003; Zhang *et al.*, 2006). The construction encoding *RPT4a::FLAG-RPT4a* and *RPT4b::FLAG-RPT4b* were transformed into plants harboring the *rpt4a-1* and *rpt4b-2* insertions, while the construction encoding *UBQ10::HA-PAP1* was transformed into *PAG1-FLAG pag1-1* plants. T1 plants were selected with the relevant plant resistance markers (Basta for both the *RPT4a* pCAMBIA3301 vector and the *RPT4b* pMDC123 vector, and hygromycin for the *PAP1* pMDC99 vector). T2, and T3 plants were generated by self-pollination. T3 progeny analysis of herbicide/antibiotic-resistant segregation was used to identify homozygous T2 plants. Expression of the FLAG-RPT4a/b transgenes was confirmed by SDS-PAGE followed by immunoblot analysis of total cell extracts with either the M2 anti-FLAG antibodies (Sigma Catalogue Number (Cat. No.) F1804) or anti-RPT4 antibodies generated by Kwang-Hee Lee (University of Wisconsin - Madison). Total cell extracts were prepared from plants were grown for 7-days on 0.8% agar plates containing standard plant growth medium (GM medium, containing MS salts, and Gamborg's B5 vitamins (Gamborg *et al.*, 1968; Julio and Jose, 2006; Murashige and Skoog, 1962)), and 2% sucrose). Expression of HA-PAP1 was confirmed by SDS-PAGE of total extracts (prepared as above) followed by immunoblot analysis with anti-HA antibodies (Sigma anti-HA antibody, Cat. No. H6908).

3.4.2. Sequence and Phylogenetic Analyses

Plant protein sequences related to *Arabidopsis* PBAC1-4 and PAP1 were obtained using the Basic Local Alignment Search Tool (BLAST) against plant genomes (*Zea mays*, *Oryza sativa*, *Medicago truncatula*, and *Physcomitrella patens*) from Phytozome v8 (Goodstein *et al.*, 2012). Sequences for Pba1-4 (*Saccharomyces cerevisiae*) and PAC1-4 (*Homo sapiens*) were obtained from literature searches. The plant PBAC nomenclature is used for consistency. Related animal, and yeast sequences were obtained using top hits in each respective proteome from iterative PSI-BLAST searches using the BLOSUM62 substitution matrix, and hits above the default expectation value (E-Value) threshold of 0.005 in each iteration (Altschul *et al.*, 1997). Sequence alignments and identity matrices were determined using the Clustal Omega, which selects the appropriate Gonnet 40, 80, 120, 160, 250 and 350 protein substitution matrices in its alignment algorithm (Gonnet *et al.*, 1992; Sievers and Higgins, 2014; Sievers *et al.*, 2011). Alignments were visualized with BOXSHADE (v3.2.3, <https://sourceforge.net/projects/boxshade/>). Phylogenetic trees were generated with MrBayes (v3.2.2) using 1 million generations and a relative burn-in of 25% (Ronquist *et al.*, 2012). The analysis was run until convergence, and the resulting consensus tree was visualized in FigTree (v1.4.2, <http://tree.bio.ed.ac.uk/software/figtree/>).

3.4.3. Proteasome Affinity Purifications using PAG1-FLAG and FLAG-RPT4a/b

26S proteasomes and various subcomplexes were affinity purified based on integrated FLAG epitopes as described previously with a minor modification in the primary buffering agent, which was switched from Tris to HEPES, as it has a more stable pK_a (Book *et al.*, 2010; Marshall *et al.*, 2017). Plant seedlings were grown for 10 days under constant 150 μmol/m²/s light in liquid GM medium containing 2% sucrose at 21-23°C with gentle shaking (90 rpm). For MG132 treatments the growth medium was replaced at day 9 with fresh medium containing 50 μM MG132 (Sell-eckChem) prepared from a 50 mM MG132 solution dissolved in DMSO, or fresh medium containing an equivalent volume of DMSO, and harvested after 16 hours. Frozen seedlings (2.5 - 5g) were pulverized by mortar and pestle at liquid nitrogen temperatures, extracted at 4°C with 1.25 mL/g of fresh weight of buffer A.1 (50 mM HEPES (pH7.5), 50 mM NaCl, 10 mM MgCl₂, and 10% (v/v) glycerol, 2 μM chymostatin, 2mM phenylmethane sulfonyl fluoride (PMSF), 5mM dithiothreitol (DTT), with or without 20 mM ATP), filtered through Miracloth (EMD Millipore), and clarified at 30,000 × g for 30 min. Clarified protein extracts were applied to 50 μL of M2 anti-flag affinity resin (Sigma, Cat. No. A2220) (pre-equilibrated by washing with 1 mL buffer A.1 three times), washed three times with 40 column volumes (2 mL) of buffer A.1. Proteasomes, or their sub-complexes were eluted

in approximately 250 μ L of 500 ng / μ L FLAG peptide (DYKDDDDK synthesized at the University of Wisconsin Biotech Center, Madison WI) in A.1 buffer without PMSF and chymostatin. Samples were stored at -80°C.

3.4.4. SDS-PAGE and Native-PAGE Analysis and Silver Staining

SDS-PAGE utilized a Tris-glycine buffering system and was prepared according to the protocol originally established by Laemmli (Laemmli, 1970) and were performed as described in (Marshall *et al.*, 2017). Gels were made using 40% (w/v) acrylamide/bis-acrylamide solution (29:1, Bio-Rad), SDS-PAGE resolving buffer (375 mM Tris-HCl (pH 8.8) and 0.125% SDS), SDS-PAGE stacking buffer (125 mM Tris-HCl (pH 6.8) and 0.125% SDS), 0.1% ammonium persulphate (APS) and polymerized with N,N,N',N'-tetramethylethylenediamine (TEMED) with a final acrylamide concentration of 11% and 3.5% for the resolving and stacking gels, respectively. The gels were cast using a Hoeffer Gel casting system into 18 x 16 cm glass plates using 0.75 mm spacers. Proteins were separated with a Hoeffer SE600 electrophoresis unit using Tris-glycine SDS running buffer (25 mM Tris (pH 8.3), 192 mM glycine, 0.1% SDS; Bio-Rad) at a constant current of 15 mA for 6 hours. Approximately 32 μ L out of 250 μ L of each affinity-purified proteasome preparation was loaded for SDS-PAGE with Laemmli sample buffer. Native-PAGE gels were

prepared according to Book *et al.* (Book *et al.*, 2010), and performed exactly as described in (Marshall *et al.*, 2017). A discontinuous Native-PAGE system was used. The lower gel contained 4.6% acrylamide (w/v), 0.12% Bis (w/v), 2.3% sucrose (w/v), Tris-Borate-EDTA (TBE) buffer (89 mM Tris-HCl (pH 8.3), 89 mM boric acid, 2 mM EDTA, Bio-Rad), 0.1% APS, 5 mM MgCl₂, 1 mM ATP. The upper gel contained 2.5% acrylamide (w/v), 0.62% Bis (w/v), 2.3% sucrose, TBE, 0.1% APS, 5 mM MgCl₂, 1mM ATP. Gels were cast and run as described above. Gels were run using a Hoeffer SE600 electrophoresis unit with TBE buffer supplemented with 0.5 mM ATP instead of Tris-glycine SDS running buffer. Approximately 32 µL of each proteasome affinity purification was supplemented with 5 µL of 0.03% (w/v) xylene cyanol to visualize loading. Gels were run at 4°C at 50V for 16 or 32 (extended Native-PAGE) hours. Silver-staining of all gels was accomplished using a MS-safe short silver nitrate staining protocol (Chevallet *et al.*, 2006).

3.4.5. MS/MS Sample Preparation

All chemicals for MS/MS analyses were obtained from Sigma-Aldrich, unless otherwise noted. All solvents used were obtained from Thermo-Fisher and were LC/MS grade or J.T Baker® grade unless otherwise noted. For in-solution digests, 200 µL of eluent from each proteasome affinity purification was dried down in a

Speed Vac centrifugal evaporator (Savant Instruments, model number SUC100H) until approximately 25 µL of volume remained, then resuspended in 100 µL of 8 M urea. The samples were reduced with 10 mM DTT (1 hour RT), alkylated with 40 mM iodoacetamide (1 hour RT in the dark), quenched with 40 mM DTT (10 min RT) and then digested with approximately 1 µg of trypsin overnight at 37°C. The samples were acidified with 10% trifluoroacetic acid (TFA) until the pH was 2, desalted on C18 Bond Elut OMIX Tips (Agilent) as according to manufacturer's instructions, and eluted in 75% acetonitrile (ACN), 0.1% acetic acid. The samples were then dried in a Speed Vac centrifugal evaporator and resuspended in 30 µL of 5% acetonitrile, 0.1% formic acid and stored at -80°C until analyzed by MS/MS.

For digestion of samples run on a native-PAGE gel, silver-stained protein bands were cut into 1-mm³ pieces with a clean razorblade, washed with MilliQ (EMD Millipore) water, and destained with a 1:1 solution of freshly prepared 100 mM sodium thiosulfate (Na₂S₂O₃) and 30 mM potassium ferricyanide (K₃Fe(CN)₆) until colorless. Silver-ions were then removed from the gel slices by washing twice with MilliQ water. The slices were dehydrated with a 50% ACN in 25 mM ammonium bicarbonate (NH₄HCO₃) solution, then dried in a Speed Vac. They were rehydrated with a reducing solution (25 mM DTT in 50 mM ammonium bicarbonate) for 20 minutes at 56°C, to cleave disulfide bonds. The gel pieces were then treated with

an alkylating solution (55 mM iodoacetamide in 50 mM ammonium bicarbonate) in the dark for 20 minutes at RT to alkylate free cysteines. The alkylating solution was then removed, the gel slices were washed with MilliQ water, dehydrated in 50% ACN in 25 mM ammonium bicarbonate solution, and then dried in a Speed Vac. The ProteaseMax (Promega) in-gel digest protocol was then used to aide in digestion and peptide extraction from gel slices (Saveliev *et al.*, 2013). Samples were then desalted with self-packed C18 stage-tips (Rappsilber *et al.*, 2003) using a spin protocol (Yu *et al.*, 2014), dried, and resuspended as described above.

3.4.6. MS/MS Analyses

In-solution-digested peptide mixtures were separated using a Dionex U3000 nano-flow ultra-high performance liquid chromatography system (nUHPLC, Thermo Scientific). The peptides were separated using an Acclaim® PepMap RSLC C18 column, 2 μm particle size, 100 \AA pore size, 75 $\mu\text{m} \times 15 \text{ cm}$ (Thermo Scientific) using a 90 minute linear gradient from 5% ACN to 33% ACN with 0.1% formic acid. Electro-spray ionization (ESI) of separated peptides was performed with a NanoSpray Flex ion source and ionized peptides were subsequently analyzed with a Thermo Q-Exactive Plus high-resolution, accurate-mass Orbitrap MS/MS. A data-dependent acquisition program was run with an MS1 mass range of 380-1500 m/z

at 70,000 mass resolution, with an automatic gain control (AGC) target of 1e6 ion intensity. The top 15 peaks were chosen for higher energy collisional dissociation (HCD) fragmentation at 28% normalized collision energy with a dynamic exclusion set to 10 s. Product MS2 scans were acquired with a 200-2000 *m/z* mass range at 17,500 mass resolution, with an AGC target of 8e3. All peaks were recorded in profile mode. All raw files associated with these experiments will be deposited in the PRoteomics IDEntification (PRIDE) database.

Peptide samples obtained from native-gel slices were instead analyzed using a nUHPLC (nanoAcuity, Waters Corporation) connected online to an electrospray ionization LTQ Velos Orbitrap mass spectrometer (Thermo Scientific). Separation of the peptide mixture employed a 100 x 365 μm fused silica capillary micro-column packed with 20 cm of 1.7 μm -diameter, 130- \AA pore size, C18 beads (Waters BEH), with an emitter tip pulled to approximately 1 μm using a laser puller (Sutter Instruments). Peptides were separated with a 2% to 30% acetonitrile gradient in 0.1% formic acid. Full-mass scans were performed in the FT Orbitrap with a mass range of 300-1500 *m/z* at a resolution of 60,000, followed by ten MS/MS high energy C-trap dissociation scans of the ten highest intensity parent ions at 42% normalized collision energy and 7,500 resolution, with a mass range starting at 100 *m/z*. Dynamic exclusion was enabled with a repeat count of two over the duration of 30 s and an

exclusion window of 120 s.

3.4.7. MS/MS Data Processing

For the proteasome associated protein studies, raw spectra files were processed with MaxQuant version 1.5.3.30 (Cox and Mann, 2008) using the Andromeda search engine against a TAIR10_pep_20101214_updated.fasta file obtained from The *Arabidopsis* Information Resource (TAIR) version 10 (Berardini *et al.*, 2015).

Andromeda searches were performed with MS1 mass tolerance of 20 ppm and an MS2 mass tolerance of 10 ppm. Up to two missed cleavages were allowed, along with the fixed modifications of carbamidomethyl on cysteine, and variable modifications including oxidation of methionine, acetylation of protein N-termini, and GlyGly (ubiquitylation remnant after cleavage with trypsin) on lysines for possible ubiquitylation events, were specified. Peptides and protein groups were identified with a 0.01 false discovery rate (FDR).

Label-free-quantitative analyses were performed using MaxQuant's MaxLFQ (Cox *et al.*, 2014) algorithm (generating the MaxLFQ values output) with default settings except that only unique peptides were used for quantification so that protein isoforms could be discriminated. Matching was set to "match from and to samples" to increase the number of quantifiable proteins per run, using the

'MaxQuant's match between runs' feature. MaxQuant's proteinGroups.txt file was then processed in Perseus (Tyanova *et al.*, 2016), a software suite for analyzing mass spectrometry data. Ten proteins were marked as likely contaminants including nitrilases, cruciferins, and seed storage albumins (full list available in Table 3.S2). MaxLFQ values were averaged for both technical replicates, and missing MaxLFQ quantification values were imputed with a 1.8-fold downshifted normal distribution with a width of 0.3. Volcano plots were generated comparing each proteasome affinity purification against its respective wild-type control using Perseus's volcano plot function using a two-sided t-test, 0.01 FDR, S0 value of 2, and 250 permutations. The resulting volcano plots were plotted using the Seaborn python graphing library (Tyanova *et al.*, 2016; Waskom, 2016), and edited in Adobe Illustrator. Proteins that were called by Perseus as statistically significant were included in Tables 3.S5 -3.S10. KDE estimations for CP and RP subunits were performed using the Seaborn graphing library, with the median test and Kolmogorov-Smirnov (KS) tests for enrichment performed using scipy.stats. For the bar graphs in Figure 3.3, missing values were imputed with a $\text{Log}_2(\text{MaxLFQ value})$ of 15 so that a "zero value" could be more easily viewed in log scale.

For the MS/MS analyses of native gel slices the data were searched with the Morpheus search engine (Wenger and Coon, 2013) against same protein database

used above. The recommended settings of MS1 search tolerance of 2.1 Da, MS2 search tolerance of 0.01 Da, and 1% maximum FDR were used. Fixed modifications of carbamidomethyl on cysteine, and variable modifications of oxidation on methionine, and GlyGly on lysine with up to two missed cleavages were used. LFQ was completed with Morpheus Spectral Counter (Gemperline *et al.*, 2016) to calculate dNSAF values as described in Chapter 2. The data were plotted in R using the heatmap.2 function, with centroided hierarchical clustering based on Pearson's correlation as the distance function.

3.4.8. Yeast-two hybrid, bimolecular fluorescent complementation, and HA-PAP1 immunopurification

Assays for protein-protein interactions between PBAC1-4 and PAP1 were performed using the ProQuest® two-hybrid system which uses the pDEST22 and pDEST32 vectors, and the *Saccharomyces cerevisiae* MaV203 strain (Vidal *et al.*, 1996) all available from Thermo Fisher Scientific. PBAC1-4 and PAP1 were amplified from *Arabidopsis* cDNA with primers (P20 -P29) and transformed into the pDONR221 entry vector (Thermo Fisher Scientific). Genes were then recombined using Gateway LR Clonase II (Thermo Fisher Scientific) into vectors pDEST22 and pDEST32. Yeast strain MaV203 was transformed with pairwise gene combinations in pDEST22

and pDEST32, or the empty vector controls. Interactions were tested by growing transformants for 2 days at 30°C on synthetic complete medium lacking leucine, tryptophan, and histidine, with added 30mM 3-amino-1,2,4-triazole.

For BiFC assays, PBAC1-3 and PAP1 cDNA sequences in pDONR221 were recombined using Gateway LR Clonase II (Thermo Fisher Scientific) into either pSITE-N-EYFP-C1 or pSITE-C-EYFP-C1 vectors (Martin *et al.*, 2009) (ABRC stocks CD3-1648 for NYFP or CD3-1649 for cYFP respectively), and subsequently transformed into *Agrobacterium tumefaciens* strain GV3101 (Gelvin, 2003). Overnight cultures were diluted in resuspension buffer (10 mM MgCl₂, 10 mM MES (pH 5.7)) to a final OD₆₀₀ of 0.5 and then infiltrated by syringe into 4-6 week old *Nicotiana benthamiana* leaves. After 36 h, fluorescence in the infiltrated regions was visualized using a Zeiss 510 Meta confocal laser scanning microscope, and images were analyzed with Zeiss LSM image browser v3.5 and ImageJ.

The immunopurification of PAP1 to determine if PAP1 could co-purify with 26S proteasome subunits was performed as follows. HA-PAP1-expressing tissue in the *PAG1-FLAG pag1-1* background were grown in liquid culture as described previously (Book *et al.*, 2010). Approximately 2 g of tissue were ground using liquid nitrogen chilled mortar and pestles, and soluble proteins were extracted in 2.5 mL of buffer A.1. 25 µL of HA-EZview (Sigma-Aldrich) beads were incubated overnight

with 1.75 mL of protein extracts in 2 mL Eppendorf tubes at 4°C with constant rotation. The HA beads were washed with 3 x 1mL of buffer A.1, and then eluted with 100µL of 2x protein sample buffer ((Laemmli, 1970)).

3.5. Results

3.5.1. *Arabidopsis* Proteasomes Can Be Effectively Affinity-Purified Using Epitope-Tagged Versions of Either Isoform of the Regulatory Particle Subunit RPT4

As a first step toward purifying *Arabidopsis* 26S proteasome via the RP, and exploring the possibility that plants assemble unique proteasome isotypes, we sought out candidate subunits amenable to epitope tagging for which useful mutants were available for genetic replacement. One of the best candidates was *RPT4*, which is encoded by two isoforms (*RPT4a* and *RPT4b*) with 96% amino-acid sequence identity. This level of identity is near average for shared sequence identity among RPT subunits (Book *et al.*, 2010); RPT2a and RPT2b share the most identity at 99%, and RPT5a and RPT5b share the least identity at 93%. Based on the structure of the yeast 26S proteasome (PDB ID: 4CR2) (Beck *et al.*, 2012), we expected that the N-terminus of RPT4 would be solvent accessible, thus making it an attractive candidate for attaching an tag that would be exposed and minimally disrupt particle

assembly. Furthermore, we identified several T-DNA insertion mutations within *RPT4a* and *RPT4b* in the Col-0 background that might generate null alleles suitable for genetic replacement (Figure 3.1 A). DNA sequence analysis of the *rpt4a/b* mutants determined the insertion sites to be at nucleotides 681, -278, and -24 for *rpt4a-1*, *rpt4a-3* and *rpt4a-4* and 151, 409, and 558 for *rpt4b-1*, *rpt4b-2*, and *rpt4b-3*, from the ATG start codon respectively. The *rpt4a-1* and *rpt4b-2* alleles were particularly notable because the T-DNA insertions sites disrupted the coding regions at 88 and 101 residues upstream of the AAA-ATPase cassette (Figure 3.S1) for each polypeptide which would likely block accumulation of the full length transcript (Figure 3.1 B). This possibility was confirmed by RT-PCR analysis of RNA extracted from seedlings homozygous for either *rpt4a-1* or *rpt4b-2* showing that the plants failed to accumulate the corresponding full-length transcripts (Figure 3.1 B, Primers P1 + P5, and Primers P6 + P9 respectively). However, low levels of the 5' region was amplified using primers that were 5' to the insertion site in both *rpt4a-1* and *rpt4b-2* plants (Primers P1 + P2, and Primers P6 + P7) suggesting that a transcript for the 5' region still accumulates. Additionally, a small amount of product relative to wild type was also detected 3' to the insertion site for *rpt4a-1* suggesting some mRNA from this region can also accumulate. At the protein level, an immunoblot analysis of SDS-PAGE-separated proteins from *rpt4a-1*, *rpt4b-2* and wild type Col-0




Figure 3.1: *rpt4a-1* and *rpt4b-2* mutants rescued with flag tagged variants. (A) Organization of the *Arabidopsis* *RPT4a* and *RPT4b* genes with insertion points shown for T-DNA alleles of *RPT4a* and *RPT4b*. **(B)** RT-PCR analysis of the *RPT4a* and *RPT4b* transcripts in wild type and *rpt4a* and *rpt4b* mutant seedlings showing aberrant *RPT4a* transcripts in *rpt4a-1* and abnormal *RPT4b* mRNAs in *rpt4b-2* plants. **(C)** Homozygous *rpt4* mutants show a phenotype similar to wild-type control. The top panel shows plants grown for 10-days in a 16-hour photoperiod, while the lower panel shows plants grown for 42 days in an 8-hour photoperiod. **(D)** Replacing the wild-type *RPT4a* and *RPT4b* with their respective flag-tagged variants show plants with normal growth when grown for 7 days on GM plates and then grown in soil for two weeks. **(E)** Immunoblots using anti-RPT4 and anti-FLAG antibodies confirm the presence of the tag and an increased size due to the FLAG tag, which is reactive to the anti-FLAG antibody. A loading control for histone-3 is shown.

seedlings showed reduction in RPT4 protein in *rpt4a-1* compared to wild type, but little to no change in *rpt4b-2* (Figure 3.1 E). While RPT4a is the dominant isoform with 169 expressed sequence tags as compared to 39 for RPT4b (Berardini *et al.*, 2015), our data detected a stronger reduction in RPT4a levels, as compared to RPT4b. There were no differences in RPT4 immunoreactivity between RPT4a and RPT4b (communications with Kwang-Hee Lee), therefore a more complex regulation of global RPT4a/b expression than anticipated may occur. The mutants were phenotypically indistinguishable to wild-type when grown on agar plates for 10 days in a long day (LD) photoperiod (16 h light / 8 h dark) or when transferred to unfertilized soil and grown until 42 days old in a short-day photoperiod (8 h light / 16 h dark) (Figure 3.1 C). Taken together these data suggest that the two RPT4 isoforms are not essential by themselves.

Plants containing the mutant alleles *rpt4a-1* and *rpt4b-2* were transformed with their respective 2XFLAG-tagged genomic variants and selected for stable transgene insertion. Each construct included codons for a 2XFLAG tag and the genomic region including introns that bracketed the *RPT4a/b* coding regions with expression drive by the promoter and 5' UTR regions of each locus (see Experimental Procedures). The initial transformants were self-fertilized and the homozygous lines for both the *rpt4* T-DNA insertion and the transgene were identified in the Basta

resistant progeny by genomic PCR. Immunoblot analysis of proteins extracted from *RPT4a/b::FLAG-RPT4a/b*-transformed plants detected accumulation of the FLAG-RPT4 proteins, which were larger than the corresponding native RPT4 polypeptides when detected with anti-RTP4 antibodies, and could be detected with anti-FLAG antibodies (Figure 3.1 E). The lower bands in the anti RPT4 immunoblot were likely RPT4b in the case of the *rpt4a-1* mutant, and RPT4a in the case of the *rpt4b-2* mutant. The transformed plants were phenotypically indistinguishable from wild type, implying that ectopic expression of either isoform bearing the 2X-FLAG tag was not detrimental to growth and development (Figure 3.1 D). Taken together these data showed that transgenic plants expressing their respective FLAG-tagged RPT4 variants displayed no gross morphological defects.

Using the strategy first described by Book *et al.* (Book *et al.*, 2010), we exploited the homozygous *FLAG-RPT4a/b rpt4a/b* seedlings to affinity purify 26S proteasomes via the RP. Briefly, frozen seedlings were homogenized in a tissue extraction buffer (see Experimental Procedures), clarified by centrifugation, and subjected to anti-FLAG chromatography, using FLAG peptide for gentle elution. To enrich for the 26S particle, 20 mM ATP was included in the indicated buffers to stabilize the CP/RP association (Book *et al.*, 2010; Liu *et al.*, 2006). To enrich for just the RP (using FLAG-RPT4a/b) or the CP (using PAG1-FLAG) the same protocol was used

except that ATP was omitted from the buffers. As a control, we also attempted to purify 26S proteasomes from wild-type tissues. Under conditions where 20 mM ATP was included in the purification buffer (indicated with a + in Figure 3.2 A), the FLAG-RPT4a/b affinity purified samples showed a comparable protein banding pattern after SDS-PAGE as compared to the PAG1-FLAG affinity-purified samples suggesting a similar complement of proteins were present (Figure 3.2 A). The only notable exception was the observation of a band that likely represented the FLAG-tagged PAG1 subunit (indicated with a black arrow) in the PAG1-FLAG samples (Book *et al.*, 2010). The size-shifted RPT4 subunits seemed to be masked by other subunits and were not readily identifiable via silver staining alone. Importantly, the mock affinity-purified samples from wild-type tissue showed very few contaminating proteins as judged by protein staining with silver, which was important for down-stream tandem mass-spectrometry (MS/MS) analyses. Overall, these affinity purification strategies based on either RPT4a or RPT4b successfully purified 26S proteasomes from whole *Arabidopsis* seedlings. When ATP was omitted from the purification buffer, there was clear enrichment for the CP in PAG1-FLAG samples, and clear enrichment for the RP in both the FLAG-RPT4a and FLAG-RPT4b samples, as judged by the size and intensity of the protein bands in Figure 3.2 A. Immunoblot analysis of these samples showed clear enrichment for two CP subunits PAG1 and




Figure 3.2: Protein composition of affinity-purified proteasomes from FLAG-RPT4a-, FLAG-RPT4b-, and PAG1-FLAG-expressing plants and from mock wild-type controls in the presence (+) or absence (-) of 20mM ATP. (A) Silver stained gel of each preparation shows clear enrichment over the negative control (Col-0) with minimal contaminants, both in the presence and absence of ATP during purification. Omitting ATP from the preparations for the CP-based PAG1-FLAG pulldown shows a decreased amount of RP subunits, while both RP-based subunit pulldowns show a decrease in the amount of CP subunits. (B) Immunoblots for various proteasome subunits for both the RP and the CP show a clear enrichment of proteasome subunits as compared to the wild type mock affinity purification control. As shown in the silver stain, omitting ATP from the preparations in each affinity purification causes a decrease in association of the alternative sub-particle (decrease in RP for CP pulldowns, and decrease in CP for RP pulldowns).

PBA1 in the CP-based samples affinity-purified in the absence of ATP, as compared to the RP subunits RPT4, RPN5, and RPN12 (Figure 3.2 B). There was also clear enrichment for the three RP subunits in the RP-based affinity purified samples when ATP was omitted from the purification as compared to the CP subunits that were immunoblotted. Taken together, these data show that 26S proteasomes can be easily purified based on a RP affinity tag, and that both isoforms for a single subunit will work. By omitting ATP from the buffer, we were able to enrich for subunits of the CP or the RP utilizing affinity purifications from PAG1-FLAG and FLAG-RPT4a/b expressing tissues, respectively.

3.5.2. MS/MS Analyses of RPT4a/b Affinity-Purified Proteins Identifies a Similar Complement of Proteins as PAG1-FLAG Affinity Purifications

Previous MS/MS analyses of affinity preparations based on PAG1-FLAG identified the fundamental subunits that make up the 26S proteasome, including the CP α -subunits (PAA-PAG), the CP β -subunits (PBA-PBG), the RP AAA-ATPases (RPT) (RPT1-6), and the RP non-ATPase subunits (RPN) (RPN1-3 and RPN5-12). Most subunits are encoded as a paralogous gene pair (named 1 and 2 for CP subunits, and a and b for RP subunits) and most isoforms were found to be incorporated in the complex (Book *et al.*, 2010). We wanted to verify that our affinity purifications

based on RPT4a and RPT4b also contained a similar set of proteins as in Book *et al.* (Book *et al.*, 2010). To do this we used label-free quantitative tandem mass spectrometry (LFQ-MS/MS) to analyze samples obtained from affinity purifications for FLAG-RPT4a, FLAG-RPT4b, PAG1-FLAG, and a mock wild-type control, in the presence and absence of ATP resulting in MaxLFQ values (Cox *et al.*, 2014) for each protein analyzed. Importantly, we identified all subunit isoforms previously identified by Book *et al.* (Book *et al.*, 2010) showing that our purifications were comparable (Figure 3.3, CP α -ring, CP β -ring, RPT, and RPN plots). Additionally, we identified a suite of key accessory proteins found to be statistically enriched as compared to control at a false discovery rate (FDR) of 0.01 for both the CP and RP (Figure 3.3 Associated Proteins).

The CP α - and β -subunits show higher MaxLFQ values in the PAG1-FLAG samples as compared to either FLAG-RPT4a/b sample under both plus and minus ATP conditions (Figure 3.3, CP α -ring, CP β -ring). This suggests some free CP was purified even in the presence of ATP, and that the complexes purified were not exclusively RP-CP proteasomes. In the PAG1-FLAG samples, the CP α and β -subunits showed higher MaxLFQ values when ATP was omitted from the affinity purification buffer as compared to when ATP was present (Figure 3.3, CP CP α -ring, CP β -ring). This increase is consistent with the preferential purification of the CP




Figure 3.3: Label-free quantitative MS/MS analysis of affinity-purified proteins in the presence (+) or absence (-) of ATP. (CP Alpha Ring) α -subunits are enriched in samples that were affinity purified via the CP subunit PAG1-FLAG. This enrichment increases even further when ATP is omitted from the preparations as shown on the left side of the graph. (CP Beta Ring) β -subunits are enriched at a higher level when purified via the CP subunit PAG1-FLAG. This enrichment increases even further when ATP is omitted from the preparations as shown on the left side of the graph.




Figure 3.3: Label-free quantitative MS/MS analysis of affinity-purified proteins in the presence (+) or absence (-) of ATP. (RPT Subunits) RPT subunits are enriched at a higher level when purified via RP subunits RPT4a or RPT4b. There are clear increases in the level of enrichment of the RPT4a subunit for the RPT4a affinity purifications, and clear increases in enrichment of the RPT4b subunit for the RPT4b affinity purifications. These respective increases are greater than the isoform unbiased PAG1 pulldowns. The respective enrichments of each isoform also increases when ATP is omitted, due to a decrease in the possibility of binding the alternative subunit isoform under ATP limiting conditions. **(RPN Subunits)** RPN subunits are enriched at a higher level when purified via RP subunits RPT4a and RPT4b. This enrichment increases even further when ATP is omitted from the preparations as shown on the left side of the graph.




Figure 3.3: Label-free quantitative MS/MS analysis of affinity-purified proteins in the presence (+) or absence (-) of ATP. (Associated Proteins) Key proteins identified as proteasome interacting proteins are specifically enriched in pulldowns based on the CP subunit PAG1, and others are specifically enriched for pulldowns based on either RP subunits RPT4a or RPT4b. Please refer to Figure 3.4 for statistical analyses of CP and RP distributions in affinity-purified samples, and for statistical analyses that identify candidates as associated proteins.

over the RP in the absence of ATP when using the PAG1-FLAG epitope tag.

Likewise, both affinity purifications that target the RP through FLAG-RPT4a/b, had higher MaxLFQ values for RP subunits as compared to the CP-based PAG1-FLAG samples under both plus and minus ATP conditions. When ATP was omitted from the purification, the FLAG-RPT4a/b samples had increased MaxLFQ values for RP subunits as compared to when ATP was included; however, this increase was considerably less than that observed for the CP subunits in the PAG1-FLAG samples (Figure 3.3, RPT, and RPN plots). FLAG-RPT4a samples had higher MaxLFQ values for RPT4a when compared to FLAG-RPT4b and PAG1-FLAG (Figure 3.3, RPT plot). Conversely, FLAG-RPT4b samples had higher MaxLFQ values for RPT4b when compared to FLAG-RPT4a and PAG1-FLAG (Figure 3.3, RPT plot). These data suggest that, as expected, RPT4a was enriched in the FLAG-RPT4a samples, and that RPT4b was enriched in the FLAG-RPT4b samples.

3.5.3. Statistical Analysis of Affinity Enrichments for FLAG-RPT4a Show Little to No Difference When Compared to FLAG-RPT4b

While mammals and yeast can form only a few types of proteasomes, *Arabidopsis*, with its many subunit paralogs, can potentially form a much wider array of proteasome isotypes. By comparing samples enriched for RPT4a and RPT4b us-

ing a Significant Analysis of Microarray (SAM) test at 0.01 FDR, we determined if any additional subunit isoforms (or other associated proteins) are incorporated preferentially, which would suggest that *Arabidopsis* can form proteasome isotypes. Surprisingly, this statistical analysis revealed that only when ATP was omitted from the purifications were the samples statistically enriched for either RPT4a or RPT4b, respectively (Figure 3.S2 B). While RPT4 and RPT4b were enriched in the samples in which ATP was included in the purification, they were not statistically significant by the SAM test at 0.01 FDR (Figure 3.S2 A). Additionally, the 26S proteasome subunits have larger MaxLFQ values in the FLAG-RPT4a sample suggesting a slightly more efficient purification for the proteasome with the RPT4a isoform. Furthermore, no significant enrichment of any other subunit isoforms was observed. As such these data suggest that there are no major differences in protein composition between affinity purifications for FLAG-RPT4a or FLAG-RPT4b other than statistically enriching for each individual isoform when ATP was omitted from the purification. This indicates that plants may assemble their proteasome randomly with respect to isoform incorporation under the conditions tested.

3.5.4. Statistical Analyses of LFQ-MS/MS Data Show Affinity Preparations Targeting Either the CP or RP Specifically Enrich for Their Respective Sub-complexes

The LFQ-MS/MS data allowed us to perform a more formal statistical test for the enrichment of the RP over the CP in samples affinity purified from the RP subunits FLAG-RPT4a/b, and for the enrichment of the CP over the RP in samples affinity purified from CP subunit PAG1-FLAG. We estimated the distribution of RP and CP subunits in these samples using kernel density estimations (KDE) of MaxLFQ values as compared to our wild-type mock-affinity purifications as a negative control (Figure 3.4 below each volcano plot, and larger plots in Figure 3.S3). As the KDE distributions were not normal, two non-parametric statistical tests, Kolmogorov-Smirnov (KS) and median test, were used to determine if there was specific enrichment of particular proteasome sub-complexes (RP vs CP) from the different affinity purifications. The KS test determines if the two distributions are different in shape or position, and the median test determines if the distribution's medians are statistically different.

The KDE distribution for the CP as compared to the RP from samples affinity purified in the absence of ATP using PAG1-FLAG showed both the KS and the median *p*-values being much less than 0.01, at 1.74E-10 and 4.3E-10, respectively




Figure 3.4: Volcano Plots comparing proteasome affinity purifications under both plus and minus ATP conditions relative to WT controls, allow identification of statistically significant interacting partners for the proteasome. Volcano plots comparing affinity purifications for PAG1-FLAG (A and B), FLAG-RPT4a (C and D), and FLAG-RPT4b (E and F) against their respective WT mock affinity purification control are shown. The top panels (A, C, and E) show comparisons for samples purified in the absence of ATP, while the lower panels (B, D, and F) show comparisons for samples purified in the presence of ATP. Two-dimensional Kernel Density Estimations (KDE) are shown for CP subunits in red and RP subunits in blue. Larger KDE plots are available in Figure 3.S3. CP subunits are labeled as red squares, while RP subunits are labeled as blue squares. Proteins outside the dashed lines are statistically significantly differentially represented in purified samples relative to control as judged by a significant analysis of microarray or (SAM) test at 0.01 false discovery rate. Proteins likely involved in CP assembly are shown as green squares, while proteins likely involved in RP assembly are shown as yellow squares. Known proteasome associated proteins or PAPs are labeled as black squares, while unknown PAPs are labeled in grey squares. Proteins identified that did not reach SAM significance are labeled as grey circles.




Figure 3.4: Volcano Plots comparing proteasome affinity purifications under both plus and minus ATP conditions relative to WT controls, allowing identification of statistically significant interacting partners for the proteasome. Volcano plots comparing affinity purifications for PAG1-FLAG (A and B), FLAG-RPT4a (C and D), and FLAG-RPT4b (E and F) against their respective WT mock affinity purification control are shown. The top panels (A, C, and E) show comparisons for samples purified in the absence of ATP, while the lower panels (B, D, and F) show comparisons for samples purified in the presence of ATP. Two-dimensional Kernel Density Estimations (KDE) are shown for CP subunits in red and RP subunits in blue. Larger KDE plots are available in Figure 3.S3. CP subunits are labeled as red squares, while RP subunits are labeled as blue squares. Proteins outside the dashed lines are statistically significantly differentially represented in purified samples relative to control as judged by a significant analysis of microarray or (SAM) test at 0.01 false discovery rate. Proteins likely involved in CP assembly are shown as green squares, while proteins likely involved in RP assembly are shown as yellow squares. Known proteasome associated proteins or PAPs are labeled as black squares, while unknown PAPs are labeled in grey squares. Proteins identified that did not reach SAM significance are labeled as grey circles.

(Figure 3.4 and 3.S3 A). This suggests that the distributions are different, and that their medians were different with a greater than 20-fold increase in CP over RP. Conversely the KDE distributions for the samples affinity purified in the presence of ATP using PAG1-FLAG (Figure 3.4 and 3.S3 B) overlap to a higher degree, with both the KS and Median *p*-values > 0.01 at 0.401 and 0.256, respectively. This indicates that in samples affinity purified in the presence of ATP that the RP and CP distributions were not significantly different, and that their means were similar. Although not statistically significant, a 1.5 median fold-change was observed for the CP over the RP in samples affinity purified in the presence of ATP using PAG1-FLAG suggesting that there may be some CP enriched as compared to the RP. These statistical analyses showed that, as expected, we were able to successfully enrich for the CP over the RP when affinity purifying from PAG1-FLAG expressing tissue.

Conversely, in samples using the RP-based FLAG-RPT4a/b purified in the absence of ATP, the KDE distributions showed enrichment for the RP over the CP, with a fold change of 3.43 and 3.41 for the RPT4a and RPT4b subunits respectively (Figures 3.4 and Figures 3.S3 C and E, respectively). Both the KS (3.303E-5 for RPT4a and 5.55E-5 for RPT4b) and median *p*-values (7.11E5 for RPT4a and 7.11E-5 for RPT4b) were < 0.01, suggesting that the two distributions were statistically different and had different medians. Again, contrasting this with the samples

purified in the presence of ATP using FLAG-RPT4a and FLAG-RPT4b (Figures 3.4 and Figures 3.S3 D and F, respectively), the KDE distributions overlap with both the KS (0.017 for RPT4a and 0.058 for RPT4b) and median *p*-values (0.089 for RPT4a and 0.023 for RPT4b) were > 0.01 , suggesting that the RP and CP distributions are similar. Although not statistically significant, we still observed a two-fold-change enrichment in both FLAG-RPT4a/b samples purified in the presence of ATP for the RP over the CP. In summary, these KDE distributions and statistical analyses provide further support for enrichment of either the CP or the RP, as expected, when samples were affinity purified in the absence of ATP via PAG1-FLAG, or FLAG-RPT4a/b, respectively.

3.5.5. Statistical Analysis of Affinity Purifications Enriched for the RP or CP Identifies Proteasome-Associated Proteins and Putative Assembly Chaperones Specific to Each Subcomplex

The ability to affinity-enrich both the CP and the RP from *Arabidopsis* under gentle non-denaturing conditions without the need for high salt washes may enable identification of associated proteins, particularly those which may be more loosely bound. Additionally, targeting the RP or CP enables the identification of proteins that may be associating with either or both subcomplexes specifically. The previous

data represented in Figure 3.3 showed the raw MaxLFQ values for some of the relevant associated proteins; however, a formal statistical test against mock affinity purifications was important given the high number of proteins identified and quantified via LFQ-MS/MS. To better understand the suite of proteins that are statistically enriched in samples that target the CP or RP specifically we compared the LFQ-MS/MS data obtained from our affinity purifications against their respective wild-type controls using volcano plots and SAM tests at 0.01 FDR.

Affinity purifications enriching for the CP via PAG1-FLAG, performed either in the presence or absence of ATP (Figure 3.4 A and 3.4 B respectively) show statistical enrichment for two orthologs of the UMP1 CP assembly chaperone, UMP1a and UMP1b. The *Arabidopsis* genome encodes a third *UMP1* ortholog, *UMP1c*; however, we have no experimental evidence that UMP1c associates with the proteasome, even though it has several tryptic peptides from an extended C-terminal domain that could differentiate it from UMP1a/b (Figure 3.S4). UMP1a, and UMP1b are considerably higher expressed at 90 and 55 ESTs as compared to UMP1c which had only 7 ESTs (Berardini *et al.*, 2015), suggesting that UMP1c may be lowly expressed, or possibly a pseudogene.

Intriguingly, these samples also show statistical enrichment for a suite of putative CP assembly chaperones (Proteasome Biogenesis Associated Chaperone)

PBAC1-4 (Figure 3.4 A and 3.4 B). Although their annotation in The *Arabidopsis* Information Resource v10 (TAIR10) was incomplete, an InterProScan of their respective sequences supported these identifications (Table 3.S3). While PBAC1 was annotated as having a PBAC2 domain, the expectation value of 3.4E-6 was much lower than the expectation value of 1E-29 for the PBAC2 domains in the potential PBAC2 homologue. Additionally, PBAC1 and PBAC2 were previously identified by iterative PSI-BLAST with the human PAC1 and PAC2 orthologs in two separate studies (Kusmierczyk *et al.*, 2011; Le Tallec *et al.*, 2007) which aids in our assignment of PBAC1 and PBAC2. This PSI-BLAST based analysis was repeated using human PAC1-4, which identified PBAC1-4 as the top hits for these proteins in *Arabidopsis* (Table 3.S4). Protein sequence alignments for PBAC1-4 show that these proteins are conserved across plants but that they show little conservation with orthologs outside the plant kingdom with < 15% protein sequence identity (Figure 3.S5-3.S8). All statistically significant interactors from PAG1-FLAG affinity purifications performed in both the absence or presence of ATP are available in Tables 3.S5 and 3.S6.

As shown in Figure 3.4 C-F, UMP1a/b and PBAC1-4 were not significantly enriched in our affinity purifications targeting the RP, suggesting that these proteins interact with the CP specifically. However, affinity purifications that targeted the

RP through either FLAG-RPT4a or FLAG-RPT4b showed statistically significant enrichment over controls for the putative orthologs of RP assembly chaperones HSM3, NAS2, and NAS6 (Figure 3.4 C-F). Importantly, these putative RP assembly chaperones were also absent from our affinity purifications that targeted the CP, suggesting that these proteins are RP specific interactors. All statistically significant interactors from FLAG-RPT4a and FLAG-RPT4b affinity purified in the presence or absence of ATP are available in Tables 3.S7 - 3.S10. Several accessory proteins are common to both CP and RP purifications including the alternate capping particle PA200 (Book *et al.*, 2010), ECM29 that is involved in stabilizing the CP-RP association (Lehmann *et al.*, 2010), and the recently identified proteasome inhibitor 31 related protein PTRE1, which is involved in modulating proteasome activity in response to auxin (Yang *et al.*, 2016).


3.5.6. A Novel Plant Proteasome-Associated Protein, PAP1, Interacts with a Putative CP Assembly Chaperone

An intriguing plant-specific protein, named here Proteasome Associated Protein 1 or PAP1 was identified and significantly enriched in affinity preparations that targeted the CP (Figure 4). While PAP1 had no domains of known function, as determined by InterProScan (Table 3.S3), it did contain a C-terminal HbYX (hydrophobic amino

acid, followed by a tyrosine, followed by any amino acid) motif. This HbYX motif is commonly found in proteins that intercalate between the α -rings of the CP (Kusmierczyk *et al.*, 2011). This motif is found in both the plant specific PAP1, and in the assembly chaperone PBAC1 (Figure 3.5 A) as well as the conserved yeast assembly chaperones Pba1 and Pba2 (Kusmierczyk *et al.*, 2011). However, Pba2/PBAC2 orthologs outside of yeast typically contain a C-terminal HbF motif, with the only notable exception being the moss *Physcomitrella patens* which instead contains a HbYX motif. To confirm whether PAP1 did indeed interact with the proteasome, we generated transgenic plants expressing HA-tagged PAP1 in the *PAG1::PAG1-FLAG pag1-1* background. Figure 3.5 B shows immunoblots of SDS-PAGE separated samples from an affinity purification of HA-PAP1 utilizing anti-HA beads. HA-PAP1 expressed at very low levels in transgenic plants (Figure 3.5 B, input). However, upon immunopurification and detection with anti-HA antibodies we were able to detect HA-PAP1, the FLAG-tagged CP α -subunit PAG1, and the FLAG-less PAG1 product that likely resulted from cleavage of the FLAG tag during overnight incubation (Book *et al.*, 2010). Low levels of both the CP β -subunit PBA1, and RPN1 subunit were also detected in these samples.

To identify which putative *Arabidopsis* assembly chaperones might interact, we performed a yeast two-hybrid (Y2H) interaction assay for all pairwise combinations

[Figure 3.5 caption follows on next page]



of PBAC1-4, and PAP1 (Figure 3.5 C). We hypothesized that PAP1 may be a CP assembly chaperone and that it may interact with one or more of the other putative assembly chaperones. The Y2H showed pairwise interaction for PBAC1 and PBAC2, and pairwise interaction for PBAC3 and PBAC4 regardless of whether they were attached to the GAL4 activating or binding domains (Figure 3.5 C). These interactions were consistent with how the CP assembly orthologs in other organisms interact (Murata *et al.*, 2009). Taken together these data suggested these proteins might form heterodimeric pairs, like other CP assembly chaperone orthologs. Importantly,

Figure 3.5 (preceding page): PAP1 interacts with the proteasome and with putative assembly chaperone PBAC1. (A) C-terminal sections of protein sequence alignments for PAP1, PBAC1 and PBAC2, show conserved HbYX / HbY/F motifs underlined in red (full alignments in Figures 3.S9, 3.S5, and 3.S6) (B) An anti-HA immunopurification of HA-PAP1 expressed in the PAG1-FLAG background shows affinity enrichment for both PAP1 and the FLAG-tagged PAG1 α -subunit of the CP. A very low level of PBA1 is also observed in the elution. (C) A yeast-two-hybrid analysis of putative plant assembly chaperones shows interaction between PBAC1 and PBAC2 independently of whether PBAC1 is used as bait or prey. Similarly, interaction is also detected between PBAC3 and PBAC4 independently of which protein is used as bait. Interaction between PAP1 PBAC1 is also observed. (D) Bi-molecular Fluorescent Complementation (BiFC) with indicated combinations of PAP1, PBAC1, and PBAC2 fused at their N-terminus with either N or C -terminal halves of YFP (N-YFP, C-YFP). BiFC results confirm the results obtained via Y2H in (B) and show that these interactions also occur *in planta* between PBAC1 and PBAC2, and between PBAC1 and PAP1 in either orientation. A control is shown in this figure documenting a lack of auto activation for N-YFP and C-YFP. Additional controls combining N-YFP and C-YFP with their respective N and C terminal PBAC1, PBAC2, PBAC3, and PAP1 controls are shown in Figure 3.S10 along with the full panel from (D) for comparison.

we saw no interaction from PBAC1 or PBAC2 to PBAC3 or PBAC4 pairs in this Y2H, suggesting these putative assembly chaperones form separate complexes like their orthologs in mammals and yeast. Interestingly, we observed PAP1 interacting with PBAC1, which suggested that it may be part of the plant assembly chaperone PBAC1/2 complex (Figure 3.5 C).

Because these interactions were analyzed in yeast, which contain their own assembly chaperones that might interfere with interactions between plant subunits, we wanted to verify that PAP1 interacted with PBAC1 *in planta*. To do this we tested several pairwise interactions by bimolecular fluorescent complementation (BiFC) via transient expression in *Nicotiana benthamiana* leaves. Figure 3.5 D shows confocal images of transiently expressed PAP1, PBAC1, PBAC2, and PBAC3 with either the N- or C-terminal half of YFP (NYFP and CYFP respectively) attached as a translational fusion to the N-terminus of each putative assembly chaperone. As controls, PAP1 and PBAC1-3 were tested for auto-activation against NYFP or CYFP alone with no strong auto-activation being observed (Figure 3.S10 B). We again observed interactions between PBAC1 and PBAC2, regardless of their attachment to the N or C terminal halves of YFP based on their strong fluorescent intensity (Figure 3.5 D). Additionally, we saw an interaction between PAP1 and PBAC1, showing that PAP1 can interact with PBAC1 *in planta*. Consistent with our Y2H data, there

was no evidence of strong interaction between PAP1 and PBAC3 (Figure 3.5 D).

3.5.7. Inhibition of the Proteasome Results in Formation of Distinct Proteasome Sub-Species Containing Putative CP Assembly Chaperones

Upon inhibition with the proteasomal inhibitor MG132, most proteasome genes are coordinately upregulated by a pair of NAC transcription factors, NAC53 and NAC78, as the plants attempt to synthesize more proteasomes in an effort to increase degradation capacity (Gladman *et al.*, 2016). Additionally, after prolonged inhibition the complex becomes heavily ubiquitylated and is turned over via autophagy, likely as a clearing mechanism for inhibited complexes (Marshall *et al.*, 2015). We reasoned that treatment with MG132 might lead to an increase in the population of newly assembling proteasomes, potentially allowing us to observe proteasome assembly intermediates. As such, proteasomes were purified in the presence of ATP from *PAG1::PAG1-FLAG pag1-1* tissue treated for 16 hours with or without MG132. A silver stain of SDS-PAGE-separated proteins affinity-purified from MG132-treated tissue showed successful purification of the 26S proteasome under these conditions (Figure 3.6 A) with the only notable difference in the treated sample being an increased association of PA200, which was also observed by Book *et al.* (Book *et al.*, 2010). An anti-ubiquitin immunoblot of the same samples (Figure 3.6 B), showed

extensive ubiquitylation of the particle consistent with data observed by Marshall et al (Marshall *et al.*, 2015).

Non-denaturing native-PAGE of affinity-purified proteasomes were performed to determine if any novel complexes could be observed. Intriguingly, silver staining of samples affinity-purified from MG132-treated tissue revealed a smear below the canonical CP band (Figure 3.6 B). Additionally, extra bands were observed above the canonical CP band, which may represent stalled intermediates with the alternate capping particle PA200 (Li *et al.*, 2007). To identify the proteins contained in the bands shown in Figure 3.6B, we cut the corresponding gel slices, trypsin-digested the proteins, and then analyzed the resulting peptides via MS/MS. A LFQ-MS approach was used to quantify the proteins present in each sample (Gemperline *et al.*, 2016). Proteins identified in gel slices obtained from both MG132-treated and untreated control samples were clustered based on relative abundance using hierarchical clustering. The clusters were then visualized via a heat map showing that the RP and CP proteins cluster away from one another in these samples (Figure 3.6 C), consistent with their separation by native-PAGE. Furthermore, an outgroup that includes the putative assembly chaperones PBAC1-4, Ump1, and PAP1, plus the accessory factors PA200, PTRE1, as well as ubiquitin, is found in the CP cluster (Figure 3.6 C).




Figure 3.6: Protein composition of PAG1-FLAG affinity-purified proteasomes from tissues treated with and without MG132. (A) A silver stain of proteasomes affinity purified from *Arabidopsis* tissue treated for 24 hours with 50 μ M MG132 (left panel). An increase in PA200 is observed in these samples as judged by silver stain. An anti-ubiquitin immunoblot shows increased ubiquitylation of the complex on the right panel. (B) Native-PAGE analysis of 26S proteasomes affinity purified from MG132-treated (+) and untreated control (-) *Arabidopsis* tissues (left panel). A different banding pattern is observed between MG132-treated and untreated controls, with a smear below, and two bands (4 and 5) above the canonical CP band. An extended gel is shown on the right to better visualize and cut out the indicated bands for mass spectrometry. Cut bands indicated in red were recovered. The proteins contained within these bands were trypsin-digested and subjected to MS/MS analyses.




Figure 3.6: Protein composition of PAG1-FLAG affinity-purified proteasomes from tissues treated with and without MG132. (C) A heat map showing LFQ MS/MS analysis (using dNSAF values calculated with Morphewus Spectral Counter (Gemperline *et al.*, 2016)) of the gel slices shown in (B), which were hierarchically clustered using a Pearson's correlation, and ordered by gel slice. Clear grouping of the CP and RP is observed. Fractions containing the unassembled α -rings and putative assembly chaperones are seen from the native gel smears shown in (B). Fractions two and four contain the putative assembly chaperones PBAC1, PBAC2, and PAP1 as well as CP subunits and likely represent pre-assembly intermediates of the CP. PBAC3 and PBAC4 are only seen in the smear.

The proteins identified in gel slices below the canonical CP band in both MG132-treated and untreated control samples (Figure 3.6 B) contained a large abundance of CP α -subunits, with very few β -subunits (Figure 3.6 C). In addition to this set of α -subunits, the previously mentioned CP-specific putative assembly chaperones PBAC1-4 were also observed, as was the plant-specific PAP1 protein and UMP1b. PBAC3 and PBAC4 were only identified in the slice below the canonical CP band. Slices 1-5 contained mostly CP subunits with very few, if any, RP subunits (Figure 3.6 C). Slice 2, above the canonical CP band, from both treated and untreated samples also contained PBAC1, PBAC2, and PAP1, suggesting that this band may represent an assembly intermediate of the plant CP. All CP subunits and the alternate capping particle PA200 were identified in slice 3, suggesting that this band represents CP capped with PA200. Slices 4 and 5 were observed only in proteasomes purified from MG132 treated tissue (Figure 3.6 B, and Table 3.S11), and interestingly, slice 4 contained both the PA200 alternate capping particle, and the assembly chaperones PBAC1, PBAC2, and PAP1 along with most of the CP subunits, suggesting that this may be a PA200 containing assembly intermediate. Slice 5 likely represented a doubly capped PA200-CP-PA200 species, as no other putative assembly chaperones were present, and a strong signal for PA200, and the various CP subunits was observed.

In contrast to slices 1-5, most RP subunits were identified in slices 6-10. Based on size, slices 6-7 likely represented free RP complexes that have some associating α -subunits, while slices 8-9 likely represent doubly or singly capped RP-CP complexes (Book *et al.*, 2010). It was difficult to discriminate between singly and doubly capped proteasomes in these native gel assays, and we did not observe discrete bands for these complexes. Slice 10 likely represented the holo-complex of the 26S proteasome (Book *et al.*, 2010), and notably is the only slice in which the CP-RP stabilizing protein ECM29 was identified; however, it was detected at low abundance. Very low levels of the putative assembly chaperones PBAC1, PBAC2, and PAP1 were detected in these higher slices. Increased amounts of ubiquitin were detected in slices 1-5 for proteasomes that were affinity-purified from MG132-treated tissue, which also contained the putative assembly chaperones PBAC1, PBAC2, and PAP1. No ubiquitin receptors were identified in these slices, as expected, which suggests that ubiquitin is covalently attached; however we were unable to identify any canonical ubiquitylation footprints (GlyGly) (Wang *et al.*, 2006) on peptides corresponding to any proteasome subunits in these samples. Taken together, these data suggest that the novel upper novel slices and lower smears observed might represent stalled assembly intermediates, and that they may be actively ubiquitylated and turned over to prevent the formation of aberrant complexes.

3.6. Discussion

Here we have shown that we can affinity purify proteasomes from plants expressing FLAG-tagged RP subunits RPT4a and RPT4b, and that these preparations are of sufficient purity to enable analysis by MS/MS. Using a LFQ-MS/MS strategy, we were able to show that these RP-based affinity purifications identify the core components of the 26S proteasome, and are comparable to our previously developed CP-based affinity purification. Interestingly the only major difference between the RPT4a and RPT4b purifications was the statistical enrichment for each respective isoform when purifications were performed in the absence of ATP. Even with this enrichment, no other isoforms or associated proteins were preferentially enriched. For RPT4a/b purifications performed in the presence of ATP, enrichment for each respective subunit was observed; however, this difference was not statistically significant. These data imply that plants do not assemble particular proteasome subunit isoforms into specific sub-particles or isotypes at the young-seedling stage. While most plant proteasome subunits are encoded as paralogs, mammals instead have only a few subunit paralogs. $\beta 1$, $\beta 2$, and $\beta 5$ are replaced in immune tissue with $\beta 1i$, $\beta 2i$, and $\beta 5i$ to form the immunoproteasome isotype (Nandi *et al.*, 1996), and in thymus tissue the immunoproteasome contains $\beta 5t$ instead of $\beta 5i$ forming the thymoproteasome isotype (Murata *et al.*, 2007). Outside of these catalytic subunits,

the only other isoform switch involves replacing the $\alpha 4$ subunit with a variant only expressed in testes forming a testes specific proteasome isotype (Belote *et al.*, 1998; Uechi *et al.*, 2014). While there is no evidence for *Arabidopsis* assembling proteasomes into isoform specific sub-types, duplication of the RPT family has also occurred in monocots, with rice having duplicated subunits for RPT1, 2, 4 and 5. This suggests that there may be some evolutionary benefit to having multiple copies of the RPT family (Shibahara *et al.*, 2004). Interestingly, proteomic analysis of bran and callus rice tissue showed some small differences in isoform expression suggesting that analysis of different *Arabidopsis* developmental stages may provide additional insight into potential isoform-specific functions (Shibahara *et al.*, 2004). In conclusion, we did not find proteasome isotypes at the young seedling stage that are specific to either RPT4 isoform. In other organisms, proteasome sub-types seem to be found in specific tissues or cell-types (Belote *et al.*, 1998; Murata *et al.*, 2007; Nandi *et al.*, 1996; Uechi *et al.*, 2014). Therefore, affinity purifications of proteasomes from different plant tissues and developmental stages may be an intriguing next step to explore the possible existence of plant proteasome isotypes.

In conjunction with our previously developed CP affinity purification (Book *et al.*, 2010), this newly developed RP affinity purification allowed us to specifically enrich for either the CP or RP sub-complexes when purifications are performed in

the absence of ATP. LFQ-MS/MS and statistical analyses of these affinity-purified proteasome sub-complexes enabled identification of sub-complex-specific PAPs. The most predominant RP-specific PAPs present in both RPT4a and RPT4b affinity-purified samples are likely orthologs of the RP assembly chaperones NAS2, NAS6, and HSM3. We were unable to identify any proteins that were likely orthologs for the RP assembly chaperone RPN14. In yeast, RPN14 and NAS6 bind together in a complex, and only double mutants show a strong growth defect in growth, suggesting that together they perform redundant functions (Funakoshi *et al.*, 2009). These data suggest that that plant proteasome assembly has diverged from the yeast and mammalian pathways and may no longer require RPN14 to assemble the RP (Funakoshi *et al.*, 2009). There are no obvious orthologs for RPN14 in the *Arabidopsis* genome (Book *et al.*, 2010) which supports this hypothesis. The exact role of NAS2, NAS6, and HSM3 in plant proteasome assembly remains to be determined; however, their specific association with the RP suggests that they may perform similar functions in RP assembly as in mammals and yeast. It will be intriguing to test if these RP assembly orthologs form similar modules with specific RP subunits like their yeast and mammalian counterparts (Nas2 with Rpt4 and 5, Nas6 with Rpt3, Hsm3 with Rpt1,2 and Rpn1, and Rpn14 with Rpt6) (Park *et al.*, 2010).

Several PAPs identified here were enriched in affinity purifications targeting

both the RP and CP. Unsurprisingly, the alternate capping particle PA200 was enriched in our CP-specific purifications. This protein is known to interact with α -subunits of the CP (Ortega *et al.*, 2005). However, we detect PA200 in our RP affinity purifications as well, which suggests that plants form a hybrid RP-CP-PA200 complex. This hybrid complex has been observed in yeast, and is speculated to aide in release of degraded substrate peptides (Ortega *et al.*, 2005; Schmidt *et al.*, 2005; Ustrell *et al.*, 2002). In plants however, PA200 mutants display a remarkably wild-type phenotype (Book *et al.*, 2010), which suggests that this hybrid complex is not essential in *Arabidopsis*. ECM29 was also found to be enriched as compared to mock controls in both CP and RP targeted purifications. ECM29 is known in yeast to bind the CP subunit PRE10 (named PAG1 in *Arabidopsis*), and the RP subunit Rpt5 (Wani *et al.*, 2016). In yeast, ECM29 is one of the only PAPs known to bind both the RP and the CP (Lehmann *et al.*, 2010), consistent with our observations. Ecm29 has been shown to inhibit the yeast 26S proteasome, and may be a checkpoint for damaged or aberrantly assembled complexes (Lehmann *et al.*, 2010). While ECM29 likely performs a similar role in *Arabidopsis*, its function in plants remains to be investigated.

A recently characterized plant ortholog of proteasome inhibitor 31 (PI31), named Proteasome Regulator 1 or PTRE1 (Yang *et al.*, 2016), was identified as a CP-specific

PAP, providing confirmation of PTRE1's association with the CP of the 26S proteasome. However, the most predominant CP-specific PAPs identified here were putative orthologs of the CP assembly chaperones PBAC1-4, UMP1, and the plant specific protein PAP1. In our analyses, searches for PBAC1-4 in the *Arabidopsis* genome using assembly chaperones from yeast (Pba1-4) or humans (PAC1-4) failed with BLAST alone. This is not surprising given that orthologs from different kingdoms typically shared <25% identity, with fungi and animal forms of these chaperones typically sharing <15% sequence identity (Figure 3.S5 - 3.S8) despite having similar functions. However two independent groups identified orthologs to the assembly chaperones PBAC1 and PBAC2 using an iterative PSI-BLAST based approach with the human orthologs PAC1 and PAC2 (Kusmierczyk *et al.*, 2011; Le Tallec *et al.*, 2007). PSI-BLAST has several advantages over BLASTp in that it is typically more sensitive, and can be useful when identifying protein like PBAC1-4 that likely share structural homology but share little sequence homology (Altschul *et al.*, 1997; Aravind and Koonin, 1999; Kusmierczyk *et al.*, 2011; Yashiroda *et al.*, 2008). We were able to identify PBAC1 and PBAC2 as the top *Arabidopsis* PSI-BLAST hits for human PAC1 and PAC2 (Table 3.S4), respectively, repeating the finding by the both Le Tallec *et al.* (2007), Kusmierczyk *et al.* (2011). Additionally, we were able to extend this analysis, with both PBAC3 and PBAC4 being top PSI-BLAST hits for

human PAC3 and PAC4 (Table 3.S4). PSI-BLAST analysis with the yeast assembly chaperones Pba1-4 failed to identify the putative *Arabidopsis* orthologs PBAC1-4, suggesting that the plant CP assembly system may be different from the yeast system. Sequence-based InterProScan analyses showed that PBAC2-4 contained PAC2-4 domains (Table 3.S3), further supporting these assignments. InterProScan analyses showed that PBAC1 contained a PAC2 domain; however, the E-value for PBAC1 containing this domain is much lower than the assignment of this domain to PBAC2 (3.4E-6 vs 1.0E-29), and is much lower than the other assigned domains. Despite this disagreement, PBAC1 was the top hit in the *Arabidopsis* genome when searching via PSI-BLAST with human PAC1, as performed by ourselves and others, and for these reasons we have decided to use the name PBAC1 for this identified protein (Kusmierczyk *et al.*, 2011; Le Tallec *et al.*, 2007). Sequence alignments of PBAC1 and PBAC2 also showed conserved HbYX and HbY/F motifs respectively that are also found in their human counterparts. Sequence analysis alone, however, is insufficient to conclude that these are assembly chaperones. There is strong divergence in sequence identity between the plant, yeast, and animal sequences (alignments and identity matrices in Figures 3.S5 - 3.S8); therefore, additional characterizations will be necessary to determine what role if any they have in plant CP assembly. Despite this divergence, these bioinformatics analyses, along with

prior literature (Kusmierczyk *et al.*, 2011; Le Tallec *et al.*, 2007), help support our identification of these CP-specific PAPs as PBAC1-4.

Phylogenetic analysis of the putative plant assembly chaperones orthologs PBAC1-4, and plant specific protein PAP1, showed that PBAC2-4 formed clades with their respective animal counterparts (Figures 3.S11 A and B); however, PBAC2 was the only protein to form a clear clade with all sequences from animals, yeast, and plants. PBAC1, 3, and 4 from yeast did not form clades with either the plant or animal sequences, suggesting that they are more different in sequence as compared to the plant and animal versions. In this analysis PBAC1 from yeast, mammals and plants was a strong exception in that it did not form a strong clade with any of their respective orthologs (Figure 3.S11). These data may not be surprising given the low sequence homology shared between yeast and animal PBAC1, with sequence identity at <13%. This low % identity makes phylogenetic analysis of these putative PBAC1 orthologs particularly challenging, therefore we cannot draw strong conclusions about PBAC1 orthology based on this analysis alone.

In yeast and mammals, Pba1/PAC1 and Pba2/PAC2 form a heterodimeric pair. Separately, Pba3/PAC3 and Pba4/PAC4 also form a heterodimeric pair. If plant PBAC1-4 are acting similarly, we would expect that PBAC1 should interact with PBAC2 and that PBAC3 should interact with PBAC4. Indeed, our Y2H analyses

shows these interactions (Figure 3.5 C) suggesting that they each form a heterodimer like their respective orthologs in yeast and mammals (Kunjappu and Hochstrasser, 2014). Additionally, in our Y2H we failed to detect interactions between these two putative heterodimeric pairs as both PBAC1 and PBAC2 failed to interact with PBAC3 and PBAC4, which suggests that these proteins form two separate complexes. These data were also re-capitulated in our BiFC analyses (Figure 3.5 D) suggesting that these interactions occur *in planta*; however, the interaction between PBAC3 and PBAC4 was not tested via BiFC.

We were able to confirm the interaction of the CP-specific PAP1 protein with the proteasome by IP with an HA tagged version, identifying the CP subunit PAG1. The presence of a conserved HbYX motif in PAP1 (Figure 3.5 A), which was also found in the putative assembly chaperone PBAC1, suggested that PAP1 might play a role in plant proteasome assembly. In yeast, the Pba1 and Pba2 proteins both have HbYX motifs important for interacting with the CP α -ring; however orthologs of Pba2 in higher eukaryotes lack HbYX motifs, and instead have a HbY/F motif that lacks the last variable C-terminal amino acid (Kusmierczyk *et al.*, 2011). Consistent with this, PBAC1 has a HbYX motif conserved in plants, and PBAC2 has a HbY/F motif conserved in all plants analyzed, except *Physcomitrella patens* (which has a HbFX motif, Figure 3.5 A), suggesting the plant-CP-assembly pathway is more

similar to that of higher eukaryotes than that of fungi. If PAP1 is functioning as a CP assembly chaperone it might be acting in concert with PBAC1, given their interaction (Figure 3.5 C and D). In one scenario, PAP1 could replace PBAC2 in the PBAC1/2 heterodimer, or in an alternative scenario, PAP1 may be forming a novel trimeric complex with both PBAC1 and PBAC2. Despite our evidence that PAP1 contains a conserved HbYX motif, and interacts with the plant assembly chaperone PBAC1 via both Y2H and *in planta* by BiFC, we cannot rule out that PAP1 functions outside of plant proteasome assembly.

While an interaction network of these putative plant proteasome assembly chaperones provides some insight into how these proteins organize, our analysis of complexes affinity purified from PAG1-FLAG tissue treated with the potent and specific proteasome inhibitor MG132 shows the formation of novel sub-complexes that contain these putative chaperones. In general, two classes of species appeared in the sample affinity purified from tissue treated with MG132 as analyzed by native-PAGE. The first is a smear running at a faster pace below the canonical CP species which was increased in intensity upon inhibitor treatment. The second class includes discrete species migrating more slowly than the canonical CP band (Figure 3.6 B).

In the untreated sample, native-PAGE showed a very faint band (as opposed to

a smear) below the canonical CP band (Figure 3.6 B, labeled Half Barrel?). This may represent a proteasome half-barrel known as a 15S precursor complex that also typically runs below the CP band in native-PAGE analyses and contains partially assembled α -subunits along with UMP1 (Kock *et al.*, 2015). Indeed, our MS/MS analyses of this faint band identified UMP1, α -subunits and the putative assembly chaperones PBAC1-4 and PAP1. While mostly α -subunits were present, some β -subunits were still identified at lower levels. In yeast, these 15S complexes lack the β 7 subunit (Kock *et al.*, 2015). We still detected the β 7 subunit PBG1; however, it was present at a 50-fold lower level as compared to the CP band in our untreated samples (Table 3.S11). Native-PAGE analysis of the affinity preparations from MG132-treated tissue showed a smear instead of the faint band seen in the untreated samples. MS/MS analyses of this smear showed that it also contained unassembled α -subunits, UMP1, and the suite of putative assembly chaperones PBAC1-4. We could not observe a discrete band similar to the 15S species from yeast but this may be due to the fact that we are inhibiting the complex and are capturing a wider array of partially assembled products, or that these partially assembled products are less stable in plants as compared to yeast. Indeed, increased levels of ubiquitin were found in this fraction even though no ubiquitin binding proteins were detected (as expected), suggesting that these sub-species may be aberrant and are likely turned

over, either via autophagy or other active proteasomes. PBAC3, PBAC4, and UMP1 are exclusively detected in the untreated and treated smears below the CP band, which is consistent with PBAC3 and PBAC4 preferentially binding free α -subunits at early stages of proteasome assembly, as established in both yeast and mammals (Kunjappu and Hochstrasser, 2014), and with UMP1 being degraded at later stages of proteasome assembly (Ramos *et al.*, 1998).

The new species observed above the canonical CP band upon MG132 treatment contained large amounts of PA200, suggesting that these are likely singly- and doubly-capped versions of the CP (CP-PA200, and PA200-CP-PA200, respectively). The first band above the CP was previously identified as CP-PA200 by our lab (Book *et al.*, 2010); however, this doubly capped complex and additional bands observed in affinity purified complexes from MG132 inhibited tissue have not been seen before in plants. Recently, other researchers have found that human cell lines treated with the proteasome inhibitor bortezomib induced the formation of a species above the canonical CP band, which also contained PA200 (Welk *et al.*, 2016) suggesting that inhibition of the complex increases CP-PA200 binding, and that this process is conserved between animals and plants. The function of increased PA200 binding to the CP is still unclear in both plants and animals. It could be that inhibition of the complex causes PA200 to associate and help release peptides from

stalled proteasomes. Alternatively, given PA200's role in assembly (Savulescu and Glickman, 2011), it could represent increased biogenesis of the CP sub-complex in response to inhibition.

Intriguingly, high levels of PBAC1, PBAC2, and PAP1 were found in the native-PAGE bands directly above the canonical CP band (slice 2 Figure 3.6 B and Figure 3.6 C) in samples affinity purified from both untreated and MG132 treated tissues. This suggests that PBAC1, PBAC2, and PAP1 may be acting at later stages of proteasome assembly, given their association with species that run higher than mature CP. While the assembly chaperones Pba1/PAC1 and Pba2/PAC2 in yeast and mammals typically do not associate with mature forms of the CP, the related archaeal assembly chaperones PbaA and PbaB are found loosely associated with mature CP when its active sites are bound with inhibitors (Kusmierczyk *et al.*, 2011). The data gathered here suggest that PBAC1, PBAC2, and PAP1 can bind mature CP and suggest that plant CP assembly at later stages may be more similar to archaeal proteasome assembly. Taken together these larger-sized bands may represent stalled assembly intermediates of the mature proteasome still carrying their assembly chaperones. Like the previously mentioned smears, increased levels of ubiquitin are found in these fractions even though no ubiquitin binding proteins were detected, suggesting that these sub-species may also be aberrant and are likely turned over. While the

exact role of PAP1 in plant proteasome assembly, if any, still needs to be determined, the data presented here suggest that it may be involved in late-stage proteasome assembly, as it co-migrates with PBAC1 and PBAC2. Figure 3.7 illustrates a possible model of CP assembly that is consistent with the data collected here.




Figure 3.7: Putative Model of 20S Proteasome Assembly in *Arabidopsis*. A model of 20S proteasome assembly adapted from (Murata *et al.*, 2009).

In summary, the data presented herein show that we were able to efficiently purify the 26S proteasome via both the CP and RP. The newly-developed RP-based purification allowed us to identify CP and RP specific PAPs, several of which may be involved in the assembly of their respective sub-complexes. The fact that the putative CP assembly chaperones were only identified in our CP-based affinity purifications, and that the putative RP assembly chaperones were only identified in our RP-based affinity purifications, suggest that only mature RP is purified with our CP affinity preparation, and that only mature CP is purified with our RP affinity purification. Our follow up analysis builds an interaction network of PBAC1-4 and PAP1, and identifies novel sub-species that contain these putative assembly chaperones when analyzing sub-complexes affinity purified from MG132-treated tissue. Taken together, these data suggest that UMP1, PBAC3, and PBAC4 may be performing a role in the early biogenesis of the CP, while PBAC1, PBAC2, and possibly PAP1 are playing a role in later stages of CP assembly. While we have been able to identify the likely players in the plant proteasome assembly pathways for both the RP and CP, additional functional analyses of these proteins will be needed to demonstrate their roles as putative assembly chaperones, and to fully elucidate the plant proteasome assembly pathway.

3.7. Literature Cited

- Altschul, S.F., Madden, T.L., Schaffer, A.A., Zhang, J., Zhang, Z., Miller, W., and Lipman, D.J.** (1997). Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res.* **25:** 3389–402.
- Aravind, L. and Koonin, E.V.** (1999). Gleaning non-trivial structural, functional and evolutionary information about proteins by iterative database searches. *J. Mol. Biol.* **287:** 1023–40.
- Beck, F., Unverdorben, P., Bohn, S., Schweitzer, A., Pfeifer, G., Sakata, E., Nickell, S., Plitzko, J.M., Villa, E., Baumeister, W., and Förster, F.** (2012). Near-atomic resolution structural model of the yeast 26S proteasome. *Proc. Natl. Acad. Sci. U.S.A.* **109:** 14870–14875.
- Belote, J.M., Miller, M., and Smyth, K.A.** (1998). Evolutionary conservation of a testes-specific proteasome subunit gene in *Drosophila*. *Gene* **215:** 93–100.
- Berardini, T.Z., Reiser, L., Li, D., Mezheritsky, Y., Muller, R., Strait, E., and Huala, E.** (2015). The arabidopsis information resource: Making and mining the gold standard annotated reference plant genome. *Genesis* **53:** 474–485.
- Besche, H.C., Haas, W., Gygi, S.P., and Goldberg, A.L.** (2009). Isolation of mammalian 26S proteasomes and p97/VCP complexes using the ubiquitin-like domain from HHR23B reveals novel proteasome-associated proteins. *Biochemistry (Mosc.)* **48:** 2538–49.
- Book, A.J., Gladman, N.P., Lee, S.S., Scalf, M., Smith, L.M., and Vierstra, R.D.** (2010). Affinity purification of the Arabidopsis 26S proteasome reveals a diverse array of plant proteolytic complexes. *J. Biol. Chem.* **285:** 25554–25569.
- Book, A.J., Smalle, J., Lee, K.H., Yang, P., Walker, J.M., Casper, S., Holmes, J.H., Russo, L.A., Buzzinotti, Z.W., Jenik, P.D., and Vierstra, R.D.** (2009). The RPN5 subunit of the 26S proteasome is essential for gametogenesis, sporophyte development, and complex assembly in *Arabidopsis*. *Plant Cell* **21:** 460–78.
- Brukhin, V., Gheyselinck, J., Gagliardini, V., Genschik, P., and Grossniklaus, U.** (2005). The RPN1 subunit of the 26S proteasome in *Arabidopsis* is essential for embryogenesis. *Plant Cell* **17:** 2723–37.
- Chevallet, M., Luche, S., and Rabilloud, T.** (2006). Silver staining of proteins in polyacrylamide gels. *Nat. Protoc.* **1:** 1852–8.

- Clough, S.J. and Bent, A.F.** (1998). Floral dip: a simplified method for Agrobacterium-mediated transformation of *Arabidopsis thaliana*. *Plant J.* **16**: 735–43.
- Cox, J., Hein, M.Y., Luber, C.A., Paron, I., Nagaraj, N., and Mann, M.** (2014). Accurate proteome-wide label-free quantification by delayed normalization and maximal peptide ratio extraction, termed MaxLFQ. *Mol. Cell. Proteomics* **13**: 2513–26.
- Cox, J. and Mann, M.** (2008). MaxQuant enables high peptide identification rates, individualized p.p.b.-range mass accuracies and proteome-wide protein quantification. *Nat. Biotechnol.* **26**: 1367–72.
- Dange, T., Smith, D.M., Noy, T., Rommel, P.C., Jurzitz, L., Cordero, R.J., Legende, A., Finley, D., Goldberg, A.L., and Schmidt, M.** (2011). The Blm10 protein promotes proteasomal substrate turnover by an active gating mechanism. *J. Biol. Chem.* **286**: 42830–42839.
- Edelheit, O., Hanukoglu, A., and Hanukoglu, I.** (2009). Simple and efficient site-directed mutagenesis using two single-primer reactions in parallel to generate mutants for protein structure-function studies. *BMC Biotechnol.* **9**: 61.
- Elsasser, S., Chandler-Militello, D., Muller, B., Hanna, J., and Finley, D.** (2004). Rad23 and Rpn10 serve as alternative ubiquitin receptors for the proteasome. *J. Biol. Chem.* **279**: 26817–22.
- Elsasser, S., Gali, R.R., Schwickart, M., Larsen, C.N., Leggett, D.S., Muller, B., Feng, M.T., Tubing, F., Dittmar, G.A., and Finley, D.** (2002). Proteasome subunit Rpn1 binds ubiquitin-like protein domains. *Nat. Cell Biol.* **4**: 725–30.
- Farmer, L.M., Book, A.J., Lee, K.H., Lin, Y.L., Fu, H., and Vierstra, R.D.** (2010). The RAD23 family provides an essential connection between the 26S proteasome and ubiquitylated proteins in *Arabidopsis*. *Plant Cell* **22**: 124–42.
- Fatimababy, A.S., Lin, Y.L., Usharani, R., Radjacommare, R., Wang, H.T., Tsai, H.L., Lee, Y., and Fu, H.** (2010). Cross-species divergence of the major recognition pathways of ubiquitylated substrates for ubiquitin/26S proteasome-mediated proteolysis. *FEBS J.* **277**: 796–816.
- Finley, D.** (2009). Recognition and processing of ubiquitin-protein conjugates by the proteasome. *Annu. Rev. Biochem.* **78**: 477–513.

- Fu, H., Doelling, J.H., Arendt, C.S., Hochstrasser, M., and Vierstra, R.D.** (1998). Molecular organization of the 20S proteasome gene family from *Arabidopsis thaliana*. *Genetics* **149**: 677–92.
- Funakoshi, M., Tomko, R. J., J., Kobayashi, H., and Hochstrasser, M.** (2009). Multiple assembly chaperones govern biogenesis of the proteasome regulatory particle base. *Cell* **137**: 887–99.
- Gamborg, O.L., Miller, R.A., and Ojima, K.** (1968). Nutrient requirements of suspension cultures of soybean root cells. *Exp. Cell. Res.* **50**: 151–8.
- Gelvin, S.B.** (2003). Agrobacterium-mediated plant transformation: the biology behind the "gene-jockeying" tool. *Microbiol. Mol. Biol. Rev.* **67**: 16–37, table of contents.
- Gemperline, D.C., Scalf, M., Smith, L.M., and Vierstra, R.D.** (2016). Morpheus Spectral Counter: A computational tool for label-free quantitative mass spectrometry using the Morpheus search engine. *Proteomics* **16**: 920–4.
- Gladman, N.P., Marshall, R.S., Lee, K.H., and Vierstra, R.D.** (2016). The Proteasome Stress Regulon Is Controlled by a Pair of NAC Transcription Factors in *Arabidopsis*. *Plant Cell* **28**: 1279–96.
- Gonnet, G.H., Cohen, M.A., and Benner, S.A.** (1992). Exhaustive matching of the entire protein sequence database. *Science* **256**: 1443–5.
- Goodstein, D.M., Shu, S., Howson, R., Neupane, R., Hayes, R.D., Fazo, J., Mitros, T., Dirks, W., Hellsten, U., Putnam, N., and Rokhsar, D.S.** (2012). Phytozome: a comparative platform for green plant genomics. *Nucleic Acids Res.* **40**: D1178–86.
- Hirano, Y., Kaneko, T., Okamoto, K., Bai, M., Yashiroda, H., Furuyama, K., Kato, K., Tanaka, K., and Murata, S.** (2008). Dissecting beta-ring assembly pathway of the mammalian 20S proteasome. *EMBO J.* **27**: 2204–13.
- Jones, P., Binns, D., Chang, H.Y., Fraser, M., Li, W., McAnulla, C., McWilliam, H., Maslen, J., Mitchell, A., Nuka, G., Pesceat, S., Quinn, A.F., Sangrador-Vegas, A., Scheremetjew, M., Yong, S.Y., Lopez, R., and Hunter, S.** (2014). InterProScan 5 genome-scale protein function classification. *Bioinformatics* **30**: 1236–40.
- Julio, S. and Jose, S.S.J.** (2006). *Arabidopsis Protocols*, Second Edition, volume 323. Humana Press.

- Kock, M., Nunes, M.M., Hemann, M., Kube, S., Dohmen, R.J., Herzog, F., Ramos, P.C., and Wendler, P.** (2015). Proteasome assembly from 15S precursors involves major conformational changes and recycling of the Pba1-Pba2 chaperone. *Nat. Commun.* **6**: 6123.
- Kunjappu, M.J. and Hochstrasser, M.** (2014). Assembly of the 20S proteasome. *Biochim. Biophys. Acta* **1843**: 2–12.
- Kusmierczyk, A.R., Kunjappu, M.J., Kim, R.Y., and Hochstrasser, M.** (2011). A conserved 20S proteasome assembly factor requires a C-terminal HbYX motif for proteasomal precursor binding. *Nat. Struct. Mol. Biol.* **18**: 622–9.
- Laemmli, U.K.** (1970). Cleavage of Structural Proteins during the Assembly of the Head of Bacteriophage T4. *Nature* **227**: 680–685.
- Lander, G.C., Estrin, E., Matyskiela, M.E., Bashore, C., Nogales, E., and Martin, A.** (2012). Complete subunit architecture of the proteasome regulatory particle. *Nature* **482**: 186–191.
- Lasker, K., Förster, F., Bohn, S., Walzthoeni, T., Villa, E., Unverdorben, P., Beck, F., Aebersold, R., Sali, A., and Baumeister, W.** (2012). Molecular architecture of the 26S proteasome holocomplex determined by an integrative approach. *Proc. Natl. Acad. Sci. U.S.A.* **109**: 1380–1387.
- Le Tallec, B., Barrault, M.B., Courbeyrette, R., Guérois, R., Marsolier-Kergoat, M.C., and Peyroche, A.** (2007). 20S proteasome assembly is orchestrated by two distinct pairs of chaperones in yeast and mammals. *Mol. Cell* **27**: 660–674.
- Leggett, D.S., Glickman, M.H., and Finley, D.** (2005). Purification of proteasomes, proteasome subcomplexes, and proteasome-associated proteins from budding yeast. *Methods Mol. Biol.* **301**: 57–70.
- Leggett, D.S., Hanna, J., Borodovsky, A., Crosas, B., Schmidt, M., Baker, R.T., Walz, T., Ploegh, H.L., and Finley, D.** (2002). Multiple associated proteins regulate proteasome structure and function. *Mol. Cell* **10**: 495–507.
- Lehmann, A., Niewienda, A., Jechow, K., Janek, K., and Enenkel, C.** (2010). Ecm29 fulfills quality control functions in proteasome assembly. *Mol. Cell* **38**: 879–88.
- Li, X., Kusmierczyk, A.R., Wong, P., Emili, A., and Hochstrasser, M.** (2007). beta-Subunit appendages promote 20S proteasome assembly by overcoming an Ump1-dependent checkpoint. *EMBO J.* **26**: 2339–49.

- Lin, Y.L., Sung, S.C., Tsai, H.L., Yu, T.T., Radjacompare, R., Usharani, R., Fatimababy, A.S., Lin, H.Y., Wang, Y.Y., and Fu, H.** (2011). The defective proteasome, not substrate recognition function, is responsible for the null phenotypes of the Arabidopsis proteasome subunit RPN10. *Plant Cell* **23**: 2754–2773.
- Liu, C.W., Li, X., Thompson, D., Wooding, K., Chang, T.L., Tang, Z., Yu, H., Thomas, P.J., and DeMartino, G.N.** (2006). ATP binding and ATP hydrolysis play distinct roles in the function of 26S proteasome. *Mol. Cell* **24**: 39–50.
- Livneh, I., Cohen-Kaplan, V., Cohen-Rosenzweig, C., Avni, N., and Ciechanover, A.** (2016). The life cycle of the 26S proteasome: from birth, through regulation and function, and onto its death. *Cell Res.* **26**: 869–85.
- Marques, A.J., Glanemann, C., Ramos, P.C., and Dohmen, R.J.** (2007). The C-terminal extension of the beta7 subunit and activator complexes stabilize nascent 20 S proteasomes and promote their maturation. *J. Biol. Chem.* **282**: 34869–76.
- Marshall, R.S., Gemperline, D.C., and Vierstra, R.D.** (2017). Isolation of Plant Organelles and Structures, volume 1 of *Methods in Molecular Biology*. Humana Press, Methods in Molecular Biology, 1 edition.
- Marshall, R.S., Li, F., Gemperline, D.C., Book, A.J., and Vierstra, R.D.** (2015). Autophagic Degradation of the 26S Proteasome Is Mediated by the Dual ATG8/Ubiqutin Receptor RPN10 in Arabidopsis. *Mol. Cell* **58**: 1053–66.
- Martin, K., Kopperud, K., Chakrabarty, R., Banerjee, R., Brooks, R., and Goodin, M.M.** (2009). Transient expression in Nicotiana benthamiana fluorescent marker lines provides enhanced definition of protein localization, movement and interactions in planta. *Plant J.* **59**: 150–62.
- Mi, H., Lazareva-Ulitsky, B., Loo, R., Kejariwal, A., Vandergriff, J., Rabkin, S., Guo, N., Muruganujan, A., Doremieux, O., Campbell, M.J., Kitano, H., and Thomas, P.D.** (2005). The PANTHER database of protein families, subfamilies, functions and pathways. *Nucleic Acids Res.* **33**: D284–8.
- Murashige, T. and Skoog, F.** (1962). A Revised Medium for Rapid Growth and Bio Assays with Tobacco Tissue Cultures. *Physiol. Plant.* **15**: 473–497.
- Murata, S., Sasaki, K., Kishimoto, T., Niwa, S., Hayashi, H., Takahama, Y., and Tanaka, K.** (2007). Regulation of CD8+ T cell development by thymus-specific proteasomes. *Science* **316**: 1349–53.

- Murata, S., Yashiroda, H., and Tanaka, K.** (2009). Molecular mechanisms of proteasome assembly. *Nat. Rev. Mol. Cell Biol.* **10**: 104–15.
- Nandi, D., Jiang, H., and Monaco, J.J.** (1996). Identification of MECL-1 (LMP-10) as the third IFN-gamma-inducible proteasome subunit. *J. Immunol.* **156**: 2361–4.
- Ortega, J., Heymann, J.B., Kajava, A.V., Ustell, V., Rechsteiner, M., and Steven, A.C.** (2005). The axial channel of the 20S proteasome opens upon binding of the PA200 activator. *J. Mol. Biol.* **346**: 1221–7.
- Paraskevopoulos, K., Kriegenburg, F., Tatham, M.H., Rösner, H.I., Medina, B., Larsen, I.B., Brandstrup, R., Hardwick, K.G., Hay, R.T., Kragelund, B.B., Hartmann-Petersen, R., and Gordon, C.** (2014). Dss1 is a 26S proteasome ubiquitin receptor. *Mol. Cell* **56**: 453–461.
- Park, S., Tian, G., Roelofs, J., and Finley, D.** (2010). Assembly manual for the proteasome regulatory particle: the first draft. *Biochem. Soc. Trans.* **38**: 6–13.
- Pathare, G.R., Nagy, I., Bohn, S., Unverdorben, P., Hubert, A., Korner, R., Nickell, S., Lasker, K., Sali, A., Tamura, T., Nishioka, T., Forster, F., Baumeister, W., and Bracher, A.** (2012). The proteasomal subunit Rpn6 is a molecular clamp holding the core and regulatory subcomplexes together. *Proc. Natl. Acad. Sci. U.S.A.* **109**: 149–54.
- Ramos, P.C., Hockendorff, J., Johnson, E.S., Varshavsky, A., and Dohmen, R.J.** (1998). Ump1p is required for proper maturation of the 20S proteasome and becomes its substrate upon completion of the assembly. *Cell* **92**: 489–99.
- Rappsilber, J., Ishihama, Y., and Mann, M.** (2003). Stop and go extraction tips for matrix-assisted laser desorption/ionization, nanoelectrospray, and LC/MS sample pretreatment in proteomics. *Anal. Chem.* **75**: 663–70.
- Ronquist, F., Teslenko, M., van der Mark, P., Ayres, D.L., Darling, A., Hohna, S., Larget, B., Liu, L., Suchard, M.A., and Huelsenbeck, J.P.** (2012). MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Syst. Biol.* **61**: 539–42.
- Russell, J.D., Scalf, M., Book, A.J., Ladrón, D.T., Vierstra, R.D., Smith, L.M., and Coon, J.J.** (2013). Characterization and quantification of intact 26S proteasome proteins by real-time measurement of intrinsic fluorescence prior to top-down mass spectrometry. *PLoS One* **8**: e58157.

- Sadre-Bazzaz, K., Whitby, F.G., Robinson, H., Formosa, T., and Hill, C.P.** (2010). Structure of a Blm10 complex reveals common mechanisms for proteasome binding and gate opening. *Mol. Cell* **37**: 728–735.
- Saveliev, S.V., Woodroffe, C.C., Sabat, G., Adams, C.M., Klaubert, D., Wood, K., and Urh, M.** (2013). Mass spectrometry compatible surfactant for optimized in-gel protein digestion. *Anal. Chem.* **85**: 907–14.
- Savulescu, A.F. and Glickman, M.H.** (2011). Proteasome activator 200: the heat is on. *Mol. Cell. Proteomics* **10**: R110 006890.
- Schmidt, M., Haas, W., Crosas, B., Santamaria, P.G., Gygi, S.P., Walz, T., and Finley, D.** (2005). The HEAT repeat protein Blm10 regulates the yeast proteasome by capping the core particle. *Nat. Struct. Mol. Biol.* **12**: 294–303.
- Schreiner, P., Chen, X., Husnjak, K., Randles, L., Zhang, N., Elsasser, S., Finley, D., Dikic, I., Walters, K.J., and Groll, M.** (2008). Ubiquitin docking at the proteasome through a novel pleckstrin-homology domain interaction. *Nature* **453**: 548–52.
- Shi, Y., Chen, X., Elsasser, S., Stocks, B.B., Tian, G., Lee, B.H., Shi, Y., Zhang, N., de Poot, S.A., Tuebing, F., Sun, S., Vannoy, J., Tarasov, S.G., Engen, J.R., Finley, D., and Walters, K.J.** (2016). Rpn1 provides adjacent receptor sites for substrate binding and deubiquitination by the proteasome. *Science* **351**.
- Shibahara, T., Kawasaki, H., and Hirano, H.** (2004). Mass spectrometric analysis of expression of ATPase subunits encoded by duplicated genes in the 19S regulatory particle of rice 26S proteasome. *Arch. Biochem. Biophys.* **421**: 34–41.
- Sievers, F. and Higgins, D.G.** (2014). Clustal Omega, accurate alignment of very large numbers of sequences. *Methods Mol. Biol.* **1079**: 105–16.
- Sievers, F., Wilm, A., Dineen, D., Gibson, T.J., Karplus, K., Li, W., Lopez, R., McWilliam, H., Remmert, M., Soding, J., Thompson, J.D., and Higgins, D.G.** (2011). Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. *Mol. Syst. Biol.* **7**: 539.
- Stadtmueller, B.M., Kish-Trier, E., Ferrell, K., Petersen, C.N., Robinson, H., Myszka, D.G., Eckert, D.M., Formosa, T., and Hill, C.P.** (2012). Structure of a proteasome Pba1-Pba2 complex: implications for proteasome assembly, activation, and biological function. *J. Biol. Chem.* **287**: 37371–82.

- Tyanova, S., Temu, T., Sinitcyn, P., Carlson, A., Hein, M.Y., Geiger, T., Mann, M., and Cox, J.** (2016). The Perseus computational platform for comprehensive analysis of (prote)omics data. *Nat. Methods*.
- Uechi, H., Hamazaki, J., and Murata, S.** (2014). Characterization of the testis-specific proteasome subunit alpha4s in mammals. *J. Biol. Chem.* **289**: 12365–74.
- Unverdorben, P., Beck, F., led, P., Schweitzer, A., Pfeifer, G., Plitzko, J.M., Baumeister, W., and Förster, F.** (2014). Deep classification of a large cryo-EM dataset defines the conformational landscape of the 26S proteasome. *Proc. Natl. Acad. Sci. U.S.A.* **111**: 5544–5549.
- Ustrell, V., Hoffman, L., Pratt, G., and Rechsteiner, M.** (2002). PA200, a nuclear proteasome activator involved in DNA repair. *EMBO J.* **21**: 3516–25.
- VanderLinden, R.T., Hemmis, C.W., Schmitt, B., Ndoja, A., Whitby, F.G., Robinson, H., Cohen, R.E., Yao, T., and Hill, C.P.** (2015). Structural basis for the activation and inhibition of the UCH37 deubiquitylase. *Mol. Cell* **57**: 901–11.
- Verma, R., Aravind, L., Oania, R., McDonald, W.H., Yates, J.R., Koonin, E.V., and Deshaies, R.J.** (2002). Role of the Rpn11 metalloprotease in deubiquitination and degradation by the 26S proteasome. *Science* **298**: 611–615.
- Vidal, M., Braun, P., Chen, E., Boeke, J.D., and Harlow, E.** (1996). Genetic characterization of a mammalian protein-protein interaction domain by using a yeast reverse two-hybrid system. *Proc. Natl. Acad. Sci. U.S.A.* **93**: 10321–6.
- Vierstra, R.D.** (2009). The ubiquitin-26S proteasome system at the nexus of plant biology. *Nat. Rev. Mol. Cell Biol.* **10**: 385–97.
- Wang, D., Kalume, D., Pickart, C., Pandey, A., and Cotter, R.J.** (2006). Identification of protein ubiquitylation by electrospray ionization tandem mass spectrometric analysis of sulfonated tryptic peptides. *Anal. Chem.* **78**: 3681–7.
- Wani, P.S., Suppahia, A., Capalla, X., Ondracek, A., and Roelofs, J.** (2016). Phosphorylation of the C-terminal tail of proteasome subunit alpha7 is required for binding of the proteasome quality control factor Ecm29. *Sci. Rep.* **6**: 27873.
- Waskom, M.** (2016). Seaborn.
- Welk, V., Coux, O., Kleene, V., Abeza, C., Trumbach, D., Eickelberg, O., and Meiners, S.** (2016). Inhibition of Proteasome Activity Induces Formation of Alternative Proteasome Complexes. *J. Biol. Chem.* **291**: 13147–59.

- Wenger, C.D. and Coon, J.J.** (2013). A proteomics search algorithm specifically designed for high-resolution tandem mass spectra. *J. Proteome Res.* **12**: 1377–86.
- Yang, B.J., Han, X.X., Yin, L.L., Xing, M.Q., Xu, Z.H., and Xue, H.W.** (2016). Arabidopsis PROTEASOME REGULATOR1 is required for auxin-mediated suppression of proteasome activity and regulates auxin signalling. *Nat. Commun.* **7**: 11388.
- Yang, P., Fu, H., Walker, J., Papa, C.M., Smalle, J., Ju, Y.M., and Vierstra, R.D.** (2004). Purification of the Arabidopsis 26 S proteasome: biochemical and molecular analyses revealed the presence of multiple isoforms. *J. Biol. Chem.* **279**: 6401–13.
- Yao, T. and Cohen, R.E.** (2002). A cryptic protease couples deubiquitination and degradation by the proteasome. *Nature* **419**: 403–7.
- Yashiroda, H., Mizushima, T., Okamoto, K., Kameyama, T., Hayashi, H., Kishimoto, T., Niwa, S., Kasahara, M., Kurimoto, E., Sakata, E., Takagi, K., Suzuki, A., Hirano, Y., Murata, S., Kato, K., Yamane, T., and Tanaka, K.** (2008). Crystal structure of a chaperone complex that contributes to the assembly of yeast 20S proteasomes. *Nat. Struct. Mol. Biol.* **15**: 228–36.
- Yu, Y., Yu, Y., Smith, M., and Pieper, R.** (2014). A spinnable and automatable StageTip for high throughput peptide desalting and proteomics. *Protocol Exchange*.
- Zhang, X., Henriques, R., Lin, S.S., Niu, Q.W., and Chua, N.H.** (2006). Agrobacterium-mediated transformation of *Arabidopsis thaliana* using the floral dip method. *Nat. Protoc.* **1**: 641–6.




Figure 3.S1: Sequence Alignments of RPT4a and RPT4b. Sequence Alignments of *Arabidopsis* RPT4a RPT4b and related sequences was performed with ClustalW. Positions of T-DNA insertion mutants for both RPT4a and RPT4b were determined by sequencing T-DNA specific PCR products and are shown with black triangles. The AAA-ATPase domain is indicated with a black bar. The Walker A and B motifs contained in the AAA-ATPase domain are indicated in red. Species names are abbreviated as follows: At - *Arabidopsis thaliana*, Pt - *Populus trichocarpa*, Os - *Oryza sativa*, Sb - *Sorghum bicolor*, Zm- *Zea mays*, Pp - *Physcomitrella patens*, Sm - *Selaginella moellendorffii*, Dm - *Drosophila melanogaster*, Sc- *Saccharomyces cerevisiae*.




Figure 3.S2: Volcano Plots Comparing Protein Profiles of Proteasome Affinity Purifications Based on RPT4a or RPT4b subunits in the presence (+) or absence (-) of ATP. (A) In the presence of ATP, statistical analysis using significant analysis of microarray (SAM) test does not show enrichment for RPT4a in the RPT4a pulldown and RPT4b in the RPT4b pulldown. (B) In the absence of ATP, statistical analysis using a SAM test does show enrichment for RPT4a in the RPT4a pulldown and RPT4b in the RPT4b pulldown, but shows no other associations with central proteasome subunits or associated proteins. This suggests the complex assembles in a random fashion with respect to isoform incorporation.




Figure 3.S3: Kernel Density Estimates (KDE) of the CP and RP distributions from label-free quantitative mass spectrometry show enrichment for the CP when purified via PAG1-FLAG, and for the RP when purified via FLAG-RPT4a/b. The X-axis shows difference in CP (red) or RP (blue) protein levels between affinity-purified samples and their respective mock affinity purification control, while the Y-axis is unit less with the area of each density corresponding to a total area of 1. Fold change (FC) CP over RP, or RP over CP was calculated based on the median of these distributions. Two *p*-values, one for a median test, and another for a Kolmogorov-Smirnov test, are shown, which test for differences in median and overall distribution respectively. Cryo-EM structures of the sub-complexes likely to be enriched in each affinity purification are shown, with free CP (A) and RP (C and E) in the absence of ATP, and in the presence of ATP half capped and doubly capped proteasomes (B, D, and F) adapted from (Beck *et al.*, 2012) **(A)** The CP is enriched over 20-fold when performing an affinity purification with ATP omitted from the purification buffer as compared to the RP. **(B)** The CP is only enriched 1.5-fold when affinity purified in the presence of 20 mM ATP. **(C)** The RP as compared to the CP is enriched approximately 3.5-fold in samples that were purified via FLAG-RPT4a in the absence of ATP. **(D)** In the presence of 20 mM ATP this RP enrichment decreases to only 2-fold for RPT4a. **(E)** In the absence of ATP the RP is enriched approximately 3.5-fold in samples affinity purified from RPT4b. **(F)** Enrichment for the RP decreases to about 2-fold in the presence of 20 mM ATP.

UMP1_a_A_t h - - - - - MES - - EKKI AHEI GGVKNDALRFGL - - H-
 UMP1_b_A_t h - - - - - MES - - QKKI AHEI GGMKNDALRFGL - - H-
 UMP1_c_A_t h - - - - -
 UMP1_Z_ma - - - - - MA - - SGS MKREI SET - HDTLRFGL - - NA
 UMP1_O_sa - - - - - MAS - - GGVVKKEI GGN - HDVLRFGV - - ND
 UMP1_M_tr - - - - - ME E - - SKTI PHQI GGVHTDVLRFGL - - P-
 UMP1_P_pa - - - - - MD - - GI QL PHDVLGS - HDSL LRHGL - - P-
 UMP1_S_ce - - - - - MNI VPQDTFKSQVSTDQ - - DKSVLSSAVPSL PDTL RQQEGGAV
 UMP1_K_la - - - - - MPMNI VPSPDFKS AVATDK - GCE HQS NAVASL PDVFR EQV - GAR
 MCS GETTI ASATPTQTAMNI VPPSNRRSALAA - - - GDATAASAAPALPDVLREQR - AAR
 UMP1_D_ha - - - - - MSLKI VPDNNQHSSI NSTK - F - GE PAS HAP GL QDI LKTQE - GPL
 UMP1_D_me - - - - - MY - - QPSLK VQPAE - - VSVL NAT GRVGMP - - T - E - A-
 UMP1_X_la - - - - - MN - - TKAL GS QLKDSI PI GE LSTH YYGI QDTL RLTGF - T-
 UMP1_D_re - - - - - MN - - TRGL RS QLKDS VPVTGL CPQAGP YGVQDSL RRGF - S-
 UMP1_M_mu - - - - - MN - - ARGL GSEL KDSI PVAELS - ASGPFES HDL LRKGF - S-
 UMP1_H_sa - - - - - MN - - ARGL GSEL KDSI PVTELS - ASGPFES HDL LRKGF - S-
 consensus + + + + +

UMP1_a_A_t h GVKS NI GS HPLE SSYE SE KKSK EA LKRTVI AHAY GT AL PL KMDM DRQI LS RF - - -
 UMP1_b_A_t h GVKS DI LRS HPLE TAYE SGKQS C E MKR RV I THTYGA AL PL KMDL DRQI LS RF - - -
 UMP1_c_A_t h - - - - - ME P APL PL KMDL DRP I L P RY GML DQYH
 UMP1_Z_ma GVKA DL APPHPL QSSI QE EAKF WADKKKF GTE A I YGSAL NI RKDL DAQI LS RF - - -
 UMP1_O_sa SVKGDL APPHPL QASVHKDAFKWADKKRFGAE A I YGSAF NI RKDL DAQI LS KF - - -
 UMP1_M_tr GVKS DI VGA HPLE LSSL QSVRGVE EAMRKQCKVNLYGA APL KE EL DRQI LS RF - - -
 UMP1_P_pa AFKS DVL PVHP VE VI QRSS KAMQA QT KRTL E RAYGS APL PMKMDI ERQI L TRF - - -
 UMP1_S_ce PLST QL NDRHPL EST LKNMETT QQR QME QYRQI FGGIAE PMKRTM E MEI VNRT - - -
 UMP1_K_la PLNT QL NDRHPL E S RVRN MDE T QH KRL E QYRQI FGGIAE PMKRV MEL KLV QNT - - -
 UMP1_E_go PLA AEL ADRHPL E S RL RDWDACA QRT RLE QYRRI FGGIAE PVRRTM E L AL VS DA - - -
 UMP1_D_ha NLA SKI NNRHPL E GRI S NME E T QY QTR L E TY RRV F GAGE P I KRTM E L EL VE AT - - -
 UMP1_D_me NCL NQL AHV HRL RDSE L NYNE HQY NR NMQML RNHE GL GPV L KMGM E RFA AAR QV - - -
 UMP1_X_la SVQSE L MPSPHPL E LSE KNF QI NQEKVNFS T VRN I QGL HAPL KL HME F KAVKQV - - -
 UMP1_D_re SVK NELL PS HPLE LSE KNF QL NQDK MNF NT L RNI QGL HAPL KL QME YRAAK QI - - -
 UMP1_M_mu CVK NELL PS HPLE LSE KNF QL NQDK MNF ST L RNI QGL F APL KL QME F KAVQV - - -
 UMP1_H_sa CVK NELL PS HPLE LSE KNF QL NQDK MNF ST L RNI QGL F APL KL QME F KAVQV - - -
 consensus + + + + +

UMP1_a_A_t h - - - - - QRPPGP - I PSSM GL E VYT GAV DDF GFE DY L NDPR DS E T - FKP
 UMP1_b_A_t h - - - - - QRPPGP - I PSSM GL E VYT GAL DNF GFE DY L NDPR RE SET - LKP
 UMP1_c_A_t h YE I PLNS DAYS A I RYE NRP RTGP - I PSL MLD E VYT GAI DDF DFE DY L NVST P QI L - ENP
 UMP1_Z_ma - - - - - QRPPGA - L PSS L LG E ALT G S L DDF GFE DY L NMPQDSEG - FRQ
 UMP1_O_sa - - - - - QRPPGA - L PSS M L GYE ALT G S L DDF GFE DY L NL P QDSES - FRA
 UMP1_M_tr - - - - - QRPPGV - I PSS M L GLE VT G T L DHF GFE DY L NDS R E SET - F RP
 UMP1_P_pa - - - - - QRPT GL - I PSS M L GLE S LT G E L D E F GFE DFL NDPK E SDT - F VP
 UMP1_S_ce - - - - - DF NPL - - STNGS I HRDI L L NKE CS I DWE DVYPGTGL QAS T MVG
 UMP1_K_la - - - - - DF NPL - - NO S NLH KDV L MNK E AS I DWE DVY PTS DF AS GMM
 UMP1_E_go - - - - - EF APL - - GA AAL H L DV N NTE AS L DWHDV Y P D E QQ - - -
 UMP1_D_ha - - - - - DF KP QLL LGGS DS MHKDVL L NKDAS VDW E D I YRGGL E SGS - - HV
 UMP1_D_me - - - - - GRL P - F L S S SNF MDDV L T GRC DS I GFE DF MNL P E NS EH - MRQ
 UMP1_X_la - - - - - QRL P - F L P I S N I A L D T L RGT D E C I S F E D I L N D P S Q S E L - MGE
 UMP1_D_re - - - - - QRL P - F L P S S N L A L D T L R G S D D T I GFE D I L N D P V Q C E M - MGD
 UMP1_M_mu - - - - - HRL P - F L P S S N L S D L R G N D E T I GFE D I L N D P S Q S E L - MGE
 UMP1_H_sa - - - - - QRL P - F L S S S N L S D L R G N D E T I GFE D I L N D P S Q S E V - MGE
 consensus + + + + +

UMP1_a_A_t h VDF - - - - - HHGME VRL GL S KGPI - - - - - YPS - - FM - - -
 UMP1_b_A_t h VDF - - - - - HHGME VRL GL S KGPA - - - - - SPS - - FM - - -
 UMP1_c_A_t h RCS NQL T S T T S WKL GL VSPKAQL NQVPCRS KQFI LYNAI KVL KL DST CMF VQP NAGE MI
 UMP1_Z_ma PDM - - - - - HHGME VRL GL S KGPI - - - - - CPS - - FS - - -
 UMP1_O_sa PDM - - - - - HHGME VRL GL S KGPV - - - - - CPS - - FN - - -
 UMP1_M_tr LDM - - - - - HHGME VRL GL S KGPV - - - - - YPS - - II - - -
 UMP1_P_pa RDM - - - - - HES M VRL GL A KGPV - - - - - SRM - - FF - - -
 UMP1_S_ce DDV - - - - - HS K I E K QL G I - - - - -
 UMP1_K_la DDV - - - - - HT K I E K RM G I - - - - -
 UMP1_E_go P D V - - - - - H A I E R S L R L - - - - -
 UMP1_D_ha QDF - - - - - H T E M E K K M G M - - - - -
 UMP1_D_me P - - - - - H A V V E K S L G I Y Q - - - - -
 UMP1_X_la P - - - - - H L M V E Y K L G L L - - - - -
 UMP1_D_re P - - - - - H I M T E Y K L G L L - - - - -
 UMP1_M_mu P - - - - - H V M V E H K L G L L - - - - -
 UMP1_H_sa P - - - - - H L M V E Y K L G L L - - - - -
 consensus + + + +

Figure 3.S4: Multiple Sequence Alignment for UMP1.

[Figure 3.S4 caption follows on next page]

UMP1_a_A. th	- - - - -
UMP1_b_A. th	- - - - -
UMP1_c_A. th	L AAP SSSG T KTL V E F N N K S A E T E P S D V K L G L D L V S S M K E E L D L G R N A P K Y F R V D F S P K
UMP1_Z. ma	- - - - -
UMP1_O. sa	- - - - -
UMP1_M. tr	- - - - -
UMP1_P. pa	- - - - -
UMP1_S. ce	- - - - -
UMP1_K. la	- - - - -
UMP1_E. go	- - - - -
UMP1_D. ha	- - - - -
UMP1_D. me	- - - - -
UMP1_X. la	- - - - -
UMP1_D. re	- - - - -
UMP1_M. mu	- - - - -
UMP1_H. sa	- - - - -
consensus
UMP1_a_A. th	- - - - -
UMP1_b_A. th	- - - - -
UMP1_c_A. th	D H T V C L E F G I L S R E F N R A H L R D G D K E K F T L K N V H K S K Q G E E A S L P T L I N L N K R D I N
UMP1_Z. ma	- - - - -
UMP1_O. sa	- - - - -
UMP1_M. tr	- - - - -
UMP1_P. pa	- - - - -
UMP1_S. ce	- - - - -
UMP1_K. la	- - - - -
UMP1_E. go	- - - - -
UMP1_D. ha	- - - - -
UMP1_D. me	- - - - -
UMP1_X. la	- - - - -
UMP1_D. re	- - - - -
UMP1_M. mu	- - - - -
UMP1_H. sa	- - - - -
consensus




Figure 3.S4 (preceding page): Multiple Sequence Alignment for UMP1. A Clustal Omega (Sievers and Higgins, 2014) sequence alignment of UMP1 from plants, yeast, and animals visualized with the program BOXSHADE. A heat map showing each combination of protein percent identity with respective values shown in each cell obtained from the resultant Clustal Omega identity matrix. Species names are abbreviated as follows: A.th - *Arabidopsis thaliana*, Z.ma - *Zea mays*, Os - *Oryza sativa*, M.tr - *Medicago truncatula*, P.pa - *Physcomitrella patens*, S.ec- *Saccharomyces cerevisiae* K.la - *Kluyveromyces lactis*, E.go - *Eremothecium gossypii*, D.ha - *Debaryomyces hansenii*, D.me - *Drosophila melanogaster*, X.la - *Xenopus laevis*, D.re - *Danio rerio*, M.mu - *Mus musculus*, H.sa, *Homo sapiens*.

PBAC1_A.th - MEDVL-TENPPPSRFQEDLNN-FVQPPES-LPSPFIIFSNP
 PBAC1_Z.ma - MEDVI-TDVPPPSRFSPDDLNN-FAAPPQAQ-PTPIFVVSPN
 PBAC1_O.sa - MEDVV-TDAPPPSRSFSPDDLNN-FAAPPQ-PTPILVVSPA
 PBAC1_M.tr - MEDVI-TEAAPPSSRLLEEDLN-FTPSSPP-LPSPFLLFPHT
 PBAC1_P.pa MAGMPDP-LTDLPPPSSRFDPDELNA-FAASDVQ-PSPLIVWAPWAKT
 PBAC1_S.ce - MLFKQW-NDLPEPKHLLDLPEISKNL-QSLEVCPVPKVEF-PQ
 PBAC1_K.la - MLTFKQW-QNDQVPRHQI DPVVQSDP-YSLVNKTTPTVTI EPE
 PBAC1_E.go - MI F KQW- QSERLPRHQ DCL GPMDP-HSLLAGAEPOVEFSGK
 PBAC1_D.ha - MLIKPW-NEI PAPRHTLDDEVDDA-IPSPINLVELN
 PBAC1_D.re - MSCPGFGEELNI PSSRAFWDDCDEDELE-NLKPVE-KLQLKFDDDC
 PBAC1_X.la - MATFFGEVQS VFSRAVDEEEED-E DDDEEVEEEEDREI I AEELERKREV RVTWNPELTA
 PBAC1_D.re - MATFFGEVLS VYSRAVEEDEYE-DM TNNEEDEQI RREIEEKRSVEVCVLLG
 PBAC1_M.mu - MAATFFGEVVKAPCRAGTEEEEQQSRRDTPE DREV RQLARKREV RLLRRQT-ET
 PBAC1_H.sa - MAATFFGEVVKAPCRAGTEDEEEEEGRRETPE DREV RLQLARKREV RLLRRQT-KT
 consensus .+.++..+.++..+++.+++++.+.+.+++.+.++.+.+...+.

PBAC1_A.th K---PELPLRPSLLIAIASSPSIYIFH-SLPS-KTLLGSILMPEVPF
 PBAC1_Z.ma - PSPPA-PRLLVVFI SPTSLAL-ASPPPLASLLPDPL
 PBAC1_O.sa - NPSPAPRRLLLI SPTSLAL-HLPSPPPLHASLLPELPL
 PBAC1_M.tr Q---QQQPLKPTLLLIAIASSPSI SLSLFH-NLLTPQTLTSSLLPELPL
 PBAC1_P.pa L---LQS GVL QPQL ALFCVSDVGHLH-QI PN-KRIVGSI VLPEISM
 PBAC1_S.ce - DL DVPQYSTAVTTKI MNPLFPKNNQL-TSI GEIKTTLTVKSPSLPQ-S
 PBAC1_K.la - SLDI SPYNHCI VLPTSLKWL LPSI I NL-VHI GI ISGTL NMNEWESMKP-V
 PBAC1_E.go - LGERYS AAL VVPSTLSW MPP EMVQL-TPV AQL VGT QAGGGECATR-LV
 PBAC1_D.ha - QNI TS MDYI VSPKTLKV L V-ED-LPNGKAVGYI NI EYPQLADL QT ND
 PBAC1_D.re - GEAAVPESEMLVVI EGVNVTHF-ASSLLEKDAKR
 PBAC1_X.la AI ESSPGKRLPCSSVILSVGDNATGF-VSSYI LSSGSWEVA
 PBAC1_D.re - SQDGPLQCSDLII GAGPNASGF-I RAYLSTVGWTAV
 PBAC1_M.mu SLEAVLLET HPCS KFII AVGS NATAF-LSAFVMNSGVWEV
 PBAC1_H.sa SLEVSLLKE YPCSKFII AIGNNAAVF-LSFVMNSGVWEV
 consensus .+.+++.+.++..+++.++.+.+.+.++.+.+.++.+.+...+.

PBAC1_A.th - SG-NT-MEPSLEDKSCNI YSLSSADDKNLTLLV
 PBAC1_Z.ma - LP-H-AP-MRVYLH-PSGALLAAA
 PBAC1_O.sa - GR-SQ-PP-I RVYHLH-PSGTLAAA
 PBAC1_M.tr - SY-PN-NT-I NI HSLS-STI LLA
 PBAC1_P.pa - VN-NE-I EPSPHNKECYLYAVD-SQASVVFA
 PBAC1_S.ce SGKHSWNYDENFPNEVDPDQKNDT ADETVYGFSPYIYS-FGKTLFSMEE-NFI
 PBAC1_K.la - PGMYDTEEHMI N-STEQFVIHYNLWR-LEDTLIAID-NEV
 PBAC1_E.go AADQFYEAD-VPQ-AERHVRVAYQLYERGPQQLVVVEE-NDL
 PBAC1_D.ha EDDDVYDEDEQLYTAV-MRKSLLDKYGKLKIPIQVHDDRYLSVNI P
 PBAC1_D.re - VC-AI PANNSTLHMSHSKQL-LAI INED-L
 PBAC1_X.la GSVTLWNERCRDCNVRK-DFLPAPSSCTFYRSITDPT-VLLC
 PBAC1_D.re AWLSF WNERSRG SERPT-VVPGPGEPS CSVLYRLIESCPT-VLIC
 PBAC1_M.mu GCAKLWNEWCRTTDTV-RLSPTDVF CVFYQLKSDPS-VFLC
 PBAC1_H.sa GCAKLWNEWCRTTDTT-HLSSTEAFCVFYHLKSNPS-VFLC
 consensus .+.+.+.+.+.+.+.+.+.+.+.+

PBAC1_A.th SVQLPVSPERSNLVSRLLI AKDI-I PERVI ILDSI QS RNF RGRLS-PDETLAAKDE
 PBAC1_Z.ma HGA-FPAHRARAAKALV-SRL-QPEEV L DAVRS GTYRGR A-ADEPVEGKLE
 PBAC1_O.sa HGPAPVPAHRARAVARSLV-SAL-QPEEV L DAVRS GAYRGR A-ADEPVEGKLE
 PBAC1_M.tr AVQNPI PDNRAYA VAEI ILLS NQI-CPDSVITLDSI QPMNRG L S-SDEAVAFKLE
 PBAC1_P.pa SI QYAI PPESATI WT KSLF-QAI-RPERVILMDGI STL HFRGKLS-ADEPLLFLLE
 PBAC1_S.ce SISPI FG-NM-I SRSII SQLAQFSPDIIVL GTS-DK-I-A
 PBAC1_K.la KYPAI VT-NF-MTRNL AQL RDI K-DI TI LVNS-DK-I-T
 PBAC1_E.go KYPPV VH-NR-LAQKLA EKL RAVGAETTVLVNS-DK-S-T
 PBAC1_D.ha HFPNI I ANNI-LAQKL TEEL DKT I NKAW I LSPSLISSNET-I-NKLEVDFDM
 PBAC1_D.re TSSGEVAEL-LLPY-VK-LAKKV TLT LKPKVEFKSEDI QLYRDHI AVVRGIG
 PBAC1_X.la QCNCHVVAE DQL F QWCEKVF GS LEK-SSLKVTVL STCPVSEYKTPEST-YSLPVPLKA
 PBAC1_D.re QCQGFVAE DQL F QFTEKVF SCVQT-RDL NVT I L SDCSSADYKTS DYL-SGSSTPFLRCLK
 PBAC1_M.mu QCSCYI AEDQQF QWL EKVFGF QPR-KS MQVTVLTCRHI TDYKTPEST-CSLSSPFLRALK
 PBAC1_H.sa QCSCYVAE DQQY QWL EKVFGS CPR-KNMQITLTCRHTD YKTS EST-GSLPSPFRLALK
 consensus .+

Figure 3.S5: Multiple Sequence Alignment for PBAC1.

[Figure 3.S5 caption follows on next page]

PBAC1_A.th	T S A E K K A T T - H T G L V N L D Y F P S G S V I D G L I S A S L L S R C Q L K N I R G T L V I S W - P E - - - - F
PBAC1_Z.ma	T R A A R A R G G V G A A K V Q A L V P P G S V V D G L G A A V L A E Q E I R G K A A S M V V T W - P A - - - - D
PBAC1_O.sa	T R A A R A R G G V G A A R G V A A L A P P G S V V D G L G A A V L A E Q E I M R G K A A S M V V T W - P E - - - - G
PBAC1_M.tr	S F A E R K K A N E E K L L G G L D Y Y P S G S V V D G L G A A V L L G R C Q I L K I R A S L C V T W - P Q - - - - F
PBAC1_P.pa	T S A S K K G K K - - N K D S I A P Y Y P S G S L V D G V A A A L I T H Q C Q L R A L R A E L L M S W - P T - - - - A
PBAC1_S.ce	- - - - - S M K V M T E N E C T L Q P P E F I T G F I G S V L T Q L I V G P S K G L K F K C L V A P S E G P N G F
PBAC1_K.la	- - - - - E L K Q L S - - - Y L V P P E F I T G S I T N F I L A L - - - P - R K E L K V I V V P S E G P I G F
PBAC1_E.go	- - - - - G V K D L E - - - E L V P P E F I T G A L A A L V C A L - - - A P S G A G R V L V V Q S E G P H G F
PBAC1_D.ha	- - - - - Q T P D I Y A A I P S L K P P H F I T G I G A S F N S Q I S L L R - - Q P R L M S L V L R S E G Q S G F
PBAC1_D.me	- - - - - A N Q K D I T E L E A P N F I A G V A A G V A S W R D Q M E L P V T S F V I Y - - T D - - - - K
PBAC1_X.la	T S E Y - - - - R E E V P C P L L E Q P N I V D G L P A A V L S H C Q V L G I P A V F Y R C Y - - T D - - - - I
PBAC1_D.re	T S T Y - - - - T H T V T C P P L E Q P N I C S G L A A A V L S H C Q V H Q I S A V L Y Q C Y - - S D - - - - V
PBAC1_M.mu	T Q T F - - - - K D A L C C P L L E Q P N I V H D L I S A A V L S Y C O V V K I P A V L Y L C Y - - T D - - - - V
PBAC1_H.sa	T Q N F - - - - K D S A C C P L L E Q P N I V H D L P A A V L S Y C O V V K I P A I L Y L C Y - - T D - - - - V
consensus	+ + . + . + + . + + + + + + . + + . + . + . + . + +
PBAC1_A.th	G P S V I R F A G A L L K K I V P S I D V A S V N K D L E K N - - S S R T G L K K - D T W L - - D S D L Y T
PBAC1_Z.ma	S R P A E F - - - G V M R R V A A E L G V D T G K - - - - - A - A A R V S - G R P D - - L D S L Y T
PBAC1_O.sa	A R P A E F - - - G V M R R V A A E L G V D P A K - - - - - A A A G R V A - G R A E - - L D A L Y T
PBAC1_M.tr	D S D V V L L L K D L L R G L G E F - E F G F S D - - D E - - V F K F G R S K - D H V F - - Q S H L Y I
PBAC1_P.pa	D T S V V N Q L A S A L Q N V C Q Q A - - L A N N V I I D F - - - I A A A K S I A - G R R L L S D S E I Y M
PBAC1_S.ce	E K I L S L S D M G S L V D L C G Q W I G F - - E - - P S R Y S E E C - Y R L W R C D S A A I G A Q S G L Y I
PBAC1_K.la	E K Y N F P T V D A L V D E M S K L L C S Y P S Q - - T E Y Y V S E C - L K L W K L D E C T - T T Q G G L Y I
PBAC1_E.go	E K H N L T V V D A L V E R V R Q I L D - - - Q - - P P E Y V A A C - T R H W K L Q A G S - Q F Q G G L Y L
PBAC1_D.ha	E K I D T D A F V D A S F V L N E L L V G T S D Q - - G - N Y L K Q V S S T V R K I N G - - - Y S N S I G M Y I
PBAC1_D.me	L P L D - - - A T A A Q P V I K V L K A A G V A - C S S Y I P P R K E S - - - - - S Y L Y M
PBAC1_X.la	S K L D - - - S V T I K A F R P L L S S G S L S - R L A A D S A N I Q E T L R K T - V K L N E I Q S N L Y I
PBAC1_D.re	L H P D - - - S A S M Q M F A A T L S S V - - - L K L E Q S L S A D V L Q R V - T R V S E I Q S N L Y T
PBAC1_M.mu	M K L D - - - R V T V E A F K P L L S S R S L K - C L V K N I P E S T E I L K K L - M T T N E I Q S N I Y T
PBAC1_H.sa	M K L D - - - L I T V E A F K P I L S T R S L K - G L V K N I P Q S T E I L K K L - M T T N E I Q S N I Y T
consensus	. + + + . + . + . + + + + + + . + * +




Figure 3.S5 (preceding page): Multiple Sequence Alignment for PBAC1. A Clustal Omega (Sievers and Higgins, 2014) sequence alignment of PBAC1 from plants, yeast, and animals visualized with the program BOXSHADE. A heat map showing each combination of protein percent identity with respective values shown in each cell obtained from the resultant Clustal Omega identity matrix. Species names are abbreviated as follows: A.th - *Arabidopsis thaliana*, Z.ma - *Zea mays*, Os - *Oryza sativa*, M.tr - *Medicago truncatula*, P.pa - *Physcomitrella patens*, S.ec- *Saccharomyces cerevisiae* K.la - *Kluyveromyces lactis*, E.go - *Eremothecium gossypii*, D.ha - *Debaryomyces hansenii*, D.me - *Drosophila melanogaster*, X.la - *Xenopus laevis*, D.re - *Danio rerio*, M.mu - *Mus musculus*, H.sa, *Homo sapiens*.

PBAC2_A.t h MEFVV-----EEG----KLVNE DCSTL VLPALS|GNVGQL AVDL LVSS T--GAER RVGY LDD
 PBAC2_Z.nz MEFAL-----VES---ESFSPSCPTL|MPALS|GNVGQL AVDL LI SSS--RARRVAYL DE
 PBAC2_O.sa MEHAV-----VEG---ESFSPDCSTL|MPALS|GNVGQL AVDL LVSS--RARRVAYL DE
 PBAC2_M.tr MKEIP-----EEG----KHLHE DCSTL|LPAVS|GNVGQL ADL LVSSM--GSEKVGY LDD
 PBAC2_P.pa MDFVE-----LEG---GE GHL QHTGSL|VPALS|GNAGQL AVDL LI S-H-GACKSGYL DE
 PBAC2_S.ce -----MSCL VLPPLVSVGN|PQLSI DWLL NSQANEWE YLEALDS
 PBAC2_K.la -----MSTLL PLVSTGNVSQLCVDLVL HSEDEFTFVKE LDS
 PBAC2_E.go MS HLT GAERS RSDS WWP HGRQES MATL|LPLVSA GNVPQL CADVL HSTPGE FRFVRE LDG
 PBAC2_D.ha MSEDTKVF NSL PG-VI LSDLADSTL|PSI SI GNI PQL ANDL ITTL--NF NKT GSL TD
 PBAC2_D.re MLFLK-----DK--RTTLDVAEHTV|PSI CVGNA AOL ACDLL I ASK--EL RRI GSITD
 PBAC2_X.la -MFVP-----VG-SEQDFPSVSTLL|LPAIS VGVNVGQL AIDLI STL--NI PKVGYF YT
 PBAC2_D.re -MFIS-----SG-DAVPSFKGFTL|LPAVS VGVNVGQL AVDL LI STL--NM PRVGVY FHT
 PBAC2_M.mu -MFVP-----CG-ES VPDL TNFTL|LPAVS VGVNVGQL AIDL II STL--NM CKI GYF YT
 PBAC2_H.sa -MFVP-----CG-ES APDL AGFTL|LPAVS VGVNVGQL AMDLI STL-NMS KI GYF YT
 consensus +.++.....+....++++*+...+**+...+...+....+++++.

PBAC2_A.t h PNLL PCVGNDAY GPL-----P-CGEIALP LEVYE SSS ATT LAQQRSPVA-KGMM
 PBAC2_Z.nz PSVL PCAGND AF GPD-----A-VGDL ALAEAYESTSHKLGF|QQRSPV-T-TGMV
 PBAC2_O.sa PSVL PCAGND AF GPD-----A-VGDL ALAEAYESTSHKLGF|QQRSPV-T-TGMV
 PBAC2_M.tr PYVL PCVGNDAY GFP-----P-QGDL ALPEAYEDSPS NGLTVI|QQRSPV-I-KGMM
 PBAC2_P.pa PHVL PCVGNDPFGPS-----A-NGDL AVAL E VYE DAEI LASI|QQRSPV-I-KGT M
 PBAC2_S.ce KYLVEFVGPL DRPEDGS DSYKDADM--KYSS AL E VFY NKKRGL FAI|QORTPLVSVNYL
 PBAC2_K.la IWLHS F|GPLDFVEGSSKPLYHDCKGS KKQFTTPLELY FYNRRL DLYVI|QQRSPV-I-AGYE
 PBAC2_E.go QWLHPEVGPLDYVEGQESLLYRDPAGK-RVTTPELY F YCEQRLQLY|QQRSPV-V-QGYE
 PBAC2_D.ha TYLYPFASPI DTS ND-----VGQKKGI STAVEVYHNKNL KLTI|QQRSPV-I-PSFT
 PBAC2_D.re PALI PVYGPSSAYQHE-----P-KEKVS S CELYEGTEDKL LVVQFRAPW-ARHT
 PBAC2_X.la DCLVPMVGNNPYETN-----E ENAKELCT NAEVY AL P SQKL AVL QL RSL VI-KKKS
 PBAC2_D.re DCLIPMAGNNPYASS-----TE DAAQLS TS AEVY S HRDL KLA VL QI RAPI|QT KV
 PBAC2_M.mu DCLVPMVGNNPYATE-----EENS NELSI NT E VY S L P SKKL LVVL QL RST F|KY KS
 PBAC2_H.sa DCLVPMVGNNPYATT-----EGNSTELSI NAEVY S L P SRKL VAL QL RSI F|KY KS
 consensus ..+++. ++...++....++*.+...+.++*.++*+....++....++....++....++....

PBAC2_A.t h I KFAEN-IA NF AASS GKKH|VLSSLD F QRL HNL DMS RG P QVYY|LSNAES DGR DDHC ER
 PBAC2_Z.nz VSFAKN-VADFI INSI GKNHI|VIISSLD DS GKRRI DASSDMQVYY|LSSCNE DGS DPE YEK
 PBAC2_O.sa VSFAKN-VANF ISSI EKDH|VII LSSLD DS GKRRI I DASSDMQVYY|LSSCNE DGS DPK CEN
 PBAC2_M.tr LEFAKN-MADFLAGS GKKH|I LSSLD FG WKQKVDMSS GLOI YY|LSSANS NGADE NCE Q
 PBAC2_P.pa MKFSKN-LATWAAQE GVKE|VII LSGL DS GKVRQ SE-TNAMQI QY|T STANE DGS DDR CEQ
 PBAC2_S.ce NNFI VEII I LPFLSKYNI SEI CI WDSL YAME DE NGVI VRP-QEV YSL GE FYF---DDE AE
 PBAC2_K.la NQFFKEVI I PLIKSYQI ES VI VLDSAGAL DPSVPFSN ANY OHQYI ATAKCQLES LDNI TLQ
 PBAC2_E.go NEFCKEVLLP LLEELAP QRVVLL|DS VGE FETE VPVHETMESRSFSVATCNVTSI SDVAKE
 PBAC2_D.ha KTYYNEVI LPFI KEAKVKRF|LID DS DAGL VEN-----LSPGTI DFYS NEDLL N K
 PBAC2_D.re AHFOKE- LVELLK-- GARRVVI|LSSGFG EKF KRVI E-E SPWAYR-A SE NF--KEAHAAQ
 PBAC2_X.la KSF RQA-LV SW KRCAF ARVI|FLSS SCHAY HR DDKQL FGT P F RYL-VTPAL QKS VADVL SE
 PBAC2_D.re KSF RKL-MI SW KSSGF LRTV LLSSAHAY QR DDQQL HGT P L RYM|LSPS LE KEER QRF E E
 PBAC2_M.mu KSF CEK-LLAWV ESSGCARI|V LSSS HS YHRNDAQL RST P F RYL-LTPCL QKS VQN KI K S
 PBAC2_H.sa KPF CEK-LLS WVK SS GCA RVI|V LSSS HS YQR NDL QL RST P F RYL|LTPSM QKS VQN KI K S
 consensus ..+.+.++..++..+...+.+.++....++....++....++....++....++....++....

PBAC2_A.t h L-----GFGRL HEYDS E GRCW KY|LSSV F E KNS V--E E L AL P S E D E L E D I D Y Y P S L P-F A A L
 PBAC2_Z.nz L-----GWKKL E D YDPS S QKR WNC|LASL VEG GGL-S E D L D D D T D E M I N D Y Y A S L P-F A A L
 PBAC2_O.sa L-----GWKKL E E YDPS S QKR WKC|LASL VEG GGH-S E D M T G D P E E M T I N D Y Y S S L P-F A A L
 PBAC2_M.tr L-----GWKKL Q E YDPS S QKH WKY|LNDL AEGNAT-P E D T T S I E D E L E E N Y Y A S L P-F A A L
 PBAC2_P.pa L-----GW KRL HHYL PSS E A W QV|LDHQAMS-A EDSL Y S E I P Q S D D L Y F P R Q P-F A A L
 PBAC2_S.ce L-----S NL H-----LN-D-Q E S M V N N W L H F T P T S F Q D K I S V D Q P I F K I L
 PBAC2_K.la F Q-----TRL S-----I Q-DDE-H S I V N Q V F H F D T A S F Q Q N F S A D Q P V F K L F
 PBAC2_E.go F A-----E GL Q-----LD-E Q N D L S V N S T L F S F R Q G S F Q R G I S T E Q F I F K F S
 PBAC2_D.ha S LET L NI GOVDAMQ-----LG-KGNDHSS-Y-----I R S L I
 PBAC2_D.re L-----G NEE ELI K W K E H-----K G D A I Y G G G N A L Q
 PBAC2_X.la L-----E W K E M E K V S S Y P E-----LN-----D N E K K L F I P G G G F T K A F
 PBAC2_D.re L-----G W R E M E K I S V F P G-----I S-----D S E Q R L Y I P G G G V T K C L
 PBAC2_M.mu L-----N W L E M E K S R C I P E-----M S-----D S E F C I R I P G G G I T K T L
 PBAC2_H.sa L-----N W E M E K S R C I P E-----I D-----D S E F C I R I P G G G I T K T L
 consensus +.+.++..++..+...+.+.++....++....++....++....++....++....++....++....++....

Figure 3.S6: Multiple Sequence Alignment for PBAC2.

[Figure 3.S6 caption follows on next page]

PBAC2_A.th	FSAF-----KARGLKMTCLLCYCCSEGDNIPPEAFILLAEA-----SSKLNGLT	TPDKFH--GEEG
PBAC2_Z.mz	FSAC-----KAKGLKVSCVLCYCSEGDNMPESFQLAEA-----ACKL	LHGPEKFH--GNGS
PBAC2_O.sa	FSAC-----KAKGLKVTCVLCYCSEGDNMPESFQLAEA-----ACKL	LHGPEKFH--GNGS
PBAC2_M.tr	FSFL-----KAKGLKVTCVLCYCSEGDNTSDAFQLADA-----ACKLLRLSH--PNS--GI EG	G
PBAC2_P.pa	LACC-----KAQGLKVVCTLCFCAGDNVPDAFLIADG-----LHRLL	CHEKKEPQ--GSG
PBAC2_S.ce	FQILNAS-QRPKALRSIKYCSCLANE	GDNSLDSQQFLQWIISQKVIKNAPP-----I
PBAC2_K.la	HHHLLYQ---SDLNLKSTYLSKVHE	GDNSWDAFQICEK--LAEI
PBAC2_E.go	YHLLHSPLARSADFELS	YVSIFIHE
PBAC2_D.ha	TYLHTGDLSS--DFEVNV	LVSYYVE
PBAC2_D.me	FKAF-----DEQKVPVML	LFRYLL
PBAC2_X.la	YND-----CLEDLQMAVVL	KFC
PBAC2_D.re	YTDC-----CTEDVPMAVVL	I
PBAC2_M.mu	YDES-----CSKEI	QMAVVLKFV
PBAC2_H.sa	YDES-----CSKEI	QMAVVLKFV
consensus	+
PBAC2_A.th	GKWOI	PYSWKS
PBAC2_Z.mz	NGWI	VPLSWKS
PBAC2_O.sa	NGWTI	PLSWKS
PBAC2_M.tr	GKWR	PLSWMS
PBAC2_P.pa	VTWKI	PLSWTT
PBAC2_S.ce	VKFVRPI	SWQGAYGMADARDK
PBAC2_K.la	GRLTAPV	SWKGVY
PBAC2_E.go	ETLRTPI	SWKGVYGS
PBAC2_D.ha	KNW	RPRSLW
PBAC2_D.me	FKLTVPKS	WLGF
PBAC2_X.la	VKWKS	PRS
PBAC2_D.re	VQWRVPPS	WKLLFGS
PBAC2_M.mu	LPWKI	PSS
PBAC2_H.sa	SRWKI	PSS
consensus	. +++++*	+***+. +**. +.




Figure 3.S6 (preceding page): Multiple Sequence Alignment for PBAC2. A Clustal Omega (Sievers and Higgins, 2014) sequence alignment of PBAC2 from plants, yeast, and animals visualized with the program BOXSHADE. A heat map showing each combination of protein percent identity with respective values shown in each cell obtained from the resultant Clustal Omega identity matrix. Species names are abbreviated as follows: A.th - *Arabidopsis thaliana*, Z.ma - *Zea mays*, Os - *Oryza sativa*, M.tr - *Medicago truncatula*, P.pa - *Physcomitrella patens*, S.ec- *Saccharomyces cerevisiae* K.la - *Kluyveromyces lactis*, E.go - *Eremothecium gossypii*, D.ha - *Debaryomyces hansenii*, D.me - *Drosophila melanogaster*, X.la - *Xenopus laevis*, D.re - *Danio rerio*, M.mu - *Mus musculus*, H.sa, *Homo sapiens*.

PBAC3_A.th - - - - - MESLDTNFPVRHRK- - - - - VSFESKGNKTEI VI CSYEDHIL- - - - I
 PBAC3_Z.ma MEVPMNTTQS GPHF PVQHRS- - - - - LSLDI KGNKTDI VI NKYEDTFL- - - - V
 PBAC3_O.sa ME MGMNS VQPNAQFPVPHKS- - - - - LSLDI NGNKTDI VI SKYEDNFM- - - - V
 PBAC3_M.tr MNI DDNVASLTPKFPMQLN- - - - - FSI EI KGNKTEI II SSYEDHFM- - - - V
 PBAC3_P.pa - - - - - MASPLTFPMKSKQ- - - - - FVTSL QGLKTEVVQFSYDDHIMSQI LAFAFEI
 PBAC3_S.ce - - - - - MI SYEFQTHL PKGKDSLNASSENKELYVQATHFNNTIL- - - - L
 PBAC3_K.la - - - - - MRFI EFOKQDPEVFSDF- - - - - PGELGI FANHYDNKIV- - - - L
 PBAC3_E.go - - - - - MI YRELQKAIPPALWG-L- DESAQNELELTAVHFTNKTL- - - - L
 PBAC3_D.ha - - - - - MESTPEFSKSI TAT- - - F-PGDEENEFTFHKEELNDKII- - - - L
 PBAC3_D.re - - - - - MNHTANKPRQI GF DRO- - - - - FSADVNQGPTEIVFHCPFAKKWF- - - - L
 PBAC3_X.la - - - - - MPVTDTKPAVVSQKQ- - - - - AEVVI NGVATQVVVCASFNYI F- - - - V
 PBAC3_D.re - - - - - MAPQPVI RSKQ- - - - - AEKTI NGNOTQVVVCTEFSNYI F- - - - I
 PBAC3_M.mu - - - - - MEDKPLVVSQKQ- - - - - KTEVVCGVPTQVVCTAFSSHL- - - - V
 PBAC3_H.sa - - - - - MEDTPLVLSQKQ- - - - - KTEVVCGVPTQVVCTAFSSHL- - - - V
 consensus + . . + + . . + . + . + . + + +

PBAC3_A.th | ATQ|GAMGT|LHARKE- EGMSVE- - - - - PTFSVSVI
 PBAC3_Z.ma | VTQ|GCMGT|LAAKKD- ESVFSD- - - - - PTYNVSVL
 PBAC3_O.sa | MVTQ|GCMGT|LAARKD- ESVFSD- - - - - PTYNVSVL
 PBAC3_M.tr | VATQ|GAMGT|LHARKE- EGMSI S- - - - - PTFNVSVL
 PBAC3_P.pa | AVTQ|GKMGTL|LHARKE- EGYGTG- - - - - PTFNISITL
 PBAC3_S.ce | QIRLNGEMDSTYEVS SKGLNPILDI NVPLAGNL GNTGGDYDDEEEE FVRDHLSDYQVVTK
 PBAC3_K.la | QLRVNGEMDSTYEVKSKGLNDTEARFLVLD- - - - - GGDDDSDYAPVNDPLSHYDVITK
 PBAC3_E.go | HIRFNGEMDATYEVQPRGAGGLAADVGGL HGGAASSLRAEGNDEDEFVVEDHLSKYNVSMK
 PBAC3_D.ha | NVLINGVMDTTFDIPVSSKSNAYNSLNSI - S- ED- - - - - GGVEPVII
 PBAC3_D.re | VITOLGKIPGIVNVHF DVKKDERVV- - - - - P- - - - - YL
 PBAC3_X.la | VVTQYGKMGTLVSVTSLVSSDLGK- - - - - PSLTTKVL
 PBAC3_D.re | VLTQYGKIGTLVLSITPDTRSGDISV- - - - - PMYTSKVL
 PBAC3_M.mu | VVTQFGKMGTLVSLPEPSNVANDISK- - - - - PVLTTTRVL
 PBAC3_H.sa | VVTQFGKMGTLVSLPESSVASDVSK- - - - - PVLTTKVL
 consensus +++. * . ++++++. + . + + . + . + . + + +

PBAC3_A.th | FGKRDEPM- - - - - LTATAR- - - - - RLIEHISSI VPSKPLVL SGLKDHSSE
 PBAC3_Z.ma | FGKRDEPL- - - - - LLACAR- - - - - QLIEHISGS GSARSLVLSLGLKDHSQG
 PBAC3_O.sa | FGKRDEPL- - - - - LLACAR- - - - - QLIEHISGS GSARSLMISLGLKDHSQA
 PBAC3_M.tr | FGKRDEPM- - - - - LVACAR- - - - - QLIEHMTLSGVSKPLVL SGLKDHSVE
 PBAC3_P.pa | MGKRD EPL- - - - - MMACAR- - - - - QLIEKMSKTCGCSRPLI LSLGLADHSAE
 PBAC3_S.ce | LGDSADPK- - - - - VPVVCVQIAELYRRVI LPEVSGTMQDNMFQFSLLISN- - - -
 PBAC3_K.la | LGDSNDMK- - - - - IPIICTQIAQLYERVI LPTLGSEVGASVPDF- - VI TI- - -
 PBAC3_E.go | MGNSNDMK- - - - - LPVVMQIAELYERAVYPSVLGAEARAGPDGLLLVTL- - - -
 PBAC3_D.ha | IDYSNLK- - - - - I QVVASQIGKLVLVSQE- - - - - PKNLILSI- - - -
 PBAC3_D.re | HGPPVDNPEFHVSVPI TMNCCLGLDTDETRSAIQFLVNRTGLHKCPTEFVVGLGLKKIDGP
 PBAC3_X.la | LGQ-DEPL- - - - - VHVCALK- - - - - NLVTFVSEESKNKPVL LSNALDKSVD
 PBAC3_D.re | LGK-DEPL- - - - - THVYAK- - - - - NVATFVSQESGNRPI LLLGSLKDNSAE
 PBAC3_M.mu | LGQ-DEPL- - - - - HVFAK- - - - - NLVAFVSQEAGNRAVL LAAVAKDKSME
 PBAC3_H.sa | LGQ-DEPL- - - - - HVFAK- - - - - NLVAFVSQEAGNRAVL LAAVAKDKSME
 consensus +*. . +++. + . + . + + + +

PBAC3_A.th | TLKEIVATVI ENRLW- - - - -
 PBAC3_Z.ma | TLKDI TVAVI ENRLW- - - - -
 PBAC3_O.sa | TMKYI VSTI I ENRLW- - - - -
 PBAC3_M.tr | TLKGIVSAVIDKRLW- - - - -
 PBAC3_P.pa | TLKDI EVVVENKI W- - - - -
 PBAC3_S.ce | - - - - - SSKI WRATKEQ-SADDNDFGKLVFVLKCI KDMYAK
 PBAC3_K.la | - - - - - STKL WRKSQ- - - - - SNEFEKLI FLLNAVKEAYVN
 PBAC3_E.go | - - - - - SSRFWRSASAGGPGSGSDFDRLMFVLQSI REMYGI
 PBAC3_D.ha | - - - - - GSRWFKGKG- - NETDENDFNKLMFVL DNVVKQLLK-
 PBAC3_D.re | NLRALAKVLEETLF- - - - -
 PBAC3_X.la | CIKTLKDVI RSCQVW- - - - -
 PBAC3_D.re | TMKTVKDMI KSCQVW- - - - -
 PBAC3_M.mu | RLKALKEVI RL CQVW- - - - -
 PBAC3_H.sa | GLKALREVIRVCQVW- - - - -
 consensus . . + + .

Figure 3.S7: Multiple Sequence Alignment for PBAC3.

[Figure 3.S7 caption follows on next page]




Figure 3.S7 (preceding page): Multiple Sequence Alignment for PBAC3. A Clustal Omega (Sievers and Higgins, 2014) sequence alignment of PBAC3 from plants, yeast, and animals visualized with the program BOXSHADE. A heat map showing each combination of protein percent identity with respective values shown in each cell obtained from the resultant Clustal Omega identity matrix. Species names are abbreviated as follows: A.th - *Arabidopsis thaliana*, Z.ma - *Zea mays*, Os - *Oryza sativa*, M.tr - *Medicago truncatula*, P.pa - *Physcomitrella patens*, S.ec- *Saccharomyces cerevisiae* K.la - *Kluyveromyces lactis*, E.go - *Eremothecium gossypii*, D.ha - *Debaryomyces hansenii*, D.me - *Drosophila melanogaster*, X.la - *Xenopus laevis*, D.re - *Danio rerio*, M.mu - *Mus musculus*, H.sa, *Homo sapiens*.

PBAC4_A.t h METEYDQVLRSVAQVNI ----- ADQI GESDDV----- ERDGNGVKVSCFTTEVFDDVTIQFQ
 PBAC4_Z.ma MSSEE- ARASFSELVVGSPTRTEGETYSSEG----- MSSEEGAHVTCFTEDLHDVTLHFQ
 PBAC4_O.sa MASEE- LRTSFSDELVGSSRTDSQVESSGD----- PSSSEGGVQVTCFTEDLHDVIHFQ
 PBAC4_M.tr MNSEE- LGKGIQELKLN----- HK----- DNEQTQLQFTCFTELVNDVNLHFQ
 PBAC4_P.pa ----- MAVAPA----- PDVEPSSLVTFQSFSDENSTFHfq
 PBAC4_S.ce MLV----- KTI SRTI E----- SES----- GFLQPTLDVIA-TLP----- AD
 PBAC4_K.la TTL----- TFPDTW----- T----- SFE GTPTS VRCTTS PHRDPI----- VS
 PBAC4_E.go MTQT----- KTADYTTS----- TLGGDDVRLLV-TMP----- VE
 PBAC4_D.ha ----- MI VTK----- QE----- NI----- NGDDYQIVI S
 PBAC4_D.re ----- MDSVPANVVAPEKESGGFTSHCVHLAAGPSQFTVR
 PBAC4_X.la ----- MESSPRAITLYNFSEKICE----- QLVHFF
 PBAC4_D.re ----- ME AVL----- NGDCSSHASADSLSLHDFSEKFLE----- QRVYFFH
 PBAC4_M.mu ----- MEESRAAADADVSLLHNFSSARLWE----- QLVHFFH
 PBAC4_H.sa ----- MEGLVVAAGGDVSLHNFSSARLWE----- QLVHFFH
 consensus +.....+.+.+..++.

PBAC4_A.t h IIRL----- AKQI YVWVGCNSA----- KFGSLIYAAASTRP----- S----- NTVSVTSVLGGS
 PBAC4_Z.ma I VRF-SKQIYVWVGCNTA----- KF GHLYAAASTRP----- D----- NRVSVTSVLGGS
 PBAC4_O.sa I I RF-SKQIYAWVGCNTA----- KF GHLYAAASTRP----- G----- NGVSVTSVLGGS
 PBAC4_M.tr I I RF-PKQIYVWVGCNSA----- KLGQLIYAAASTRP----- N----- NVVSVTSILGGNS
 PBAC4_P.pa I LRL-TNQLYVWVGCNSA----- RMGNLYAALPTRW----- D----- KTAGVAALVGGGA
 PBAC4_S.ce - DRSKKIPIISL VVGF KQEASL NSSSSLSCYYYAIPLMRDRHI NLKGGSNVVGI PLLDTK
 PBAC4_K.la QDRKTALSIAIYLNCGDLSSGDSRTYLQGYCYGIEFVGRVQ-STPKSNEVCVADLIESN
 PBAC4_E.go LTASKKVPVSI VVSGRS DGV----- NEE LAATVYG----- VPYKGSDVTSVLLDTA
 PBAC4_D.ha IDKHDKSSINVYVNEMTNT----- SVGSVYV----- SI N----- DKKSQI YSSVLQDH
 PBAC4_D.re ALKM-QGSTMLIINPKGLE----- VFEE LAVAMPSRN----- PASSQSLSSTLGGHG
 PBAC4_X.la VMGN- EDCFLFWVGL - SP----- VL ANLISMA MCSRF----- D----- SVPLS STLVGDT
 PBAC4_D.re LMKL-SGGFFLWVGTATNP----- LS S NLA VSI LS RT----- D----- SAPLS STLVLGDT
 PBAC4_M.mu VMRL-TDSLFLWVGA- TP----- HL RNL AVAMCSRY----- D----- PI PVCTS LFGDT
 PBAC4_H.sa VMRL-TDSLFLWVGA- TP----- HL RNL AVAMCSRY----- D----- SI PVSTSLLGDT
 consensus +++++.++++.+++.+.++.+++.+.++++.+++++.

PBAC4_A.t h DNTIGSGIARR----- LVLKTGLNIILACNIPKNPLLEA----- KAEEKVLI RKLIDL
 PBAC4_Z.ma DNTIGSSMARR----- LVLKTGLNI VLACNIPKDSPI LEA----- AAERKLVEKLKG
 PBAC4_O.sa DNTIGSMARR----- LVLKTGLNI VLACNIPKDSPMLEA----- AAERKLI EKL RGL
 PBAC4_M.tr DNTIGSGIARR----- LVLKTGLNI IMACNIPKNSPMVEI----- EAEKILIKQLISL
 PBAC4_P.pa DSTIGASMARR----- LS MVI GWSIVLASNIPS NAPVLEA----- FAEQRLFQELRSM
 PBAC4_S.ce DDRI RD MARHMATI I SERFNRPCYVTVSSLPSEDPMSMLVANH----- LYI LKICL DLLKTEL
 PBAC4_K.la DHQI HDLHVKTTRILV LKYHRPCYVQIASKTAEFT-LNGADQI NMLQNTKCI DRNVK
 PBAC4_E.go DNRI R DFG STFCRVLVKKVGL PCYVSVSSVGRM----- L DS GDQ----- LAVI RKALEHLTS
 PBAC4_D.ha SNI VTE QARNLGYLVVKKYYCRPSYTNI NGQI SDYPQVL----- KVVFNLLIDESI
 PBAC4_D.re QT DSSVLA AAK----- LS KRY CRQFYVSLNLLKLDRL-MGP----- LFEKTLTVYMDH
 PBAC4_X.la DI MSTSFAQR----- LAKKT KKQVFA SVNLPTNESQML----- LIEKRI RQE MDTF
 PBAC4_D.re DTIASAL AQR----- LTKKT KKQVFVSYNLPVTDSNLL----- LVENRI KQE MELH
 PBAC4_M.mu DT TSTGL AQR----- LARKT S KQVFVSYNL SNTDSNFTL----- LVENRI KEEMETF
 PBAC4_H.sa DT TSTGL AQR----- LARKT NKQVFVSYNL QNTDSNFAL----- LVENRI KEEMEA
 consensus +.+.+.++++.++++.++++.++++.+.+.+.++++.++++.+.+

PBAC4_A.t h GYTMSKPVRAT-----
 PBAC4_Z.ma GYIKTPAVGANSSAQ-----
 PBAC4_O.sa GYVRPRAGEVNTSTAQ-----
 PBAC4_M.tr GYTKSRLLEGTS-----
 PBAC4_P.pa GYIKPASKPLNPGAPLAPI
 PBAC4_S.ce GE-----
 PBAC4_K.la -----
 PBAC4_E.go -----
 PBAC4_D.ha -----
 PBAC4_D.re LEHFA-----
 PBAC4_X.la PEKF-----
 PBAC4_D.re PDKF-----
 PBAC4_M.mu PEKF-----
 PBAC4_H.sa PEKF-----
 consensus

Figure 3.S8: Multiple Sequence Alignment for PBAC4.

[Figure 3.S8 caption follows on next page]




Figure 3.S8 (preceding page): Multiple Sequence Alignment for PBAC4. A Clustal Omega (Sievers and Higgins, 2014) sequence alignment of PBAC4 from plants, yeast, and animals visualized with the program BOXSHADE. A heat map showing each combination of protein percent identity with respective values shown in each cell obtained from the resultant Clustal Omega identity matrix. Species names are abbreviated as follows: A.th - *Arabidopsis thaliana*, Z.ma - *Zea mays*, Os - *Oryza sativa*, M.tr - *Medicago truncatula*, P.pa - *Physcomitrella patens*, S.ec- *Saccharomyces cerevisiae* K.la - *Kluyveromyces lactis*, E.go - *Eremothecium gossypii*, D.ha - *Debaryomyces hansenii*, D.me - *Drosophila melanogaster*, X.la - *Xenopus laevis*, D.re - *Danio rerio*, M.mu - *Mus musculus*, H.sa, *Homo sapiens*.

[Figure 3.S9 caption follows on next page]

PAP1_A.th	- - - - M	KVAE KI VFLW NDS DGF AAT I SDT L NPS PSS SPL RKL E E Q I QL P L D KY G V E G V E T G G
PAP1_Z.ma	- - - - M	RVAPRVLFLV RDAAGY GAAL ADAL RPPP - - GL TRES S P F E L P L GKY GL DGE KA CA
PAP1_O.sa	- - - - M	RVAPKVLVLVR DAAGY GAAL ADAL RPTP - - GL TRES STPL E L PLAKY GL DGE KA SG
PAP1_M.tr	- - - - M	KL API TVF MFRD S EG F AS A I S Q A L S S - - S S F T R L E D S F E L S L E G Y G I K DH K A SG
PAP1_P.pa	MGVAME L GE E VVLL YDG GK VGVTR FSE ALV PEA S PPL Q RVVC NL - - PWKD - P NGE VLT KS	
consensus	. . . * ++++++++ . . ++++++++ * . + . . +++. . . +++. . . +++. + . . +.	
PAP1_A.th	S I VHF VD E	N E V Y Q V S I F L L R S Y E P P V - L V C A M N E L L D L I T R G - S T L P - - T I I A P F F V A A
PAP1_Z.ma	E L L S F S D S	S C S P Q V T I F V L P D Y K P P L - A A C V V N G V L E L I S S E A T S T E R - - T L I V P Y I T R S
PAP1_O.sa	E L V N F S D S	S G D P Q V S F F V L P D Y K P P V - A A C A M N E I L A C A L V S S E A P S T Q P - - T L I V P F M T R S
PAP1_M.tr	N V L H Y L D N H R S	Y K V S I V I M Q H Y E P P V - L A C A L N E V L N K I V G G D P S S N P - - T L I V P F L V E S
PAP1_P.pa	E V V N F K D N A C L	T K V S L I V F R Q E V T S A C G R V V H D L F E Y L L S N A P I Q S P G F S L V V P A I S R M
consensus	+++ . + . * +. . + * + + + + . + . + + + . . +	
PAP1_A.th	S K L K F N N R S L E A N N R K A S L H Y V Q V A T E	T E T S R L F A S R I E K P P L S M Q I H Y E P L S C L L H L A R
PAP1_Z.ma	S S Y H H G T E - -	H A T K A S T V L H G A E I G A T T D Y T H L L V G G T T K P P T S L Q I R S E P I L C L L E M V R
PAP1_O.sa	P N Y Y H G A K - -	T G Q - - L A T L H G A E I G A T T E F T Q M L V D G T T K L P Q S L Q V R S E P I L C L L E M V R
PAP1_M.tr	S K V K G N S K S L R S D E S R A L T Y G I Q I G Q I T E T M Q T L L K K T Q E P P S S L R I Q Q E N F A C F L H F V R	
PAP1_P.pa	P G S V L N M D - -	T G N - - N A N I Y A A F F N R - T S E G S G N L D G L P K L P S T F A V R D G F L A Y V F H Y V H
consensus	+ . . . + . + . . + + . . + . + + + + + + . + + . +	
PAP1_A.th	V K R L P T A I L I G Q R S N S L T H K A L D E E L Q V I H E T G E L V A S W T G L S F S R D R I K W S - - A S K T S K	
PAP1_Z.ma	V L K V P T V L L V T S G G Q Q Q G K S S T V S D L E V L Q C L G E H L A K H T A L E F S K G T V L K R - - G E E K S R	
PAP1_O.sa	V L N I P T V I L F A S G G Q H Q G K S S T D V D L E V L Q C V G D H L A K H I N L E F S K E T V L Q R - - G E E K S P	
PAP1_M.tr	V M K L P T F F L I G Q T S Q Y L D N K S P K - Q H E A I C E I G E I L A S S T G L E F S E D R V I W N - - P K K T L K	
PAP1_P.pa	A T Q L P T L V L V S P S T E G V T A Q K - N E S N E V L L A L G D V V G R Q I G L T C S K Q E L N H V T S L I P L A S	
consensus	++++* + +* ++. . ++. . + . . + + + + + . + * + + + + . + +* + +* ++. . ++. . + + . . + + . . + +	
PAP1_A.th	E E E S P W R A L Y G	
PAP1_Z.ma	I D Q E P W R E L Y G	
PAP1_O.sa	V F Q E P W R E L Y Q	
PAP1_M.tr	E I K E P W R D L Y G	
PAP1_P.pa	S M E D D M R R L Y I	
consensus	. . . + + * * + * * +	





Figure 3.S9 (preceding page): Multiple Sequence Alignment for PAP1. A Clustal Omega (Sievers and Higgins, 2014) sequence alignment of PAP1 from plants, yeast, and animals visualized with the program BOXSHADE. A heat map showing each combination of protein percent identity with respective values shown in each cell obtained from the resultant Clustal Omega identity matrix. Species names are abbreviated as follows: A.th - *Arabidopsis thaliana*, Z.ma - *Zea mays*, Os - *Oryza sativa*, M.tr - *Medicago truncatula*, P.pa - *Physcomitrella patens*, S.ec- *Saccharomyces cerevisiae* K.la - *Kluyveromyces lactis*, E.go - *Eremothecium gossypii*, D.ha - *Debaryomyces hansenii*, D.me - *Drosophila melanogaster*, X.la - *Xenopus laevis*, D.re - *Danio rerio*, M.mu - *Mus musculus*, H.sa, *Homo sapiens*.

[Figure 3.S10 caption follows on next page]

A



B




Figure 3.S10 (preceding page): Full BiFC Panel and Controls. BiFC analysis showing both GFP channel and a merged bright-field and GFP channel. **(A)** A repeat of Figure 6D is shown for reference **(B)** Additional controls combining N-YFP and C-YFP with their respective N and C terminal YFP-fused PBAC1, PBAC2, PBAC3, and PAP1.




Figure 3.S11: Phylogenetic Analysis of PAP1 and PBAC1-4.

[Figure 3.S11 caption follows on next page]

B




Figure 3.S11 (preceding page): Phylogenetic Analysis of PAP1 and PBAC1-4. Bayesian phylogenetic analysis of PAP1 and PBAC1-4 from sequence alignments shown in Figures 3.S5-3.S9. Trees were generated with MrBayes (Ronquist *et al.*, 2012) and then visualized with FigTree. Nodes are color-coded with clade probabilities (probability of observing this grouping in the set of sampled trees), and a color key is shown on the right. Species names are abbreviated as follows: A.th - *Arabidopsis thaliana*, Z.ma - *Zea mays*, Os - *Oryza sativa*, M.tr - *Medicago truncatula*, P.pa - *Physcomitrella patens*, S.ec- *Saccharomyces cerevisiae* K.la - *Kluyveromyces lactis*, E.go - *Eremothecium gossypii*, D.ha - *Debaryomyces hansenii*, D.me - *Drosophila melanogaster*, X.la - *Xenopus Laevis*, D.re - *Danio rerio*, M.mu - *Mus musculus*, H.sa, *Homo sapiens*. **(A)** Phylogeny generated with all sequences from yeast, animals, and plants. Clear grouping is observed between the yeast, animal, and putative plant PBAC2 orthologs into a single clade. Grouping was observed for PBAC3 between the animal, and putative plant sequences; however, yeast PBAC3 was not found in this clade. Clear grouping is observed for PBAC4 between the animal, and putative plant sequences; however, yeast PBAC4 was not found in this clade. Animal, yeast, and plant PBAC1 did not clearly group, likely due to considerable sequence divergence. **(B)** Phylogeny generated when omitting the yeast sequences and only utilizing animal and plant sequences. Clear grouping is observed between the putative plant orthologs PBAC2-4 with their respective animal orthologs into separate clades. Animal PBAC1 does not group with the putative plant specific PBAC1 orthologs, likely due to considerable sequence divergence from animal PBAC1.

Table 3.S1: Oligonucleotide Primer Sequences Used in this Study. A table of primers used throughout this chapter is provided for reference.

Primers	Notes	Sequence
P1		CATAAGGAACCTCGAATCCCC TTICATTATCCAGGGTCGAAAG
P2		GATCACCTAACCCCTCCGACA
P3		CATGGATGAAATCGATGCTA
P4		GCACAAGAAAGAGTGTACTCTC
P5		TGATAAAATGCTCAGTGTCAA
P6		GCTGCCCTAAATCTCCCTAGC
P7		TGAACCTTTCCCTTAGAGTTGGGA
P8		ACTACAAACCTCAAAACAAACATTGA
P9		GGGGACACAAGTTGTACAAAAGCAGGCTTAACAAAGCTTGAAAGTATGAA
P10		GGGGACCCACTTTCTACAAAGAAAAGCTGGCTATTGATGTTGCAATTCTTTGATT
P11		cagaatccaaaggatggACTACAAGGACCGACGATGACAAGatgaccggactgtatg
P12	1xFLAG RPT4a F	catcgacgtggccatCTTGTCACTCGTCGTCCTGTAGTCatgttgtctgaatctcg
P13	1xFLAG RPT4a R	GatgGACTACAAGGACGACGATGACAAGGATTACAAAGGATGATGATAAGatgaccggactgtatg
P14	2xFLAG RPT4a F	catcgacgtggccatCTTATCACTCATCCTGTAAATCTGTACATCGTCCTGTAGTCatC
P15	2xFLAG RPT4a R	catcgatggactacaaggacgtgatgataaggactacaaggatgatgacttgcggagcgg
P16	5-Flag-RPT4B-Nco1	tccccggatgtactttccaaaatccgc
P17	3-RPT4B-Sma1	AAAAACTTGACAGgttggatacttgttttcatgt
P18	5-prRPT4B-Pst1	catcgatggattttacatgtacttgcggatgttttcatgt
P19	3-prRPT4B-Nco1	GGGGACACAAGTTGTACAAAAGCAGGCTTAATGAAAGTACGGGAGAAG
P20	PAP1 F	GGGGACCCACTTTCTACAAAGGACGATGGCTATTAGCCATAAGAGCAG
P21	PAP1 R	GGGGACACAAGTTGTACAAAAGGCTTAATGCAAGATGTACTTACC
P22	PBAC1 F	GGGGACCCACTTTCTACAAAGGACGATGGCTATTAGCTATATAGATCAGA
P23	PBAC1 R	GGGGACACAAGTTGTACAAAAGGACGCTTAATGGAGTTTGTGTTGA
P24	PBAC2 F	GGGGACCCACTTTCTACAAAGGACGATGGCTATTAAACATAGACATATC
P25	PBAC2 R	GGGGACACAAGTTGTACAAAAGGACGCTTAATGAAAAGCTGGACACTAAT
P26	PBAC3 F	GGGGACCCACTTTCTACAAAGGACGATGGCTATTACCAAAGGGATCTC
P27	PBAC3 R	GGGGACACAAGTTGTACAAAAGGACGCTTAATGGAGACTGAATACGAC
P28	PBAC 4 F	GGGGACCCACTTTCTACAAAGGACGCTTAATGGAGACTGAATACGAC
P29	PBAC 4 R	GGGGACCCACTTTCTACAAAGGACGCTTAATGGAGACTGAATACGAG

Table 3.S2: Contaminants Removed from MaxQuant's ProteinGroups.txt file.
 A list of common contaminants that were removed from MaxQuant's ProteinGroups.txt file for the proteomics experiments involving PAG1-FLAG, and FLAG-RPT4a/b affinity purified proteasomes. ATG Identifiers, Symbols and descriptions from The *Arabidopsis* Information Resource v 10 (TAIR10) are provided.

ATG Identifier	Symbols	TAIR10 Description
AT1G03880.1	CRU2, CRB	cruciferin 2
AT3G44300.1	NIT2, AtNIT2	nitrilase 2
AT3G44310.3	NIT1, ATNIT1, NIT1	nitrilase 1
AT3G44310.2	NIT1, ATNIT1, NIT1	nitrilase 1
AT4G27140.1	SESA1, AT2S1	seed storage albumin 1
AT4G27160.1	AT2S3, SESA3	seed storage albumin 3
AT4G27170.1	SESA4, AT2S4	seed storage albumin 4
AT4G28520.1	CRU3, CRC	cruciferin 3
AT5G44120.3	CRA1, ATCRA1, CRU1	RmlC-like cupins superfamily protein
AT5G54740.1	SESA5	seed storage albumin 5

Table 3.S3: InterProScan analysis of *Arabidopsis* protein sequences for PBAC1-4. InterProScan Signatures from both PFAM and Panther databases with their respective expectation-values (E-values) are shown for reference (Jones *et al.*, 2014). While PBAC1 (previously identified as a likely ortholog for PAC1 (Kus-mierczyk *et al.*, 2011)) contains a Proteasome Assembly Chaperone (PAC) 2 domain shown with a star, it is of considerably lower expectation-value (E-value) than PBAC2 which contains a Proteasome Assembly Chaperone (PAC) 2 domain. PBAC3 contains a PAC3 domain, and PBAC4 contains a PAC4 domain. PAP1 contains no domains of known function but does contain a conserved domain annotated in the Panther database (Mi *et al.*, 2005).

Protein	AGI Identifier	InterproScan Signature	Annotation	E-value	Start Residue	End Residue
PBAC1	AT3G25545	PF09754 PTHR37227	Proteasome Assembly Chaperone 2* Family Not Named	3.4E-06	60	247
PBAC2	AT3G18940	PF09754 PTHR12970	Proteasome Assembly Chaperone 2 Member 5-induced protein 1 class3	1.6E-116	2	267
PBAC3	AT5G14710	PF10178 PTHR31051	Proteasome Assembly Chaperone 3 Proteasome Assembly Chaperone 3	1.0E-29	18	249
PBAC4	AT1G48170	PF16093 PTHR33559	Proteasome Assembly Chaperone 4 Family Not Named	4.8E-102	3	172
PAP1	AT3G07640	PTHR37221	Family Not Named	1.6E-26	37	120
				2.9E-45	1	124
				1.4E-17	47	120
				9.5E-56	14	153
				8.1E-109	2	241

Table 3.S4: Iterative PSI-BLAST Analysis using Human PAC1-4 to Identify PBAC1-4. Iterative PSI-BLAST Analysis was performed on NCBI's non-redundant (nr) database using the human PAC1-4 sequences using the default parameters. PBAC1-4 were the top hits in *Arabidopsis* for their respective PAC1-4 search sequences. ATG identifiers, RefSeq identifiers, Number of PSI-BLAST iterations, and expectation values (E-value) are listed for each analysis. PBAC1 required a second iteration to be identified, in which all sequences above the default expectation threshold were included in the second search.

Human	Arabidopsis	ATG Identifier	Matching Accession	Iteration	E-value
PAC1	PBAC1	AT3G25545	NP_566770.1	2	2.0E-04
PAC2	PBAC2	AT3G18940	NP_566621.1	1	1.0E-21
PAC3	PBAC3	AT5G14710	NP_196975.1	1	5.0E-08
PAC4	PBAC4	AT1G48170	NP_564522.1	1	4.0E-06

Table 3.S5: Proteasome associated proteins identified after PAG1-based affinity purification under minus ATP Conditions. A table of proteins identified as significantly enriched at FDR 0.01 in the indicated affinity preparations. Shown are the protein's ATG identifiers, descriptions from *The Arabidopsis Information Resource version 10* (TAIR10), symbols from TAIR10, average MS/MS count, average sequence coverage, fold change compared to control using the MaxLFQ value from MaxQuant, and p-values obtained from the volcano plots in Figure 3.4 A.

ATG	TAIR10 Description	Symbol(s)	Annotation	Average		Fold Change Compared to Control	P-Value
				Proteasome Count*	Average Sequence MS/MS Coverage (%)		
AT3G18940.1	clast3-related	PBAC2		41.5	56.2	2734.5	1.21E-06
AT3G07640.1	unknown protein	PAP1		49.8	59.9	2462.4	6.05E-06
AT3G25545.1	unknown protein	PBAC1		58.2	68.4	2319.3	2.73E-07
AT3G13330.1	proteasome activating protein 200	PA200	PA200	54.3	30.1	846.7	3.02E-05
AT5G38650.1	Proteasome maturation factor UMP1	Ump1b		11.5	67.4	279.2	1.26E-05
AT1G67250.1	Proteasome maturation factor UMP1	Ump1a		8.7	56.0	277.0	1.96E-05
AT3G53970.1	proteasome inhibitor-related	PTRE1		6.3	29.4	240.6	9.99E-06
AT1G23520.1	Domain of unknown function (DUF220)			8.7	3.0	165.3	1.02E-03
AT1G01320.2	Tetratricopeptide repeat (TPR)-like superfamily protein			2.0	1.7	156.0	4.31E-07
AT1G48170.1	unknown protein	PBAC4		19.8	79.5	138.4	5.30E-02
AT5G14710.1	CONTAINS InterPro DOMAIN/s: Proteasome assembly chaperone 3 (InterPro:IPR018788)	PBAC3		7.8	39.4	126.3	5.79E-05
AT3G13090.1	multidrug resistance-associated protein 8	ATMRP8, ABCC6, MRP8		0.0	0.5	86.8	1.01E-01
AT2G47360.1	unknown protein			3.2	2.6	64.5	3.39E-03
AT2G26780.1	ARM repeat superfamily protein	ECM29		3.5	9.7	35.6	1.91E-02
AT4G17100.1	CONTAINS InterPro DOMAIN/s: Endoribonuclease XendoU (InterPro:IPR018998)			0.2	4.4	26.0	6.55E-05
AT5G27960.1	AGAMOUS-like 90			0.2	6.3	25.1	2.71E-02
AT1G21530.1	AMP-dependent synthetase and ligase family protein			0.2	3.1	24.4	1.22E-05
AT1G23780.1	F-box family protein			2.0	8.6	21.5	1.63E-04
AT3G26020.2	Protein phosphatase 2A regulatory B subunit family protein			0.8	2.7	14.5	2.53E-04
AT2G17200.1	ubiquitin family protein	DSK2		2.5	9.4	13.7	1.10E-02
ATCG00680.1	photosystem II reaction center protein B	PSBB		0.0	2.2	13.4	3.07E-04
AT5G55220.1	trigger factor type chaperone family protein			0.2	2.3	13.2	5.56E-03
AT5G13450.1	delta subunit of Mt ATP synthase	ATP5		0.0	4.5	8.5	9.55E-04
AT1G79530.1	glyceraldehyde-3-phosphate dehydrogenase of plastid 1	GAPCP-1		3.3	17.6	7.4	3.56E-04

Table 3.S6: Proteasome associated proteins identified after PAG1-based affinity purification under plus ATP Conditions. A table of proteins identified as significantly enriched at FDR 0.01 in the indicated affinity preparations. Shown are the protein's ATG identifiers, descriptions from *The Arabidopsis Information Resource version 10* (TAIR10), symbols from TAIR10, average MS/MS count, average sequence coverage, fold change compared to control using the MaxLFQ value from MaxQuant, and *p*-values obtained from the volcano plots in Figure 3.4 B.

ATG	TAIR10 Description	Symbol(s)	Proteasome Annotation	Average			
				Average MS/MS Count*	Sequence Coverage (%)	Fold Change Compared to Control	
AT3G25545.1	unknown protein	PBAC1	33.7	40.7	2130.4	8.05E-06	
AT3G18940.1	clast3-related	PBAC2	12.2	39.6	1340.9	1.45E-05	
AT2G26780.1	ARM repeat superfamily protein	ECM29	0.0	0.8	908.0	5.36E-04	
AT3G13330.1	proteasome activating protein 200	PA200	PA200	10.8	11.8	755.5	1.20E-05
AT3G07640.1	unknown protein	PAP1	21.0	27.4	446.8	1.09E-03	
AT5G38650.1	Proteasome maturation factor UMP1	Ump1b	6.2	45.4	326.9	1.47E-04	
AT1G48170.1	unknown protein	PBAC4	8.3	41.1	291.7	6.90E-05	
AT1G67250.1	Proteasome maturation factor UMP1	Ump1a	4.7	36.7	247.3	2.59E-04	
AT2G47360.1	unknown protein			2.7	2.6	197.7	1.08E-05
AT3G13090.1	multidrug resistance-associated protein 8	ATMRP8, ABCC6, MRP8		0.0	1.3	184.1	1.15E-02
AT5G14710.1	CONTAINS InterPro DOMAIN/s: Proteasome assembly chaperone 3 (InterPro:IPR018788)	PBAC3	2.8	24.5	162.2	4.32E-05	
AT3G48670.2	XH/XS domain-containing protein	IDN2, RDM12		1.5	2.2	113.5	3.33E-04
AT2G28070.1	ABC-2 type transporter family protein			0.8	1.8	105.9	2.27E-04
AT2G14170.2	aldehyde dehydrogenase 6B2	ALDH6B2	0.8	1.0	100.8	5.43E-05	
AT5G27960.1	AGAMOUS-like 90	AGL90	0.0	6.3	80.4	4.59E-04	
AT1G74260.1	purine biosynthesis 4	PUR4	0.3	0.9	62.9	1.70E-02	
AT3G53970.1	proteasome inhibitor-related		PTRE1	2.3	14.4	62.7	3.29E-03
AT5G66420.1	CONTAINS InterPro DOMAIN/s: Uncharacterised conserved protein UCP033271 (InterPro:IPR008322), TIM-barrel signal transduction protein, predicted (InterPro:IPR009215)			0.2	1.5	57.2	1.03E-03
AT3G26020.2	Protein phosphatase 2A regulatory B subunit family protein			0.5	2.3	54.7	1.55E-04
AT2G37650.1	GRAS family transcription factor			0.3	0.7	46.7	2.71E-04
AT4G25500.2	arginine/serine-rich splicing factor 35	ATRSP35, ATRSP40, AT-SRP40, RSP35, RS40, At-RS40		1.7	15.4	46.6	8.75E-04
AT2G04235.1	unknown protein			1.5	0.9	42.5	6.59E-04
AT1G23520.1	Domain of unknown function (DUF220)			3.7	3.0	42.3	1.56E-03
AT2G17200.1	ubiquitin family protein	DSK2	0.7	4.3	40.9	3.39E-04	
AT4G23680.1	Polyketide cyclase/dehydrase and lipid transport superfamily protein			1.5	30.3	34.5	3.15E-03
AT5G05180.2	unknown protein			0.8	2.5	30.6	1.62E-04
AT4G17100.1	CONTAINS InterPro DOMAIN/s: Endoribonuclease XendoU (InterPro:IPR018998)			0.0	4.9	23.8	5.73E-05
AT1G79530.1	glyceraldehyde-3-phosphate dehydrogenase of plastid 1	GAPCP-1	0.3	7.3	23.6	1.31E-04	
AT1G78860.1	D-mannose binding lectin protein with Apple-like carbohydrate-binding domain			0.2	2.8	21.2	1.21E-04
AT4G34700.1	LYR family of Fe/S cluster biogenesis protein	CIB22, AtCIB22		0.2	2.6	19.4	5.40E-03
AT5G15390.1	tRNA/rRNA methyltransferase (SpoU) family protein			0.2	4.0	16.5	7.05E-04
AT4G30160.1	villin 4	VLN4, ATVLN4		0.0	1.3	15.4	5.91E-03
AT2G36880.2	methionine adenosyltransferase 3	MAT3	0.0	7.8	12.3	2.16E-03	
AT3G28780.1	Protein of unknown function (DUF1216)			0.5	2.5	11.0	6.32E-04

Table 3.S7: Proteasome associated proteins identified after RPT4a-based affinity purification under minus ATP conditions. A table of proteins identified as significantly enriched at FDR 0.01 in the indicated affinity preparations. Shown are the protein's ATG identifiers, descriptions from *The Arabidopsis Information Resource version 10 (TAIR10)*, symbols from TAIR10, average MS/MS count, average sequence coverage, fold change compared to control using the MaxLFQ value from MaxQuant, and *p*-values obtained from the volcano plots in Figure 3.4 C.

ATG	TAIR10 Description	Symbol(s)	Proteasome Annotation	Average			Fold Change Compared to Control	<i>P</i> -Value
				Average MS/MS Count*	Sequence Coverage (%)			
AT2G26780.1	ARM repeat superfamily protein		ECM29	153.7	55.2		499.0	2.73E-03
AT2G03430.1	Ankyrin repeat family protein		NAS6	22.8	68.5		155.3	1.65E-03
AT3G13090.1	multidrug resistance-associated protein 8		ATMRP8, ABCC6, MRP8	0.0	1.3		135.8	4.78E-06
AT3G15180.1	ARM repeat superfamily protein		HSM3	54.5	51.8		110.2	4.30E-06
AT5G27960.1	AGAMOUS-like 90		AGL90	1.0	7.5		94.1	6.40E-03
AT5G57950.1	26S proteasome regulatory subunit, putative		NAS2	2.8	32.3		82.1	5.94E-06
AT3G53970.1	proteasome inhibitor-related			1.8	11.1		23.4	9.04E-05
AT2G28070.1	ABC-2 type transporter family protein			0.0	0.7		19.3	1.01E-04
AT5G49550.1	Putative homolog of mammalian BLOC-1 Subunit 2. Protein - protein interaction with BLOS1.	BLOS2		0.7	10.5		16.9	5.10E-04
AT5G21040.2	F-box protein 2		FBX2	0.5	2.6		14.8	7.77E-05
AT3G13330.1	proteasome activating protein 200	PA200	PA200	0.0	0.7		13.7	2.90E-03
AT5G13450.1	delta subunit of Mt ATP synthase		ATP5	0.7	7.5		8.8	1.42E-03

Table 3.S8: Proteasome associated proteins identified after RPT4a-based affinity purification under plus ATP conditions. A table of proteins identified as significantly enriched at FDR 0.01 in the indicated affinity preparations. Shown are the protein's ATG identifiers, descriptions from *The Arabidopsis Information Resource version 10* (TAIR10), symbols from TAIR10, average MS/MS count, average sequence coverage, fold change compared to control using the MaxLFQ value from MaxQuant, and p-values obtained from the volcano plots in Figure 3.4 D.

ATG	TAIR10 Description	Symbol(s)	Annotation	Average			P-Value
				Proteasome	Average MS/MS Count*	Sequence Coverage (%)	
AT3G15180.1	ARM repeat superfamily protein	HSM3		27.3	36.2	1024.2	7.3E-06
AT5G57950.1	26S proteasome regulatory subunit, putative	NAS2		18.0	59.9	822.1	4.0E-06
AT2G03430.1	Ankyrin repeat family protein	NAS6		35.8	63.5	542.5	2.0E-02
AT3G48670.2	XH/XS domain-containing protein	IDN2, RDM12		2.3	2.2	189.3	1.6E-04
AT2G26780.1	ARM repeat superfamily protein	ECM29		0.3	1.4	183.9	6.9E-04
AT5G27960.1	AGAMOUS-like 90	AGL90		0.3	7.5	170.6	2.2E-05
AT3G13090.1	multidrug resistance-associated protein 8	ATMRP8, ABCC6, MRP8		0.3	1.5	131.3	1.5E-02
AT3G13330.1	proteasome activating protein 200	PA200	PA200	0.8	2.0	111.6	1.1E-05
AT2G47360.1	unknown protein;BEST Arabidopsis thaliana protein match is: unknown protein (TAIR:AT1G02570.1);Has 58 Blast hits to 55 proteins in 6 species: Archae - 0;Bacteria - 0;Metazoa - 0;Fungi - 0;Plants - 58;Viruses - 0;Other Euka			2.2	2.6	79.0	3.5E-05
AT5G66420.1	CONTAINS InterPro DOMAIN/s: Uncharacterised conserved protein UCP033271 (InterPro:IPR008322), TIM-barrel signal transduction protein, predicted (InterPro:IPR009215);Has 30201 Blast hits to 17322 proteins in 780 species: Archae -			0.2	1.0	57.7	2.1E-04
AT2G04235.1	unknown protein;FUNCTIONS IN: molecular_function unknown;INVOLVED IN: biological_process unknown;LOCATED IN: cellular_component unknown;EXPRESSED IN: cultured cell;Has 713 Blast hits to 345 proteins in 122 species: Archae - 2			1.2	0.9	53.5	2.5E-05
AT2G28070.1	ABC-2 type transporter family protein			0.2	1.2	40.4	5.6E-03
AT2G17200.1	ubiquitin family protein	DSK2	DSK2	0.7	4.1	39.1	1.3E-04
AT5G05180.2	unknown protein;BEST Arabidopsis thaliana protein match is: unknown protein (TAIR:AT3G10880.1);Has 50151 Blast hits to 28081 proteins in 1838 species: Archae - 860;Bacteria - 6013;Metazoa - 25867;Fungi - 3836;Plants - 2236;			1.0	2.5	36.9	4.1E-06
AT2G14170.2	aldehyde dehydrogenase 6B2	ALDH6B2		0.8	1.2	28.1	1.7E-04
AT2G37650.1	GRAS family transcription factor			0.2	1.1	23.9	8.1E-04
AT4G25500.2	arginine-serine-rich splicing factor 35	ATRSP35, ATRSP40, AT-SRP40, RSP35, RS40, At-RS40		1.7	17.4	21.9	6.8E-04
AT5G15390.1	tRNA/rRNA methyltransferase (SpoU) family protein			0.5	4.0	20.7	3.0E-04
AT3G05070.1	CONTAINS InterPro DOMAIN/s: mRNA splicing factor, Cwf18 (InterPro:IPR013169);Has 292 Blast hits to 292 proteins in 153 species: Archae - 0;Bacteria - 0;Metazoa - 118;Fungi - 83;Plants - 38;Viruses - 11;Other Eukaryotes - 42			0.8	8.1	15.3	1.8E-04
AT3G26020.2	Protein phosphatase 2A regulatory B subunit family protein			0.3	1.8	14.7	1.1E-04

Table 3.S9: Proteasome associated proteins identified after RPT4b-based affinity purification under minus ATP conditions. A table of proteins identified as significantly enriched at FDR 0.01 in the indicated affinity preparations. Shown are the protein's ATG identifiers, descriptions from *The Arabidopsis Information Resource version 10 (TAIR10)*, symbols from TAIR10, average MS/MS count, average sequence coverage, fold change compared to control using the MaxLFQ value from MaxQuant, and *p*-values obtained from the volcano plots in Figure 3.4 E.

ATG	TAIR10 Description	Symbol(s)	Proteasome Annotation	Average			Fold Change Compared to Control	<i>P</i> -Value
				Average MS/MS Count*	Sequence Coverage (%)			
AT2G26780.1	ARM repeat superfamily protein		ECM29	153.7	55.2		499.0	2.73E-03
AT2G03430.1	Ankyrin repeat family protein		NAS6	22.8	68.5		155.3	1.65E-03
AT3G13090.1	multidrug resistance-associated protein 8		ATMRP8, ABCC6, MRP8	0.0	1.3		135.8	4.78E-06
AT3G15180.1	ARM repeat superfamily protein		HSM3	54.5	51.8		110.2	4.30E-06
AT5G27960.1	AGAMOUS-like 90		AGL90	1.0	7.5		94.1	6.40E-03
AT5G57950.1	26S proteasome regulatory subunit, putative		NAS2	2.8	32.3		82.1	5.94E-06
AT3G53970.1	proteasome inhibitor-related			1.8	11.1		23.4	9.04E-05
AT2G28070.1	ABC-2 type transporter family protein			0.0	0.7		19.3	1.01E-04
AT5G49550.1	Putative homolog of mammalian BLOC-1 Subunit 2. Protein - protein interaction with BLOS1.	BLOS2		0.7	10.5		16.9	5.10E-04
AT5G21040.2	F-box protein 2		FBX2	0.5	2.6		14.8	7.77E-05
AT3G13330.1	proteasome activating protein 200	PA200	PA200	0.0	0.7		13.7	2.90E-03
AT5G13450.1	delta subunit of Mt ATP synthase		ATP5	0.7	7.5		8.8	1.42E-03

Table 3.S10: Proteasome associated proteins identified after RPT4b-based affinity purification under plus ATP conditions. A table of proteins identified as significantly enriched at FDR 0.01 in the indicated affinity preparations. Shown are the protein's ATG identifiers, descriptions from *The Arabidopsis Information Resource version 10 (TAIR10)*, symbols from TAIR10, average MS/MS count, average sequence coverage, fold change compared to control using the MaxLFQ value from MaxQuant, and p-values obtained from the volcano plots in Figure 3.4 F

ATG	TAIR10 Description	Proteasome Symbol(s)	Annotation	Average			P-Value
				Average MS/MS Count*	Sequence Coverage (%)	Fold Change Compared to Control	
AT3G15180.1	ARM repeat superfamily protein	HSM3		27.3	36.2	1024.2	7.3E-06
AT5G57950.1	26S proteasome regulatory subunit, putative	NAS2		18.0	59.9	822.1	4.0E-06
AT2G03430.1	Ankyrin repeat family protein	NAS6		35.8	63.5	542.5	2.0E-02
AT3G48670.2	XH/XS domain-containing protein	IDN2, RDM12		2.3	2.2	189.3	1.6E-04
AT2G26780.1	ARM repeat superfamily protein	ECM29		0.3	1.4	183.9	6.9E-04
AT5G27960.1	AGAMOUS-like 90	AGL90		0.3	7.5	170.6	2.2E-05
AT3G13090.1	multidrug resistance-associated protein 8	ATMRP8, ABCC6, MRP8		0.3	1.5	131.3	1.5E-02
AT3G13330.1	proteasome activating protein 200	PA200	PA200	0.8	2.0	111.6	1.1E-05
AT2G47360.1	unknown protein;BEST Arabidopsis thaliana protein match is: unknown protein (TAIR:AT1G02570.1);Has 58 Blast hits to 55 proteins in 6 species: Archae - 0;Bacteria - 0;Metazoa - 0;Fungi - 0;Plants - 58;Viruses - 0;Other Euka			2.2	2.6	79.0	3.5E-05
AT5G66420.1	CONTAINS InterPro DOMAIN/s: Uncharacterised conserved protein UCP033271 (InterPro:IPR008322), TIM-barrel signal transduction protein, predicted (InterPro:IPR009215);Has 30201 Blast hits to 17322 proteins in 780 species: Archae -			0.2	1.0	57.7	2.1E-04
AT2G04235.1	unknown protein;FUNCTIONS IN: molecular_function unknown;INVOLVED IN: biological_process unknown;LOCATED IN: cellular_component unknown;EXPRESSED IN: cultured cell;Has 713 Blast hits to 345 proteins in 122 species: Archae - 2			1.2	0.9	53.5	2.5E-05
AT2G28070.1	ABC-2 type transporter family protein			0.2	1.2	40.4	5.6E-03
AT2G17200.1	ubiquitin family protein	DSK2	DSK2	0.7	4.1	39.1	1.3E-04
AT5G05180.2	unknown protein;BEST Arabidopsis thaliana protein match is: unknown protein (TAIR:AT3G10880.1);Has 50151 Blast hits to 28081 proteins in 1838 species: Archae - 860;Bacteria - 6013;Metazoa - 25867;Fungi - 3836;Plants - 2236;			1.0	2.5	36.9	4.1E-06
AT2G14170.2	aldehyde dehydrogenase 6B2	ALDH6B2		0.8	1.2	28.1	1.7E-04
AT2G37650.1	GRAS family transcription factor			0.2	1.1	23.9	8.1E-04
AT4G25500.2	arginine-serine-rich splicing factor 35	ATRSP35, ATRSP40, AT-SRP40, RSP35, RS40, At-RS40		1.7	17.4	21.9	6.8E-04
AT5G15390.1	tRNA/rRNA methyltransferase (SpoU) family protein			0.5	4.0	20.7	3.0E-04
AT3G05070.1	CONTAINS InterPro DOMAIN/s: mRNA splicing factor, Cwf18 (InterPro:IPR013169);Has 292 Blast hits to 292 proteins in 153 species: Archae - 0;Bacteria - 0;Metazoa - 118;Fungi - 83;Plants - 38;Viruses - 11;Other Eukaryotes - 42			0.8	8.1	15.3	1.8E-04
AT3G26020.2	Protein phosphatase 2A regulatory B subunit family protein			0.3	1.8	14.7	1.1E-04

Table 3.S11: Fold Change dNSAF values comparing Slice 1 and Smear from Figure 3.6. Fold change values were calculated for both treated and untreated samples by dividing the dNSAF values for Smear / Slice 1. #DIV/0! Is listed as a divide by zero error, in that those proteins were detected in the smear but not in Slice 1. Larger values were color coded by red in Microsoft Excel, and smaller values were color coded by blue.

Protein	Slice1		Smear		Fold Change		Fold Change
	Treated	Untreated	Treated	Untreated	Treated	Smear/CP(Slice 1)	Smear/CP(Slice 1)
PBAC2	0.0000	0.0000	0.0089	0.0108	#DIV/0!		#DIV/0!
PBAC3	0.0000	0.0000	0.0098	0.0048	#DIV/0!		#DIV/0!
PBAC4	0.0000	0.0000	0.0188	0.0184	#DIV/0!		#DIV/0!
UMP1	0.0000	0.0000	0.0007	0.0021	#DIV/0!		#DIV/0!
PAP1	0.0002	0.0000	0.0162	0.0188	73.2669		#DIV/0!
PBAC1	0.0008	0.0000	0.0220	0.0295	26.1164		#DIV/0!
UBQ	0.0025	0.0008	0.0408	0.0107	16.2543		12.7573
PA200	0.0016	0.0000	0.0067	0.0024	4.1181		#DIV/0!
PTRE1	0.0009	0.0011	0.0037	0.0023	4.0152		2.1968
PAG1	0.0664	0.0675	0.0935	0.0336	1.4069		0.4973
PAA2	0.0553	0.0532	0.0460	0.0219	0.8316		0.4119
PAD2	0.0276	0.0271	0.0149	0.0200	0.5394		0.7403
PAF1	0.0491	0.0485	0.0245	0.0256	0.4997		0.5285
PAA1	0.0436	0.0573	0.0210	0.0195	0.4811		0.3410
PAB2	0.0285	0.0330	0.0133	0.0114	0.4666		0.3457
PAB1	0.0443	0.0391	0.0199	0.0121	0.4495		0.3105
PAE2	0.0287	0.0275	0.0122	0.0151	0.4238		0.5485
PAE1	0.0446	0.0437	0.0181	0.0175	0.4045		0.4012
PAC1	0.0588	0.0551	0.0171	0.0162	0.2915		0.2932
PAD1	0.0473	0.0422	0.0093	0.0123	0.1961		0.2912
PBD2	0.0339	0.0343	0.0050	0.0041	0.1463		0.1180
PAF2	0.0162	0.0155	0.0019	0.0021	0.1174		0.1356
PBB2	0.0186	0.0170	0.0017	0.0046	0.0892		0.2713
PBB1	0.0225	0.0228	0.0017	0.0088	0.0742		0.3874
PBA1	0.0447	0.0386	0.0025	0.0036	0.0568		0.0944
PBD1	0.0234	0.0236	0.0011	0.0025	0.0466		0.1056
PBF1	0.0653	0.0774	0.0025	0.0059	0.0383		0.0761
PBG1	0.0611	0.0652	0.0004	0.0012	0.0061		0.0186
PBC1	0.0341	0.0314	0.0000	0.0113	0.0000		0.3616
PBC2	0.0150	0.0146	0.0000	0.0000	0.0000		0.0000
PBE1	0.0506	0.0538	0.0000	0.0004	0.0000		0.0083
PBE2	0.0173	0.0155	0.0000	0.0000	0.0000		0.0000

Chapter 4

FUTURE DIRECTIONS AND CLOSING REMARKS

4.1. Introduction

The Ubiquitin Proteasome System (UPS) plays a central role in protein degradation in all eukaryotes, and impacts almost all aspects of plant growth and development. The selective removal of key regulatory and aberrant proteins occurs via the covalent attachment of the polypeptide ubiquitin followed by degradation via the 26S proteasome. This central enzymatic effector of the UPS is made up of two sub-particles, the regulatory particle (RP) responsible for recognizing and unfolding ubiquitylated substrates, and the core particle (CP) which degrades the unfolded protein. The bulk of my thesis has been spent analyzing the sub-particles of the 26S proteasome and their interacting proteins, with follow-up analyses characterizing the interaction network of several CP-specific proteins that may be involved in plant proteasome assembly.

In Chapter 2, I developed the first downstream label-free quantitative analysis tool for the Morpheus mass spectrometry search engine, Morpheus Spectral Counter (MSpC) (Gemperline *et al.*, 2016). The goal was to use this software to perform analyses of the plant 26S proteasome complex with a focus on analyzing

its isoform composition. The Morpheus search engine typically identifies a greater number of peptide spectral matches (PSMs), and is considerably faster than other search engines (Wenger and Coon, 2013). MSpC combined with Morpheus outperformed other spectral counting based measures in both speed and accuracy (Gemperline *et al.*, 2016). MSpC implements the proven dNSAF algorithm that corrects for protein length and uniqueness (Zhang *et al.*, 2010). The software maps Peptide-Spectral Matches (PSMs) to Protein groups and determines if they are distinct or shared, and then corrects the spectral counts based on protein length and total number of PSMs identified in an experiment. I was able to determine the relative incorporation level for most of the CP subunit isoforms, and these results matched previously published data that utilized UV-VIS quantitative methods coupled with top-down mass spectrometry (Gemperline *et al.*, 2016; Russell *et al.*, 2013). MSpC in combination with MS/MS analyses was also used in Chapter 3 to analyze native-PAGE slices of proteasome samples affinity purified from inhibitor-treated tissue. MSpC comes with an easy to use graphic user interface and indeed, others have already started to use the software.

In Chapter 3, I developed an affinity purification of the RP based on two isoforms of RPT4, RPT4a and RPT4b. I identified all of the core 26S proteasome subunits and their isoforms in these affinity preparations, showing that this protocol was

suitable for analysis of the plant RP. These purifications coupled with a previously developed purification that targeted the CP enabled me to affinity-purify samples that were statistically enriched for either the RP or CP sub-complexes. LFQ-MS/MS of these CP and RP enriched samples, along with mock affinity controls, enabled me to confidently identify a suite of interacting proteins that were CP- and RP-specific. The most abundant CP-associated proteins identified in our samples were putative assembly chaperones PBAC1-4, while the most abundant RP-associated proteins identified were putative RP assembly chaperones NAS2, NAS6, and HSM3. Taken together these data suggest that affinity preparations targeting the CP contain only mature RP, and affinity purifications targeting the RP contain only mature CP.

Next, I used a combination of yeast-two-hybrid and split-YFP approaches to build an interaction network involving a subset of the CP-specific putative assembly chaperones (PBAC1-4), showing similar interactions as observed between their mammalian and yeast orthologs. I also show that the novel plant, CP-specific interactor, PAP1, interacts with putative assembly chaperone PBAC1, suggesting that PAP1 may have a role in plant proteasome assembly. Additionally, I also used native-PAGE coupled with MS/MS analyses to identify possible assembly intermediates of the 26S proteasome that contain these putataive, CP-specific chaperones and show that these proteins co-migrate with proteasome subunits. PBAC1-4, and

PAP1 are found to associate with fractions containing mostly α subunits and the assembly chaperone UMP1; whereas only PBAC1, PBAC2, and PAP1 associate with more mature forms of the CP. These data suggest that PBAC3 and PBAC4 may be acting early in assembly, while PBAC1 and PBAC2 may be acting later. This is reminiscent to the role played by orthologous proteins in mammalian and yeast systems (Le Tallec *et al.*, 2007). Taken together the analyses herein provide the first glimpse at plant proteasome assembly, and the likely players involved. Finally, I go on to show that there are very few, if any, differences in protein isoform composition between proteasomes affinity-purified with tagged RPT4a and RPT4b, suggesting that plants assemble their isoforms in a random fashion.

4.2. Future Directions

4.2.1. Proteasome Purifications

Now that I have developed an affinity purification strategy that is based on the RP base subunit RPT4, it may be useful to also develop an affinity purification that is based on an RP lid subunit. We now have tools in plants to affinity purify the CP via PAG1, and the RP-base via RPT4 through my work presented in Chapter 3; however, we have no tools that specifically target the plant RP-lid. Tools in yeast exist to affinity purify the CP via PRE1, the RP base via RPT1, and the RP lid via RPN11

(Leggett *et al.*, 2005, 2002). RP-lid affinity preparations of the yeast proteasome based on RPN11 were able to identify the de-ubiquitylating enzyme Ubp6, and an E3 ubiquitin ligase Hul5, which both have likely orthologs in *Arabidopsis*. Unfortunately, to date, Hul5 has not been identified in plant preparations. While I have been able to identify a suite of RP interacting proteins including likely orthologs for the assembly chaperones NAS2, NAS6, and HSM3, there may be additional proteins that associate with the plant RP lid specifically. Affinity purifications based on the RPN5 subunit may be an attractive candidate to explore, as C-terminal tags with GFP have been used successfully in the past to rescue plants deficient in RPN5 (Book *et al.*, 2009; Marshall *et al.*, 2015). Unfortunately, structural studies of RP-lid subunits in yeast have suggested that the C-termini are essential for proper RP-lid assembly (Estrin *et al.*, 2013). Therefore, an N-terminal tag may be more appropriate. I have already generated constructs that contain N-terminally FLAG-tagged variants of RPN5a, and RPN5b driven by their native promoters, that could be transformed into *Arabidopsis*, and analyzed. In my RP-based affinity preparations, I failed to identify several proteins that likely help the plant complex process substrates including obvious plant orthologs of UCH37, and RPN13. In yeast, RPN13 acts in concert with UCH37 to appropriately process substrates, with RPN13 binding ubiquitin, and UCH37 de-ubiquitylating substrates (Husnjak *et al.*,

2008; Jiao *et al.*, 2014). RPN13 binds the proteasome subunit RPN2 in a highly labile and sub-stoichiometric association (Sakata *et al.*, 2012). It may be reasonable to try cross-linking strategies coupled with MS/MS analyses to identify additional sub-stoichiometric and labile interactors of the plant proteasome.

4.2.2. CP Assembly Chaperones


The most abundant CP-specific proteasome-associated proteins (PAPs) identified, PBAC1-4, may be orthologs of the mammalian assembly chaperones PAC1-4. We have evidence that these proteins associate with the proteasome and are CP-specific, as they are not identified in affinity purifications that target the RP via RPT4. These putative chaperones form an interaction network that is reminiscent of their yeast and mammalian counterparts, and are found in putative assembly intermediates. While we have indirect evidence for protein interactions through both yeast-two-hybrid and bimolecular fluorescent complementation analysis, direct, *in vitro* binding assays should be performed to validate these indirect interactions. This would provide further support for these proteins forming an interaction network like their yeast and mammalian counterparts. *In vitro* binding assays would require expression in heterologous systems such as *E. coli*, and purification of these PBAC1-4 proteins. I have already generated expression vectors using two separate tags (6x

histidine and glutathione-s-transferase (GST)), and performed some preliminary expression in *E. coli* of PBAC1-3, and PAP1 (Figure 4.1). PBAC4 was also recently cloned into the same expression vectors, but its expression has not been tested. If post-translational modifications of these chaperones are required for their interaction, this may be a potential problem with *in vitro* binding assays, and would not be recapitulated in an *E. Coli* recombinant expression system. Another potential issue with *in vitro* expression is that others have reported that yeast Pba3 and Pba4 were completely insoluble unless co-expressed (Kusmierczyk *et al.*, 2008), so an alternative expression system may be required such as the Duet expression system, to allow expression of both PBAC3 and PBAC4 at the same time (Scheich *et al.*, 2007).

Despite these downfalls, it would be useful to generate antibodies against PBAC1-4 as this would be helpful in determining the content of plant assembly intermediates. Assuming *in vitro* expression of PBAC1-4 can be accomplished, these proteins (either expressed as heterodimeric pairs or alone) could also be used as bait to enrich for potential assembly intermediates from crude *Arabidopsis* protein extracts. Proteins that interact with these bait proteins could then be determined by MS/MS analyses.

While PBAC1-4 associate with the CP and form an interaction network, it is

[Figure 4.1 caption follows on next page]



unclear if these proteins function as assembly chaperones. Bioinformatic analyses, including PSI-BLAST and InterProScan domain analyses, are insufficient to determine that these proteins function as true CP assembly chaperones in plants. Analysis of *Arabidopsis* plants deficient in *PBAC1-4* will be useful in dissecting the plant proteasome assembly pathway. I have not been able to confirm any mutants in *PBAC1-4*; however, there may be candidates that could be tested (see Table 4.1). These genes are quite small, and identifying T-DNA insertions has proved challenging. A CRISPR based approach may be useful to precisely target these genes to generate single mutants, as well as combinations of mutants.

When yeast proteasomes are compromised, either chemically, or by genetic disruption, the Pba1 or Pba2 mutants show no phenotype; however, the double mutant exhibits a strong growth phenotype (Kusmierczyk *et al.*, 2011). This suggests that double mutant analysis may be necessary to study the putative plant assembly chaperones. Once there are sufficient mutants identified in *PBAC1-4*, it would be

Figure 4.1 (preceding page): Recombinant protein expression of PAP1 and PBAC1-3. Recombinant proteins were expressed in *E. coli* BL21 DE3 with the indicated GST or 6xHis Tags. The corresponding protein extracts were then subjected to SDS-PAGE and then stained with Coomassie Blue. Protein extracts from IPTG-induced (+IPTG) and un-induced (-IPTG) control cultures are shown. At the top of each lane, samples are labeled with the name of the expressed protein followed by a number that defines the proteins expected size (in kDa). Black arrows on the gel indicate the likely induced proteins.

Table 4.1: Candidate alleles for PBAC1-4. Candidate alleles were obtained from The Arabidopsis Information Resource (Berardini *et al.*, 2015) using T-DNA Express. SALK or SAIL identifiers are shown for PBAC1-4 with their predicted insertion sites.

Putative Chaperone	ATG identifier	Potential Insertions	Predicted Insertion Location
PBAC1	AT3G25545	SALK_112630	Exon
PBAC2	AT3G18940	SALK_104978C SAIL_598_E03 SAIL_477_D092	5'UTR Exon Intron
PBAC3	AT5G14710	SALK_092651 SALK_006045	Exon Intron
PBAC4	AT1G48170	SALK_10090 SALK_078009	Exon Exon

useful to cross these mutants into the *FLAG-PAG1* affinity purification background, so that potential assembly intermediates could be purified and analyzed. Alternatively, CRISPR mutagenesis of each putative assembly chaperone gene could be performed in the *PAG1-FLAG* background, possibly shortcutting the need to select for triple mutants in the putative assembly chaperone, the *PAG1-FLAG* allele, and the *pag1-1* allele in a standard crossing scheme.

One complication with this mutant analysis strategy is that deficiency in the mammalian assembly chaperone PAC3, is lethal in mice. Several strategies could be used to overcome this. Hypomorphs could be generated using amiRNA silencing, or heterozygous mutants could be rescued with conditional expressors (Eamens *et al.*, 2014). Conditional knockout strategies could be used by driving a Cre promoter using tissue or developmental time point-specific promoters to excise a LoxP flanked rescue cassette (Louwerse *et al.*, 2007). Given the conserved nature of the HbYX

motif found in PBAC1, and the HbYX-like motif found in PBAC2 it will be useful to study the role of these domains in plant proteasome assembly. HbYX-less variants should be generated to see if they can rescue their respective mutants (assuming a phenotype for each (or both) is found).

An alternative, and possibly faster strategy for confirming that these PBAC1-4 proteins may be involved in proteasome assembly would be to try cross-species mutant rescue experiments. I have already obtained yeast deficient in Pba1 and Pba2 from the Hochstrasser lab (Kusmierczyk *et al.*, 2011), and plans are underway to rescue these yeast mutants with their putative *Arabidopsis* orthologs PBAC1 and PBAC2 with and without HbYX motifs. It may be possible to rescue these deficient yeast lines with *Arabidopsis* *PBAC1*, and/or *PBAC2*; however, yeast deficient in the assembly chaperone Ump1, cannot be rescued with the human variant, suggesting that this strategy may still be difficult, especially considering the poor sequence conservation between yeast and *Arabidopsis* (Burri *et al.*, 2000). Another complicating factor is that PAP1 may be involved in plant proteasome assembly, as it interacts with PBAC1. If PAP1 is involved in CP assembly, it would suggest that the plant system has diverged considerably from all known CP assembly systems, and this may make it difficult to rescue a yeast system with a plant ortholog.

4.2.3. PAP1

The plant-specific protein PAP1 was localized to the CP specifically, found to interact with PBAC1, and also contained a conserved HbYX motif. This suggested that PAP1 may be involved in plant proteasome assembly; however, additional analyses will be necessary to determine PAP1's role, if any, in construction of the plant particle. While PAP1 interacts with PBAC1 indirectly, it would be useful to test this interaction *in vitro*. Like above with *PBAC1-3*, I have already cloned *PAP1* into an expression vector and shown that it can be expressed in *E. coli* (Figure 4.1). While, like above, there could be issues with protein solubility or required post-translation modifications for the PAP1 interaction, it would still be useful to test the PAP1-PBAC1 interaction. I have already identified several mutations that disrupt *PAP1* (Figure 4.2), with *pap1-3* in particular causing a significant reduction in the *PAP1* transcript; however, plants homozygous for *pap1-3* display a remarkably normal phenotype. These preliminary data suggest that *PAP1* is not essential for plant growth and development. However, initial analyses of total protein extracted from *pap1-3* plants using glycerol gradient centrifugation showed a small shift in CP subunits PAG1, and PBA1 (Figure 4.3) suggesting that the CP may be destabilized. This analysis should be repeated to confirm the small shift in CP subunits. While the gross morphology of *pap1-3* seemed normal, glycerol gradient centrifugation

may be a way to show that the CP is slightly destabilized in the *pap1-3* background. I have already crossed *pap1-3* into the *PAG1-FLAG pag1-1* line and once triple mutants are isolated, it may be possible to affinity purify assembly intermediates using our standard anti-FLAG affinity protocol. It would be useful to generate antibodies against PAP1 so that we can confirm that *pap1-3* is a null allele and that no protein product is produced. Additionally, antibodies would allow us to determine if PAP1 is present in any assembly intermediates identified.

If PAP1 is involved in CP assembly, it could replace PBAC2 in a PBAC1/2 heterodimer, forming PBAC1-PAP1 complex, or PAP1 could form a trimeric complex with PAP1, PBAC1, and PBAC2. If the above expression system can be used to purify these proteins, this model should be tested directly using *in vitro* binding assays. An alternative FLAG tag could be used with PAP1, and we should test for simultaneous co-IP with both PBAC1 and PBAC2. Additionally, the formation of a stable trimeric complex with PAP1, PBAC1, and PBAC2 using native-PAGE co-migration assays would support the hypothesis of a trimeric complex. In the alternative model of PAP1 replacing PBAC2 in a PBAC1/2 heterodimer, if stable PBAC1 and PBAC2 binding can be achieved as described above, competition assays could be used to see if PAP1 can competitively bind PBAC1 and cause release of PBAC2.





Figure 4.2: Insertion alleles for *PAP1* and qRT-PCR analysis. (A) A gene diagram showing 5' and 3' untranslated regions in black, with exons in green, and introns indicated as a black line. The positions of T-DNA insertions (as determined by sequencing the products of PCR amplification overlapping with the corresponding T-DNA borders) for *pap1-1*, *pap1-2*, and *pap1-3* are shown in red. **(B)** qRT-PCR analysis of *PAP1* expression in wild type (Col-0), *pap1-1*, *pap1-2*, and *pap1-3* seedlings using primers that spanned the full length of *PAP1*. For each mutant, expression levels were normalized relative to wild type levels (set at 100%) of the average of two reference genes, *PP2A* and, *ACT2*. Error bars indicate standard deviation for 3 different biological replicates.

[Figure 4.3 caption follows on next page]



Besides *in vitro* binding assays, we could also test for PAP1 cross-species mutant rescue. Given that PAP1 binds with PBAC1 and contains a HbYX motif, it may be possible for PAP1 to rescue yeast deficient in Pba1 and Pba2. PAP1 with and without its HbYX motif should be assayed to determine if it can successfully rescue yeast mutants deficient in Pba1 and Pba2.

It is also possible that PAP1 acts in plant proteasome assembly, but indirectly. In mammals, the PAC1/PAC2 proteasome assembly chaperone heterodimer interacts with and is regulated by iRhom1 (Lee *et al.*, 2015). iRhom1 is a member of the Rhomboid protease family that is generally located in the endoplasmic reticulum (ER); however, iRhom1 lacks protease catalytic activity, and is thought to inhibit translocation of epithelial growth factor ligand family members to the Golgi, by targeting them and bringing them to the proteasome (Lee *et al.*, 2015). A recent study

Figure 4.3 (preceding page): Glycerol gradient centrifugation analysis of *pap1-3*. Total protein extracted from *pap1-3* plants was fractionated by glycerol gradient centrifugation as described previously (Book *et al.*, 2010). Fraction numbers indicate 0.5 mL fractions, with the first fraction being the least dense at 10% glycerol, and the last fraction being the densest at 40% glycerol. 10 µL of each fraction was subjected to SDS-PAGE analyses followed by immunoblot analyses with antibodies for the CP subunits PAG1 and PBA1, and the RP subunits RPT2, RPT4, and RPN1 (defined to the left of each panel). A small shift is observed in *pap1-3* samples relative to wild type for fractions 7 and 8 when immunoblotting for the CP subunits PAG1 and PBA1 which may be due to a slight disruption in CP stability, or possibly due to the appearance of complexes containing fewer subunits that might represent assembly intermediates. The RP distribution does not seem to be different between wild-type Col-0 and *pap1-3*

showed that reduced levels of iRhom1 resulted in reduced proteasome capacity; whereas increased levels of iRhom1 stimulated proteasome activity (Lee *et al.*, 2015). Intriguingly, iRhom1, was found to increase when cells were treated with the ER stressor tunicamycin and also with other protein synthesis inhibitors, hygromycin and cycloheximide (Lee *et al.*, 2015). In iRhom1 knockdown cells PAC1 was found to be rapidly degraded, and both endogenous PAC1 and PAC2 levels were reduced (Lee *et al.*, 2015). The authors then go on to show that iRhom1 has a role in stabilizing the PAC1/2 heterodimer (Lee *et al.*, 2015). A model emerges in which iRhom1 regulates proteasome assembly by stabilizing this PAC1/2 heterodimer, and plays a role in ER stress response (Lee *et al.*, 2015). Because PAP1 interacts with the putative PAC1 ortholog PBAC1, and could be performing a similar role to iRhom1, *pap1-3* mutants should be assayed for phenotypes under tunicamycin treatments that induce ER stress. Additionally, GFP-tagged PAP1 lines should be developed using the *PAP1* native promoter to see if PAP1 localizes to the ER. If PAP1 is performing a similar function as iRhom1, and antibodies become available for PBAC1, it would also be useful to test PBAC1 stability in the *pap1-3* background. While PAP1 may be playing a similar role as iRhom1, PAP1 does not seem to share any sequence homology with iRhom1, and iRhom1 does not contain a HbYX motif, whereas PAP1 does. Additionally, a potential ortholog for iRhom1, ATRBL1 (Rhomboid-like

1) can be identified by protein BLAST (expectation value 3E-19). ATRBL1 may be an interesting follow up candidate as potential regulator of PBAC1 and PBAC2 activity.

It is possible that PAP1 is not involved in plant proteasome assembly. PAP1 could act as a shuttling factor or adapter to help bring other proteins in close proximity to the CP, or PAP1 could be used to anchor the CP to a specific sub-cellular localization. *pap1-3* mutants could be crossed to a PAG1-GFP reporter line (Marshall *et al.*, 2015) to determine if the CP is miss-localized when deficient in *PAP1*.

4.2.4. RP Assembly Chaperones

While I have been able to identify RP-specific interactors that are putative orthologs for NAS2, NAS6, and HSM3, further analyses will be necessary to determine if these function in RP assembly. Mutants should be identified in *NAS2*, *NAS6*, and *HSM3*. Once plants defective in these putative chaperones are obtained, they could then be crossed into the FLAG-RPT4a/b lines to determine if any assembly intermediates could be identified in affinity preparations that are based on the RP.

4.2.5. Identifying CP Assembly Intermediates

MS/MS analyses of proteasomes affinity purified from tissue treated with the proteasome inhibitor MG132 fractionated by native-PAGE have been useful in iden-

tifying potential assembly intermediates. In addition to these studies, I performed sample preparation for a similar analysis along a time course of 0, 4, 8, and 16 hours of MG132 treatment. Unfortunately, MS/MS analyses of these samples did not identify all proteasome subunits, and we did not identify a large number of PSMs for the proteasome subunits that were identified suggesting that there were issues, either with the sample preparation, or with sensitivity of the MS.

Additional strategies, such as those used in yeast and mammals, could be used to induce the formation of assembly intermediates. Overexpression of the β 7 subunit, along with a variant of β 5 lacking its propeptide induced the formation of 15S half-proteasome intermediates (Li *et al.*, 2007). A similar strategy using the β 5 (PBE1/2) and β 7 (PBG1) subunits may be useful to induce the formation of assembly intermediates in *Arabidopsis*.

As described above, recombinant PBAC1-4 and/or PAP1 could also be used to enrich for potential CP assembly intermediates. A similar strategy could be used to identify RP-assembly intermediates using recombinant NAS2, NAS6, or HSM3 in co-IP experiments to determine if they form complexes ([Hsm3 with Rpn1, Rpt1, and Rpt2], [Nas2 with Rpt4, and Rpt5], [Nas6 with Rpt3], and [Rpn14 with Rpt6]), like their yeast and mammalian counterparts by MS/MS analyses.

4.2.6. Isoform-Specific Proteasomes

MS/MS analyses of affinity preparations enriched for either RPT4a or PRT4b showed a similar complement of proteasome subunit isoforms suggesting that plants assemble proteasomes randomly with respect to isoform incorporation, and do not form proteasome isotypes. However, I have only performed these affinity enrichments in young seedlings, at a single developmental time point. Data from rice suggests that RPT isoforms may differ in expression in different tissue types (Shibahara *et al.*, 2004). Therefore, proteasomes should be affinity purified using the FLAG-RPT4a and FLAG-RPT4b lines from different tissue types and/or developmental time points to determine if *Arabidopsis* assembles developmental or tissue-specific proteasome isotypes. Additionally, it may be possible that isoform subunit incorporation changes when plants are exposed to different environmental conditions, or experience different stressors. Indeed, transcriptional analyses of *Arabidopsis* proteasome subunit isoforms revealed that typically only one isoform is upregulated in response to MG132 (Gladman *et al.*, 2016), and therefore this would be an interesting stressor to assay for differential isoform incorporation. The PAG1-FLAG line could also be used for these analyses, with a focus on CP subunit incorporation.

It may be difficult to obtain enough tissue to perform these experiments. If this

is the case, an alternative strategy using targeted proteomics of total protein extracts could be performed utilizing the Skyline software suite (MacLean *et al.*, 2010). Based on the data obtained from our affinity purifications, we could identify tryptic peptides specific to each subunit isoform and then use this targeted proteomics approach to assay total protein extracts for subunit isoform abundance. One downside to this approach, is that it would only give us an idea of protein levels for each individual isoform present in total protein extracts, and we would not be able to tell if these subunits are effectively incorporated into the proteasome.

4.3. Concluding Remarks

My thesis has contributed to the overall understanding of the *Arabidopsis* UPS with a focus on identifying proteasome associated proteins, many of which may be likely players in the biogenesis of both the CP and RP of the plant proteasome. Along the way, I developed label-free quantitative mass spectrometry software that was useful in assaying proteasome samples for specific isoform distributions, and indeed, others have already started using this software to perform their own label-free quantitative analyses.

I developed an affinity purification method for purifying the 26S proteasome via the RP utilizing two different isoforms of the RPT4 subunit, RPT4a and RPT4b.

These affinity preparations enabled me to explore the possibility that plants generate specific proteasome isotypes, and I was able to show that the incorporation of specific subunit isoforms likely occurs in a random fashion. These affinity purifications that targeted the RP, in conjunction with previously developed CP-based affinity purifications enabled me to identify a suite of RP and CP-specific interactors. Several of these CP-specific interactors may be orthologs of CP assembly chaperones, while several of the RP-specific interactors are likely orthologs of RP assembly chaperones. Follow up characterization of the CP-specific interactors PBAC1-4 show that they form an interaction network that may be similar to their putative yeast and mammalian counterparts. A novel plant-specific protein PAP1 interacts with PBAC1 both via yeast-two-hybrid analyses, and *in planta* using split-YFP assays suggesting that PAP1 may be involved in the biogenesis of the CP. Finally, I show that PBAC1-4 and PAP1 co-migrate with the plant proteasome, and are found in possible assembly intermediates of the CP. While PBAC3 and PBAC4 and UMP1 are found exclusively in fractions that are similar to the 15S half-barrel assembly intermediate, PAP1, PBAC1, and PBAC2 seem to associate with this half-barrel, and with more mature forms of the CP. Taken together these data suggest that UMP1, PBAC3, and PBAC4 are involved in early stages of CP biogenesis, while PAP1, PBAC1, and PBAC2 are likely involved at both early, and later stages.

Together these data provide the first glimpse at the machinery that may drive proteasome biogenesis in plants. Continued analyses of this complex process should yield insights that are central to the entire UPS. As this system is responsible for modulating most if not all aspects of plant growth and development, a general understanding of plant proteasome regulation, and in particular an understanding of proteasome biogenesis, will have impacts in bioenergy research, and hopefully agriculture.

4.4. Literature Cited

- Berardini, T.Z., Reiser, L., Li, D., Mezheritsky, Y., Muller, R., Strait, E., and Huala, E.** (2015). The arabidopsis information resource: Making and mining the gold standard annotated reference genome. *Genesis* **53**: 474–485.
- Book, A.J., Gladman, N.P., Lee, S.S., Scalf, M., Smith, L.M., and Vierstra, R.D.** (2010). Affinity purification of the Arabidopsis 26S proteasome reveals a diverse array of plant proteolytic complexes. *J. Biol. Chem.* **285**: 25554–25569.
- Book, A.J., Smalle, J., Lee, K.H., Yang, P., Walker, J.M., Casper, S., Holmes, J.H., Russo, L.A., Buzzinotti, Z.W., Jenik, P.D., and Vierstra, R.D.** (2009). The RPN5 subunit of the 26S proteasome is essential for gametogenesis, sporophyte development, and complex assembly in Arabidopsis. *Plant Cell* **21**: 460–78.
- Burri, L., Hockendorff, J., Boehm, U., Klamp, T., Dohmen, R.J., and Levy, F.** (2000). Identification and characterization of a mammalian protein interacting with 20S proteasome precursors. *Proc. Natl. Acad. Sci. U.S.A.* **97**: 10348–53.
- Eamens, A.L., McHale, M., and Waterhouse, P.M.** (2014). The use of artificial microRNA technology to control gene expression in *Arabidopsis thaliana*. *Methods Mol. Biol.* **1062**: 211–24.

- Estrin, E., Lopez-Blanco, J., Chacón, P., and Martin, A.** (2013). Formation of an Intricate Helical Bundle Dictates the Assembly of the 26S Proteasome Lid. *Structure* **21**: 1624–1635.
- Gemperline, D.C., Scalf, M., Smith, L.M., and Vierstra, R.D.** (2016). Morpheus Spectral Counter: A computational tool for label-free quantitative mass spectrometry using the Morpheus search engine. *Proteomics* **16**: 920–4.
- Gladman, N.P., Marshall, R.S., Lee, K.H., and Vierstra, R.D.** (2016). The Proteasome Stress Regulon Is Controlled by a Pair of NAC Transcription Factors in Arabidopsis. *Plant Cell* **28**: 1279–96.
- Husnjak, K., Elsasser, S., Zhang, N., Chen, X., Randles, L., Shi, Y., Hofmann, K., Walters, K.J., Finley, D., and Dikic, I.** (2008). Proteasome subunit Rpn13 is a novel ubiquitin receptor. *Nature* **453**: 481–488.
- Jiao, L., Ouyang, S., Shaw, N., Song, G., Feng, Y., Niu, F., Qiu, W., Zhu, H., Hung, L.W., Zuo, X., Eleonora Shtykova, V., Zhu, P., Dong, Y.H., Xu, R., and Liu, Z.J.** (2014). Mechanism of the Rpn13-induced activation of Uch37. *Protein Cell* **5**: 616–30.
- Kusmierczyk, A.R., Kunjappu, M.J., Funakoshi, M., and Hochstrasser, M.** (2008). A multimeric assembly factor controls the formation of alternative 20S proteasomes. *Nat. Struct. Mol. Biol.* **15**: 237–244.
- Kusmierczyk, A.R., Kunjappu, M.J., Kim, R.Y., and Hochstrasser, M.** (2011). A conserved 20S proteasome assembly factor requires a C-terminal HbYX motif for proteasomal precursor binding. *Nat. Struct. Mol. Biol.* **18**: 622–9.
- Le Tallec, B., Barrault, M.B., Courbeyrette, R., Guérois, R., Marsolier-Kergoat, M.C., and Peyroche, A.** (2007). 20S proteasome assembly is orchestrated by two distinct pairs of chaperones in yeast and mammals. *Mol. Cell* **27**: 660–674.
- Lee, W., Kim, Y., Park, J., Shim, S., Lee, J., Hong, S.H., Ahn, H.H., Lee, H., and Jung, Y.K.** (2015). iRhom1 regulates proteasome activity via PAC1/2 under ER stress. *Sci. Rep.* **5**: 11559.
- Leggett, D.S., Glickman, M.H., and Finley, D.** (2005). Purification of proteasomes, proteasome subcomplexes, and proteasome-associated proteins from budding yeast. *Methods Mol. Biol.* **301**: 57–70.

- Leggett, D.S., Hanna, J., Borodovsky, A., Crosas, B., Schmidt, M., Baker, R.T., Walz, T., Ploegh, H., and Finley, D.** (2002). Multiple Associated Proteins Regulate Proteasome Structure and Function. *Mol. Cell* **10**: 495–507.
- Li, X., Kusmierczyk, A.R., Wong, P., Emili, A., and Hochstrasser, M.** (2007). beta-Subunit appendages promote 20S proteasome assembly by overcoming an Ump1-dependent checkpoint. *EMBO J.* **26**: 2339–49.
- Louwerse, J.D., van Lier, M.C.M., van der Steen, D.M., de Vlaam, C.M.T., Hooykaas, P.J.J., and Vergunst, A.C.** (2007). Stable Recombinase-Mediated Cassette Exchange in *Arabidopsis* Using *Agrobacterium tumefaciens*. *Plant Physiol.* **145**: 1282–1293.
- MacLean, B., Tomazela, D.M., Shulman, N., Chambers, M., Finney, G.L., Frewen, B., Kern, R., Tabb, D.L., Liebler, D.C., and MacCoss, M.J.** (2010). Skyline: an open source document editor for creating and analyzing targeted proteomics experiments. *Bioinformatics* **26**: 966–968.
- Marshall, R.S., Li, F., Gemperline, D.C., Book, A.J., and Vierstra, R.D.** (2015). Autophagic Degradation of the 26S Proteasome Is Mediated by the Dual ATG8/U-biquitin Receptor RPN10 in *Arabidopsis*. *Mol. Cell* **58**: 1053–66.
- Russell, J.D., Scalf, M., Book, A.J., Ladror, D.T., Vierstra, R.D., Smith, L.M., and Coon, J.J.** (2013). Characterization and quantification of intact 26S proteasome proteins by real-time measurement of intrinsic fluorescence prior to top-down mass spectrometry. *PLoS One* **8**: e58157.
- Sakata, E., Bohn, S., Mihalache, O., Kiss, P., Beck, F., Nagy, I., Nickell, S., Tanaka, K., Saeki, Y., Forster, F., and Baumeister, W.** (2012). Localization of the proteasomal ubiquitin receptors Rpn10 and Rpn13 by electron cryomicroscopy. *Proc. Natl. Acad. Sci. U.S.A.* **109**: 1479–84.
- Scheich, C., Kummel, D., Soumailakakis, D., Heinemann, U., and Bussow, K.** (2007). Vectors for co-expression of an unrestricted number of proteins. *Nucleic Acids Res.* **35**: e43.
- Shibahara, T., Kawasaki, H., and Hirano, H.** (2004). Mass spectrometric analysis of expression of ATPase subunits encoded by duplicated genes in the 19S regulatory particle of rice 26S proteasome. *Arch. Biochem. Biophys.* **421**: 34–41.
- Wenger, C.D. and Coon, J.J.** (2013). A proteomics search algorithm specifically designed for high-resolution tandem mass spectra. *J. Proteome Res.* **12**: 1377–86.

Zhang, Y., Wen, Z., Washburn, M.P., and Florens, L. (2010). Refinements to Label Free Proteome Quantitation: How to Deal with Peptides Shared by Multiple Proteins. *Anal. Chem.* **82**: 2272–2281.

COLOPHON

This document was typeset with L^AT_EX . It is based on the University of Wisconsin dissertation template created by William C. Benton (available at <https://github.com/willb/wi-thesis-template>).