

Mathematical Stories

Mathematical Stories

Lecture Notes

July 15, 2026

Contents

Introduction	2
Purpose and Philosophy	2
How the Course Works	2
How to Prepare Mathematical Notes	4
1 Euclidean Geometry and Cartesian Coordinates	8
1.1 The Euclidean Geometry Available Before Coordinates	8
1.2 Constructing Cartesian Coordinates from Euclidean Geometry	17
2 Vectors, Bases, and Coordinate Systems	23
2.1 Vectors from Coordinate Displacements	23
2.2 Changing the Basis: New Coordinates for the Same Vector	36
2.3 Beyond a Change of Basis: Polar Coordinates	41
3 Lines and Planes in the Plane and in Space	45
3.1 Lines in the Cartesian Plane	45
3.2 Lines in Three-Dimensional Space	53
3.3 Planes in Three-Dimensional Space	56
4 Second-Degree Equations, Conic Sections, and Quadric Surfaces	59
4.1 Second-Degree Equations and Conic Sections	59
4.2 Second-Degree Surfaces in Three-Dimensional Space	63
Chapter 5	68
Chapter 6	69
Chapter 7	70
Chapter 8	71
Chapter 9	72
Chapter 10	73
Chapter 11	74
Chapter 12	75

Introduction

Purpose and Philosophy

These notes are an invitation to look at mathematics in a different way. For many students, mathematics has long appeared as a collection of formulas, rules, and mnemonic patterns: one recognises the type of problem, chooses a remembered procedure, performs the required operations, and moves on. Such skills have their place, but they are not the heart of mathematics. Mathematics begins when we stop for a moment and ask what we are really looking at, what kind of object has appeared, what question is being asked, what is known, what is unknown, and why a proposed step is justified.

The purpose of this course is therefore not only to practise calculations. We will learn how to read mathematical expressions, formulate precise questions, distinguish an object from its notation, notice structure, and build simple but meaningful chains of reasoning. A solved problem is valuable not only because it produces an answer, but also because it can reveal a method of thought: how to begin, what to observe, which assumptions matter, what can be transformed, what remains unchanged, and how a conclusion follows from earlier facts.

We will try to study this language from the inside. A definition should not be treated as a sentence to memorise, but as a way of introducing a new kind of object. A theorem should not be treated as a decorative statement, but as an answer to a clear question under precise assumptions. An example should not be treated as a template to copy blindly, but as a small experiment in reasoning. Even a simple calculation can show something important if we ask why it works and what it tells us.

For this reason, some habits of thought will deliberately return in different places. The same question of what is fixed and what is variable appears when we study functions, vectors, equations, and transformations. The same need for precision appears when we define a line, compute a derivative, solve a system of equations, or interpret an integral. A mathematical question can often be seen algebraically, geometrically, and conceptually. This repetition is intentional: it helps the reader recognise patterns of reasoning rather than only patterns of exercises.

The number of answers produced is not, by itself, a measure of progress. What matters is the quality of the work. Did we understand the question? Did we identify the object correctly? Did we choose notation consciously? Did we know why each important step was valid? Did we interpret or verify the result? These questions are not an addition to mathematics; they are mathematics being done carefully.

Mathematics is useful in physics, engineering, computer science, data analysis, modelling, and many other fields, but its value is not only practical. It is also a way of seeing. It teaches precision, discipline, imagination, and the ability to move from simple observations to general conclusions. These notes are meant to show a piece of that beauty, strength, and richness. The goal is not merely to survive mathematical techniques, but to begin to appreciate mathematics as the art of asking good questions and giving well-reasoned answers.

How the Course Works

This book is part of a weekly learning process rather than a collection of material to be read only before an examination. Each of the twelve thematic chapters is designed to support one week of study. A chapter provides the mathematical structure for the lecture, a complete narrative to which the student can return, and a starting point for individual problem solving and classroom discussion.

In a fifteen-week semester, the twelve thematic weeks establish the main rhythm of the course. The remaining time can be used to introduce the method of work, connect topics, revise important ideas, and present more substantial solutions.

The Weekly Learning Cycle

A student should not leave a topic behind when the lecture ends. A typical week consists of the following connected stages:

1. **Attend the lecture.** The lecture introduces the topic, motivates the central questions, and demonstrates how the relevant mathematical objects and methods should be read.
2. **Return to the chapter.** Read the corresponding chapter carefully after the lecture. The PDF is not merely a summary of slides. It is an organised narrative containing definitions, explanations, examples, interpretations, and mathematical arguments that can be examined more than once.
3. **Ask questions and clarify the material.** Mark every place where a definition, transition, calculation, or conclusion is not yet clear. Use the text, your own examples, classroom discussion, and appropriate digital tools to resolve these questions.
4. **Solve the complete assigned problem set.** Move from following an explanation to constructing arguments independently. Work on every assigned problem and prepare a separate, self-contained note for each one.
5. **Prepare professional mathematical notes.** Use precise notation, explanatory sentences, justified transitions, and a clear conclusion. The note should show not only the answer, but also how it follows from the given information, definitions, and earlier results.
6. **Present and discuss your work.** During exercise classes, be prepared to explain a solution, answer questions about its steps, and describe what you learned while preparing it.

These stages form one continuous process. Each stage prepares the next, and the presentation and feedback lead back to further reading, revision, and improved work.

Exercise Classes: Presentation, Feedback, and Revision

Presenting mathematics is part of learning mathematics. It reveals whether an argument has genuinely been understood and whether the student can communicate it clearly. During a presentation, the student should be able to read the notation, explain the role of each important formula, justify the method, and respond to questions without relying on AI to speak on the student's behalf.

Feedback given during exercise classes is intended for the whole group, not only for the student whose work is currently being presented. Everyone should follow the argument and listen carefully to the discussion. By comparing different examples, students learn what a complete solution looks like, which explanations make an argument clear, which steps require justification, and which weaknesses require revision.

Students should not focus only on comments addressed directly to their own work. Listening to feedback on the work of others provides a broader set of examples and develops an operational understanding of the course standards: what makes mathematical work clear, justified, and complete, and what still needs improvement.

The feedback concerns the mathematical work and its results, not the personal worth of the student. What is examined is the quality of the note, the understanding demonstrated in the

argument, and the clarity of the presentation. Feedback should identify what already works and what should be changed in order to produce a stronger solution.

The first submitted version does not always have to be the final version. Students should learn to receive comments, reconsider their choices, correct errors, add missing explanations, and improve both the written note and its presentation. Revision is not evidence of failure. It is a normal part of serious intellectual work.

This process is important throughout university study and later in professional life. Students and professionals are given tasks, asked to solve problems, expected to present results, and required to respond to feedback and critical review. Learning to improve an initial solution is therefore not an additional classroom exercise. It is a fundamental habit of responsible academic and professional work.

Progress and Assessment

Progress is demonstrated by the quality and increasing independence of the student's work. By the end of a weekly cycle, the student should be able to identify the main mathematical objects, state the essential definitions, solve representative problems, justify the principal steps, verify or interpret the result, and communicate the argument clearly.

Assessment in the exercise component focuses primarily on how the student works throughout the semester. Evidence of serious engagement includes:

- regular preparation and completion of the assigned problem sets,
- documented attempts to solve problems rather than bare answers,
- active participation in presentations and classroom discussion,
- separate, self-contained notes for individual problems,
- careful revision of work after feedback,
- increasing clarity, independence, and mathematical precision.

A student is not defined by the quality of the first solution or presentation. Some students may initially find mathematical notation, formal writing, or public explanation difficult. A weak beginning does not prevent a very good final grade for the exercise component. A student who works conscientiously, responds to feedback, corrects earlier weaknesses, and demonstrates substantial development may finish the exercise component with a very good grade.

This approach does not lower the standard of mathematical work. It recognises that reaching the standard through sustained effort and revision is itself a meaningful achievement. The exercise-component grade reflects a process of learning, not merely a snapshot taken at the beginning of the semester.

Systematic work according to these guidelines provides the basis for earning a passing grade for the exercise component. The examination is assessed separately and tests the student's mathematical knowledge and ability to use it independently. Regular work does not replace the examination, but it develops the understanding and skills that substantially increase the likelihood of successfully completing the course.

How to Prepare Mathematical Notes

A mathematical note is not merely a place where an answer is recorded. Its purpose is to show how a problem is understood, how relevant information is selected, how a method is chosen, and how a conclusion follows from definitions, assumptions, and earlier results.

Mathematical expressions are compact. A single formula may contain several ideas at once: objects, operations, conditions, and logical relationships. Reading a formula therefore means more than pronouncing its symbols. A student should be able to explain what the objects represent, what operation is being performed, why the expression is meaningful, and what information the formula provides.

The same principle applies to writing. A sequence of formulas without explanatory sentences rarely communicates a complete mathematical argument. Formulas should be connected by words that explain their role. A good solution tells the reader why a calculation is being performed, which definition or theorem is being used, whether its assumptions are satisfied, and how the result answers the original question.

Make the Note Self-Contained

Prepare one separate, self-contained note for each assigned problem, including all of its subparts. Begin by writing the problem or a complete and faithful formulation of it. The reader should not need to search through another document to discover what is being solved. After several weeks, you should still be able to open the note and understand both the question and the solution.

Identify what is given and what must be found, proved, or explained. Introduce the relevant notation and determine what kinds of mathematical objects appear in the problem. Before calculating, ask what information is available and what would count as a satisfactory answer.

Make the Reasoning Visible

Present the principal idea or method before, or while, carrying out the calculation. Important transitions should be explained with complete sentences. For example:

- Because the determinant is non-zero, the matrix is invertible.
- We may therefore multiply both sides by the inverse matrix.
- By the definition of the derivative, we consider the following limit.
- Since the direction vector is non-zero, it determines a line.
- Substituting the result into the original equation verifies the solution.

Such sentences are not decoration around the mathematics. They are part of the mathematics because they show the logical relationship between successive formulas.

It is not necessary to describe every elementary arithmetic operation. The amount of explanation should be proportional to the problem. A simple exercise may require only a few sentences, while a more substantial problem may require a longer argument. The goal is not to make every note long. The goal is to make every essential step understandable and justified.

Write as briefly as possible, but as completely as necessary.

A Complete Solution

A complete note should make the following five stages visible:

1. **Understand the problem.** Record the problem, identify the given information, and state clearly what must be found, proved, or explained.
2. **Choose a method.** Select the definitions, theorems, or techniques that connect the available information with the goal.
3. **Construct the argument.** Carry out the necessary steps and connect formulas with sentences that explain why each important transition is valid.

4. **Verify the result.** Check calculations, assumptions, domain restrictions, or geometric meaning whenever appropriate.
5. **State the conclusion.** Answer the original question clearly and in its proper context.

A result without its context is not a complete solution. A bare final answer, an unexplained sequence of formulas, or a few unconnected calculations will be treated as incomplete work and should be revised.

Quality cannot be replaced by quantity, but quality does not remove the obligation to complete the assigned work. Students are expected to prepare separate, complete solutions for the entire assigned problem set. Producing many unexplained answers is insufficient, but preparing one polished solution while ignoring the remaining problems is also insufficient.

Working with AI

We are studying in an age in which AI tools are widely available. The free versions of these tools are sufficient for the kinds of support expected in this course: asking questions, clarifying notation, generating simpler examples, checking reasoning, identifying missing steps, and improving the organisation of mathematical writing. Access to expensive software is not required.

AI should be used as an individual learning assistant. It may help transform rough calculations into clear mathematical prose, but it does not replace the student's mathematical work. Receiving a well-written answer is not the same as understanding it. The student remains responsible for checking every definition, calculation, assumption, conclusion, and formula included in the submitted note.

Use AI with a focused context. Work on one problem at a time and prepare one complete note for that problem. Do not paste an entire problem set into a single conversation and request all answers at once. Handling many unrelated tasks together encourages brief, generic, and mathematically weak notes. Focused work makes it easier to question each step, identify missing justifications, and produce a solution that can be understood and presented.

The one-problem-at-a-time principle does not mean that only one problem from the set should be prepared. Students are expected to repeat this focused process for every problem in the complete assigned set.

A polished AI-generated note is insufficient if the student cannot read its formulas, explain its terminology, justify its transitions, or present the argument independently. When using AI, ask not only for a solution, but also questions that deepen understanding:

- What does this formula mean?
- What mathematical object does each symbol represent?
- Why is this step valid?
- Which definition or theorem is being used?
- Are the assumptions of the theorem satisfied?
- Is there a simpler way to explain this transition?
- How can the result be checked?
- What should I be able to explain during class?

Preparing notes in this way is part of learning mathematics. It develops the ability to read notation, recognise mathematical objects, organise information, construct arguments, and communicate ideas precisely. Repeating this process for the complete problem set builds the habits of systematic and conscientious work that should accompany the student throughout university study and later professional life.

Student GitHub Workspace

The current instructions, problem templates, and technical workflow for student work are maintained in the public repository [dchorazkiewicz/Math_Problems_Repo](#). Students should create a public fork of this repository and commit their notes to their own fork. The course repository is the common template; it is not the place where individual student solutions should be submitted.

The repository contains a short map of the working process:

- `README.md` explains the purpose and structure of the workspace and provides the weekly index;
- `SOURCE_MATERIAL.md` links the current lecture PDF and maps each week to the corresponding LaTeX problem source used by an AI assistant;
- `NOTE_GUIDELINES.md` defines the standard of a complete mathematical note;
- `AI_WORKFLOW.md` describes the one-problem-at-a-time workflow with AI, review, commits, presentation, feedback, and revision;
- `AGENTS.md` gives a connected AI assistant the operational rules for working safely in the student's fork;
- the `docs/` directory contains twelve weekly folders and one Markdown file for each problem.

The workspace uses Markdown for student notes, Git and GitHub for version history and public forks, MathJax for mathematical typesetting, MkDocs for the web presentation, and GitHub Actions for automatic validation and deployment. These instructions and templates may be updated independently of this PDF, so students should consult the repository for the current working procedure. The repository links back to this PDF and to its LaTeX sources. Students use the PDF as the readable course material containing the theory, definitions, examples, and problem sets. A connected AI assistant uses the current mapped LaTeX file as the exact machine-readable source of a selected problem statement.

Chapter 1

Euclidean Geometry and Cartesian Coordinates

1.1 The Euclidean Geometry Available Before Coordinates

Before introducing coordinates, we recall the geometric world that is already available. At this stage there are no ordered pairs, no axes, and no vectors. There are points, straight lines, segments, circles, and angles, together with relations such as incidence, betweenness, congruence, perpendicularity, and parallelism.

Remark

Coordinates will later translate this geometry into arithmetic. They do not create the geometry. We can already construct and compare geometric objects before assigning numbers to all points and angles.

1.1.1 Three Different Levels Must Not Be Confused

We shall distinguish three kinds of statements.

1. *Euclid's postulates* describe the geometric world being assumed.
2. *Straightedge-and-compass rules* describe which elementary drawings and intersections may be used in a construction.
3. *Derived constructions* must be built from those elementary operations and justified by a proof.


In particular, a perpendicular bisector, a perpendicular through a point, an angle bisector, or a parallel through a point will not be treated as primitive construction postulates. They will be constructed.

1.1.2 Euclid's Five Postulates

The following are the five postulates used in the first book of Euclid's *Elements*. The diagrams illustrate their meaning; they are not proofs.

Postulate I: Joining Two Points

A straight segment can be drawn from any point to any other point.



a segment is drawn from A to B

Postulate II: Extending a Segment


A finite straight segment can be extended continuously in the same straight line.



AB is extended beyond B

Postulate III: Drawing a Circle

A circle can be drawn with any chosen centre and any chosen radius.




centre O and radius r determine a circle

This postulate is what permits a length to be copied geometrically without first assigning a number to it.

Postulate IV: Equality of Right Angles

All right angles are congruent.




every right angle represents the same geometric magnitude

This postulate does not say that a perpendicular may simply be drawn without construction. It says that once right angles have been constructed, they are all equal.

Postulate V: The Parallel Postulate

If a transversal meets two straight lines and the interior angles on one side sum to less than two right angles, then the two lines, when extended, meet on that side.



the angle condition determines on which side ℓ and m meet

A familiar modern equivalent is Playfair's formulation: through a point outside a given line there passes exactly one parallel to that line. We shall use that form later, after the relation between the two formulations has been made clear.

1.1.3 Elementary Rules for Straightedge-and-Compass Work

For explicit constructions we use the following operational rules.

1. Through two already constructed distinct points, draw the straight line.
2. With an already constructed point as centre and an already constructed segment as radius, draw a circle.
3. Retain intersection points of two constructed lines, a line and a circle, or two circles whenever such intersections exist.

The first two rules are the operational forms of Euclid's first three postulates; the third explains how new constructible points are obtained. These rules are the complete elementary toolkit used in the constructions below.

1.1.4 Constructions Derived from the Toolkit

The Perpendicular Bisector and the Midpoint

Let A and B be distinct points. Draw two circles with the same radius, one centred at A and one centred at B , choosing the radius larger than half of $|AB|$. Let their intersection points be P and Q .


Because P lies on both circles,

$$|PA| = |PB|,$$

and similarly,

$$|QA| = |QB|.$$

Thus both P and Q are equidistant from A and B . The line PQ is the perpendicular bisector of AB .



The intersection of PQ with AB is the midpoint of AB . Thus midpoint and perpendicularity have been constructed from equal radii and intersections.

A Perpendicular Through a Point on a Line

Let P lie on a line ℓ .

Step 1. Draw a circle centred at P . It meets ℓ at two points A and B . Since A and B lie on the same circle,

$$|PA| = |PB|.$$

Thus P is the midpoint of AB .


Step 2. Draw two circles with the same radius, one centred at A and one centred at B , choosing the radius larger than $|AP|$. Let their intersection points be Q and R . Then

$$|AQ| = |BQ| \quad \text{and} \quad |AR| = |BR|.$$

Both Q and R are therefore equidistant from A and B , so the line QR is the perpendicular bisector of AB . Because P is the midpoint of AB , the line QR passes through P . Since AB lies on ℓ ,

$$QR \perp AB, \quad \text{hence} \quad QR \perp \ell.$$

The required perpendicular is the line $m = QR$.



The perpendicular has now been obtained entirely from the permitted straightedge-and-compass operations.

A Perpendicular Through a Point Outside a Line

Let P lie outside a line ℓ .

Step 1. Draw a circle centred at P with radius large enough to meet ℓ at two points A and B . Since A and B lie on the same circle,

$$|PA| = |PB|.$$

Step 2. Draw two circles of radius $|PA|$: one centred at A and one centred at B . The two circles meet at P and at a second point Q . Therefore


$$|AP| = |BP| \quad \text{and} \quad |AQ| = |BQ|.$$

Thus both P and Q are equidistant from A and B . The line PQ is therefore the perpendicular bisector of AB .

Because A and B lie on ℓ , the segment AB is contained in ℓ . Consequently,

$$PQ \perp AB, \quad \text{hence} \quad PQ \perp \ell.$$

The required perpendicular is the line $m = PQ$.



The perpendicular has now been obtained entirely from the permitted straightedge-and-compass operations: drawing circles, retaining their intersection points, and drawing the line through two constructed points.

Bisecting an Angle

Let two rays with common endpoint O form an angle. Draw a circle centred at O meeting the rays at A and B . Then draw equal-radius circles centred at A and B , and let one of their intersections inside the angle be P . Since

$$|OA| = |OB|, \quad |AP| = |BP|, \quad |OP| = |OP|,$$

the triangles OAP and OBP are congruent. Therefore

$$\angle AOP = \angle POB.$$

The ray OP is the angle bisector.

Constructing a Parallel Through a Point

Let P lie outside a line ℓ . First construct a line m through P perpendicular to ℓ . Then construct a line n through P perpendicular to m .

The transversal m forms equal corresponding right angles with n and ℓ . By the converse of the corresponding-angles theorem, the two lines are parallel:

$$n \parallel \ell.$$

The fifth postulate, equivalently Playfair's postulate, guarantees that this is the unique parallel to ℓ through P .

1.1.5 Parallel Lines and Proportional Segments

A central fact of Euclidean geometry is that parallel lines preserve ratios. This is the content of Thales' theorem in the form needed here.


Theorem

Let D lie on AB and E lie on AC in a triangle ABC . If

$$DE \parallel BC,$$

then

$$\frac{AD}{AB} = \frac{AE}{AC} = \frac{DE}{BC}.$$



The smaller triangle ADE and the larger triangle ABC have the same three angles. Their corresponding sides differ by one common scale factor. This is the basic phenomenon of similarity.


Definition

Two triangles are similar when their corresponding angles are equal. In that case their corresponding side lengths are proportional.

Similarity lets us change the size of a figure while preserving its shape. It is the reason ratios of corresponding lengths can depend only on angles.

1.1.6 Sine and Cosine Before Coordinates

Take a right triangle containing an acute angle α . Let h denote the hypotenuse, let a be the leg adjacent to α , and let b be the leg opposite α .



$$\frac{a'}{a} = \frac{b'}{b} = \frac{h'}{h}, \quad \frac{a'}{h'} = \frac{a}{h}, \quad \frac{b'}{h'} = \frac{b}{h}.$$

For an acute geometric angle α , define

$$\cos \alpha = \frac{a}{h}, \quad \sin \alpha = \frac{b}{h}.$$

These ratios are independent of the chosen right triangle because all right triangles containing the same acute angle are similar.

For angles occurring in an arbitrary Euclidean triangle, we also need the right and obtuse cases. We set

$$\cos(\text{right angle}) = 0.$$

If α is obtuse and β is its acute supplement, so that together they form a straight angle, define

$$\cos \alpha = -\cos \beta.$$

Thus the geometric cosine is defined for every angle between 0 and a straight angle. For acute angles it is positive, for a right angle it is zero, and for obtuse angles it is negative.

The essential role of Thales' theorem and similarity is that these values depend only on the angle, not on the size of the triangle used to represent it.

For an acute angle, the Pythagorean theorem gives

$$a^2 + b^2 = h^2.$$

Dividing by h^2 yields

$$\left(\frac{a}{h}\right)^2 + \left(\frac{b}{h}\right)^2 = 1,$$

and hence

$$\cos^2 \alpha + \sin^2 \alpha = 1.$$

Thus the fundamental trigonometric identity is a direct reformulation of the Pythagorean theorem.


1.1.7 Angles and Their Numerical Measure

An angle exists geometrically before it is assigned a number. We can compare, copy, add, and bisect angles without degrees or radians. A numerical angle measure requires an additional choice of unit.

Definition

Let r_1 and r_2 be two rays with the same initial point O . The rays, together with a choice of one of the two regions between them, determine an angle α with vertex O .

The angle is the chosen geometric region. It is neither the small arc drawn near the vertex nor a number. The arc is only a graphical mark indicating which region has been selected.



Why circles give a consistent measure

Draw a circle centred at O . The rays cut from the circle an arc corresponding to the chosen angle.

Theorem

For every Euclidean circle, the ratio of circumference C to diameter D is the same constant:

$$\frac{C}{D} = \pi.$$

This constant is called π . Since $D = 2r$, every circle of radius r has circumference

$$C = 2\pi r.$$

The constancy of C/D follows from similarity: enlarging a circle by a factor λ multiplies every length, including its diameter, circumference, and corresponding arc lengths, by λ .

If the same two rays meet two circles of radii r_1 and r_2 and cut arc lengths s_1 and s_2 , then

$$\frac{s_1}{s_2} = \frac{r_1}{r_2},$$

and therefore

$$\frac{s_1}{r_1} = \frac{s_2}{r_2}.$$

Thus the quotient s/r depends only on the angle, not on the chosen circle.

Definition

Let s be the length of the selected arc on a circle of radius r centred at the vertex. The radian measure of α is

$$m_{\text{rad}}(\alpha) = \frac{s}{r}.$$

For a full turn, $s = 2\pi r$, so

$$m_{\text{rad}}(\text{full turn}) = 2\pi.$$

A half-turn has radian measure π , and a quarter-turn has radian measure $\pi/2$.

Degrees are a second numerical scale. A full turn is assigned 360° , so

$$2\pi \text{ radians} = 360^\circ$$

and

$$m_{\text{deg}}(\alpha) = \frac{180^\circ}{\pi} m_{\text{rad}}(\alpha).$$

In particular, a constructible right angle has the numerical measures

$$m_{\text{rad}}(\text{right angle}) = \frac{\pi}{2}, \quad m_{\text{deg}}(\text{right angle}) = 90^\circ.$$

The angle is still the geometric region; $\pi/2$ and 90° are two numbers assigned to it by two different measurement systems.

Sine and cosine as functions of a number

The trigonometric ratios were first defined for acute geometric angles. We now extend them to a full turn. For $t \in [0, 2\pi)$, let $\beta \in [0, \pi/2]$ be the acute reference angle between the direction of t and the nearest horizontal ray. The magnitudes $\sin \beta$ and $\cos \beta$ come from the right-triangle definition; their signs are determined by the quadrant:

	$\cos t$	$\sin t$
$0 \leq t \leq \pi/2$	$+\cos \beta$	$+\sin \beta$
$\pi/2 \leq t \leq \pi$	$-\cos \beta$	$+\sin \beta$
$\pi \leq t \leq 3\pi/2$	$-\cos \beta$	$-\sin \beta$
$3\pi/2 \leq t < 2\pi$	$+\cos \beta$	$-\sin \beta$

At the axes this gives the familiar values 0, 1, and -1 . For arbitrary real t , define sine and cosine periodically:

$$\sin(t + 2\pi k) = \sin t, \quad \cos(t + 2\pi k) = \cos t, \quad k \in \mathbb{Z}.$$


If $t = m_{\text{rad}}(\alpha)$, then $\cos t$ is the same value as the geometric cosine $\cos \alpha$. This explains the notation $\cos(m_{\text{rad}}(\alpha))$.

The Euclidean law of cosines

Directly measuring an arc is often difficult. A triangle provides another way to recover the measure of an angle. Choose points

$$A \in r_1 \setminus \{O\}, \quad B \in r_2 \setminus \{O\}.$$

The points form the Euclidean triangle OAB .



Theorem

If $\alpha = \angle AOB$, then the side lengths of the Euclidean triangle satisfy

$$|AB|^2 = |OA|^2 + |OB|^2 - 2|OA||OB|\cos \alpha.$$

Equivalently, the last factor may be written as $\cos(m_{\text{rad}}(\alpha))$.

The theorem belongs to Euclidean geometry; it does not require Cartesian coordinates. Solving it for the angle measure gives

$$m_{\text{rad}}(\alpha) = \arccos \left(\frac{|OA|^2 + |OB|^2 - |AB|^2}{2|OA||OB|} \right).$$

Later, coordinates will provide a convenient arithmetic method for calculating the three Euclidean lengths $|OA|$, $|OB|$, and $|AB|$.

Remark

At this point no vector has been introduced. We have only Euclidean objects, postulates, constructions, similarity, trigonometric ratios, numerical angle measure, and the Euclidean law of cosines. Vectors will appear only in the next chapter, after the Cartesian coordinate system has been constructed.

The next step is to choose two perpendicular lines, an origin, directions, and a unit length. Those choices will convert the Euclidean plane into a Cartesian coordinate plane.

1.2 Constructing Cartesian Coordinates from Euclidean Geometry

We begin with a Euclidean plane, not with ordered pairs and not with a coordinate system that has already been drawn. Points, straight lines, right angles, perpendicularity, congruent segments, and the Pythagorean theorem belong to the geometry available before coordinates are introduced. Our task is to construct, step by step, the additional choices that turn this plane into a Cartesian coordinate plane.

Remark

Coordinates do not create Euclidean geometry. They arise after we choose and construct a particular system for assigning numerical labels to points of an already existing Euclidean plane.

1.2.1 The Plane Before Coordinates

At first there is only a Euclidean plane, with no distinguished line, origin, directions, or numerical scale.



a Euclidean plane with no chosen coordinates

1.2.2 First Construction: A Straight Line


Choose two distinct points of the plane and construct the unique straight line through them. At this stage it is only a chosen Euclidean line. It is not yet an axis, because there is not yet a second line, an origin, a direction, or a scale.



the first constructed line

1.2.3 Second Construction: A Point on the Line

Choose an arbitrary point O on the constructed line. This point will later become the common zero of both coordinate axes, but it becomes an origin only after the rest of the coordinate structure has been constructed.




a point chosen on the first line

1.2.4 Third Construction: The Perpendicular Line

Through O , construct the unique line perpendicular to the first line. The two constructed lines now intersect at a right angle. Only now do we have the geometric framework from which two coordinate axes can be made.

We call the first line the x -axis, the perpendicular line the y -axis, and their intersection O the origin.




The axes are perpendicular because this was required in their geometric construction. No coordinate formula and no numerical angle measure has been used.

1.2.5 Choosing Positive and Negative Directions

Each axis is divided by O into two opposite rays. Choose one ray on the x -axis as positive and one ray on the y -axis as positive. The opposite rays are then called negative.

These choices are not forced by the Euclidean plane. Reversing one of them would produce a different coordinate system on the same geometric plane.




1.2.6 Constructing a Common Unit Length

Choose a positive segment length and call it one unit. Using a compass, transfer this same length from O repeatedly along both axes and in both directions. The first marks are labelled 1 and -1 ; repetition constructs the integer marks

$$\dots, -3, -2, -1, 0, 1, 2, 3, \dots$$

on each axis.



Remark


The same unit must be used on both axes. A segment labelled as one unit on the x -axis must be congruent to a segment labelled as one unit on the y -axis. This common scale is what later allows the Pythagorean theorem to produce the standard coordinate formula for Euclidean distance.

1.2.7 Refining the Scale by Bisection

The compass and straightedge construction does not stop at integer marks. Bisecting each unit segment constructs halves. Bisecting again constructs quarters, then eighths, and so on. Repeating this process on every integer interval and on both sides of the origin gives all dyadic marks

$$\frac{k}{2^n}, \quad k \in \mathbb{Z}, \quad n \in \mathbb{N}.$$

successive bisections of one unit



The dyadic marks are dense on the axis: every interval, however small, contains such marks. They therefore provide arbitrarily fine numerical approximations to positions on the line.

Definition


Continuity assumption. Once an origin, a positive direction, and a unit length are fixed on a Euclidean line, every point of the line corresponds to exactly one real number, and every real number corresponds to exactly one point of the line.

This assumption is additional to the finite straightedge-and-compass operations. Those operations construct many particular positions, including all dyadic marks, but not every real position. Continuity supplies the complete real scale used by coordinates. After this step, every point of each axis has a unique signed real label.

1.2.8 Assigning Coordinates by Orthogonal Projection

Only after the axes, directions, common unit, integer marks, refined scale, and real labels have been constructed can an arbitrary point of the plane be given coordinates.

Let P be any point of the plane. Through P , construct the line perpendicular to the x -axis. It meets the x -axis at a unique point P_x . Through P , construct the line perpendicular to the y -axis. It meets the y -axis at a unique point P_y .



If the signed real labels of P_x and P_y are x and y , respectively, we assign to P the ordered pair

$$P \longleftrightarrow (x, y).$$

The construction is unambiguous because the perpendicular through a given point to a given line is unique. Conversely, given real numbers x and y , locate the corresponding points on the two axes and construct through them lines perpendicular to their respective axes. Those two lines meet in exactly one point of the plane.

Theorem

Once the two perpendicular axes, their positive directions, and a common unit are fixed, orthogonal projection gives a one-to-one correspondence

$$\text{points of the Euclidean plane} \longleftrightarrow \text{pairs } (x, y) \in \mathbb{R}^2.$$

The ordered pair is the numerical address of a geometric point in the chosen coordinate

system. It is not the point itself and it is not the source of the geometry.

1.2.9 The Coordinate Formula for Euclidean Distance

The Euclidean distance between two points is the length of the straight segment joining them. Coordinates now allow this geometric length to be calculated.


For points

$$P = (x_1, y_1), \quad Q = (x_2, y_2),$$

their projections show that the horizontal and vertical separations have lengths

$$|x_2 - x_1| \quad \text{and} \quad |y_2 - y_1|.$$

Together with the segment PQ , these two perpendicular segments form a right triangle. Here “right” refers to the Euclidean right angle already used in the construction; no numerical measure of that angle is required.



By the Pythagorean theorem,

$$d_E(P, Q)^2 = (x_2 - x_1)^2 + (y_2 - y_1)^2,$$

and therefore

$$d_E(P, Q) = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}.$$

The formula does not create a new notion of distance. It is the numerical coordinate expression of the Euclidean segment length already present in the plane.

Summary

The construction proceeds in the following order:

1. begin with a Euclidean plane containing no chosen coordinates;
2. construct one straight line;
3. choose a point O on that line;
4. construct through O the perpendicular line, and only then name the two lines as coordinate axes and O as their origin;
5. choose positive directions on both axes;
6. choose one common unit length and transfer it with a compass in both positive and negative directions to construct the integer marks;

7. repeatedly bisect intervals to construct the dense family of dyadic marks;
8. use Euclidean continuity to identify every axis position with a real number;
9. assign to every point its unique pair of orthogonal projections;
10. derive the coordinate distance formula from the resulting right triangle and the Pythagorean theorem.

Chapter 2

Vectors, Bases, and Coordinate Systems

2.1 Vectors from Coordinate Displacements

A coordinate pair tells us where a point is. It does not yet tell us how one point is related to another. To describe motion from one point to another we need a new object: not another position, but a record of the change of position.

Choose

$$A = (1, 2), \quad B = (4, 6).$$

To travel from A to B , the horizontal coordinate must increase by 3 and the vertical coordinate by 4. The pair

$$(3, 4)$$

therefore records an instruction:

move three units in the positive x direction and four units in the positive y direction.

This instruction can be carried out from A , but it can also be carried out from any other point. The starting point may change while the displacement remains the same. That distinction is the idea from which vectors arise.

The development below follows a fixed question-driven order. We first decide when two arrows represent the same displacement. We then ask how successive displacements combine, what it means to repeat or reverse one, what addition and scaling can build, how to measure length and direction, and how two displacements can be compared geometrically. The coordinate formulas will be consequences of those questions, not isolated rules introduced for memorisation.

2.1.1 A Vector Is a Displacement, Not a Place

Let

$$A = (x_1, y_1), \quad B = (x_2, y_2).$$

The point A is a position and the point B is another position. The arrow from A to B asks a different question:

what change of coordinates carries the starting point to the endpoint?

The horizontal change is $x_2 - x_1$ and the vertical change is $y_2 - y_1$. Therefore

$$\overrightarrow{AB} = (x_2 - x_1, y_2 - y_1).$$

This subtraction is not an arbitrary convention. It is forced by the equations

$$x_1 + \Delta x = x_2, \quad y_1 + \Delta y = y_2.$$

Solving them gives

$$\Delta x = x_2 - x_1, \quad \Delta y = y_2 - y_1.$$

Definition

A vector in the Cartesian plane is a coordinate displacement

$$\mathbf{v} = (v_1, v_2).$$

It instructs us to change the first coordinate by v_1 and the second coordinate by v_2 .


A vector is therefore not tied to one starting point. If $P = (p_1, p_2)$ and we apply the same displacement $\mathbf{v} = (v_1, v_2)$, then the endpoint must be

$$Q = (p_1 + v_1, p_2 + v_2),$$

and indeed

$$\overrightarrow{PQ} = (v_1, v_2) = \mathbf{v}.$$

Two arrows drawn in different places represent the same vector when they have the same coordinate changes. Their positions may differ; their displacement is the same.



The zero vector

$$\mathbf{0} = (0, 0)$$

means no change at all. Starting anywhere and applying $\mathbf{0}$ leaves the point fixed.

The opposite vector

$$-\mathbf{v} = (-v_1, -v_2)$$

undoes the original displacement. If $\mathbf{v}$ carries A to B , then $-\mathbf{v}$ carries B back to A . Thus

$$\overrightarrow{BA} = -\overrightarrow{AB}.$$

2.1.2 Two Journeys Become One: Why Vectors Add

Suppose a point is moved twice. The first displacement is

$$\mathbf{u} = (u_1, u_2),$$

and the second is

$$\mathbf{v} = (v_1, v_2).$$

We want one vector that has exactly the same final effect as these two moves performed successively.

Start at an arbitrary point (x, y) . After applying $\mathbf{u}$ we are at

$$(x + u_1, y + u_2).$$


Applying $\mathbf{v}$ from there gives

$$(x + u_1 + v_1, y + u_2 + v_2).$$

Relative to the original point, the total horizontal change is $u_1 + v_1$ and the total vertical change is $u_2 + v_2$. Therefore the single displacement with the same effect is forced to be

$$\boxed{\mathbf{u} + \mathbf{v} = (u_1 + v_1, u_2 + v_2)}.$$

The formula is simply bookkeeping for a journey performed in two stages. It adds changes of the same kind: horizontal change to horizontal change and vertical change to vertical change.



The diagram contains two routes to the same endpoint:

$$\mathbf{u} \text{ then } \mathbf{v}, \quad \mathbf{v} \text{ then } \mathbf{u}.$$

They agree because addition of real numbers is commutative:

$$u_1 + v_1 = v_1 + u_1, \quad u_2 + v_2 = v_2 + u_2.$$

Hence

$$\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}.$$

Geometrically this is the parallelogram rule. Algebraically it is the fact that the order of adding coordinate changes does not affect the final position.

Subtraction answers a different practical question: what displacement must be performed after $\mathbf{v}$ in order to have the same effect as $\mathbf{u}$? We seek a vector $\mathbf{w}$ satisfying

$$\mathbf{v} + \mathbf{w} = \mathbf{u}.$$

Solving coordinate by coordinate gives

$$\boxed{\mathbf{u} - \mathbf{v} = (u_1 - v_1, u_2 - v_2)}.$$

In particular, the displacement from A to B is the position of B minus the position of A :

$$\overrightarrow{AB} = B - A.$$

The expression does not subtract geometric points as physical objects. It subtracts their coordinate addresses to recover the change carrying one to the other.

2.1.3 Stretching, Shrinking, and Reversing a Displacement

Addition combines different displacements. A second operation appears when the same displacement is repeated.

If

$$\mathbf{v} = (v_1, v_2),$$

then performing it twice gives

$$\mathbf{v} + \mathbf{v} = (2v_1, 2v_2).$$

Performing it three times gives


$$3\mathbf{v} = (3v_1, 3v_2).$$

The coordinates are multiplied by the same number because both components must be stretched by the same factor if the direction is to remain unchanged.

This suggests the general definition

$$\boxed{\lambda\mathbf{v} = (\lambda v_1, \lambda v_2), \quad \lambda \in \mathbb{R}}.$$

The number λ is called a scalar. It does not provide a new direction; it changes how much of the old displacement is taken.



The picture separates the possible cases:

- if $\lambda > 1$, the vector is stretched;
- if $0 < \lambda < 1$, it is shortened;
- if $\lambda = 0$, it collapses to the zero vector;
- if $\lambda < 0$, it is also reversed.

In every case the result lies on the same line through the origin as $\mathbf{v}$. This gives a first geometric meaning of scalar multiples: nonzero vectors $\mathbf{u}$ and $\mathbf{v}$ are parallel precisely when one is a scalar multiple of the other.

Scalar multiplication also explains why fractions and real factors are natural. The vector $\frac{1}{2}\mathbf{v}$ is the displacement that must be repeated twice to produce $\mathbf{v}$, while $\sqrt{2}\mathbf{v}$ has the same direction and $\sqrt{2}$ times the length. The operation extends the simple idea of repetition to continuous scaling.


2.1.4 Linear Combinations: What Addition and Scaling Can Build

Addition and scalar multiplication were introduced as two separate operations, but they are most useful when they are combined. Given vectors $\mathbf{u}$ and $\mathbf{v}$ and real numbers a and b , the vector

$$a\mathbf{u} + b\mathbf{v}$$

is called a *linear combination* of $\mathbf{u}$ and $\mathbf{v}$.

The phrase has a direct geometric meaning. First scale $\mathbf{u}$ by a and $\mathbf{v}$ by b ; then add the two resulting displacements. Thus a linear combination is not a new operation. It is a controlled use of the two operations already understood.



A single nonzero vector can generate only one line through the origin:

$$\{\lambda \mathbf{v} : \lambda \in \mathbb{R}\}.$$

Changing λ changes the length and possibly the orientation, but never the underlying line of motion.

Two vectors behave differently. If $\mathbf{u}$ and $\mathbf{v}$ are parallel, then every linear combination still lies on one line. If they are not parallel, then their linear combinations fill the whole plane. Geometrically, the two independent directions allow us to reach any endpoint by a suitable parallelogram.

Theorem

Two nonparallel vectors $\mathbf{u}$ and $\mathbf{v}$ determine every vector in the plane uniquely in the form

$$\mathbf{w} = a\mathbf{u} + b\mathbf{v}.$$

The numbers a and b record how much motion occurs in each chosen direction.

This observation is the bridge to the later notion of a basis. A basis is not an unrelated new object: it is a pair of directions whose linear combinations can describe every vector uniquely.

2.1.5 How Much Motion and in Which Direction?


The coordinate pair of a vector records two changes, but it does not immediately answer two basic geometric questions:

1. How long is the displacement?
2. In which direction does it point?

Let

$$\mathbf{v} = (v_1, v_2)$$

be drawn from the origin. Its endpoint, together with its horizontal and vertical coordinate changes, forms a right triangle. The legs have lengths $|v_1|$ and $|v_2|$, while the vector itself is the hypotenuse.



By the Pythagorean theorem,

$$\|\mathbf{v}\|^2 = v_1^2 + v_2^2,$$

so the nonnegative length of the vector is

$$\boxed{\|\mathbf{v}\| = \sqrt{v_1^2 + v_2^2}}.$$

The double bars distinguish the vector from the number measuring its length. In particular,

$$\|\mathbf{0}\| = 0,$$

and every nonzero vector has positive length.

This formula also reconnects vectors with the Euclidean distance between points. Since

$$\overrightarrow{AB} = B - A,$$

the length of the displacement from A to B is

$$\boxed{d_E(A, B) = \|B - A\|}.$$

A vector therefore packages both the direction from one point to another and the distance travelled.

Now assume $\mathbf{v} \neq \mathbf{0}$ and let α be its angle of inclination, measured from the positive x -axis. The right triangle gives

$$\cos \alpha = \frac{v_1}{\|\mathbf{v}\|}, \quad \sin \alpha = \frac{v_2}{\|\mathbf{v}\|}.$$

Rearranging,

$$v_1 = \|\mathbf{v}\| \cos \alpha, \quad v_2 = \|\mathbf{v}\| \sin \alpha.$$

Hence

$$\boxed{\mathbf{v} = \|\mathbf{v}\|(\cos \alpha, \sin \alpha)}.$$

This formula separates the two pieces of information carried by a nonzero vector:

- $\|\mathbf{v}\|$ says how much displacement occurs;
- $(\cos \alpha, \sin \alpha)$ is a unit vector saying in which direction.

Indeed,

$$\|(\cos \alpha, \sin \alpha)\| = \sqrt{\cos^2 \alpha + \sin^2 \alpha} = 1.$$

Thus every nonzero vector is a positive length multiplied by a unit direction. This decomposition will make comparison of two vectors by their angle possible in the next subsection.

2.1.6 One Angle Calculation, Two Geometric Tests

Take two arbitrary nonzero vectors

$$\mathbf{u} = (u_1, u_2), \quad \mathbf{v} = (v_1, v_2).$$

Let their angles of inclination be α and β . Then


$$\mathbf{u} = \|\mathbf{u}\|(\cos \alpha, \sin \alpha),$$

$$\mathbf{v} = \|\mathbf{v}\|(\cos \beta, \sin \beta).$$

The difference $\beta - \alpha$ measures the directed turn from $\mathbf{u}$ to $\mathbf{v}$. The ordinary angle between the vectors will be denoted by

$$\theta = |\beta - \alpha|$$

when the smaller angle is intended.



Parallel directions arise from the sine of the angle difference

Two nonzero vectors determine the same unoriented direction precisely when the turn from one to the other is an integer multiple of a half-turn:

$$\beta - \alpha \equiv 0 \pmod{\pi}.$$

Equivalently,

$$\sin(\beta - \alpha) = 0.$$

Using the sine-difference identity,

$$\sin(\beta - \alpha) = \sin \beta \cos \alpha - \cos \beta \sin \alpha.$$

Now substitute

$$\cos \alpha = \frac{u_1}{\|\mathbf{u}\|}, \quad \sin \alpha = \frac{u_2}{\|\mathbf{u}\|},$$

$$\cos \beta = \frac{v_1}{\|\mathbf{v}\|}, \quad \sin \beta = \frac{v_2}{\|\mathbf{v}\|}.$$

We obtain

$$\begin{aligned} \sin(\beta - \alpha) &= \frac{v_2}{\|\mathbf{v}\|} \frac{u_1}{\|\mathbf{u}\|} - \frac{v_1}{\|\mathbf{v}\|} \frac{u_2}{\|\mathbf{u}\|} \\ &= \frac{u_1 v_2 - u_2 v_1}{\|\mathbf{u}\| \|\mathbf{v}\|}. \end{aligned}$$

The denominator is nonzero. Therefore

$$\sin(\beta - \alpha) = 0 \iff u_1 v_2 - u_2 v_1 = 0.$$

We have discovered the coordinate test

$$\mathbf{u} \parallel \mathbf{v} \iff u_1 v_2 - u_2 v_1 = 0.$$

The same condition implies that one vector is a scalar multiple of the other. Indeed, if $v_1 \neq 0$, set $\lambda = u_1/v_1$. The equality

$$u_1v_2 - u_2v_1 = 0$$

gives $u_2 = \lambda v_2$, and hence

$$\mathbf{u} = \lambda \mathbf{v}.$$

If $v_1 = 0$, then $v_2 \neq 0$ and the same argument uses $\lambda = u_2/v_2$. Thus

$$\mathbf{u} \parallel \mathbf{v} \iff \mathbf{u} = \lambda \mathbf{v} \quad \text{for some } \lambda \neq 0.$$

This scalar-multiple criterion is a conclusion of the angle calculation, not a rule imposed in advance.

The cosine of the same difference reveals perpendicularity

The vectors are perpendicular precisely when their angle difference is a quarter-turn modulo a half-turn:

$$\beta - \alpha \equiv \frac{\pi}{2} \pmod{\pi}.$$

Equivalently,

$$\cos(\beta - \alpha) = 0.$$

The cosine-difference identity gives

$$\cos(\beta - \alpha) = \cos \beta \cos \alpha + \sin \beta \sin \alpha.$$

Substituting the coordinate ratios gives

$$\begin{aligned} \cos(\beta - \alpha) &= \frac{v_1}{\|\mathbf{v}\|} \frac{u_1}{\|\mathbf{u}\|} + \frac{v_2}{\|\mathbf{v}\|} \frac{u_2}{\|\mathbf{u}\|} \\ &= \frac{u_1v_1 + u_2v_2}{\|\mathbf{u}\| \|\mathbf{v}\|}. \end{aligned}$$

Since the denominator is nonzero,

$$\mathbf{u} \perp \mathbf{v} \iff u_1v_1 + u_2v_2 = 0.$$

The coordinate expression has emerged from the geometry. Only now do we give it a name.

Definition

For vectors in the plane, their *dot product* is

$$\mathbf{u} \cdot \mathbf{v} = u_1v_1 + u_2v_2.$$

In three-dimensional space,

$$(u_1, u_2, u_3) \cdot (v_1, v_2, v_3) = u_1v_1 + u_2v_2 + u_3v_3.$$

The preceding calculation proves

$$\mathbf{u} \perp \mathbf{v} \iff \mathbf{u} \cdot \mathbf{v} = 0.$$

It proves more than the right-angle case. Because cosine is even and $\theta = |\beta - \alpha|$,

$$\cos(\beta - \alpha) = \cos \theta.$$

Therefore

$$\boxed{\mathbf{u} \cdot \mathbf{v} = \|\mathbf{u}\| \|\mathbf{v}\| \cos \theta}$$

for every pair of nonzero vectors in the plane.

Example

Take

$$\mathbf{u} = (3, 2), \quad \mathbf{v} = (-2, 3).$$

Then

$$\mathbf{u} \cdot \mathbf{v} = 3(-2) + 2(3) = 0.$$

The coordinate calculation says that

$$\cos \theta = 0,$$

so the angle between the vectors is a right angle.

2.1.7 Why the Dot Product Can Be Expanded

The dot product was discovered from the cosine of the angle between two vectors. Before using it in longer calculations, we must verify how it behaves under addition and scalar multiplication.

Let

$$\mathbf{u} = (u_1, u_2), \quad \mathbf{v} = (v_1, v_2), \quad \mathbf{w} = (w_1, w_2).$$

Then

$$\begin{aligned} (\mathbf{u} + \mathbf{w}) \cdot \mathbf{v} &= (u_1 + w_1)v_1 + (u_2 + w_2)v_2 \\ &= (u_1v_1 + u_2v_2) + (w_1v_1 + w_2v_2) \\ &= \mathbf{u} \cdot \mathbf{v} + \mathbf{w} \cdot \mathbf{v}. \end{aligned}$$

Thus the dot product distributes over vector addition. Similarly,

$$(\lambda \mathbf{u}) \cdot \mathbf{v} = \lambda(\mathbf{u} \cdot \mathbf{v}).$$

The symmetry of multiplication of real numbers also gives

$$\mathbf{u} \cdot \mathbf{v} = \mathbf{v} \cdot \mathbf{u}.$$

Theorem

For vectors $\mathbf{u}, \mathbf{v}, \mathbf{w}$ and a real number λ ,

$$\begin{aligned} (\mathbf{u} + \mathbf{w}) \cdot \mathbf{v} &= \mathbf{u} \cdot \mathbf{v} + \mathbf{w} \cdot \mathbf{v}, \\ (\lambda \mathbf{u}) \cdot \mathbf{v} &= \lambda(\mathbf{u} \cdot \mathbf{v}), \\ \mathbf{u} \cdot \mathbf{v} &= \mathbf{v} \cdot \mathbf{u}. \end{aligned}$$

Moreover,

$$\mathbf{u} \cdot \mathbf{u} = \|\mathbf{u}\|^2.$$

The last identity follows immediately:

$$\mathbf{u} \cdot \mathbf{u} = u_1^2 + u_2^2 = \|\mathbf{u}\|^2.$$

These properties justify every later expansion involving expressions such as $(\mathbf{u} - \mathbf{v}) \cdot (\mathbf{u} - \mathbf{v})$. They are not formal rules borrowed from elsewhere; they follow from the coordinate definition of the dot product.

2.1.8 The Angle Between Two Vectors

The difference of inclination angles $\beta - \alpha$ is a directed turn and is understood modulo a full turn. The ordinary angle between two nonzero vectors is a different object: it is the smaller, unoriented separation between their directions.

Definition

For nonzero vectors $\mathbf{u}$ and $\mathbf{v}$, their angle is the unique number

$$\theta \in [0, \pi]$$

satisfying

$$\cos \theta = \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{u}\| \|\mathbf{v}\|}.$$

The interval $[0, \pi]$ chooses the smaller unoriented angle. Thus directions with inclinations 10° and 350° are separated by 20° , not 340° .

The definition agrees with the earlier geometric interpretation:

$$\theta = 0 \iff \text{same direction}, \quad \theta = \frac{\pi}{2} \iff \text{perpendicular}, \quad \theta = \pi \iff \text{opposite directions}.$$

The Cauchy–Schwarz inequality proved later guarantees that the quotient inside the cosine lies between -1 and 1 .

2.1.9 The Cosine Theorem Emerges Immediately

Draw $\mathbf{u}$ and $\mathbf{v}$ from a common point. The displacement from the endpoint of $\mathbf{v}$ to the endpoint of $\mathbf{u}$ is

$$\mathbf{u} - \mathbf{v}.$$

Thus these three vectors determine a triangle with side lengths

$$\|\mathbf{u}\|, \quad \|\mathbf{v}\|, \quad \|\mathbf{u} - \mathbf{v}\|.$$

We calculate the square of the third side:

$$\begin{aligned} \|\mathbf{u} - \mathbf{v}\|^2 &= (\mathbf{u} - \mathbf{v}) \cdot (\mathbf{u} - \mathbf{v}) \\ &= \mathbf{u} \cdot \mathbf{u} - \mathbf{u} \cdot \mathbf{v} - \mathbf{v} \cdot \mathbf{u} + \mathbf{v} \cdot \mathbf{v} \\ &= \|\mathbf{u}\|^2 + \|\mathbf{v}\|^2 - 2\mathbf{u} \cdot \mathbf{v} \\ &= \|\mathbf{u}\|^2 + \|\mathbf{v}\|^2 - 2\|\mathbf{u}\| \|\mathbf{v}\| \cos \theta. \end{aligned}$$

If

$$a = \|\mathbf{u}\|, \quad b = \|\mathbf{v}\|, \quad c = \|\mathbf{u} - \mathbf{v}\|,$$

then


$$c^2 = a^2 + b^2 - 2ab \cos \theta.$$

Theorem

For a triangle with side lengths a , b , and c , where θ is the angle between the sides of lengths a and b ,

$$c^2 = a^2 + b^2 - 2ab \cos \theta.$$

The cosine theorem is not inserted as an external formula. It is the coordinate expansion of the displacement between the endpoints of two arbitrary vectors.




2.1.10 Projection Is Forced by the Right-Angle Condition

Let $\mathbf{v} \neq \mathbf{0}$. We seek a multiple $\lambda\mathbf{v}$ such that the remainder

$$\mathbf{u} - \lambda\mathbf{v}$$

is perpendicular to $\mathbf{v}$. Geometrically, this means decomposing $\mathbf{u}$ into a component parallel to $\mathbf{v}$ and a component perpendicular to $\mathbf{v}$:



The point H is chosen on the line spanned by $\mathbf{v}$. The vector from O to H is the parallel component $\lambda\mathbf{v}$, while the vector from H to the endpoint of $\mathbf{u}$ is the perpendicular remainder. The right-angle condition determines the value of λ .

The criterion just derived requires

$$(\mathbf{u} - \lambda\mathbf{v}) \cdot \mathbf{v} = 0.$$

Expanding gives

$$\mathbf{u} \cdot \mathbf{v} - \lambda(\mathbf{v} \cdot \mathbf{v}) = 0,$$

so

$$\lambda = \frac{\mathbf{u} \cdot \mathbf{v}}{\mathbf{v} \cdot \mathbf{v}}.$$

Thus the parallel component is

$$\text{proj}_{\mathbf{v}} \mathbf{u} = \frac{\mathbf{u} \cdot \mathbf{v}}{\mathbf{v} \cdot \mathbf{v}} \mathbf{v}.$$

The formula is the solution of a geometric requirement, not a rule to be memorised without explanation.

2.1.11 Orthogonal Decomposition and Its Consequences

The projection formula splits $\mathbf{u}$ into two geometrically different parts:

$$\mathbf{u} = \text{proj}_{\mathbf{v}} \mathbf{u} + (\mathbf{u} - \text{proj}_{\mathbf{v}} \mathbf{u}).$$

The first term is parallel to $\mathbf{v}$ and the second is perpendicular to $\mathbf{v}$. This is the orthogonal decomposition of $\mathbf{u}$ relative to the direction of $\mathbf{v}$.

Because the two components are perpendicular, the Pythagorean theorem gives

$$\|\mathbf{u}\|^2 = \|\text{proj}_{\mathbf{v}} \mathbf{u}\|^2 + \|\mathbf{u} - \text{proj}_{\mathbf{v}} \mathbf{u}\|^2.$$

Scalar component and vector projection

The signed scalar component of $\mathbf{u}$ in the direction of nonzero $\mathbf{v}$ is

$$\text{comp}_{\mathbf{v}} \mathbf{u} = \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{v}\|}.$$

It measures how much signed length of $\mathbf{u}$ lies along the direction of $\mathbf{v}$. The vector projection is obtained by multiplying this number by the unit vector in that direction:

$$\text{proj}_{\mathbf{v}} \mathbf{u} = \text{comp}_{\mathbf{v}} \mathbf{u} \frac{\mathbf{v}}{\|\mathbf{v}\|}.$$

The Cauchy–Schwarz inequality

The length of the parallel component cannot exceed the length of the entire vector:

$$|\text{comp}_{\mathbf{v}} \mathbf{u}| \leq \|\mathbf{u}\|.$$

Substitution gives

$$|\mathbf{u} \cdot \mathbf{v}| \leq \|\mathbf{u}\| \|\mathbf{v}\|.$$

Therefore

$$-1 \leq \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{u}\| \|\mathbf{v}\|} \leq 1,$$

which guarantees that the angle formula using arccos is meaningful.

The triangle inequality

The direct displacement $\mathbf{u} + \mathbf{v}$ cannot be longer than travelling first by $\mathbf{u}$ and then by $\mathbf{v}$. Algebraically,

$$\begin{aligned}\|\mathbf{u} + \mathbf{v}\|^2 &= \|\mathbf{u}\|^2 + 2\mathbf{u} \cdot \mathbf{v} + \|\mathbf{v}\|^2 \\ &\leq \|\mathbf{u}\|^2 + 2\|\mathbf{u}\| \|\mathbf{v}\| + \|\mathbf{v}\|^2 \\ &= (\|\mathbf{u}\| + \|\mathbf{v}\|)^2.\end{aligned}$$

Hence

$$\|\mathbf{u} + \mathbf{v}\| \leq \|\mathbf{u}\| + \|\mathbf{v}\|.$$

This is the algebraic form of the geometric fact that the straight segment is the shortest route between two points.

2.1.12 From the Plane to Three-Dimensional Vectors

Adding a third coordinate does not change the basic meaning of a vector. A vector still records a displacement, but now it records three independent coordinate changes:

$$\mathbf{v} = (v_1, v_2, v_3).$$

Addition and scalar multiplication remain componentwise, because successive changes in each coordinate direction still combine independently.

Applying the Pythagorean theorem first to the first two coordinate changes and then to the third gives

$$\|\mathbf{v}\| = \sqrt{v_1^2 + v_2^2 + v_3^2}.$$

The dot product extends in exactly the same way:

$$\mathbf{u} \cdot \mathbf{v} = u_1v_1 + u_2v_2 + u_3v_3.$$

Its geometric role is unchanged:

$$\mathbf{u} \cdot \mathbf{v} = \|\mathbf{u}\| \|\mathbf{v}\| \cos \theta.$$

Consequently,

$$\mathbf{u} \perp \mathbf{v} \iff \mathbf{u} \cdot \mathbf{v} = 0.$$

Thus the dot product detects perpendicularity in both two and three dimensions.

A second product available in three dimensions

In three dimensions there is another natural question. Given two vectors $\mathbf{u}$ and $\mathbf{v}$, can we construct a new vector perpendicular to both of them?

For

$$\mathbf{u} = (u_1, u_2, u_3), \quad \mathbf{v} = (v_1, v_2, v_3),$$

define

$$\mathbf{u} \times \mathbf{v} = (u_2v_3 - u_3v_2, u_3v_1 - u_1v_3, u_1v_2 - u_2v_1).$$

This formula is useful only because of what it guarantees. Direct substitution gives

$$(\mathbf{u} \times \mathbf{v}) \cdot \mathbf{u} = 0, \quad (\mathbf{u} \times \mathbf{v}) \cdot \mathbf{v} = 0.$$

Therefore $\mathbf{u} \times \mathbf{v}$ is perpendicular to both original vectors.


The statement is stronger. Every linear combination

$$a\mathbf{u} + b\mathbf{v}$$

is also perpendicular to $\mathbf{u} \times \mathbf{v}$, because

$$\begin{aligned} (\mathbf{u} \times \mathbf{v}) \cdot (a\mathbf{u} + b\mathbf{v}) &= a(\mathbf{u} \times \mathbf{v}) \cdot \mathbf{u} + b(\mathbf{u} \times \mathbf{v}) \cdot \mathbf{v} \\ &= a \cdot 0 + b \cdot 0 \\ &= 0. \end{aligned}$$

We do not yet turn this family of linear combinations into a theory of planes. That geometric construction belongs to the later chapter on lines and planes. For now, the important fact is simply that one cross product produces a vector perpendicular to every combination of its two inputs.



The cross product detects parallelism

If $\mathbf{u}$ and $\mathbf{v}$ are parallel, then one is a scalar multiple of the other. Let $\mathbf{v} = \lambda\mathbf{u}$. Substitution into the coordinate formula makes every component of the cross product vanish, so

$$\mathbf{u} \times \mathbf{v} = \mathbf{0}.$$

Conversely, if two nonzero vectors have zero cross product, then their directions are parallel. Hence

$$\mathbf{u} \parallel \mathbf{v} \iff \mathbf{u} \times \mathbf{v} = \mathbf{0}$$

for nonzero vectors.

This gives two complementary geometric tests:

$$\begin{array}{ll} \mathbf{u} \cdot \mathbf{v} = 0 & \iff \mathbf{u} \perp \mathbf{v}, \\ \mathbf{u} \times \mathbf{v} = \mathbf{0} & \iff \mathbf{u} \parallel \mathbf{v} \quad (\mathbf{u}, \mathbf{v} \neq \mathbf{0}). \end{array}$$

Order and orientation

There are two opposite directions perpendicular to both nonparallel vectors. To choose one consistently, we use the right-hand rule: curl the fingers of the right hand from $\mathbf{u}$ toward $\mathbf{v}$; the thumb indicates the direction of $\mathbf{u} \times \mathbf{v}$.

Reversing the order reverses the result:

$$\mathbf{v} \times \mathbf{u} = -(\mathbf{u} \times \mathbf{v}).$$

Thus the cross product is not commutative.

Its length satisfies

$$\|\mathbf{u} \times \mathbf{v}\| = \|\mathbf{u}\| \|\mathbf{v}\| \sin \theta.$$

This formula explains the parallelism test: for parallel vectors, $\theta = 0$ or $\theta = \pi$, so the sine and therefore the cross product vanish. The later geometry chapters will use the same quantity when areas, lines, and planes are constructed.

Summary

The two vector products answer different geometric questions:

operation	output	main test
$\mathbf{u} \cdot \mathbf{v}$	number	$\mathbf{u} \perp \mathbf{v} \iff \mathbf{u} \cdot \mathbf{v} = 0$
$\mathbf{u} \times \mathbf{v}$	vector	$\mathbf{u} \parallel \mathbf{v} \iff \mathbf{u} \times \mathbf{v} = \mathbf{0}$

Moreover, $\mathbf{u} \times \mathbf{v}$ is perpendicular to $\mathbf{u}$, to $\mathbf{v}$, and to every linear combination $a\mathbf{u} + b\mathbf{v}$. These facts are developed here as properties of vectors. They will become tools for constructing and analysing lines and planes only in the following chapter.

2.2 Changing the Basis: New Coordinates for the Same Vector

The preceding section developed the basic geometry and algebra of vectors: addition, scalar multiplication, linear combinations, length, angle, the dot product, parallelism, perpendicularity, and orthogonal projection. We now use that entire toolkit to study a new question: how the coordinate description of one fixed geometric vector changes when the basis is changed.

Up to this point every vector has been described with respect to the horizontal and vertical coordinate directions. That choice was natural because the Cartesian axes had already been constructed, but it was still only a choice. The geometric vector itself does not contain a preferred pair of coordinate axes.

We now ask a new question:

What numbers describe the same vector when we choose two different coordinate directions?

This is the problem of changing basis. The vector will remain fixed. Only the language used to describe it will change.

A basis is a pair of independent directions

Definition

Two nonparallel vectors $\mathbf{b}_1$ and $\mathbf{b}_2$ form a basis of the plane. This means that every vector $\mathbf{u}$ can be written in one and only one way as

$$\mathbf{u} = c_1 \mathbf{b}_1 + c_2 \mathbf{b}_2.$$

The numbers c_1 and c_2 are called the coordinates of $\mathbf{u}$ in the basis

$$\mathcal{B} = (\mathbf{b}_1, \mathbf{b}_2).$$

We write


$$[\mathbf{u}]_{\mathcal{B}} = (c_1, c_2).$$

The basis vectors need not be perpendicular and need not have length one. The only forbidden situation is that they are parallel, because then both vectors point along the same line and cannot describe arbitrary displacements in the plane.

Geometrically, the equation

$$\mathbf{u} = c_1 \mathbf{b}_1 + c_2 \mathbf{b}_2$$

means the following. Starting at the origin, move first by $c_1 \mathbf{b}_1$, then by $c_2 \mathbf{b}_2$. The endpoint must coincide with the endpoint of $\mathbf{u}$.



The coefficients are read from the oblique parallelogram. They are not lengths of perpendicular shadows. They tell us how many signed copies of each basis vector are needed.

The same vector on two different grids

Let the ordinary Cartesian basis be

$$\mathcal{E} = (\mathbf{e}_1, \mathbf{e}_2), \quad \mathbf{e}_1 = (1, 0), \quad \mathbf{e}_2 = (0, 1).$$

Choose also the slanted basis

$$\mathcal{B} = (\mathbf{b}_1, \mathbf{b}_2), \quad \mathbf{b}_1 = (2, 1), \quad \mathbf{b}_2 = (1, 2).$$

Consider the geometric vector

$$\mathbf{u} = (5, 4)$$

when it is described in the Cartesian basis.


On the rectangular grid, the endpoint is found by moving five units in the $\mathbf{e}_1$ direction and four units in the $\mathbf{e}_2$ direction. Hence

$$[\mathbf{u}]_{\mathcal{E}} = (5, 4).$$

On the slanted grid, we must find numbers c_1, c_2 such that

$$\mathbf{u} = c_1 \mathbf{b}_1 + c_2 \mathbf{b}_2.$$

The two pictures below show the same arrow. Only the grid used to read its coordinates has changed.



On the rectangular grid we read the coordinates by projecting parallel to the coordinate axes. On the slanted grid we do the analogous construction, but the lines through the endpoint are drawn parallel to the other basis vector. This produces an oblique parallelogram rather than a rectangle.

Finding the new coordinates by a system of equations

The geometric picture tells us what must be true, but it does not yet calculate c_1 and c_2 . Write the basis vectors in ordinary Cartesian coordinates:

$$\mathbf{b}_1 = (a_1, a_2), \quad \mathbf{b}_2 = (b_1, b_2), \quad \mathbf{u} = (x, y).$$

The equation

$$\mathbf{u} = c_1 \mathbf{b}_1 + c_2 \mathbf{b}_2$$

means equality of the two Cartesian components. Therefore

$$\begin{cases} a_1 c_1 + b_1 c_2 = x, \\ a_2 c_1 + b_2 c_2 = y. \end{cases}$$

This ordinary system of two linear equations is the complete algebraic content of changing from the Cartesian basis to the basis $\mathcal{B}$.

For our example,

$$\mathbf{b}_1 = (2, 1), \quad \mathbf{b}_2 = (1, 2), \quad \mathbf{u} = (5, 4),$$

so

$$2c_1 + c_2 = 5,$$

$$c_1 + 2c_2 = 4.$$

From the first equation,

$$c_2 = 5 - 2c_1.$$

Substituting into the second gives

$$c_1 + 2(5 - 2c_1) = 4,$$

so

$$-3c_1 = -6, \quad c_1 = 2.$$

Then

$$c_2 = 5 - 2 \cdot 2 = 1.$$

Thus

$$\boxed{[\mathbf{u}]_{\mathcal{B}} = (2, 1)}.$$

The calculation agrees with the oblique parallelogram in the figure.

A general algebraic formula in the plane

The same system can be solved symbolically. Starting from

$$a_1c_1 + b_1c_2 = x,$$

$$a_2c_1 + b_2c_2 = y,$$

multiply the first equation by b_2 and the second by b_1 :

$$a_1b_2c_1 + b_1b_2c_2 = xb_2,$$

$$a_2b_1c_1 + b_1b_2c_2 = yb_1.$$

Subtracting eliminates c_2 :

$$(a_1b_2 - a_2b_1)c_1 = xb_2 - yb_1.$$

Similarly, eliminating c_1 gives

$$(a_1b_2 - a_2b_1)c_2 = a_1y - a_2x.$$

Hence

$$\boxed{c_1 = \frac{xb_2 - yb_1}{a_1b_2 - a_2b_1}, \quad c_2 = \frac{a_1y - a_2x}{a_1b_2 - a_2b_1}}.$$

The denominator

$$a_1b_2 - a_2b_1$$

is nonzero precisely because the basis vectors are not parallel. If it were zero, the two chosen directions would collapse onto one line and the coordinates would not be uniquely determined.

Changing directly from one nonstandard basis to another

Suppose

$$\mathcal{B} = (\mathbf{b}_1, \mathbf{b}_2) \quad \text{and} \quad \mathcal{C} = (\mathbf{c}_1, \mathbf{c}_2)$$

are two bases. Assume that

$$[\mathbf{u}]_{\mathcal{B}} = (p, q).$$

Then first reconstruct the geometric vector:

$$\mathbf{u} = p\mathbf{b}_1 + q\mathbf{b}_2.$$

If

$$\mathbf{b}_1 = (a_1, a_2), \quad \mathbf{b}_2 = (b_1, b_2),$$

then

$$\mathbf{u} = (pa_1 + qb_1, pa_2 + qb_2).$$

Now seek numbers r, s such that

$$\mathbf{u} = r\mathbf{c}_1 + s\mathbf{c}_2.$$

Writing

$$\mathbf{c}_1 = (d_1, d_2), \quad \mathbf{c}_2 = (e_1, e_2),$$

we solve

$$\begin{aligned} d_1r + e_1s &= pa_1 + qb_1, \\ d_2r + e_2s &= pa_2 + qb_2. \end{aligned}$$

The solution (r, s) is exactly $[\mathbf{u}]_{\mathcal{C}}$.

Thus a change from one basis to another is not a mysterious new operation. It is two familiar steps:

$\text{old coordinates} \longrightarrow \text{the geometric vector} \longrightarrow \text{new coordinates}.$

Why perpendicular projection is special

For the Cartesian basis, the coordinate directions are perpendicular unit vectors. Therefore the horizontal and vertical coordinates can be read from ordinary perpendicular projections.

For a slanted basis, this is no longer true. Projecting $\mathbf{u}$ perpendicularly onto $\mathbf{b}_1$ and $\mathbf{b}_2$ produces two shadows, but those shadows generally do not add back to $\mathbf{u}$. The correct geometric construction uses lines parallel to the other basis vector, exactly as in the oblique grid above.

Remark

Later, after matrices have been introduced, the same systems of equations will be written much more compactly. Matrix notation will not create a new method; it will only abbreviate the elimination calculations performed here.

Summary

A basis is a pair of nonparallel coordinate directions. The same geometric vector may have different coordinate pairs in different bases. To find the coordinates in a slanted basis, write

$$\mathbf{u} = c_1\mathbf{b}_1 + c_2\mathbf{b}_2$$

and solve the resulting two scalar equations. Rectangular coordinates use perpendicular projections; slanted coordinates use an oblique parallelogram whose sides are parallel to the basis vectors.

2.2.1 A Negative Coordinate Means Motion in the Opposite Direction

The example $[\mathbf{u}]_{\mathcal{B}} = (2, 1)$ used positive coefficients. That is convenient for drawing a parallelogram, but basis coordinates need not be positive.

Keep

$$\mathbf{b}_1 = (2, 1), \quad \mathbf{b}_2 = (1, 2),$$

and consider


$$\mathbf{w} = 2\mathbf{b}_1 - \mathbf{b}_2.$$

Calculating with ordinary coordinate pairs gives

$$\mathbf{w} = 2(2, 1) - (1, 2) = (4, 2) - (1, 2) = (3, 0).$$

Therefore

$$[\mathbf{w}]_{\mathcal{E}} = (3, 0), \quad [\mathbf{w}]_{\mathcal{B}} = (2, -1).$$



The coefficient -1 does not mean a negative length. It means one copy of $\mathbf{b}_2$ taken in the opposite direction. Basis coordinates are signed instructions for reconstructing the vector.

2.3 Beyond a Change of Basis: Polar Coordinates

The preceding section changed the basis of the plane. In every such change we kept two fixed, nonparallel vector directions and described a vector by the signed amounts of those two directions. The coordinate grid could become slanted, but it still consisted of two families of parallel straight lines.

There is a more radical possibility. We may keep the same Euclidean plane and describe a point by quantities that are not coefficients of two fixed basis vectors. Polar coordinates provide the first important example.

Remark

A change from one vector basis to another is a *linear* change of coordinates. Polar coordinates are not obtained from a new vector basis. Their coordinate curves are rays and circles, and the directions used to read the coordinates vary from point to point.

Two different questions about the same point

The Cartesian description of a point uses two signed coordinate values obtained from projections parallel to the coordinate directions:

$$P = (x, y).$$

This is natural when two fixed straight directions are important. We may instead ask:

1. How far is the point from the origin?
2. In which direction do we move from the origin to reach it?

These questions lead to the polar pair

$$(r, \theta),$$

where, for a point different from the origin,

$$r > 0, \quad 0 \leq \theta < 2\pi.$$

The number r is the Euclidean distance from the origin, and θ is the angle measured counter-clockwise from the positive x -axis.

The origin is handled separately. In the usual non-unique polar notation every pair $(0, \theta)$ represents the origin. In this course we choose the single representative


$$O = (0, 0)$$

so that every point has exactly one allowed polar pair. Thus


$$\mathcal{P} = \{(0, 0)\} \cup ((0, \infty) \times [0, 2\pi)).$$

Rectangular and polar grids

The two descriptions below refer to the same Euclidean plane. What changes is the family of curves used to locate a point.



Cartesian grid
two families of parallel lines



polar grid
rays and concentric circles

In the Cartesian grid, keeping x fixed gives a vertical straight line and keeping y fixed gives a horizontal straight line. In the polar grid, keeping θ fixed gives a ray from the origin, while keeping r fixed gives a circle. This is the visible reason polar coordinates are not merely a slanted basis.

Converting between the descriptions

Let P have polar coordinates (r, θ) . The right triangle determined by P gives

$$\boxed{x = r \cos \theta, \quad y = r \sin \theta}.$$

Thus the position vector of P may be written as

$$\overrightarrow{OP} = r(\cos \theta, \sin \theta).$$

This resembles a scalar multiple of a unit vector, but the direction $(\cos \theta, \sin \theta)$ depends on the coordinate θ . There are not two fixed basis vectors whose constant coefficients are r and θ .

Conversely, if $(x, y) \neq (0, 0)$, then

$$r = \sqrt{x^2 + y^2}.$$

The polar angle is the unique number $\theta \in [0, 2\pi)$ satisfying

$$\cos \theta = \frac{x}{r}, \quad \sin \theta = \frac{y}{r}.$$

These two equations determine both the quadrant and the exact direction. The single expression $\arctan(y/x)$ is insufficient because it loses quadrant information and is undefined when $x = 0$. In computation the correct angle is usually obtained with the two-argument function $\text{atan2}(y, x)$, with a negative output shifted by 2π .

Example

For

$$(r, \theta) = \left(2, \frac{\pi}{3}\right),$$

we obtain

$$x = 2 \cos \frac{\pi}{3} = 1, \quad y = 2 \sin \frac{\pi}{3} = \sqrt{3}.$$

Thus the same point is written as

$$P = (1, \sqrt{3})$$

in Cartesian coordinates and as

$$P = \left(2, \frac{\pi}{3}\right)$$

in polar coordinates.

Why the transformation is nonlinear

For a change of vector basis, multiplying a vector by λ multiplies each of its coordinates by λ , and the coordinates of a sum are obtained by adding coordinate pairs. Polar coordinates do not behave this way.

For example, the points with polar pairs

$$(1, 0) \quad \text{and} \quad \left(1, \frac{\pi}{2}\right)$$

have Cartesian position vectors $(1, 0)$ and $(0, 1)$. Their vector sum is $(1, 1)$, whose polar coordinates are

$$\left(\sqrt{2}, \frac{\pi}{4}\right),$$

not

$$\left(2, \frac{\pi}{2}\right).$$

Therefore polar coordinate pairs cannot be added componentwise to add vectors. The formulas

$$x = r \cos \theta, \quad y = r \sin \theta$$

contain trigonometric functions and describe a genuinely nonlinear change of coordinates.

A circle becomes simple

Polar coordinates are useful when distance from the origin is the main geometric feature. Let

$$C_R = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 = R^2\}, \quad R > 0.$$

Substituting

$$x = r \cos \theta, \quad y = r \sin \theta$$

gives

$$r^2 \cos^2 \theta + r^2 \sin^2 \theta = R^2.$$


Since $\cos^2 \theta + \sin^2 \theta = 1$, this reduces to

$$\boxed{r = R}.$$

Thus

$$x^2 + y^2 = R^2 \quad \text{and} \quad r = R$$

describe the same geometric circle. The set has not changed; only the coordinate language has changed.



Summary

A vector basis uses fixed straight directions and leads to linear coordinate changes. Polar coordinates instead use a distance and an angle:

$$(r, \theta) \longleftrightarrow (r \cos \theta, r \sin \theta).$$

Their coordinate curves are circles and rays, their directions vary across the plane, and vector addition is not performed by adding polar coordinate pairs. This is our first example of a nonlinear coordinate system.

Chapter 3

Lines and Planes in the Plane and in Space

3.1 Lines in the Cartesian Plane

A line is one of the simplest geometric subsets of the plane. It can be drawn, described by a condition on coordinates, or defined explicitly as a subset of $\mathbb{R}^2$. We will keep these descriptions separate and explain how they are connected.

3.1.1 A Line as a Set of Points

Consider the condition

$$y = 2x + 1.$$

This is a relation between the coordinates x and y . The corresponding geometric object is the set

$$L = \{(x, y) \in \mathbb{R}^2 : y = 2x + 1\}.$$

Definition

A non-vertical line in the Cartesian plane is a set of the form

$$L_{m,b} = \{(x, y) \in \mathbb{R}^2 : y = mx + b\},$$

where $m, b \in \mathbb{R}$ are fixed.


Every value of x determines exactly one value of y . For example, for $y = 2x + 1$ we obtain

x	-2	-1	0	1	2
y	-3	-1	1	3	5

and therefore all of the points

$$(-2, -3), \quad (-1, -1), \quad (0, 1), \quad (1, 3), \quad (2, 5)$$

belong to L .



Remark

A finite list of points only gives examples of elements of a line. The complete line is the entire set of points satisfying its defining condition.

3.1.2 Slope, Angle, and Vertical Change

Take two distinct points on a non-vertical line. Let their horizontal and vertical changes be

$$\Delta x = x_2 - x_1, \quad \Delta y = y_2 - y_1.$$

Because the line is not vertical, $\Delta x \neq 0$.

Definition

The *slope* of a non-vertical line is


$$m = \frac{\Delta y}{\Delta x}.$$

It measures vertical change per unit of horizontal change.

For a line described by $y = mx + b$, replacing x by $x + \Delta x$ changes y by

$$\Delta y = m \Delta x.$$

Thus the coefficient m in the equation is exactly the geometric slope.



The line and the positive horizontal direction form an angle α . In the right triangle shown above, the tangent of α is defined as

$$\tan \alpha = \frac{\text{opposite side}}{\text{adjacent side}} = \frac{\Delta y}{\Delta x}.$$

Theorem

If α is an angle of inclination of a non-vertical line, then

$$m = \tan \alpha.$$

Thus slope is both a coordinate ratio and a description of direction.

The sign of m has a direct geometric meaning:

- $m > 0$: the line rises from left to right;
- $m < 0$: the line falls from left to right;
- $m = 0$: the line is horizontal.

3.1.3 The Constant Term and Axis Intersections

For the line

$$y = mx + b,$$

setting $x = 0$ gives $y = b$.

Theorem

The line

$$L_{m,b} = \{(x, y) \in \mathbb{R}^2 : y = mx + b\}$$

meets the y -axis at

$$(0, b).$$

To find an intersection with the x -axis, set $y = 0$ and solve the resulting equation for x .

Example

For

$$L = \{(x, y) \in \mathbb{R}^2 : y = -\frac{1}{2}x + 3\},$$

the y -axis intersection is $(0, 3)$. Setting $y = 0$ gives

$$0 = -\frac{1}{2}x + 3,$$

so $x = 6$. Hence the x -axis intersection is $(6, 0)$.


3.1.4 Vertical Lines

A vertical line cannot be written as $y = mx + b$, because one value of x is paired with every real value of y .

Definition

For a fixed $c \in \mathbb{R}$, the vertical line through $x = c$ is

$$V_c = \{(x, y) \in \mathbb{R}^2 : x = c\}.$$



Remark

A vertical line has no real-valued slope. For two distinct points on it, $\Delta x = 0$, so the quotient $\Delta y / \Delta x$ is undefined.

3.1.5 The General Linear Equation

Definition

Let $a, b, c \in \mathbb{R}$, where a and b are not both zero. The set

$$L_{a,b,c} = \{(x, y) \in \mathbb{R}^2 : ax + by = c\}$$

is a line in the Cartesian plane.

If $b \neq 0$, solving for y gives

$$y = -\frac{a}{b}x + \frac{c}{b}.$$

Therefore

$$\boxed{m = -\frac{a}{b}}, \quad \boxed{b_{\text{intercept}} = \frac{c}{b}}.$$

If $b = 0$, then $a \neq 0$ and the condition becomes

$$x = \frac{c}{a},$$

which describes a vertical line.

Example

For

$$L = \{(x, y) \in \mathbb{R}^2 : 2x + 3y = 6\},$$

solving for y gives

$$y = -\frac{2}{3}x + 2.$$

Thus the slope is $-2/3$, the y -axis intersection is $(0, 2)$, and the x -axis intersection is $(3, 0)$.

3.1.6 The Line Through Two Points

Let

$$P = (x_1, y_1), \quad Q = (x_2, y_2)$$

be distinct points.


Theorem

If $x_1 \neq x_2$, the unique line through P and Q has slope

$$m = \frac{y_2 - y_1}{x_2 - x_1}$$

and can be defined by

$$L = \{(x, y) \in \mathbb{R}^2 : y - y_1 = m(x - x_1)\}.$$



The second formula follows because every point (x, y) on the line must produce the same vertical-to-horizontal change ratio from P .

If $x_1 = x_2$, the two points determine the vertical line

$$L = \{(x, y) \in \mathbb{R}^2 : x = x_1\}.$$

Example

The points $P = (1, 2)$ and $Q = (4, 8)$ give

$$m = \frac{8 - 2}{4 - 1} = 2.$$

Hence

$$y - 2 = 2(x - 1),$$

which simplifies to $y = 2x$. Therefore

$$L = \{(x, y) \in \mathbb{R}^2 : y = 2x\}.$$

3.1.7 Parallel Lines

Definition

Two distinct lines L_1 and L_2 are *parallel* when they have no common point:

$$L_1 \neq L_2 \quad \text{and} \quad L_1 \cap L_2 = \emptyset.$$

Consider

$$L_1 = \{(x, y) \in \mathbb{R}^2 : y = m_1x + b_1\}, \quad L_2 = \{(x, y) \in \mathbb{R}^2 : y = m_2x + b_2\}.$$

A common point would have to satisfy

$$m_1x + b_1 = m_2x + b_2.$$

Equivalently,

$$(m_1 - m_2)x = b_2 - b_1.$$

If $m_1 \neq m_2$, this equation has exactly one solution for x , so the lines intersect. If $m_1 = m_2$, then either $b_1 = b_2$ and the lines are identical, or $b_1 \neq b_2$ and no common point exists.

Theorem

Two non-vertical lines are parallel precisely when

$$m_1 = m_2 \quad \text{and} \quad b_1 \neq b_2.$$

The equality of slopes is therefore a consequence of the geometric definition, not the definition itself.

Two distinct vertical lines are also parallel, because conditions $x = c_1$ and $x = c_2$ with $c_1 \neq c_2$ cannot hold at the same point.

3.1.8 Perpendicular Lines

Definition

Two intersecting lines are *perpendicular* when the angle formed at their intersection is a right angle:

$$\frac{\pi}{2}.$$

Let two non-vertical, non-horizontal lines have inclination angles α_1 and α_2 . They are perpendicular exactly when their directions differ by $\pi/2$. Thus we may write

$$\alpha_2 = \alpha_1 + \frac{\pi}{2}$$

up to reversing the direction along a line.

From the previously established relation $m = \tan \alpha$,

$$m_1 = \tan \alpha_1, \quad m_2 = \tan \alpha_2.$$

Using

$$\tan\left(\alpha + \frac{\pi}{2}\right) = -\frac{1}{\tan \alpha},$$

we obtain

$$m_2 = -\frac{1}{m_1}.$$

Theorem

For two non-vertical, non-horizontal lines,

$$L_1 \perp L_2 \iff m_1 m_2 = -1.$$

This criterion is derived from the geometric right-angle condition and the relation $m = \tan \alpha$.

A horizontal line and a vertical line form the remaining perpendicular case. The product formula is not applicable there because the vertical line has no real-valued slope.

Remark

Two lines may intersect without being perpendicular. Intersection only says that they have a common point. Perpendicularity additionally prescribes the angle at that point.

Summary

The central relations in this section are

$$L_{m,b} = \{(x, y) \in \mathbb{R}^2 : y = mx + b\},$$

$$m = \frac{\Delta y}{\Delta x} = \tan \alpha,$$

$$L_1 \parallel L_2 \iff m_1 = m_2 \text{ and } b_1 \neq b_2,$$

$$L_1 \perp L_2 \iff m_1 m_2 = -1 \quad (\text{non-vertical, non-horizontal case}).$$

Each algebraic criterion has been obtained from a geometric definition.

3.1.9 The Parametric Equation of a Line

The equations used so far describe a line by a condition that its coordinates must satisfy. A parametric equation describes the same line differently: we start from one point and move along a fixed direction.

Let

$$P_0 = (x_0, y_0)$$

be a point of the line, and let

$$(u, v) \neq (0, 0)$$

be a direction pair. Multiplying the direction pair by a real number t changes only the distance and orientation of the displacement, not its direction. Starting at P_0 therefore gives the points

$$(x, y) = (x_0, y_0) + t(u, v), \quad t \in \mathbb{R}.$$

Definition


The line through $P_0 = (x_0, y_0)$ with direction pair $(u, v) \neq (0, 0)$ is parametrised by

$$L = \{(x_0 + tu, y_0 + tv) : t \in \mathbb{R}\}.$$

Equivalently, its coordinates satisfy

$$\begin{cases} x = x_0 + tu, \\ y = y_0 + tv, \end{cases} \quad t \in \mathbb{R}.$$

When $t = 0$, the parametrisation gives the initial point P_0 . Positive and negative values of t move from P_0 in the two opposite directions along the line. Since t ranges over all real numbers, every point of the line is obtained.



If $u \neq 0$, eliminating the parameter gives

$$t = \frac{x - x_0}{u}$$

and hence

$$y - y_0 = \frac{v}{u}(x - x_0).$$

Thus the slope is

$$m = \frac{v}{u}.$$

If $u = 0$, then $v \neq 0$ and the parametrisation becomes

$$x = x_0, \quad y = y_0 + tv,$$

which describes a vertical line. The parametric form therefore treats non-vertical and vertical lines in one uniform way.

For two distinct points

$$P = (x_1, y_1), \quad Q = (x_2, y_2),$$

the difference

$$Q - P = (x_2 - x_1, y_2 - y_1)$$

is a direction pair for their line. Consequently,

$$L = \{P + t(Q - P) : t \in \mathbb{R}\},$$

or in coordinates,

$$\begin{cases} x = x_1 + t(x_2 - x_1), \\ y = y_1 + t(y_2 - y_1), \end{cases} \quad t \in \mathbb{R}.$$

Example

The line through $P = (1, 2)$ with direction pair $(3, 6)$ has the parametrisation

$$\begin{cases} x = 1 + 3t, \\ y = 2 + 6t, \end{cases} \quad t \in \mathbb{R}.$$

Because $3 \neq 0$, we have

$$t = \frac{x - 1}{3}.$$

Substitution into the second equation gives

$$y = 2 + 6 \frac{x - 1}{3} = 2x.$$

Thus the parametric and Cartesian equations describe the same line.

Remark

A parametrisation of a line is not unique. Replacing (u, v) by any non-zero multiple $\lambda(u, v)$ changes the parameter but not the set of points on the line.

3.2 Lines in Three-Dimensional Space

A point in three-dimensional Cartesian space has coordinates

$$P = (x, y, z) \in \mathbb{R}^3.$$

A line is determined by one point and one nonzero direction.

Definition

Let

$$P_0 = (x_0, y_0, z_0)$$

and let

$$\mathbf{v} = (a, b, c) \neq (0, 0, 0).$$

The line through P_0 with direction vector $\mathbf{v}$ is

$$L = \{P_0 + t\mathbf{v} : t \in \mathbb{R}\}.$$


Equivalently,

$$(x, y, z) = (x_0, y_0, z_0) + t(a, b, c).$$

In coordinates this becomes the system

$$\begin{cases} x = x_0 + at, \\ y = y_0 + bt, \\ z = z_0 + ct, \end{cases} \quad t \in \mathbb{R}.$$

As t varies, the point moves along the line. Positive and negative values of t describe the two opposite directions along the same geometric line.



3.2.1 The Line Through Two Points in Space

Let

$$P = (x_1, y_1, z_1), \quad Q = (x_2, y_2, z_2)$$

be distinct points. Their difference

$$Q - P = (x_2 - x_1, y_2 - y_1, z_2 - z_1)$$

is a nonzero direction vector of the line through them.

Theorem

The unique line through two distinct points $P, Q \in \mathbb{R}^3$ is

$$L = \{P + t(Q - P) : t \in \mathbb{R}\}.$$

In coordinates,

$$\begin{cases} x = x_1 + t(x_2 - x_1), \\ y = y_1 + t(y_2 - y_1), \\ z = z_1 + t(z_2 - z_1). \end{cases}$$

Example

For

$$P = (1, 0, 2), \quad Q = (3, 4, 1),$$

a direction vector is

$$Q - P = (2, 4, -1).$$

Therefore the line through P and Q is

$$(x, y, z) = (1, 0, 2) + t(2, 4, -1),$$

or

$$\begin{cases} x = 1 + 2t, \\ y = 4t, \\ z = 2 - t. \end{cases}$$

3.2.2 Symmetric Form

If a , b , and c are all nonzero, the parameter can be eliminated from

$$x = x_0 + at, \quad y = y_0 + bt, \quad z = z_0 + ct.$$

This gives

$$\frac{x - x_0}{a} = \frac{y - y_0}{b} = \frac{z - z_0}{c}.$$

If one component of the direction vector is zero, the corresponding coordinate is constant. For example, if $c = 0$, then $z = z_0$, while the remaining two coordinates may still be written in symmetric form.

3.2.3 Parallel and Intersecting Lines in Space

Consider

$$L_1 = P_1 + t\mathbf{v}_1, \quad L_2 = P_2 + s\mathbf{v}_2.$$

The lines are parallel when their direction vectors are proportional:

$$\mathbf{v}_2 = \lambda\mathbf{v}_1$$

for some nonzero λ .

To test whether two nonparallel lines intersect, one solves

$$P_1 + t\mathbf{v}_1 = P_2 + s\mathbf{v}_2$$

for the two parameters t and s . This is a system of three scalar equations. It may have one solution or no solution.

Definition

Two lines in $\mathbb{R}^3$ are *skew* when they are neither parallel nor intersecting.

Skew lines have no analogue in the plane. In two dimensions, two distinct nonparallel lines must intersect; in three dimensions, one line may pass above or beside another without meeting it.

Summary

A line in $\mathbb{R}^3$ is described by

$$L = \{P_0 + t\mathbf{v} : t \in \mathbb{R}\}, \quad \mathbf{v} \neq \mathbf{0}.$$

Two distinct points determine the direction $Q - P$. Parallel lines have proportional direction vectors, while nonparallel lines may either intersect or be skew.

3.3 Planes in Three-Dimensional Space

A line in $\mathbb{R}^3$ is determined by one point and one direction. A plane requires one point and two independent directions. Equivalently, it may be described by one point and a direction perpendicular to the whole plane.

3.3.1 A Plane Through a Point with Two Direction Vectors

Let

$$P_0 = (x_0, y_0, z_0)$$

be a point and let

$$\mathbf{u} = (u_1, u_2, u_3), \quad \mathbf{v} = (v_1, v_2, v_3)$$

be non-parallel vectors.

Definition


The plane through P_0 generated by $\mathbf{u}$ and $\mathbf{v}$ is

$$\Pi = \{P_0 + s\mathbf{u} + t\mathbf{v} : s, t \in \mathbb{R}\}.$$

In coordinates,

$$\begin{cases} x = x_0 + su_1 + tv_1, \\ y = y_0 + su_2 + tv_2, \\ z = z_0 + su_3 + tv_3. \end{cases}$$

The parameters s and t allow independent motion in two directions. Because $\mathbf{u}$ and $\mathbf{v}$ are not parallel, these motions fill a plane rather than a single line.



3.3.2 The Normal Vector and the Cartesian Equation

A nonzero vector

$$\mathbf{n} = (a, b, c)$$

is called a normal vector to a plane when it is perpendicular to every direction contained in that plane.

For a point $P = (x, y, z)$ on the plane, the displacement from P_0 to P is

$$P - P_0 = (x - x_0, y - y_0, z - z_0).$$

This displacement lies in the plane and is therefore perpendicular to $\mathbf{n}$. Hence

$$\mathbf{n} \cdot (P - P_0) = 0.$$

Theorem

A plane through $P_0 = (x_0, y_0, z_0)$ with normal vector $\mathbf{n} = (a, b, c) \neq (0, 0, 0)$ has equation

$$a(x - x_0) + b(y - y_0) + c(z - z_0) = 0.$$

After collecting constants, this becomes

$$ax + by + cz = d$$

for some $d \in \mathbb{R}$.

Thus a first-degree equation in three variables describes a plane, just as a first-degree equation in two variables describes a line.

Example

The plane through $P_0 = (1, -1, 2)$ with normal vector $\mathbf{n} = (2, 1, -3)$ satisfies

$$2(x - 1) + (y + 1) - 3(z - 2) = 0.$$

Therefore

$$2x + y - 3z = -5.$$

3.3.3 A Plane Through Three Points

Let P , Q , and R be three points that do not lie on one line. Then

$$\mathbf{u} = Q - P, \quad \mathbf{v} = R - P$$

are non-parallel direction vectors in the plane.

Theorem

Three non-collinear points determine exactly one plane. A parametric equation is

$$\Pi = \{P + s(Q - P) + t(R - P) : s, t \in \mathbb{R}\}.$$

A normal vector may be obtained from the cross product

$$\mathbf{n} = (Q - P) \times (R - P).$$

Example

For

$$P = (1, 0, 0), \quad Q = (0, 1, 0), \quad R = (0, 0, 1),$$

we have

$$Q - P = (-1, 1, 0), \quad R - P = (-1, 0, 1).$$

Their cross product is

$$(Q - P) \times (R - P) = (1, 1, 1).$$

Using the point P , the plane equation is

$$(x - 1) + y + z = 0,$$

so

$$x + y + z = 1.$$

3.3.4 Relative Positions of Planes

Consider

$$\Pi_1 : a_1x + b_1y + c_1z = d_1, \quad \Pi_2 : a_2x + b_2y + c_2z = d_2,$$

with normal vectors

$$\mathbf{n}_1 = (a_1, b_1, c_1), \quad \mathbf{n}_2 = (a_2, b_2, c_2).$$

Theorem

Two distinct planes are parallel precisely when their normal vectors are parallel. If the normal vectors are not parallel, the planes intersect in a line.

The planes are perpendicular when their normal vectors are perpendicular:

$$\mathbf{n}_1 \cdot \mathbf{n}_2 = 0.$$

3.3.5 A Line as the Intersection of Two Planes

In $\mathbb{R}^3$, two non-parallel planes usually meet in a line. Thus a line may also be described by a system

$$\begin{cases} a_1x + b_1y + c_1z = d_1, \\ a_2x + b_2y + c_2z = d_2. \end{cases}$$

A direction vector for the intersection line is perpendicular to both normal vectors, so one may take

$$\mathbf{v} = \mathbf{n}_1 \times \mathbf{n}_2.$$

Summary

A plane in $\mathbb{R}^3$ may be described in two fundamental ways:

$$P = P_0 + s\mathbf{u} + t\mathbf{v}$$

with two non-parallel direction vectors, or

$$ax + by + cz = d$$

with a nonzero normal vector (a, b, c) . Three non-collinear points determine one plane, and two non-parallel planes intersect in a line.

Chapter 4

Second-Degree Equations, Conic Sections, and Quadric Surfaces

4.1 Second-Degree Equations and Conic Sections

Lines arise from equations in which the coordinates occur only to the first power. The next natural class consists of equations in which terms of total degree two are allowed.

Definition

A *second-degree equation* in the Cartesian plane has the form

$$Ax^2 + Bxy + Cy^2 + Dx + Ey + F = 0,$$

where $A, B, C, D, E, F \in \mathbb{R}$ and at least one of A, B , or C is nonzero. Its solution set is the collection of all points $(x, y) \in \mathbb{R}^2$ whose coordinates satisfy the equation.

The terms

$$Ax^2 + Bxy + Cy^2$$

form the quadratic part. The remaining terms affect the position and size of the curve, while the quadratic part largely determines its geometric type.

4.1.1 A First Classification

For the equation

$$Ax^2 + Bxy + Cy^2 + Dx + Ey + F = 0,$$

define

$$\Delta = B^2 - 4AC.$$

Theorem

For a non-degenerate real conic, the sign of Δ gives the following classification:

$$\begin{array}{l|l} \Delta < 0 & \text{ellipse type, including circles,} \\ \Delta = 0 & \text{parabola type,} \\ \Delta > 0 & \text{hyperbola type.} \end{array}$$

This classification is preserved by rotations and translations of the coordinate system. A rotation can remove the mixed term Bxy , while a translation can move the centre or vertex to a simpler position.

Remark

The sign of Δ identifies the conic type only after degenerate cases are separated. A second-degree equation may also describe a single point, no real points, one line, or two lines. These possibilities are discussed below.

4.1.2 Circle

A circle with centre (h, k) and radius $r > 0$ is described by

$$(x - h)^2 + (y - k)^2 = r^2.$$


After expansion,

$$x^2 + y^2 - 2hx - 2ky + h^2 + k^2 - r^2 = 0.$$

Here $A = C = 1$ and $B = 0$, so

$$\Delta = 0^2 - 4 \cdot 1 \cdot 1 = -4 < 0.$$

A circle is therefore a special ellipse-type conic in which the coefficients of x^2 and y^2 are equal and there is no mixed term after a suitable rotation.



4.1.3 Ellipse

The standard ellipse centred at the origin has equation

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1, \quad a > 0, \quad b > 0.$$


Equivalently,

$$b^2x^2 + a^2y^2 - a^2b^2 = 0.$$

Thus $A = b^2$, $B = 0$, and $C = a^2$, so

$$\Delta = -4a^2b^2 < 0.$$

The parameters a and b are the semi-axis lengths. When $a = b$, the ellipse becomes a circle.



4.1.4 Parabola

A standard vertical parabola has equation

$$y = ax^2, \quad a \neq 0,$$

or equivalently

$$ax^2 - y = 0.$$


Here $A = a$, $B = 0$, and $C = 0$, hence

$$\Delta = 0.$$

The sign of a determines whether the parabola opens upward or downward. A translated form is

$$y - k = a(x - h)^2,$$

whose vertex is (h, k) .



4.1.5 Hyperbola

A standard hyperbola centred at the origin has equation

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1, \quad a > 0, \quad b > 0.$$

Equivalently,


$$b^2x^2 - a^2y^2 - a^2b^2 = 0.$$

Here $A = b^2$, $B = 0$, and $C = -a^2$, so

$$\Delta = 4a^2b^2 > 0.$$

Its asymptotes are

$$y = \frac{b}{a}x, \quad y = -\frac{b}{a}x.$$



4.1.6 The Mixed Term and Rotation

The term Bxy indicates that the natural axes of the conic may be rotated relative to the chosen Cartesian axes. For example,

$$x^2 + 2xy + y^2 = 1$$

can be written as

$$(x + y)^2 = 1,$$

so it is not a non-degenerate ellipse: it describes the two parallel lines

$$x + y = 1, \quad x + y = -1.$$

For a genuine rotated conic, one introduces new coordinates (u, v) obtained by a rotation through an angle θ :

$$x = u \cos \theta - v \sin \theta, \quad y = u \sin \theta + v \cos \theta.$$

Choosing θ appropriately removes the uv term. When $A \neq C$, one may choose θ so that

$$\tan(2\theta) = \frac{B}{A - C}.$$

The resulting equation can then be classified in the new coordinates using the standard forms above.

4.1.7 Degenerate Second-Degree Equations

Not every second-degree equation describes a non-degenerate conic. Factorisation or completing squares may reveal a simpler set.

Example

The equation


$$x^2 - y^2 = 0$$

factors as

$$(x - y)(x + y) = 0.$$

Its solution set is the union of two intersecting lines:

$$y = x \quad \text{or} \quad y = -x.$$



Other common degenerate cases include

$$x^2 + y^2 = 0,$$

which gives only the point $(0,0)$,

$$x^2 + y^2 + 1 = 0,$$

which has no real solutions, and

$$x^2 = 0,$$

which describes the single line $x = 0$ counted algebraically twice.

Summary

Second-degree equations have the general form

$$Ax^2 + Bxy + Cy^2 + Dx + Ey + F = 0.$$

For non-degenerate real conics, the discriminant

$$\Delta = B^2 - 4AC$$

separates the three principal types:

$\Delta < 0$ gives ellipse type, $\Delta = 0$ gives parabola type, $\Delta > 0$ gives hyperbola type.

Translations and rotations reduce many equations to standard forms, while factorisation and completing squares expose degenerate cases.

4.2 Second-Degree Surfaces in Three-Dimensional Space

The three-dimensional analogue of a conic section is a *quadric surface*. Instead of an equation in two coordinates, we now allow a second-degree equation in three coordinates.

Definition

A second-degree equation in $\mathbb{R}^3$ has the form

$$Ax^2 + By^2 + Cz^2 + Dxy + Exz + Fyz + Gx + Hy + Iz + J = 0,$$

where at least one quadratic coefficient is nonzero. Its solution set is called a quadric surface when it forms a non-degenerate surface.

Translations move the centre or vertex of the surface. Rotations of the coordinate system remove mixed terms such as xy , xz , and yz . After these changes of coordinates, the principal quadric surfaces have simple standard forms.


4.2.1 Ellipsoid

An ellipsoid centred at the origin has equation

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1, \quad a, b, c > 0.$$

Its intersections with the coordinate planes are ellipses. When $a = b = c = r$, the ellipsoid becomes the sphere

$$x^2 + y^2 + z^2 = r^2.$$




4.2.2 Elliptic Paraboloid

An elliptic paraboloid has standard equation

$$z = \frac{x^2}{a^2} + \frac{y^2}{b^2}, \quad a, b > 0.$$

Horizontal sections $z = k > 0$ are ellipses, while vertical sections through the z -axis are parabolas. The surface has one vertex and opens in one direction.




4.2.3 Hyperbolic Paraboloid

A hyperbolic paraboloid has standard equation

$$z = \frac{x^2}{a^2} - \frac{y^2}{b^2}.$$

Sections parallel to the xz -plane and yz -plane are parabolas opening in opposite directions. Horizontal sections are hyperbolas, except at $z = 0$, where the section consists of two intersecting lines. The surface is commonly called a saddle.




4.2.4 Hyperboloid of One Sheet

A hyperboloid of one sheet has standard equation

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} - \frac{z^2}{c^2} = 1.$$

Horizontal sections are ellipses. Vertical sections through the z -axis are hyperbolas. The surface is connected and has a narrow central waist.




4.2.5 Hyperboloid of Two Sheets

A hyperboloid of two sheets has standard equation

$$-\frac{x^2}{a^2} - \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1.$$

The condition forces $|z| \geq c$, so the surface has two disconnected parts. Horizontal sections are ellipses, while suitable vertical sections are hyperbolas.




4.2.6 Elliptic Cone

An elliptic cone has equation

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} - \frac{z^2}{c^2} = 0.$$

It consists of two nappes meeting at the origin. Horizontal sections away from the origin are ellipses. Vertical sections through the axis are pairs of intersecting lines.



4.2.7 Cylinders as Second-Degree Surfaces


If one coordinate does not occur in the equation, the corresponding planar curve is extended parallel to that coordinate axis. For example,

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$

describes an elliptic cylinder parallel to the z -axis, while

$$y = x^2$$

describes a parabolic cylinder parallel to the z -axis.



4.2.8 Degenerate Quadrics

As in the planar case, a second-degree equation in three variables may be degenerate. It may describe a point, a line, a plane, two planes, or no real points. For example,

$$x^2 + y^2 + z^2 = 0$$

describes only the origin, while

$$x^2 - y^2 = 0$$

factors as

$$(x - y)(x + y) = 0$$

and therefore describes the union of the two planes

$$x - y = 0 \quad \text{and} \quad x + y = 0.$$

Summary

Second-degree equations in three variables produce quadric surfaces. After translations and rotations, the main non-degenerate types are ellipsoids, elliptic and hyperbolic paraboloids, hyperboloids of one or two sheets, and elliptic cones. Equations missing one coordinate give cylinders, while factorisation may reveal degenerate unions of planes or smaller sets.

Chapter 5

Chapter 6

Chapter 7

Chapter 8

Chapter 9

Chapter 10

Chapter 11

Chapter 12