

# Evolving a Single Puncture Using the Moving Puncture Method\*

Ann Author<sup>†</sup> and Second Author<sup>‡</sup>

*Authors' institution and/or address*

*This line break forced with \\*

(MUSO Collaboration)

Charlie Author<sup>§</sup>

*Second institution and/or address*

*This line break forced and*

*Third institution, the second for Charlie Author*

Delta Author

*Authors' institution and/or address*

*This line break forced with \\*

(CLEO Collaboration)

(Dated: October 6, 2023)

In this paper, we investigate the evolution of a single puncture within the framework of the Moving Puncture Method. The Moving Puncture Method has proven to be a powerful technique in numerical relativity for evolving black hole spacetimes. In our study, we focus on its application to the evolution of a single puncture, which can serve as a fundamental building block in simulating more complex spacetime scenarios.

**Usage:** Secondary publications and information retrieval purposes.

**Structure:** You may use the `description` environment to structure your abstract; use the optional argument of the `\item` command to give the category of each item.

## I. INTRODUCTION

Since the first successful simulations of merging black-hole binaries for computing accurate gravitational waveforms, various formulations and techniques have been employed. These include the BSSN (*Baumgarte-Shapiro-Shibata-Nakamura*) formulation [1–3], conformal decompositions [4, 5], the Bowen-York approach [6], and the moving puncture gauge [7]. These methods have been in use until recently.

Evolving a Schwarzschild black hole serves as a valuable benchmark for assessing the performance of numerical codes, given that we possess its analytical solutions.

In this paper, we present a method for obtaining analytic solutions for a Schwarzschild black hole under time-independent conditions. We provide numerical setups and results for comparison with these analytical solutions.

## II. INITIAL DATA

### A. Spatial Metric

We need to specify the spatial metric  $\gamma_{ij}$  and the extrinsic curvature  $K_{ij}$  which satisfy the Hamiltonian constraint

$$R + K^2 - K_{ij}K^{ij} = 16\pi\rho, \quad (1)$$

and the momentum constraint

$$D_j(K^{ij} - \gamma^{ij}K) = 8\pi j^i. \quad (2)$$

To construct a single puncture initial data, we adopt York-Lichnerowicz conformal decompositions [4, 5]. The spatial metric is decomposed into a conformal factor  $\psi$  multiplying an conformally related metric:

$$\gamma_{ij} = \psi^4 \bar{\gamma}_{ij}. \quad (3)$$

With this, (1) turns into

$$\bar{D}^2\psi - \frac{\psi}{8}\bar{R} + \frac{\psi^5}{8}(K_{ij}K^{ij} - K^2) = -2\pi\psi^5\rho, \quad (4)$$

where we define the conformal Laplace operator

$$\bar{D}^2\psi \equiv \bar{\gamma}^{ij}\bar{D}_i\bar{D}_j\psi. \quad (5)$$

We assume

1) *vacuum spacetime* ( $\rho = 0, j^i = 0$ ),

2) *time symmetry* ( $\partial_t\gamma_{ij} = 0, \beta^i = 0$ ),

so the extrinsic curvature

$$K_{ij} = \frac{1}{2\alpha}(\mathcal{L}_\beta\gamma_{ij} - \partial_t\gamma_{ij}) \quad (6)$$

vanishes and also  $K = 0$ . Therefore, (2) satisfied identically. If we assume *conformal flatness*, the conformally related metric to be flat,

$$\bar{\gamma}_{ij} = \eta_{ij}, \quad (7)$$

---

\* A footnote to the article title

<sup>†</sup> Also at Physics Department, XYZ University.

<sup>‡</sup> Second.Author@institution.edu

<sup>§</sup> <http://www.Second.institution.edu/~Charlie.Author>

(4) reduces to the simple Laplace equation

$$\bar{D}^2 \psi = 0, \quad (8)$$

which has the solution

$$\psi = A + \frac{B}{r}. \quad (9)$$

To satisfy asymptotically flat as  $r \rightarrow \infty$ , and ADM mass to be  $M$ , we should choose  $A = 1$  and  $B = \frac{M}{2}$ :

$$\psi = 1 + \frac{M}{2r}. \quad (10)$$

TwoPUNCTURES [8] uses the Bowen-York approach [6] and the method presented in [7]. Since there is a single puncture with neither spin nor momentum, the result should be the same as in (10). We set TP\_epsilon to  $10^{-6}$  and TP\_Tiny to 0, resulting in

$$\psi = 1 + \frac{M}{2\tilde{r}}, \quad (11)$$

where

$$\tilde{r} = (r^4 + 10^{-24})^{1/4}. \quad (12)$$

The initial metric set by TwoPUNCTURES exactly takes the form

$$\gamma_{ij} = \left(1 + \frac{M}{2\tilde{r}}\right)^4 \eta_{ij}, \quad (13)$$

for all grid points. Fig. 1 shows  $\gamma_{xx}$  along the  $x$ -axis at the initial time. Each marker represents a grid point for a respective refinement level. The non-diagonal components of the spatial metric and all components of the extrinsic curvature have explicit values of 0.




FIG. 1.  $\gamma_{xx}$  along the  $x$ -axis at the initial time. Only 3 of the 7 refinement levels are plotted.

## B. Constraints

From (1) and (2), we can define

$$\mathcal{H} \equiv R + K^2 - K_{ij} K^{ij} - 16\pi\rho, \quad (14)$$

$$\mathcal{M}^i \equiv D_j (K^{ij} - \gamma^{ij} K) - 8\pi j^i, \quad (15)$$

and it should satisfy that  $\mathcal{H} = 0$  and  $\mathcal{M}^i = 0$ . Fig. 2 shows  $\log_{10} |\mathcal{H}|$  on the  $xy$  plane at the initial time. Every value at each grid point is less than  $10^{-7}$ . In the case of  $\mathcal{M}^i$ , all of the values are explicitly 0.




FIG. 2.  $\log_{10} |\mathcal{H}|$  on the  $xy$  plane at the initial time.

Fig. 3 shows the Hamiltonian constraint violation  $\mathcal{H}$  along the  $x$  axis. The data at  $x = 0$  has been cut since its value is on the order of 1 at the late time.  $\mathcal{H}$  increased as time passed, and most peak point's  $x$  coordinate also increased. But it still remains on the order of  $10^{-6}$  at  $t = 115.2M$ .




FIG. 3. The Hamiltonian constraint violation along the  $x$  axis.

### III. EVOLUTION

#### A. Gauge Condition

The 1+log slicing with advection term is given by

$$\partial_t \alpha = -2\alpha K + \beta^i \partial_i \alpha, \quad (16)$$

and the hyperbolic gamma driver condition for the shift with advection term is defined as

$$\partial_t \beta^i = \frac{3}{4} B^i + \beta^j \partial_j \beta^i, \quad (17)$$

$$\partial_t B^i = \partial_t \bar{\Gamma}^i - B^i + \beta^j \partial_j B^i, \quad (18)$$

which is typically employed in the moving puncture method [9, 10]. Sometimes we can drop the advection term, simplifying (16), (17), and (18) to

$$\partial_t \alpha = -2\alpha K, \quad (19)$$

$$\partial_t \beta^i = \frac{3}{4} B^i, \quad (20)$$

$$\partial_t B^i = \partial_t \bar{\Gamma}^i - B^i, \quad (21)$$

which we adopt in this paper.

We used the twopunctures-averaged initial lapse, which is given by

$$\alpha = \frac{1}{2}(1 + \alpha'), \quad (22)$$

where

$$\alpha' = \frac{1 - \frac{M}{2r}}{1 + \frac{M}{2r}}, \quad (23)$$

ensuring that the lapse satisfies  $0 \leq \alpha \leq 1$ .

When the simulation progresses sufficiently, and the solution becomes time-independent, we have  $\partial_t \alpha = 0$ , implying  $K = 0$ . Since we choose the hyperbolic gamma driver condition, it has a special solution (see [11, 12])

$$\alpha = \sqrt{1 - \frac{2M}{r_s} + \frac{27M^4}{16r_s^4}}, \quad (24)$$

where  $r_s$  is an areal radius related to the isotropic radius  $r$  as

$$r = \frac{2r_s + M + (4r_s^2 + 4Mr_s + 3M^2)^{1/2}}{4} \times \left( \frac{(4 + 3\sqrt{2})(2r_s - 3M)}{8r_s + 6M + 3(8r_s^2 + 8Mr_s + 6M^2)^{1/2}} \right)^{1/\sqrt{2}}. \quad (25)$$

The conformal factor is given by  $\psi = \left(\frac{r_s}{r}\right)^{1/2}$ , and when substituted into (25), we obtain

$$\psi = \left( \frac{4r_s}{2r_s + M + (4r_s^2 + 4Mr_s + 3M^2)^{1/2}} \right)^{1/2} \left( \frac{8r_s + 6M + 3(8r_s^2 + 8Mr_s + 6M^2)^{1/2}}{(4 + 3\sqrt{2})(2r_s - 3M)} \right)^{1/2\sqrt{2}}. \quad (26)$$

Fig. 4 shows  $\alpha$  along the  $x$  axis at several times. After  $t = 49.152M$ ,  $\alpha$  remains almost constant. The black solid line represents the analytic solution given in (24).

Fig. 5 shows  $\gamma_{xx}$  along the  $x$  axis. The black solid line indicates the analytic solution  $\gamma_{ij} = \psi^4 \eta_{ij}$ , where  $\psi$  is given by (26).

#### B. BSSN

The 3+1 ADM evolution equations are given as:

$$(\partial_t - \mathcal{L}_\beta) \gamma_{ij} = -2\alpha K_{ij}, \quad (27)$$

$$\begin{aligned} (\partial_t - \mathcal{L}_\beta) K_{ij} = & -D_i D_j \alpha + \alpha (R_{ij} + K K_{ij} - 2K_{ik} K^k_j) \\ & + 4\pi \alpha M_{ij}, \end{aligned} \quad (28)$$

However, this set of partial differential equations is only weakly hyperbolic and is therefore not suitable for stable numerical evolution. To address this, we adopt the BSSN (*Baumgarte-Shapiro-Shibata-Nakamura*) formulation [1–3].

In the BSSN formulation, the spatial metric  $\gamma_{ij}$  is decomposed into a conformally related metric as in (3), with  $\det(\tilde{\gamma}_{ij}) = 1$ . The extrinsic curvature is also decomposed into its trace and traceless parts, and we conformally transform the traceless part as follows:

$$K_{ij} = e^{4\phi} \tilde{A}_{ij} + \frac{1}{3} \gamma_{ij} K. \quad (29)$$




FIG. 4.  $\alpha$  along the  $x$  axis at several times. Only 5 refinement levels are plotted. The time unit is divided by  $M$ .




FIG. 5.  $\gamma_{xx}$  along the  $x$  axis at several times. Only refinement level 4 is plotted. The time unit is divided by  $M$ .

We then promote the following variables to evolution variables:

$$\phi = \ln \psi = \frac{1}{12} \ln \gamma, \quad (30)$$

as well as the conformal connection functions:

$$\bar{\Gamma}^i = \bar{\gamma}^{jk} \bar{\Gamma}_{jk}^i = -\partial_j \bar{\gamma}^{ij}. \quad (31)$$

The evolution equation for  $\gamma_{ij}$  splits into two equations:

$$\partial_t \phi = -\frac{1}{6} \alpha K + \beta^i \partial_i \phi + \frac{1}{6} \partial_i \beta^i, \quad (32)$$

$$\begin{aligned} \partial_t \bar{\gamma}_{ij} = & -2\alpha \bar{A}_{ij} + \beta^k \partial_k \bar{\gamma}_{ij} \\ & + \bar{\gamma}_{ik} \partial_j \beta^k + \bar{\gamma}_{jk} \partial_i \beta^k - \frac{2}{3} \bar{\gamma}_{ij} \partial_k \beta^k. \end{aligned} \quad (33)$$

The evolution equation for  $K_{ij}$  also splits into two equations:

$$\begin{aligned} \partial_t K = & -D^i D_i \alpha + \alpha (\bar{A}_{ij}^{ij} + \frac{1}{3} K^2) \\ & + \beta^i \partial_i K + 4\pi \alpha (\rho + S), \end{aligned} \quad (34)$$

$$\begin{aligned} \partial_t \tilde{A}_{ij} = & e^{-4\phi} [-D_i D_j \alpha + \alpha (R_{ij} - 8\pi S_{ij})]^{TF} \\ & + \alpha (K \tilde{A}_{ij} - 2 \tilde{A}_{ik} \tilde{A}^k_j) + \beta^k \partial_k \tilde{A}_{ij} \\ & + \tilde{A}_{ik} \partial_j \beta^k + \tilde{A}_{jk} \partial_i \beta^k - \frac{2}{3} \tilde{A}_{ij} \partial_k \beta^k, \end{aligned} \quad (35)$$

where the superscript  $TF$  denotes the trace-free part of a tensor. The Ricci tensor is also split into:

$$\begin{aligned} R_{ij} = & \bar{R}_{ij} - 2(\bar{D}_i \bar{D}_j \phi + \bar{\gamma}_{ij} \bar{\gamma}^{lm} \bar{D}_l \bar{D}_m \phi) \\ & + 4((\bar{D}_i \phi)(\bar{D}_j \phi) - \bar{\gamma}_{ij} \bar{\gamma}^{lm} (\bar{D}_l \phi)(\bar{D}_m \phi)) \\ \equiv & \bar{R}_{ij} + R_{ij}^\phi. \end{aligned} \quad (36)$$

The  $\bar{\Gamma}^i$  are now treated as independent functions that satisfy their own evolution equations:

$$\begin{aligned} \partial_t \bar{\Gamma}^i = & 2\alpha \left( \bar{\Gamma}^i_{jk} \bar{A}^{kj} - \frac{2}{3} \bar{\gamma}^{ij} \partial_j K - 8\pi \bar{\gamma}^{ij} S_j + 6 \bar{A}^{ij} \partial_j \phi \right) \\ & - 2 \bar{A}^{ij} \partial_j \alpha + \beta^j \partial_j \bar{\Gamma}^i - \bar{\Gamma}^j \partial_j \beta^i \\ & + \frac{2}{3} \bar{\Gamma}^i \partial_j \beta^j + \frac{1}{3} \bar{\gamma}^{il} \partial_l \partial_j \beta^j + \bar{\gamma}^{jl} \partial_j \partial_l \beta^i. \end{aligned} \quad (37)$$

We use the variable:

$$W = \gamma^{-1/6} = e^{-2\phi} \quad (38)$$

instead of  $\phi$ . The evolution equation for  $W$  is:

$$\partial_t W = \frac{1}{3} W (\alpha K - \partial_i \beta^i) + \beta^i \partial_i W. \quad (39)$$

The other choice, which evolves  $\phi$ , can lead to crashes due to numerical singularities.

We use McLachlan [13–15] and adopt the set of equations (33), (34), (35), (37), (39) for evolution and the gauge conditions (19), (20).

## IV. NUMERICAL SETUP

### A. Mesh Refinement

We employ mesh refinement [16] for the puncture center with 7 levels. The basic grid resolution is  $3.072M$ , and the refinement factor is set to 2. Therefore, the minimum resolution is  $3.072M \times 2^{-6} = 0.048M$ .

We utilize the EINSTEIN TOOLKIT [17] to simulate the evolution of a single puncture.

## V. RESULT

### A. Gravitational Waves

For wave extraction, we utilized the WEYLSCAL4 and MULTIPOLE thorns [18]. These thorns extract the Weyl scalar and decompose it into the spin-weighted spherical harmonics. To calculate the radiated energy, we considered modes up to  $l = 8$ , which are considered credible because higher modes are dominated by numerical noise [19].

Since the Schwarzschild metric has spherical symmetry, there should be no gravitational wave radiation. Figure 6 displays the real and imaginary parts of  $r_{\text{ex}}\psi_4$ . There is no notable difference between  $r_{\text{ex}} = 50M$ . Furthermore, all values of  $r_{\text{ex}}\psi_4$  are on the order of  $10^{-14}$ , which can be regarded as numerical noise.




FIG. 6. Real and imaginary parts of  $r_{\text{ex}}\psi_4$ . The red dashed line indicates  $r = r_{\text{ex}} = 50M$ .

### B. Apparent Horizon

We measured apparent horizon at each time using AHFINDERDIRECT [20, 21]. It finds an apparent horizon by numerically solving equation

$$\Theta \equiv D_i n^i + K_{ij} n^i n^j - K = 0, \quad (40)$$

where  $n^i$  is the outward-pointing unit normal to the apparent horizon, and  $D_i$  is the covariant derivative operator associated with the 3-metric in the slice.

We don't fixed puncture location at the origin, but there was no puncture moving and the puncture still at the origin. Fig. 7 shows expansion of apparent horizon as function of time. Fig. 8 shows irreducible mass as function of time.




FIG. 7. Expansion  $\Theta$  of apparent horizon as function of time.




FIG. 8. Irreducible mass  $M_{\text{irr}}$  as function of time.

- plement **90**, 1 (1987), <https://academic.oup.com/ptps/article-pdf/doi/10.1143/PTPS.90.1/5201911/90-1.pdf>.
- [2] M. Shibata and T. Nakamura, Evolution of three-dimensional gravitational waves: Harmonic slicing case, *Phys. Rev. D* **52**, 5428 (1995).
  - [3] T. W. Baumgarte and S. L. Shapiro, On the numerical integration of Einstein's field equations, *Phys. Rev. D* **59**, 024007 (1999), [arXiv:gr-qc/9810065](https://arxiv.org/abs/gr-qc/9810065).
  - [4] J. W. York, Gravitational degrees of freedom and the initial-value problem, *Phys. Rev. Lett.* **26**, 1656 (1971).
  - [5] A. Lichnerowicz, On completely harmonic Riemannian spaces, *Bull. Soc. Math. Fr.* **72**, 146 (1944).
  - [6] J. M. Bowen and J. York, James W., Time asymmetric initial data for black holes and black hole collisions, *Phys. Rev. D* **21**, 2047 (1980).
  - [7] B. Brügmann, J. A. González, M. Hannam, S. Husa, U. Sperhake, and W. Tichy, Calibration of moving puncture simulations, *Phys. Rev. D* **77**, 024027 (2008).
  - [8] M. Ansorg, B. Brügmann, and W. Tichy, A single-domain spectral method for black hole puncture data, *Phys. Rev. D* **70**, 064011 (2004), [arXiv:gr-qc/0404056](https://arxiv.org/abs/gr-qc/0404056).
  - [9] M. Campanelli, C. O. Lousto, P. Marronetti, and Y. Zlochower, Accurate evolutions of orbiting black-hole binaries without excision, *Phys. Rev. Lett.* **96**, 111101 (2006), [arXiv:gr-qc/0511048](https://arxiv.org/abs/gr-qc/0511048).
  - [10] M. Alcubierre, B. Brügmann, P. Diener, M. Koppitz, D. Pollney, E. Seidel, and R. Takahashi, Gauge conditions for long term numerical black hole evolutions without excision, *Phys. Rev. D* **67**, 084023 (2003), [arXiv:gr-qc/0206072](https://arxiv.org/abs/gr-qc/0206072).
  - [11] T. W. Baumgarte and S. G. Naculich, Analytical representation of a black hole puncture solution, *Phys. Rev. D* **75**, 067502 (2007).
  - [12] M. Hannam, S. Husa, F. Ohme, B. Brügmann, and N. Ó Murchadha, Wormholes and trumpets: Schwarzschild spacetime for the moving-puncture generation, *Phys. Rev. D* **78**, 064020 (2008).
  - [13] J. D. Brown, P. Diener, O. Sarbach, E. Schnetter, and M. Tiglio, Turduckening black holes: an analytical and computational study, *Phys. Rev. D* **79**, 044023 (2009), [arXiv:0809.3533 \[gr-qc\]](https://arxiv.org/abs/0809.3533).
  - [14] Kranc, Kranc: Kranc assembles numerical code.
  - [15] McLachlan, McLachlan, a public BSSN code.
  - [16] Carpet, Carpet: Adaptive Mesh Refinement for the Cactus Framework.
  - [17] L. Werneck, S. Cupp, T. Assumpção, S. R. Brandt, C.-H. Cheng, P. Diener, J. Doherty, Z. Etienne, R. Haas, T. P. Jacques, B. Karakaş, K. Topolski, B.-J. Tsao, M. Alcubierre, D. Alic, G. Allen, M. Ansorg, M. Babiuc-Hamilton, L. Baiotti, W. Bender, E. Bentivegna, S. Bernuzzi, T. Bode, G. Bozzola, B. Brendal, B. Bruegmann, M. Campanelli, F. Ciolletta, G. Corvino, R. D. Pietri, A. Dima, H. Dimmelmeier, R. Doolley, N. Dorband, M. Elley, Y. E. Khamra, J. Faber, G. Ficarra, T. Font, J. Friebe, B. Giacomazzo, T. Goodale, C. Gundlach, I. Hawke, S. Hawley, I. Hinder, E. A. Huerta, S. Husa, T. Ikeda, S. Iyer, L. Ji, D. Johnson, A. V. Joshi, H. Kalyanaraman, A. Kankani, W. Kastaun, T. Kellermann, A. Knapp, M. Koppitz, N. Kuo, P. Laguna, G. Lanferman, P. Lasky, L. Leung, F. Löffler, H. Macpherson, J. Masso, L. Menger, A. Merzky, J. M. Miller, M. Miller, P. Moesta, P. Montero, B. Mundim, P. Nelson, A. Nerozzi, S. C. Noble, C. Ott, L. J. Papenfort, R. Paruchuri, D. Pollney, D. Price, D. Radice, T. Radke, C. Reisswig, L. Rezzolla, C. B. Richards, D. Rideout, M. Ripeanu, L. Sala, J. A. Schewtschenko, E. Schnetter, B. Schutz, E. Seidel, E. Seidel, J. Shalf, K. Sible, U. Sperhake, N. Stergioulas, W.-M. Suen, B. Szilagyi, R. Takahashi, M. Thomas, J. Thornburg, C. Tian, M. Tobias, A. Tonita, S. Tootle, P. Walker, M.-B. Wan, B. Wardell, A. Wen, H. Witek, M. Zilhão, B. Zink, and Y. Zlochower, The einstein toolkit (2023), to find out more, visit <http://einstein toolkit.org>.
  - [18] J. G. Baker, M. Campanelli, and C. O. Lousto, The Lazarus project: A Pragmatic approach to binary black hole evolutions, *Phys. Rev. D* **65**, 044001 (2002), [arXiv:gr-qc/0104063](https://arxiv.org/abs/gr-qc/0104063).
  - [19] D. Pollney, C. Reisswig, E. Schnetter, N. Dorband, and P. Diener, High accuracy binary black hole simulations with an extended wave zone, *Phys. Rev. D* **83**, 044045 (2011), [arXiv:0910.3803 \[gr-qc\]](https://arxiv.org/abs/0910.3803).
  - [20] J. Thornburg, Finding apparent horizons in numerical relativity, *Phys. Rev. D* **54**, 4899 (1996), [arXiv:gr-qc/9508014](https://arxiv.org/abs/gr-qc/9508014).
  - [21] J. Thornburg, A Fast Apparent-Horizon Finder for 3-Dimensional Cartesian Grids in Numerical Relativity, *Class. Quantum Grav.* **21**, 743 (2004), [arXiv:gr-qc/0306056](https://arxiv.org/abs/gr-qc/0306056).