

# *Artemisia* terpene diversity across scales: biosynthetic evolution and bounded antiparasitic evidence from a living scoping review

## Abstract

The genus *Artemisia* contains a chemically diverse collection of volatile terpenes, nonvolatile terpenoids, and sesquiterpene lactones. This review develops a source-backed framework for comparing that chemistry across species while avoiding a common error: treating an essential-oil profile, a pathway gene, or an in-vitro parasite assay as a genus-wide pharmacological conclusion. Across 538 registered sources and 553 source-linked evidence records, we synthesize monoterpenes, sesquiterpenes, diterpenes, triterpenes, and sesquiterpene lactones; evaluate biosynthetic compartmentalization and gene-family evolution; and connect candidate pathway changes to chemotype and antiparasitic evidence. The best-resolved pathway is artemisinin biosynthesis in *A. annua*, but recent *A. argyi* results illustrate why gene presence/absence claims require assembly-, ploidy-, and annotation-aware reconciliation. Malaria and artemisinin form the primary validated case study, while wormwood (*A. absinthium*) provides a separate, preparation-specific veterinary vermifuge/anthelmintic case. Essential-oil, extract, whole-leaf, and purified-compound assays provide phenotype anchors across protozoa, helminths, and vectors, yet constituent attribution remains unresolved in most mixture studies. We propose a reproducible, phylogeny-aware research program combining voucher-linked metabolomics, secretory-tissue transcriptomics, comparative genomics, enzyme assays, and fractionated parasite testing.

Keywords: *Artemisia*; terpenes; sesquiterpene lactones; artemisinin; chemotype; comparative genomics; essential oil; antiparasitic activity

## 1. Introduction

*Artemisia* is a large genus with substantial ecological, morphological, ploidy, and phytochemical diversity. Its volatile chemistry is concentrated especially in leaves and flowers, but reported profiles vary with genotype, chemotype, locality, tissue, ontogeny, harvest, drying, and extraction. Consequently, “the terpene profile of *Artemisia*” is not a single biological object; the appropriate unit of comparison is a documented specimen or accession with analytical context.

This review has two aims. First, it organizes the major terpene classes reported from *Artemisia* and summarizes the biosynthetic logic connecting precursor supply, synthase diversification, oxidation, lactonization, storage, and regulation. Second, it evaluates whether comparative evolutionary genomics can connect that diversity to antiparasitic phenotypes without overstating causality.

The evidence extraction unit is a specimen, accession, or explicitly bounded experimental preparation. Representative records are provided in `chemotype-table.csv`; they are not species means. The review follows the protocol in `review-protocol.md` and the source identifiers in `sources.json`.

### 1.1 What previous reviews establish-and what remains open

Yes, *Artemisia* terpene and terpenoid reviews have been published. The principal precedents include a genus-level essential-oil review, a 2012-2017 essential-oil update, a dedicated sesquiterpene-lactone analysis and methods review, a review of edible species and selected sesquiterpene lactones, a 2024 genus-wide phytochemistry, pharmacology, and toxicology review, a 2024 synthesis of STL biosynthesis/regulation/genomics, a 2024 evolution-focused STL review, a recent review focused on *A. herba-alba* and *A. judaica* chemistry and omics, a 1990 review of *A. annua* constituents and antimalarial pharmacology, a 2022 review of *A. annua* artemisinin biotechnology, a 2023 malaria-focused review of isolated *A. annua* and *A. afra* metabolites and PK/PD, a 2025 medicinal-chemistry review of *Artemisia* antimalarials, and an August 2026 systematic review of whole-plant *A. annua* tea (Abad et al. 2012; Pandey and Singh 2017; Ivanescu et al. 2015; Trendafilova et al. 2021; Hussain et al. 2024; Frey et al. 2024; Agatha et al. 2024; Al-Khayri et al. 2022; Woerdenbag et al. 1990; Shinyuy et al. 2023; Diab et al. 2026; Rauf et al. 2025; Munyangi Wa Nkola et al. 2026). These reviews establish the breadth of reported chemistry and biological claims, but they generally do not provide one specimen-level, class-balanced, phylogeny-aware evidence model connecting terpene chemistry, evolutionary genomics, tissue context, antiparasitic stage, and safety. The present package is designed as an auditable update and integration, not as a claim that prior reviews were absent or noncomprehensive. Two adjacent reviews are particularly useful for scoping the parasite sections. Sharma’s 2025 review surveys *Artemisia*, artemisinin, semisynthetic derivatives, and derived oligomers across protozoal, helminth, bacterial, viral, and fungal contexts, while Jamil and colleagues’ 2025/2026 review situates *A. absinthium* within botanical anthelmintics for ruminants (Sharma 2025; Jamil et al. 2025/2026). We use both as discovery and framing sources. Their broad mechanism and efficacy summaries are not substituted for specimen-level chemistry, parasite-stage assays, host counter-screens, or primary clinical and veterinary evidence. A species-focused wormwood review provides a complementary synthesis of *A. absinthium* constituents, pharmacological reports, anthelmintic context, pharmacokinetics, and toxicity (Batiha et al. 2020). A veterinary network meta-analysis likewise places *A. absinthium* among botanical anthelmintic candidates, but its cross-plant and heterogeneous evidence base requires caution when interpreting any species-specific or terpene-specific effect (Calzetta et al. 2020). These reviews strengthen the rationale for the wormwood section while leaving preparation, constituent attribution, and human translation as primary-study questions.

## 1.2 Literature-search and evidence strategy

The literature review was organized around five linked evidence domains: (i) compound occurrence and analytical identity; (ii) enzyme, pathway, and compartment evidence; (iii) genotype, population, tissue, developmental, and environmental variation; (iv) comparative genomics and gene-family evolution; and (v) antiparasitic phenotype, target, and safety evidence. Searches combined *Artemisia* taxon terms with terpene-class, analytical, biosynthetic, genomic, and parasite terms. The reproducible search strings, dates, database results, de-duplication outputs, and title/abstract decisions are archived in `search-log.md`, `search-results.json`, and the screening CSV files; the PRISMA-ScR mapping is in `prisma-scr-checklist.md`. A source-linked full-text eligibility queue covers 110 unique antiparasitic source records representing 129 evidence rows, with malaria and *A. absinthium* vermifuge records prioritized. Full-text verification now covers 81 publicly accessible or publisher-full-text sources and 96 linked evidence rows; 29 queue sources remain unverified, and none is treated as quantitatively pooled until the full-text gate is complete. The working corpus contains both primary studies and prior reviews, but mechanistic and quantitative statements are preferentially anchored to primary studies. Because full-text screening remains incomplete, this report is a living scoping review and evidence map rather than a completed systematic review.

Evidence was graded by what an experiment actually established. Isolation with NMR or a matched analytical standard supports compound identity in the tested material; GC-MS library matching supports a lower-confidence assignment unless retention-index and standard data are also available. Heterologous enzyme assays support catalytic capacity under the assay conditions, not native flux. Transcript abundance and co-expression prioritize candidates but do not prove product identity. Parasite growth inhibition supports a preparation-specific phenotype; causal target claims require biochemical, structural, genetic, or activity-based target-engagement evidence. This hierarchy allows broad coverage without converting association into mechanism.

The review is intentionally class-balanced. Essential-oil surveys dominate the searchable literature, but the synthesis separately tracks nonvolatile sesquiterpene lactones, diterpenes, triterpenes, and cuticular metabolites so that analytical convenience does not become a biological conclusion. Likewise, *A. annua* is treated as the best-resolved reference rather than a proxy for the genus. Evidence from *A. argyi*, *A. absinthium*, *A. herba-alba*, *A. afra*, *A. campestris*, and other species is retained with its own voucher, population, tissue, and preparation boundaries wherever those data are reported.

## 2. Chemical scope and terminology

Terpenes are assembled from C5 isoprenoid units. Monoterpenes are principally C10, sesquiterpenes C15, diterpenes C20, and triterpenes C30; oxygenated and rearranged products are more precisely terpenoids. Sesquiterpene lactones are C15 terpenoids containing a lactone ring and are generally treated as extractable specialized metabolites rather than ordinary essential-oil constituents.

Representative volatile classes include pinene, limonene, myrcene, camphor, 1,8-cineole, borneol, thujones, germacrene D,  $\beta$ -caryophyllene,  $\alpha$ -humulene, davanone, spathulenol, and caryophyllene oxide. Important nonvolatile or less volatile chemistry includes artemisinin and related compounds, guaianolide and germacranolide sesquiterpene lactones, and triterpenoids associated with plant surfaces and defense.

### 2.1 Evidence map by class

Table 1. Major terpene classes, analytical contexts, and interpretation limits in *Artemisia*.

| Class                                                    | Representative <i>Artemisia</i> chemistry                                                                         | Main analytical context                                                   | Review interpretation                                                                      |
|----------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------|--------------------------------------------------------------------------------------------|
| Monoterpenes and oxygenated monoterpenes                 | $\alpha$ -/ $\beta$ -pinene, limonene, myrcene, camphor, 1,8-cineole, borneol, thujones, artemisia ketone         | GC-MS/GC-FID of volatile oils and headspace fractions                     | Often abundant, but highly sensitive to chemotype, tissue, stage, drying, and distillation |
| Sesquiterpene hydrocarbons and oxygenated sesquiterpenes | germacrene D, $\beta$ -caryophyllene, $\alpha$ -humulene, davanone, spathulenol, caryophyllene oxide              | GC-MS of oils; targeted volatile profiling                                | Useful markers of population chemistry, not universal species signatures                   |
| Sesquiterpene lactones                                   | artemisinin, arteannuin B, artemisinic acid, santonin, guaianolides, germacranolides, absinthin-related compounds | LC-MS/HPLC, NMR, isolation; GC is often unsuitable without derivatization | Usually nonvolatile; structural and stereochemical confirmation is critical                |

| Class                   | Representative <i>Artemisia</i> chemistry                                                  | Main analytical context                                                       | Review interpretation                                                                                                |
|-------------------------|--------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------|
| Diterpenes              | species-specific diterpenoid constituents and chlorophyll-associated phytol derivatives    | Solvent extraction, LC-MS, NMR; often reported in broad phytochemical surveys | Less comprehensively sampled than C10/C15 chemistry; presence should be linked to isolation or validated MS evidence |
| Triterpenes and sterols | $\beta$ -amyrin/ $\alpha$ -amyrin-, friedelin-, cycloartane- and sterol-associated reports | Solvent extraction, LC-MS, GC-MS after derivatization, NMR                    | Surface and storage chemistry; not part of ordinary essential-oil comparisons                                        |

The unevenness of this evidence map is itself a result: volatile mono- and sesquiterpenes are overrepresented because they are accessible to routine GC analysis, whereas nonvolatile diterpenes, triterpenes, and lactones require extraction workflows that are less directly comparable across studies.

Recent structure-confirmed work expands the under-sampled C10 and C15 space without supplying a shortcut to pharmacology. Aerial parts of *A. persica* yielded ten undescribed monoterpenoids, including four enantiomeric pairs, characterized by NMR, high-resolution mass spectrometry, and ECD; none exceeded 50% activity at the tested concentrations (Umaraliev et al. 2026). The same species also yielded eleven undescribed bisabolane-type sesquiterpenoids, with only limited anti-inflammatory activity reported for two compounds (Umaraliev et al. 2025). These records add authenticated chemical space and a useful negative/limited-activity control, but neither study establishes antiparasitic activity, native biosynthetic genes, or species-wide abundance. They support a core rule for the comparative dataset: a newly named terpene is an occurrence record until its pathway, ecological context, and parasite phenotype are independently linked.

Postharvest handling can create an apparent chemotype shift even when genotype is unchanged. Five weeks of extended natural shade drying reduced *A. monosperma* essential-oil yield by 23.7% and altered its constituent profile, although beta-pinene and selected major constituents were comparatively stable (Mohammed and Alfazi 2026). In three *A. argyi* cultivars, drying method also changed the volatile balance: 80 °C hot-air drying/fixation reduced oxygenated monoterpenes such as eucalyptol and thujone while increasing selected sesquiterpenes including caryophyllene and neointermedeol (Ming et al. 2025). Any parasite assay using dried plant material should therefore record drying duration, temperature, cultivar, oil yield, and full chromatographic composition before attributing activity to a terpene or comparing studies.

An additional volatile comparator is *A. serotina*, or winter wormwood. Its aerial-part oil was dominated by the irregular monoterpene santolina alcohol (34.6%) and included santolinane-, artemisane-, and lavandulane-type constituents; the oil showed fungicidal activity against *Candida albicans* in the reported assay (Kadyrbay et al. 2025). This record broadens the irregular-monoterpene space but does not establish malaria, helminth, or vermifuge activity. It also illustrates why common names such as “wormwood” should not be used as chemical surrogates for *A. absinthium* or for a particular thujone exposure.

Wormwood stereochemistry and postharvest history add two distinct sources of chemical uncertainty. VCD/DFT and NMR assigned the absolute configurations of two naturally occurring ocimene monoterpenoids from domesticated *A. absinthium* oils, converting tentative C10 identities into stereochemically resolved records (Julio et al. 2017). In a seasonal Iranian collection, 44 oil constituents and total yield changed across preflowering, flowering, and post-flowering stages, with shifts among alpha-phellandrene, beta-pinene, chamazulene, and related volatiles (Mohammadi et al. 2014). A separate storage model showed that drying and storage alter wormwood profiles through evaporation and parent-to-daughter chemical transformations (Blagojević et al. 2015). These findings make stereochemical confirmation, harvest stage, drying, and storage mandatory fields when comparing wormwood vermifuge preparations.

Primary studies show that the under-sampled classes are not merely theoretical. In *A. annua*, seed isolation recovered fourteen sesquiterpenes, three monoterpenes, and one diterpene (Brown et al. 2003). Separate trichome transcriptome and heterologous-expression work functionally characterized the oxidosqualene cyclase OSC2 and P450 CYP716A14v2 in aerial-cuticle triterpenoid production (Moses et al. 2015), while a diterpene-synthase study identified ten *A. annua* diTPS genes in a glandular-trichome and stress-resilience context (Chen et al. 2021). These are class-specific anchors, not evidence that all *Artemisia* species share the same diterpene or triterpene repertoire.

## 2.2 Monoterpenes and volatile sesquiterpenes

Chiral stereochemistry is an additional authentication variable for volatile antiparasitic candidates. Four *A. herba-alba* oils were modeled with infrared/VCD spectra and compared with chiral HPLC, resolving variable proportions and prevailing enantiomers of (-)-alpha-thujone, (+)-beta-thujone, and (-)-camphor (Said Mel et al. 2016). This approach provides a practical route to distinguish nominally identical GC-MS peaks that differ in absolute configuration. It is not evidence that *A. herba-alba* chemistry represents

*A. absinthium*, or that any one enantiomer is a vermifuge; it shows why chiral identity should accompany abundance, voucher, and parasite-stage metadata.

Monoterpenes are the most method-accessible *Artemisia* terpenoids. Across oils, recurrent chemistries include pinene and limonene hydrocarbons, camphor, borneol, 1,8-cineole, thujones, and irregular-monoterpene products such as artemisia ketone. Volatile sesquiterpenes add germacrene D,  $\beta$ -caryophyllene,  $\alpha$ -humulene, caryophyllene oxide, davanone, spathulenol, and related oxygenated products. These names should be treated as reported identities with confidence levels, not interchangeable biomarkers: retention-index agreement, authentic-standard co-injection, mass-spectrum library matching, and NMR confirmation provide different evidentiary strength.

The biological unit is often a glandular-trichome or aerial-tissue oil, not a purified constituent. In *A. annua*, three recombinant monoterpene synthases generated distinct product spectra, while wounding and hormone treatment changed transcript levels. This connects volatile chemistry to enzyme specificity and inducible regulation, but it does not show that the induced oil is antiparasitic. In *A. frigida* and *A. absinthium*, population, geography, and phenology alter the relative abundance of volatile compounds. A cross-species comparison should therefore model presence/absence, relative abundance, and analytical detectability separately.

A dedicated review of *A. annua* essential oil is useful as a calibration source because it documents both the breadth and the comparability problem. Reported major-constituent maxima included camphor up to 48%, germacrene D up to 18.9%, artemisia ketone up to 68%, and 1,8-cineole up to 51.5%; the review also summarizes antibacterial and antifungal assays performed with non-equivalent methods and organisms (Bilia et al. 2014). These values are literature maxima rather than species means, and the review does not establish parasite activity. Its strongest contribution to the present synthesis is methodological: oil composition, extraction, assay organism, endpoint, and concentration must remain linked before a volatile profile is compared with malaria or wormwood evidence.

Enzyme studies reveal why one-gene/one-compound annotation is often inadequate. Recombinant *A. annua* epi-cedrol synthase produced epi-cedrol as its major oxygenated C15 product but also generated several olefinic sesquiterpenes and showed lower activity with the C10 precursor GPP (Mercke et al. 1999). Two additional *A. annua* sesquiterpene-cyclase cDNAs and later transient-expression experiments expanded the functionally tested synthase repertoire (Van Geldre et al. 2000; Kanagarajan et al. 2012). Product multiplicity means that an expanded TPS clade can increase chemical opportunity without predicting a unique metabolite, while heterologous expression can reveal catalytic potential that may be constrained by native substrate supply or cell type.

The underrepresented irregular-monoterpene space is also phylogenetically informative. Structure-confirmed isolation from *A. tridentata* subsp. *vaseyana*, *A. cana* subsp. *viscidula*, and *A. tridentata* subsp. *spiciformis* yielded 26 regular and irregular monoterpenes, including artemisia triene, trans-chrysanthemal, lavandulol, and a lavandulyl-skeletal compound reported as new to the Anthemideae family (Gunawardena et al. 2002). The result shows that C10 diversity is not exhausted by routine pinene/camphor/thujone profiling, but isolated occurrence in selected western taxa does not establish abundance, pathway orthology, ecological function, or antiparasitic activity. It is a useful target for phylogeny-stratified metabolomics and enzyme assays.

Functional divergence can occur even among close homologs. An AaADS-like enzyme from *A. maritima* cyclized FPP to 4-amorphen-11-ol rather than amorph-4,11-diene, and the product was recovered from engineered yeast (Muangphrom et al. 2018). This is a useful evolutionary experiment because it separates sequence homology from product identity: an ADS-like copy may preserve the substrate class while changing product outcome. The result is heterologous catalytic evidence, not proof of native abundance, artemisinin production, or parasite activity.

Downstream enzymes further diversify volatile products. A glandular-trichome-specific *A. annua* alcohol dehydrogenase oxidized monoterpene alcohol substrates in recombinant assays, demonstrating that volatile profiles reflect not only cyclization but subsequent redox chemistry (Polichuk et al. 2010). At the regulatory level, AabHLH112 perturbation changed  $\beta$ -caryophyllene, epi-cedrol, and  $\beta$ -farnesene-associated sesquiterpene metabolism, connecting a transcription factor to several non-artemisinin branches rather than a single endpoint (Xiang et al. 2022). Conversely, suppression of  $\beta$ -caryophyllene synthase increased artemisinin-pathway output in transgenic plants, providing direct evidence that FPP-consuming branches can compete for precursor under at least some engineered conditions (Chen et al. 2011).

Population studies show an orthogonal source of complexity. Oils of *A. campestris* and *A. herba-alba* varied among natural populations, and their non-parasitic bioassay profiles varied with composition (Younsi et al. 2017). In *A. herba-alba*, analysis of 80 individuals from eight Tunisian populations resolved four oil chemotypes and found substantial within-population marker diversity; global chemical divergence was not reducible to geographic or bioclimatic distance alone (Younsi et al. 2018). A controlled comparison of five Iranian species further illustrates the scale and limits of cross-species variation: 20-34 compounds were assigned by GC-MS per species, artemisinin ranged from 0.12 to 0.41 mg/g dry weight, and dominant volatiles differed markedly, including camphor-rich *A. fragrans*, beta-thujone-rich *A. absinthium*, and beta-farnesene-rich *A. biennis* (Jamshidi et al. 2024). The same study found no uniform correlation between artemisinin content and the relative expression of the surveyed pathway genes. Because the plants were grown under a common controlled regime and GC-MS assignments were not all confirmed by isolation, this is an accession- and method-bounded comparison rather than a species-wide chemotype or evolutionary gain/loss test.

These observations argue for individual- or population-level sampling and against using a single oil chromatogram as a species diagnostic.

Spatial analysis adds an ecological-scale counterpart to accession-level comparisons. A province-level analysis of Chinese *A. annua* reported positive spatial autocorrelation of artemisinin content, with high-value clusters in southwestern regions and sunshine, temperature, and precipitation identified as major explanatory factors (Zhang et al. 2017). Because the record is an observational spatial analysis with region-level aggregation, it cannot separate genotype, seed source, agronomy, processing, and microclimate or establish a causal environmental response. It nonetheless supports explicitly modeling geography and climate when testing whether pathway-gene variation predicts malaria-relevant chemistry.

A fifteen-population study of *A. vulgaris* makes the safety implication concrete: davanone, germacrene D, 1,8-cineole, camphor, thujone, and cis-chrysanthenyl acetate each dominated different Lithuanian oils, and brine-shrimp toxicity varied across the profiles (Judzentiene and Garjonyte 2016). The result supports chemotype-stratified toxicology, but brine-shrimp LC0 values are not mammalian therapeutic windows and do not identify a single toxic constituent. Similarly, oral *A. argyi* oil produced rapid rat absorption and distribution of eight volatile constituents, including eucalyptol, alpha-thujone, camphor, borneol, bornyl acetate, beta-caryophyllene, and caryophyllene oxide, with prominent tissue exposure followed by prompt elimination (Hou et al. 2021). This is disposition evidence for a defined oil preparation, not proof of antiparasitic efficacy or a safe human dose.

Newer wormwood analyses reinforce the need to treat geography, phenology, and authentication as causal design variables rather than metadata. *A. absinthium* oils shifted among oxygenated monoterpene-, oxygenated sesquiterpene-, and monoterpene-hydrocarbon-dominated profiles across sites and flowering stages (Malaspina et al. 2025). A separate low-thujone oil was dominated by trans-sabinyl acetate, with trans-thujone only 1.2 +/- 0.5% in the reported sample (Judzientienė and Būdienė 2026). Conversely, authenticated and commercial oils formed distinct thujone/camphor/ocimene-epoxide clusters, and alpha- and beta-thujone produced tephritid behavioral or toxicity endpoints (Tabanca et al. 2025). These records support lot-specific chemical authentication and safety testing; they do not permit the species name “wormwood” to stand in for a defined thujone exposure or a vermifuge dose.

An additional Indian *A. absinthium* leaf oil shows how strongly the dominant class can change with provenance. The Western Ghats sample contained 64 identified constituents and was rich in oxygenated monoterpenes (39.7–41.1%), with borneol, methyl hinokiate, isobornyl acetate, beta-gurjunene, and caryophyllene oxide among the major compounds (Joshi 2013). Its tested phenotype was antibacterial and antifungal, not an anthelmintic assay. The record therefore strengthens the chemical case for regional chemotypes while weakening any attempt to infer vermifuge activity from the common name, traditional use, or a single dominant volatile.

An agricultural-invertebrate assay adds a useful but deliberately separate wormwood comparator. A Moroccan *A. absinthium* oil was rich in 3,3,5-trimethylcyclohexene and camphor and produced stage-dependent toxicity against the potato tuber moth *Phthorimaea operculella* (Mahboub et al. 2025). This supports a lot-specific volatile profile and an insect-control phenotype, but it is not evidence against helminths and does not identify a causal constituent; the result should not be used to strengthen a wormwood vermifuge claim.

Extraction technology is another source of apparent chemotype variation. In flowering *A. annua* aerial material, supercritical CO2 extraction yielded 0.62–1.92% and produced artemisia ketone, camphor, 1,8-cineole, arteannuin B, and beta-selinene profiles that differed from hydrodistilled and steam-distilled oils (Mandić et al. 2025). This provides a controlled process comparison across mono- and sesquiterpene fractions, but it does not identify the best therapeutic preparation, establish antimalarial activity, or make the resulting profiles comparable to a species-wide oil mean.

Recent comparative assays extend the same logic from chemotype to arthropod control. Five *Artemisia* oils contained only 9–24 identified constituents per species and separated into distinct multivariate profiles; *A. ordosica* and *A. tangutica* were the strongest contact-toxic oils against the two tested stored-product insects (Zhang et al. 2022). In natural *A. scoparia* populations, altitude was associated with wide ranges of beta-myrcene, gamma-terpinene, and trans-beta-caryophyllene and with differences in mosquito repellency and larvicidal activity (Parveen et al. 2024). A camphor-rich *A. vulgaris* oil produced time- and concentration-dependent fumigant, contact, and repellent activity against the gall mite *Aceria pongamiae* (Kunnathattil et al. 2025). These records support environment- and species-stratified vector experiments, but the oils remain mixtures and the reported insect or mite endpoints cannot be used as proxies for helminth or malaria efficacy.

Full-text extraction makes the chemotype/phenotype relationship more quantitative. Six-hour hydrodistillation of authenticated aerial material from five Chinese species produced oils with 0.02–0.53% yields and markedly different profiles: *A. tanacetifolia* was rich in 3-carene (45.98%), *A. sieversiana* in neryl propanoate (22.88%), *A. dalai-lamae* in cineol (32.62%), and *A. ordosica* in caryophyllene (17.81%), with PCA explaining 62.45% of variance. *A. ordosica* had the lowest reported *Liposcelis bostrychophila* contact-toxicity LD50 (52.11 microgram/cm2), whereas *A. tangutica* was strongest against *Tribolium castaneum* (17.42 microgram/adult); low-dose *A. tangutica* and *A. sieversiana* could be attractive rather than repellent. In five Pakistani *A. scoparia* populations, steam-distilled oils varied from 0.15–0.57% yield, capillene from 9.6–31.8%, and beta-myrcene from 6.8–21.5%; the high-altitude Swat oil retained repellency against *Aedes aegypti* for up to 165 minutes at the higher tested dose, while the Attock

oil had larvicidal LC50 values of 55.5 and 43.5 microgram/mL at 24 and 48 hours. These are replicated, GC-MS-linked mixture results, not single-terpene causal tests (Zhang et al. 2022; Parveen et al. 2024).

The *A. vulgaris* mite study adds a complementary route-specific control. A Kerala oil obtained by four-hour hydrodistillation yielded 0.86 +/- 0.14% and was reported as 45.20% oxygenated monoterpenes, 21.29% monoterpene hydrocarbons, and 21.88% aromatics, with camphor at 28.94%, alpha-pinene at 6.61%, eucalyptol at 6.39%, fenchol at 6.03%, and a substantial non-terpene 4-tert-butylaniline assignment. Against *Aceria pongamiae*, fumigant LC50 fell from 1.29 to 0.43 microliter/mL air and contact LC50 from 37.37 to 4.56 microliter/mL between 24 and 72 hours; repellency reached 86.11% at 72 hours. The reported non-terpene peak and library-based chemical identification reinforce the need for orthogonal standards and non-target counterscreens before attributing activity to camphor or another terpene (Kunnathattil et al. 2025).

The chemical reference space is still expanding through structure-confirmed isolation. *A. tournefortiana* aerial parts yielded two novel sesquiterpenes and one new monoterpene, with NMR, HRESIMS, and calculated ECD used to assign structures and antibacterial testing showing only moderate activity for two compounds (Wu et al. 2024). A recent *A. argyi* leaf-oil study was dominated by 1,8-cineole, beta-caryophyllene, and longifolene and reported insecticidal activity against *Callosobruchus maculatus* at high oil concentration (Kačániová et al. 2026). These studies add authenticated C10/C15 occurrence and a contemporary *A. argyi* chemotype, but neither identifies a native pathway allele, a purified active, or a parasite target.

An earlier process benchmark similarly extracted artemisinin and artemisinic acid from six *A. annua* samples using supercritical CO2 with 3% methanol at 50 degrees C and 15 MPa, with rapid recovery and no reported analyte decomposition under the tested conditions (Kohler et al. 1997). This is useful for reproducing analyte recovery and comparing extraction selectivity, but process yield remains distinct from native pathway flux, biological activity, and clinical suitability.

Older population and phenology studies remain valuable controls for the apparent stability of an oil profile. In the narrow endemic *A. molinieri*, 69 aerial-part volatiles were identified; ascaridole ranged from 19–76%, alpha-terpinene from trace levels to 36%, and germacrene D from 0.6–15% across season, phenological stage, and plant age (Masotti et al. 2003). A geographically matched *A. absinthium* comparison likewise separated French oils dominated by (Z)-epoxyocimene and chrysanthenyl acetate from Croatian oils dominated by (Z)-epoxyocimene and beta-thujone, with additional changes before and after anthesis (Juteau et al. 2003). The reported yeast assays in these studies are not parasite endpoints. Their main contribution is design logic: “wormwood oil” must be tied to geography, stage, organ, and a complete chromatographic fingerprint before a vermifuge experiment can be compared across lots.

### 2.3 Diterpenes and triterpenes

Diterpene evidence is comparatively sparse and often mixed into broad phytochemical surveys. The *A. annua* seed study provides isolation-level evidence for one diterpene among three monoterpenes and fourteen sesquiterpenes, while a dedicated *A. annua* study identified ten diterpene synthases in relation to glandular trichome formation, artemisinin production, and stress resilience. A multi-platform *A. absinthium* oil analysis additionally reported diterpene- and homoditerpene-associated constituents alongside GC and NMR confirmation (Judzentiene et al. 2009). These findings support a real C20 biosynthetic dimension but not a genus-wide diterpene chemotype. Diterpene records should preserve seed versus leaf or trichome tissue, extraction polarity, and whether the compound was isolated or only assigned by MS.

Triterpenes and sterols are especially important for avoiding a false “essential-oil-only” view of *Artemisia*. OSC2 and CYP716A14v2 were functionally connected to  $\alpha$ -,  $\beta$ -, and  $\delta$ -amyrin-related cuticular triterpenoids in *A. annua*, and triterpenoids have also been isolated from *A. igniaria*. Their cuticle or surface-associated localization suggests roles in barrier and stress biology, but current records do not support transferring those roles to antiparasitic action. Because triterpenes are less volatile, a GC-only survey can systematically under-report them.

The current diterpene and triterpene literature remains too sparse and methodologically heterogeneous for a genus-wide abundance synthesis. A defensible survey should distinguish a structurally isolated C20 or C30 metabolite from a tentative LC-MS feature, and it should separate pathway-gene discovery from measured products. The *A. annua* diterpene-synthase study is unusually informative because it combined gene discovery with developmental and physiological perturbation, whereas the OSC2/CYP716A14v2 work paired trichome comparisons with recombinant functional tests. These studies provide reference loci for comparative genomics, but their products must be re-measured in homologous tissues before copy-number differences are interpreted as ecological or antiparasitic adaptations.

A compound-resolved infectious comparator reinforces the need to retain chemical structure and assay context. Bio-guided fractionation of field wormwood, *A. campestris* subsp. *glutinosa*, isolated the pseudoguaianolide sesquiterpene lactone damsine and the steroidal triterpene lactone canrenone. In MT-2-cell HIV-1 assays, reported IC50 values were 0.84 micromolar for damsine and 18.87 micromolar for canrenone, with different cytotoxicity boundaries (Apaza-Ticona et al. 2020). This is stronger constituent attribution than an unfractionated oil assay and broadens the authenticated STL/triterpene space, but it is antiviral rather than antiparasitic evidence; the cellular mechanism cannot be relabeled as a parasite-protein interaction or transferred to *A. absinthium*.

Targeted primary isolations confirm that the sparse C20/C30 literature reflects under-sampling rather than an absence of chemistry. *A. sacrorum* yielded an ent-kauranoid diterpene diglycoside and two known ent-kauranoids; *A. caruifolia* yielded a new triterpene with co-isolated lignans; and whole-plant *A. sphaerocephala* yielded a new oleanane-type triterpenoid saponin (Li et al. 1990; Ma et al. 2001; Li et al. 2008). Petroleum-ether fractionation of *A. argyi* further identified nordammarane and cycloartane triterpenoids, while *A. vulgaris* yielded both new sesquiterpenoids and triterpenoids with structure confirmation (Zhang et al. 2020; Liu et al. 2022). These records establish chemical space and analytical identity; their cytotoxic, alpha-glucosidase, or anti-inflammatory endpoints do not establish antiparasitic activity. A genome-wide *A. annua* OSC survey identified 24 genes across CAS, LAS, LUS, and unknown subfamilies and reported differential expression under UV-B, but its bioinformatic and expression evidence still requires product-level and recombinant-enzyme validation (Guo et al. 2025).

An enzyme-resolved *A. argyi* example shows why C30 evolution should be studied at the active-site level. AaOSC-22030 converted 2,3-oxidosqualene into diverse triterpene skeletons; multiscale QM/MM analysis and mutagenesis implicated S728 in relaxed intermediate accommodation and catalysis, whereas S728Y made dammarendiol II the exclusive reported product and the N369/E371 dyad controlled terminal hydration of pentacyclic products (Liu et al. 2026). This is strong sequence-to-function evidence for catalytic plasticity in one recombinant enzyme, but engineered specificity and computationally resolved transition-state hypotheses do not establish native abundance, lineage-wide selection, or antiparasitic activity. It supplies a concrete comparative test: map active-site residues onto common-annotation OSC gene trees, then verify product spectra in matched tissues.

The class-balanced tranche adds further structure-confirmed evidence. Aerial parts of *A. suksdorfii* yielded five new polyol monoterpenes and seven new sesquiterpene lactones, with high-field NMR and X-ray support for one lactone (Ahmed et al. 2004). Across *A. giraldii* var. *giraldii*, *A. mongolica*, and *A. vestita*, isolation work recovered a new antifungal monoterpene, a new germacranolide, alpha-amyrin, and sterol-associated constituents (Tan et al. 1999). In *A. igniaria*, four triterpenoids were structurally assigned from an extract, including beta-glutanol as a first natural occurrence report (Hu and Feng 2000). These studies materially widen the C10/C15/C30 inventory, but they remain specimen- and extraction-bounded: they do not supply genus-wide abundance estimates, shared pathway orthology, or antiparasitic efficacy. The antifungal results are retained as bioactivity context rather than recoded as malaria or helminth evidence.

An earlier cross-species phytochemical study makes the same class boundary explicit. *A. sieversiana* aerial parts yielded a new guaianolide, while antifungal fractions of *A. annua* contained arteannuin B, artemisinin, oleanolic acid, beta-sitosterol, stigmasterol, and several flavones; arteannuin B was active against the tested *Candida* strain at 100 microgram/mL, whereas most other isolates were inactive (Tang et al. 2000). This is structure- and activity-resolved evidence for C15 and C30 chemical space, but the fungal endpoint does not establish malaria, helminth efficacy, or a shared *Artemisia* pathway. It also illustrates why non-terpene co-isolates and sterols should remain visible in the evidence table rather than being forced into a terpene-only interpretation.

Historical records add depth to the C10/C15 literature while also showing where the evidence is thin. An indexed *A. filifolia* study explicitly addressed new monoterpene structures, synthesis, rearrangements, and biosynthesis, providing a foundational C10 reference for structure-function comparison, although detailed product extraction still requires full-text recovery (Torrance and Steelink 1974). Seasonal sampling is represented by a primary *A. feddei* study of sesquiterpene-lactone constitution in material collected in Kochi prefecture; because its abstract is unavailable in the current retrieval, it is retained as a phenology lead rather than a quantitative seasonal result (Matsueda et al. 1973). A separate *A. feddei* isolation study reported five sesquiterpene lactones, including two new guaianolides, from aerial parts (Tan et al. 1992). *A. sieversiana* aerial parts likewise yielded one known and three new guaianolides alongside lignans and sterols, with extensive spectral characterization; antifungal testing in that study is not an antiparasitic endpoint (Tan et al. 1998). Together these records support a broad structural and phenological search space, but the title-level and isolation-level evidence should not be converted into native pathway evolution, abundance, or malaria/vermifuge claims.

Structure-confirmed isolation from *A. sieberi* provides another class-balanced STL anchor. Two new sesquiterpene lactones and two known compounds were assigned by NMR, mass spectrometry, and X-ray diffraction; the tested activities were antimicrobial or antifungal rather than antiparasitic (Mohamed et al. 2017). The study therefore expands authenticated C15 lactone space and supplies a candidate set for future parasite assays, but it does not support a claim that these compounds are vermifuges or that their antimicrobial endpoints predict mammalian selectivity.

An earlier *A. annua* isolation study adds a distinct C30 anchor: friedelin and friedelan-3-beta-ol were isolated alongside artemisinin-related terpenoids, and friedelane-type triterpenoids were reported from the plant for the first time (Zheng 1994). The associated endpoint was cytotoxicity in human tumor cell lines, so this record expands nonvolatile triterpene coverage without supporting parasite activity or a safe therapeutic window. In a separate *A. herba-alba* fractionation study, a new monoterpene dimer, a known monoterpene, and three sesquiterpene lactones were characterized by multidimensional NMR and HRMS (Abou El-Hamd et al. 2013). The two records together illustrate why class-balanced sampling must include both structure-confirmed nonvolatile constituents and unusual terpene architectures, not only routine essential-oil percentages.

Sesquiterpene engineering supplies a complementary functional anchor. Coexpression of valencene synthase and valencene oxidase in *A. annua* produced nootkatone, and plastid-targeted constructs reached 12.11-47.80 microgram/g fresh weight without changing artemisinin production (Gou et al. 2021). The result demonstrates that substrate compartment and enzyme choice can

redirect C15 flux in an engineered plant, but it is not evidence that nootkatone is a native *A. annua* chemotype or an antiparasitic constituent.

The first committed artemisinin-pathway step has also been demonstrated with unusually direct sequence-to-product evidence. An amorpho-4,11-diene synthase cDNA from *A. annua* produced amorpho-4,11-diene from FPP in recombinant *E. coli*, and transformed tobacco accumulated the product at 0.2–1.7 ng/g fresh weight (Wallaart et al. 2001). This anchors catalytic capacity and heterologous plant production, but it does not establish native flux, tissue localization, or conservation of the exact product across *Artemisia*. It is therefore a useful positive control for comparative ADS-like sequence-function analysis rather than a genus-wide artemisinin marker.

## 2.4 Sesquiterpene lactones

Structure-confirmed dimeric guaianolides extend the nonvolatile wormwood inventory beyond the monomeric lactones usually emphasized in pharmacognosy. Aerial parts of *A. absinthium* yielded five new absinthins (A–E) and seven known dimeric guaianolides; absinthin C and isoabsinthin inhibited LPS-induced nitric-oxide production in BV-2 cells at 1.52 and 1.98  $\mu\text{M}$ , respectively (Turak et al. 2014). The NMR, HRESIMS, and X-ray evidence supports authenticated C15-dimer chemical space, while the immune-cell endpoint remains distinct from helminth or malaria activity. These molecules are therefore priority standards for fractionated *A. absinthium* parasite assays, not evidence that “absinthin” is the active vermifuge principle.

Isolation from in-vitro shoot culture expands the non-artemisinin lactone reference set. *A. aucheri* yielded the eudesmanolide artemin (2,5-dihydroxy-12,6-eudesmanolide-4(15)-ene), reported from this plant for the first time; artemin showed micromolar cytotoxicity in prostate and breast cancer cell lines (Abbaspour et al. 2019). This is isolation-level sesquiterpene-lactone evidence, not antiparasitic efficacy, and the tissue-culture origin means native abundance and ecological function remain untested.

Structure-confirmed isolations widen the STL repertoire beyond the artemisinin branch. *A. anomala* aerial parts yielded a dimeric guaianolide, a glaucolide, a seco-guaiaretic acid, additional guaianolides, and known STLs (Wen et al. 2010). *A. dubia* yielded eight new artemdubolides and nineteen known STLs, while aerial parts of *A. myriantha* yielded ten guaianolides including artemyriantholide E, arglabin, dehydroleucodin, and related structures (Huang et al. 2010; Zan et al. 2018). In *A. argyi*, HPLC activity profiling followed by ECD/VCD and NMR resolved the absolute configuration of a diverse STL set, and a separate leaf study added new guaianolide and eudesmane structures (Reinhardt et al. 2019; Zhang et al. 2021). These records establish specimen-bounded chemical diversity and the need for stereochemical authentication; cytotoxic or immune-cell endpoints in several of them are not antiparasitic evidence. They also provide a tractable comparative resource: common-annotation gene trees can now be paired with authenticated STL profiles and recombinant oxidase/lactonization assays rather than inferred from compound names alone. The structure space continues into dimeric guaianolides and absinthin-related chemistry. *A. mongolica* aerial parts yielded seven undescribed and seventeen known sesquiterpenes, including artemongolide dimers, artemanomalides, absinthin-related compounds, and other guaianolides; NMR, HRESIMS, ECD, and first X-ray determinations supported the assignments (Zhu et al. 2024). A *A. gorgonum* study used santonin-derived photochemistry to prepare and revise seven seco-guaianolide/guaianolide structures, with X-ray-supported absolute configurations; the strongest reported endpoint was phytotoxicity in wheat coleoptiles rather than parasite activity (Macías et al. 2012). These studies are valuable for stereochemical and defense-chemistry coverage, but they should not be used to infer anthelmintic or antimalarial action. They also reinforce a practical design rule: dimeric and highly oxygenated STL features should be searched by LC-MS/MS and structure-guided isolation, not assumed from volatile-oil fingerprints.

A newer structure-confirmed tranche shows that this dimeric and oxygenated space is still expanding. *A. myriantha* yielded 19 new sesquiterpenolides spanning 13 germacranolides, four guaianolides, and two eudesmanolides, including uncommon cis-fused 10/5 systems and an unusual (7S) configuration (Tang et al. 2020). *A. dubia* yielded 20 previously undescribed guaiane-type dimers: A–Q were proposed as [4+2] Diels-Alder adducts of monomeric guaianolides, whereas R–T were ester-linked (Gao et al. 2022). *A. princeps* added 13 new dimeric sesquiterpenoids, also assigned a Diels-Alder origin, with X-ray/ECD-supported configurations (Su et al. 2023). In *A. porrecta*, 27 STLs covered eudesmanolide, germacranolide, melampolide, and heliangolide classes, while *A. myriantha* var. *pleiocephala* supplied eight new guaiane-type sesquiterpenoids plus 14 known compounds (Abdullajanov et al. 2025; Wang et al. 2025). The associated endpoints were cytotoxicity, anti-inflammatory activity, or structural elucidation—not malaria or helminth assays. These records therefore expand the candidate library and suggest experimentally testable assembly hypotheses, but dimer assignment, native biosynthetic provenance, parasite activity, and host selectivity still require LC-MS/MS, NMR/X-ray confirmation, pathway-enzyme assays, and matched parasite testing.

The next queue tranche further argues against treating named STL families as a closed inventory. *A. sacrorum* yielded 23 highly oxygenated guaiane-type lactones, including acetoxyated  $\alpha$ -methylene- $\gamma$ -lactones and 1,10-rearranged structures; whole-plant *A. mongolica* yielded 12 new and seven known STLs with multiple X-ray assignments (He et al. 2024; Zhu et al. 2022). *A. leucophylla* added 12 new and 13 known eudesmane-type sesquiterpenoids, including unprecedented 1,2-seco-1-nor-eudesmane skeletons, and *A. heptapotamica* supplied both monomeric and dimeric lactones (Wang et al. 2023; Zhamilya et al. 2019). Dimeric chemistry was also reported from *A. selengensis* through proposed [2+2] and [4+2] couplings, from *A. myriantha* through additional Diels-Alder-like dimers, and from *A. myriantha* var. *pleiocephala* through a second variety-specific series (Chen et al. 2025; Wang et al. 2025; Wang et al. 2024). Finally, *A. ordosica* yielded cadinane-monoterpene heterodimers and unusual sesquiterpenoids, while *A. annua* yielded 14 cadinane-involved dimers spanning several coupling modes (Wang et al. 2025; He et al. 2024).



These findings are valuable for comparative chemical evolution and target-library design, but their tumor-cell, inflammatory-cell, or computational endpoints do not establish malaria, helminth, or other parasite efficacy. A publishable synthesis should therefore report them as authenticated chemical-space expansions and explicitly preserve the gap between structural novelty, native pathway reconstruction, and antiparasitic mechanism.

Sesquiterpene lactones (STLs) include guaianolide, germacranolide, eudesmanolide, and related structural families. Their  $\alpha$ -methylene- $\gamma$ -lactone or other electrophilic motifs can react with nucleophiles, which provides a chemically plausible basis for bioactivity but not a demonstrated parasite target. In *Artemisia*, STL reports include artemisinin-related amorphane chemistry, santonin, and numerous guaianolide and germacranolide constituents. Analytical workflows differ from oil profiling: LC-MS, isolation, multidimensional NMR, and stereochemical assignment are often needed, while direct GC percentages are not a suitable universal denominator.

Wormwood provides an unusually useful analytical reference for this class. A validated HPLC-DAD method separated and quantified five sesquiterpene lactones, two lignans, and a polymethoxylated flavonoid in *A. absinthium* material and commercial preparations; HPLC-MS and HPLC-SPE-NMR also characterized absinthin degradation products and showed strong stability under several storage conditions (Aberham et al. 2010). This is important for the vermifuge question because measured lactone exposure depends on extraction and storage, but the analytical record does not by itself establish anthelmintic activity or a safe dose.

Santonin adds a historically important but safety-limited branch to the wormwood question. A validated HPLC-UV method measured santonin in *A. cina* extracts and leaves from eight Kazakhstan *Artemisia* taxa, including *A. absinthium*, providing a practical lot-specific control for a compound formerly used as an anthelmintic (Sakipova et al. 2017). In a separate *A. cina* fractionation study, high-speed countercurrent chromatography removed santonin from a CO<sub>2</sub> subcritical extract while the santonin-free preparation retained antinociceptive and anti-inflammatory activity (Sakipova et al. 2020). Because the latter endpoints were not parasite assays, these studies do not prove that santonin or santonin-free extract is a vermifuge. They do show why historical anthelmintic reputation, measured santonin exposure, preparation-level activity, and human safety must be analyzed as separate variables.

A particularly relevant attribution lead comes from a parasite-oriented analytical study of *A. absinthium*. HPLC peak deconvolution and reconstructed spectra tentatively identified the sesquiterpene lactones alpha-santonin and ketopelenolid-A in methanolic wormwood extracts; the authors presented alpha-santonin as a possible active principle based on its established antiparasitic reputation (Perez-Souto et al. 1992). This links a named wormwood preparation to named lactone candidates, but the identification was tentative and the record does not provide a matched purified-compound assay, direct parasite-protein experiment, or standardized extract exposure. The result therefore motivates targeted LC-MS/MS confirmation and fractionation rather than proving that alpha-santonin explains any modern wormwood phenotype. Historical indexing also records treatment of ascariasis with santonin prepared from *A. transiliensis* (Guseinov 1954). Because that record is title- and metadata-level in the current retrieval, it documents historical vermifuge context only; it cannot support a dose, efficacy magnitude, product-quality, or current clinical claim.

The malaria literature has been repeatedly synthesized, but review recurrence should not be mistaken for new primary evidence. A foundational endoperoxide review documented the early chemical and antimalarial development of artemisinin, while a recent review revisited artemisinin chemistry, extraction, photochemistry, mechanisms, and translational applications across *Artemisia* sources (Ziffer et al. 1997; Das et al. 2024). These reviews justify the historical and mechanistic framing of the malaria section; the present synthesis retains primary plant, parasite-stage, pharmacokinetic, and safety records as the basis for quantitative or clinical statements.

Historical pharmacognosy also links santonin to an *A. maritima* source in India (Chopra and Chandler 1924). Because the indexed record is historical and not a modern standardized composition or controlled parasite study, it strengthens provenance for santonin's vermifuge reputation only. It cannot establish that contemporary *A. absinthium* preparations contain an equivalent dose, retain equivalent activity, or have an acceptable safety margin.

The best-defined biosynthetic model for a non-artemisinin STL branch begins with FPP cyclization to germacrene A, oxidation to germacrene A acid, and downstream P450-dependent oxidation and lactonization. The presence of a germacranolide-like metabolite is therefore compatible with several levels of evidence-from structural analogy to a fully demonstrated enzyme sequence-and those levels must not be collapsed. STL abundance should also be reported separately from volatile sesquiterpene abundance because extraction and storage can partition them differently. In *A. annua*, field measurements of glandular-trichome dynamics across leaf development and senescence, together with dead-versus-green leaf precursor profiles, show why tissue age and precursor conversion must be retained as explicit variables (Lommen et al. 2006; Lommen et al. 2007). Enzyme evolution adds a testable explanation for structural diversification: a germacrene-A oxidase accepted all seven alternate sesquiterpene substrates tested in yeast, whereas amorphadiene oxidase was comparatively rigid; engineered active-site substitutions increased artemisinic-acid activity per enzyme but reduced stability (Nguyen et al. 2019). This supports catalytic promiscuity as an evolutionary opportunity, not a claim that every GAO homolog makes a native lactone. A 2024 functional reanalysis adds an important correction to the AMO/GAO boundary. One *A. annua* CYP71AV1-like copy, termed AMOHAP, had catalytic-region features and a divergent transcript context more consistent with germacrene-A oxidase; recombinant protein assays demonstrated GAO activity rather than a canonical amorphadiene-oxidase interpretation (Chen et al. 2024). This result supports a pathway-evolution model in which a new

artemisinin branch coexists with, or replaces, an ancestral costunolide-related branch through both catalytic specialization and expression-level divergence. It also supplies a concrete comparative test: common-annotation CYP71/GAO trees should be paired with cis-regulatory, tissue-expression, and matched germacrene-A/amorphadiene enzyme assays before assigning orthology or pathway completion in other *Artemisia* species.

Pathway organization is another experimentally testable layer between gene inventory and metabolite abundance. A Molecular Plant report explicitly investigated promoting artemisinin biosynthesis in *A. annua* by substrate channeling, but the current indexed retrieval lacks an abstract; it is therefore retained as a full-text-recovery lead rather than assigned a quantitative flux or compartment claim (Han et al. 2016). In an open-access transformation study, ADS plus the viral antisilencing factor P19 increased measured artemisinin to 0.18% in transformed leaves and 0.095% in transformed hairy roots, whereas control hairy roots had no detected artemisinin (Elfahmi et al. 2020). These engineered outcomes demonstrate production potential in *A. annua*, not native cross-species conservation or parasite efficacy. Environment can further separate transcript abundance from terminal product: dynamic cold-stress profiling found transient jasmonate-associated induction and coordinated PAL/C4H/ADS/DBR2 responses, while prolonged exposure increased dihydroartemisinic and artemisinic acids as terminal conversion was inhibited (He et al. 2024). This duration dependence is a useful design variable for comparative experiments and a reason not to encode “stress-induced artemisinin” as a single invariant trait.

A complementary combinatorial-engineering study showed that AaDBR2 and a *Tanacetum parthenium* CYP71CB1 could act on multiple sesquiterpene-lactone pathway substrates, enabling heterologous production of dihydrocostunolide, dihydroparthenolide, and hydroxylated derivatives (Beyraghdar Kashkooli et al. 2019). Such catalytic plasticity offers a plausible route by which duplicated or recruited downstream enzymes can enlarge lactone chemical space, but it also warns against inferring a unique product from a gene-family label: substrate access, stereochemistry, compartment, and downstream turnover must be experimentally matched.

Evolutionary sequence-function work provides a complementary explanation for terpene diversity outside the lactone branch. By recombining naturally occurring residues between cyclic ADS and linear beta-farnesene synthase, Salmon et al. identified a small epistatic network in which one dominant substitution could activate cyclization and additional residues could suppress or restore it; the corresponding network also affected a *Citrus* homolog (Salmon et al. 2015). This is direct evidence that a few interacting amino-acid changes can alter TPS product topology in a heterologous assay. It motivates gene-tree- and structure-aware tests across *Artemisia*, but does not by itself identify ancestral states, native selection, or a parasite-relevant product.

Recent structure-confirmed isolations show that the genus contains substantially more STL architecture than the artemisinin branch alone suggests. *A. vulgaris* yielded twelve new and ten known guaianolides, with NMR, electronic-circular-dichroism, and crystallographic support for stereochemistry; *A. verlotorum* yielded seven new and nineteen known sesquiterpenoids, including uncommon bicyclic, iphionane, and oxygen-bridged eudesmanolide structures (Chen et al. 2022; Wang et al. 2023). Basin big sagebrush leaf fractionation and *A. nitrosa* isolation add geographically distinct, structure-confirmed lactone records, including a new germacrene-type lactone and known sesquiterpene series (Anibogwu et al. 2024; Kazymbetova et al. 2022). The associated macrophage or cytotoxicity assays are useful for structure-activity context, but they are not antiparasitic evidence. These studies strengthen the evolutionary hypothesis that repeated germacrene, guaiane, and eudesmane diversification creates a large chemical search space; demonstrating lineage-specific pathway evolution still requires common annotations, gene trees, expression, and direct enzyme assays.

The artemisinin branch demonstrates how metabolite isolation, enzyme biochemistry, and nonenzymatic chemistry can be integrated. Dihydroartemisinic acid and its hydroperoxide were isolated as candidate late intermediates before genetic perturbation established a broader branch architecture (Wallaart et al. 1999a; Wallaart et al. 1999b). CYP71AV1 disruption subsequently redirected metabolites toward arteannuin X and supported a major role for spontaneous oxidative chemistry in terminal artemisinin formation. The evidence therefore supports a hybrid pathway model: enzyme-controlled scaffold formation and oxidation establish the precursor pool, while storage environment and post-biosynthetic oxidation help determine terminal products.

The chemical activation step has an independent physical-chemistry basis. EPR experiments detected carbon-centered radicals after iron-catalyzed decomposition of artemisinin and related endoperoxides, supporting a reactive-intermediate model for the peroxide bridge (Butler et al. 1998). In a parasite assay, extracellular heme at pathologic concentrations enhanced artemisinin activity against *P. falciparum* and increased lipid-peroxidation products; vitamin E reduced the enhancement, whereas the same heme manipulation did not potentiate quinoline drugs (Tangnitipong et al. 2012). These results connect iron/heme context to activation and phenotype, but neither identifies one necessary parasite receptor nor validates an unstandardized *Artemisia* preparation. The appropriate comparative-genomics question is therefore whether a plant accession changes the amount and exposure of an activatable peroxide-bearing metabolite, not whether an ADS-like gene alone predicts parasite killing.

Full-text review of the J-STAGE study makes the heme result more quantitatively explicit. Synchronized 10–15-hour ring-stage *P. falciparum* TM267R parasites were exposed to artemisinin with 0.1 or 1 micromolar heme for 24 hours. Artemisinin IC<sub>50</sub> fell from 7.0 +/- 1.1 nM without heme to 3.1 +/- 0.3 and 1.9 +/- 0.5 nM, respectively; severe-malaria plasma heme was 3.5 +/- 0.7 micromolar versus 0.3 +/- 0.2 micromolar in normal subjects (six per group). Heme increased TBARs in parasite media and parasitized erythrocytes, and vitamin E attenuated the artemisinin-associated oxidative signal, while heme did not potentiate

chloroquine, quinine, or mefloquine. This is a controlled purified-drug activation/redox experiment—not an infected-host intervention, a botanical-exposure study, or proof that extracellular heme is the dominant in-vivo activator (Tangnitipong et al. 2012).

Non-artemisinin lactones broaden both structural and pharmacological space. *A. gorgonum* yielded isolated guaianolide and germacranolide compounds with antiplasmodial testing, while *Artemisia*-derived or *Artemisia*-associated lactones such as dehydroleucodine have been tested against trypanosomatids. Dehydroleucodine inhibited cultured *T. cruzi* epimastigote growth and altered parasite physiology in a defined-compound study (Breggio et al. 2000). Ten additional germacrane-type sesquiterpenoids, artemyrianosins A-J, were isolated and structurally characterized from aerial parts of *A. myriantha*; selected compounds showed 43.7-89.3 microM cytotoxicity against human hepatoma cell lines (Zhang et al. 2022). That study expands confirmed STL structural space, but its endpoint is anticancer cytotoxicity and should not be recoded as antiparasitic activity. Such studies are stronger for constituent attribution than an oil assay, but species provenance, parasite stage, exposure, and host-cell selectivity still prevent direct translation to therapeutic efficacy.

### 3. Biosynthesis and compartmentalization

Production is also shaped by nutrient state and cultivation context. In cloned greenhouse plants, potassium deficiency increased artemisinin concentration from 0.93% to 1.62%, but total per-plant yield was not significantly different from complete nutrition, illustrating the difference between concentration and biomass-normalized output (Ferreira 2007). In a fed-batch hairy-root bioreactor, model-based sucrose feeding reached 1.0 mg/g artemisinin after 16 days, showing process control rather than native field flux (Patra and Srivastava 2015). AMF responses were similarly conditional: *Rhizophagus irregularis* increased artemisinin and essential-oil measures in semi-open soils, but open-field artemisinin differences were not detected and beta-farnesene/germacrene D shifted; a separate phosphate gradient found an inverse artemisinin response and lower leaf artemisinin with *Funneliformis mosseae* (Domokos et al. 2020; Todeschini et al. 2022). These studies make nutrient, symbiont, soil, and cultivation regime explicit covariates for comparative chemotype experiments.

Common-environment and plant-microbe experiments help separate genetic, ecological, and physiological components of this variation. In a Korean panel, 103 individuals grown under the same conditions formed six oil chemotype labels (artemisia-ketone, camphor, beta-cubebene, eucalyptol, alpha-pinene, and beta-selinene), demonstrating substantial intraspecific variation beyond immediate growth conditions (Hong et al. 2023). In contrast, AM colonization of *A. annua* did not change total terpene emission, but increased limonene and artemisia-ketone emission and altered selected leaf sesquiterpenes, emphasizing that volatile release and tissue accumulation are related but non-identical phenotypes (Rapparini et al. 2008). Geographic HPLC-DAD and LC-QTOF-MS/MS profiling likewise separated regional groups by artemisinin, arteannuin B, and artemisinic acid content (Fu et al. 2020). Together, these results justify common-environment designs followed by genotype, expression, and metabolite linkage rather than treating a location-level oil profile as a fixed species trait.

The plastidial MEP pathway and cytosolic mevalonate pathway supply IPP and DMAPP, which are condensed into prenyl diphosphates. GPPS and related enzymes provide C10 substrates for monoterpene synthases; FPPS provides the C15 substrate FPP for sesquiterpene synthases, including germacrene A synthase and ADS. Cytochrome P450s, reductases, dehydrogenases, oxidases, acyltransferases, glycosyltransferases, and spontaneous reactions expand the scaffold space.

Functionally characterized *A. annua* monoterpene synthases illustrate why copy number alone is insufficient: recombinant AaTPS2, AaTPS5, and AaTPS6 produced different product spectra, and transcript abundance changed after wounding and hormone treatments (Ruan et al. 2016). Product specificity, inducibility, and tissue localization therefore need to be modeled together in evolutionary comparisons.

The artemisinin pathway in *A. annua* is the strongest gene-to-metabolite reference. ADS converts FPP to amorphadiene, a foundational enzyme result established in early biochemical work (Bouwmeester et al. 1999); CYP71AV1 oxidizes pathway intermediates; DBR2 and ALDH1 support the route toward dihydroartemisinic acid, after which terminal chemistry includes nonenzymatic steps. Glandular secretory trichomes provide a relevant biosynthetic and storage context. These statements are pathway anchors, not proof that homologous genes in another species produce artemisinin or confer antiparasitic activity.

Sequence-function experiments provide the bridge from comparative gene trees to testable catalytic hypotheses. A structure-based combinatorial protein-engineering workflow developed active-site and focused libraries for plant sesquiterpene synthases, using defined residue combinations to expose epistasis and alternative product spectra (Dokarry et al. 2012). Applied to *Artemisia* TPS and oxidase clades, this approach can distinguish a lineage-specific sequence change that alters product topology from a copy-number difference that merely expands catalytic opportunity. It should be paired with authentic-standard product detection, tissue expression, and substrate-matched assays; library activity alone cannot establish native evolution, in-planta flux, or parasite relevance.

Quantitative analytical validation helps connect pathway genes to measured chemistry. Rapid chloroform immersion followed by LC-ESI-Q-TOF-MS/MS recovered more than 97% of artemisinin, arteannuin B, artemisitene, and artemisinic acid from fresh *A. annua* material across a 0.1–3.0 microgram/mL calibration range (Van Nieuwerburgh et al. 2006). The authors used the recovery pattern to support localization in the subcuticular gland space, but analytical recovery is not itself proof of complete tissue local-

ization or in-vivo pathway flux. In a historical vegetation-period study, artemisinin was highest immediately before flowering, one growth regulator increased content by approximately 30%, and most tested non-*annua* material lacked technically high concentrations (Liersch et al. 1986). These records support a stage- and method-aware sampling design rather than a universal preflowering rule.

Regulatory studies add a causal layer between stress signaling, trichome development, and pathway output. AaWRKY1 activated *ADS* and *CYP* promoter reporters, and overexpression increased pathway-gene expression and artemisinin in *A. annua* (Jiang et al. 2016). AaNAC1 was induced by dehydration, cold, salicylic acid, and methyl jasmonate; its overexpression increased artemisinin and dihydroartemisinic acid while improving drought and *Botrytis* tolerance (Lv et al. 2016). At the post-translational level, AaPP2C1 physically interacted with and dephosphorylated AaAPK1, reducing AabZIP1-mediated activation of artemisinin-biosynthesis genes (Zhang et al. 2019). Finally, exogenous beta-ocimene increased juvenile-plant artemisinin together with glandular-secretory-trichome size and density (Xiao et al. 2020). These are strong *A. annua* regulatory anchors, but overexpression, reporter, and elicitation results do not establish conserved function in other *Artemisia* lineages or increased parasite potency.

Developmental timing and culture conditions provide important negative and process controls. Constitutive *fpf1* expression advanced flowering in transgenic *A. annua* by about 20 days, yet artemisinin content did not differ significantly from the comparison plants, arguing against flowering time as a sufficient proxy for pathway output (Wang et al. 2004). In transformed hairy roots, autoclaving and the choice of filter-sterilized sugar changed biomass and artemisinin yield even at constant total carbon; glucose stimulated artemisinin production while sucrose supported better growth than glucose (Weathers et al. 2004). These studies strengthen the genotype-by-development-by-culture model and caution against interpreting a production increase as a conserved regulatory mechanism or as evidence of greater parasite activity.

Selection and engineering provide complementary tests of heritable versus inducible production. Four recurrent-selection cycles increased *A. annua* artemisinin from 0.15% to 1.16% and identified a molecular marker associated with high-yielding genotypes, linking standing genetic variation to a reproducible chemotype shift (Paul et al. 2010). In a separate transgenic experiment, co-overexpression of *CYP71AV1* and its reductase partner *CPR* increased artemisinin by up to 38% in one line (Shen et al. 2012). These results establish genetic and enzymatic leverage within *A. annua*, but neither identifies a conserved cross-species mechanism nor shows that a higher plant titer improves parasite selectivity or clinical efficacy.

Trichome regulators and plant-microbe interactions add another layer. The trichome-specific AP2/ERF factor AaORA positively regulated *ADS*, *CYP71AV1*, and *DBR2*; perturbation changed artemisinin and dihydroartemisinic acid, and disease-response phenotypes were also reported (Lu et al. 2013). Co-cultivation of *A. annua* shoots with *Piriformospora indica* increased biomass and artemisinin from 0.80% in controls to 1.30% in one treatment (Sharma and Agrawal 2013). These studies support a regulatory and symbiotic context for pathway output, while retaining the necessary boundary between production enhancement, native ecology, and antiparasitic phenotype.

The pathway was assembled through convergent evidence rather than a single experiment. *ADS* cloning and bacterial expression established the first committed cyclization; a trichome-enriched cDNA strategy identified *CYP71AV1* and demonstrated sequential oxidation of amorpha-4,11-diene and related intermediates; and *DBR2* cloning showed reduction at the branch that favors dihydroartemisinic-acid-derived chemistry (Chang et al. 2000; Teoh et al. 2006; Zhang et al. 2008). Stable-isotope feeding in intact growing plants supplied an independent carbon-flow perspective (Schramek et al. 2010). Together, these data are considerably stronger than pathway reconstruction from annotation alone, yet even this reference pathway remains sensitive to developmental state, precursor competition, and terminal nonenzymatic conversion.

The early pathway evidence also illustrates the value of preserving intermediate-level observations. Trichome and leaf analyses detected oxygenated amorpha-diene derivatives, while measured amorpha-4,11-diene hydroxylase, alcohol/dehydroaldehyde dehydrogenase, and related activities supported a proposed sequence toward dihydroartemisinic acid (Bertea et al. 2005). Recombinant *CYP71AV1* independently oxidized amorpha-4,11-diene, artemisinic alcohol, and artemisinic aldehyde (Teoh et al. 2006). The combined record is strong step-level evidence in *A. annua*, but a pathway diagram should still distinguish detected intermediates, measured activities, inferred order, and demonstrated in-vivo flux.

A 2025 enzyme study adds a distinct branch to the production model. AaDHAADH catalyzed bidirectional conversion between artemisinic acid and dihydroartemisinic acid; a P26L variant increased activity toward dihydroartemisinic acid production, and engineered yeast reached 3.97 g/L DHAA in a 5-L bioreactor (Guo et al. 2025). This is strong enzyme and metabolic-engineering evidence, but it does not demonstrate that AaDHAADH controls native *A. annua* flux in planta or that homologs in other *Artemisia* species have the same directionality or physiological importance. A production-perturbation study provides an additional tissue-level control: foliar application of the strigolactone analog GR24 increased growth, glandular-trichome attributes, and artemisinin content and yield in the tested *A. annua* plants (Wani et al. 2022). This is useful evidence that hormone signaling can alter the secretory capacity associated with a sesquiterpene-lactone phenotype, but it is not a conserved cross-species mechanism, a field recommendation, or evidence that increased artemisinin production necessarily increases parasite potency.

Chemotype regulation and precursor competition provide complementary constraints on pathway interpretation. In high- versus low-artemisinin *A. annua* chemotypes, *DBR2L* promoter variants, *DBR2L* expression, sesquiterpene profiles, and recombinant

reduction activity were associated with different artemisinin yields (Wang et al. 2023). Separately, loss of an AaBPS-associated locus reduced camphor and other monoterpenes while increasing FPP and selected sesquiterpenes, but not artemisinin (Olofsson et al. 2022). Together these observations argue that chemotype evolution can act through both downstream branch control and precursor allocation; neither study alone establishes a universal natural tradeoff or a parasite mechanism.

Compartment evidence is equally important. Immunolocalization placed key artemisinin enzymes in apical cells of glandular secretory trichomes, and a trichome transcriptome enriched the candidate space for isoprenoid and downstream-modification genes (Olsson et al. 2009; Wang et al. 2009). Tissue-expression studies showed that terpene-metabolism genes are not uniformly expressed across roots, stems, leaves, and reproductive structures (Olofsson et al. 2011). A grafting study further implicated root-derived regulation of shoot trichomes and artemisinin-related metabolites, indicating that the producing cell is embedded in a whole-plant signaling system (Wang et al. 2016).

Transport and developmental regulation add two distinct layers to that compartment model. AaPDR3 was localized mainly to T-shaped trichomes of old leaves and roots; RNAi reduced beta-caryophyllene, overexpression increased it, and transformed yeast accumulated beta-caryophyllene preferentially over germacrene D and beta-farnesene (Fu et al. 2017). In parallel, AaMIXTA1 perturbation changed glandular-trichome initiation, cuticle-related transcription, and artemisinin content (Shi et al. 2018). These studies show that a metabolite phenotype can depend on compound-selective transport and secretory-cell development, not only on synthase copy number. They also provide experimentally tractable cross-species tests: orthologous transporter activity, trichome-cell expression, and matched leaf-surface metabolomics.

Promoter-reporter evidence sharpens this compartment model. A 1,151-bp *CYP71AV1* promoter drove GUS expression in glandular secretory trichomes of young leaves and flower buds, shifted toward T-shaped trichomes in older leaves, and differed among *A. annua* varieties; methyl jasmonate increased reporter activity (Wang et al. 2013). This supports age-, tissue-, and genotype-dependent regulation, but reporter localization is not a measurement of native protein abundance, pathway flux, or conserved promoter function in another species.

These tissue and redox relationships were synthesized early in the artemisinin literature, which emphasized the coupled roles of trichomes, roots, reactive oxygen species, and regulatory enzymes (Nguyen et al. 2011; Wen and Yu 2011). The reviews remain useful for organizing hypotheses, but pathway-step and quantitative claims are anchored here to primary enzyme, perturbation, and isotope-tracing studies.

Precursor supply and branch competition have been tested genetically. Bidirectional perturbation of AaHDR1 altered artemisinin and other terpenoids, supporting a role for MEP-pathway supply while also showing class-specific responses rather than a uniform increase (Ma et al. 2017). Suppression of  $\beta$ -caryophyllene synthase shifted carbon away from one sesquiterpene branch, and a T-DNA insertion line produced a broader change in terpenoid composition and crude-extract bioactivity (Chen et al. 2011; Karaket et al. 2014). These perturbations support network competition, but engineered responses should not be assumed to describe standing natural variation.

Regulation is multilayered. Jasmonate-responsive AaERF1/AaERF2, glandular-trichome-specific WRKY1, AaMYB108, and the AaGSW1-AaYABBY5/AaWRKY9 complex have each been connected to artemisinin-pathway expression or metabolite accumulation through transgenic, interaction, and expression experiments (Yu et al. 2012; Chen et al. 2017; Liu et al. 2023; Kayani et al. 2023). Cold response adds another regulatory layer: AaHLH112 is induced at low temperature, is enriched in glandular secretory trichomes, and promotes artemisinin biosynthesis indirectly by binding the AaERF1 promoter and activating the associated pathway program (Xiang et al. 2019). The emerging model is therefore not a simple linear pathway but a tissue-localized, environment-responsive network in which developmental programs and stress signals act on precursor supply, synthase branches, oxidation, chromatin state, and trichome capacity. The uncaptured production and stress literature makes the environmental term more quantitative. Hairy-root lines selected from hundreds of transformants differed in growth and artemisinin accumulation, showing that culture context and clonal background can generate large production differences even within one taxon (Liu et al. 1998). In field-grown *A. annua*, plant density changed the relationship between artemisinin and essential-oil yield under a defined North-Indian site and season (Ram et al. 1997). Soil copper produced a nonlinear, genotype-dependent response: a low treatment increased artemisinin content whereas higher doses reduced accumulation and impaired growth (Zehra et al. 2020). Hydrogen sulfide partially rescued copper-associated injury while increasing glandular-trichome traits and artemisinin in the tested material (Nomani et al. 2022). These studies support genotype-by-environment and canopy-by-tissue interaction terms, but they are not evidence that stress manipulation improves parasite control or that a production phenotype is conserved across *Artemisia*.

An earlier Ri-plasmid study independently showed that transformed *A. annua* roots produced artemisinin and that pH, sucrose, auxin, and gibberellin altered growth or product accumulation (Cai et al. 1995). In a complementary stress experiment, a nitric-oxide donor improved photosynthetic and redox traits, trichome attributes, and artemisinin under cadmium exposure (Wani et al. 2023). Both studies strengthen the case for culture- and environment-aware sampling, but neither establishes native aerial chemotype evolution, a field intervention, or antiparasitic efficacy. Defense and post-transcriptional regulation provide complementary causal anchors. AaWRKY17 bound W-box motifs in the *ADS* promoter, increased artemisinin in transgenic *A. annua*, and reduced susceptibility to *Pseudomonas syringae* (Chen et al. 2021). A small-RNA, degradome, and perturbation study found that miR160 suppressed glandular-trichome formation and artemisinin biosynthesis, whereas ARF1 activated *AaDBR2* (Guo et

al. 2023). Together these results connect defense signaling, trichome development, and a pathway branch point, while leaving control of non-artemisinin terpenes, natural selection, cross-species orthology, and antiparasitic activity unresolved. An allele-aware *A. annua* perspective is therefore useful as a genome-interpretation frame, not as a substitute for primary abundance or enzyme evidence (Xu et al. 2022). An additional jasmonate perturbation adds a negative-regulatory node: AabHLH5 and two JAZ proteins were identified using transformation, yeast two-hybrid, and bimolecular-fluorescence assays, and AabHLH5 was verified as a negative regulator of jasmonate-linked artemisinin biosynthesis (Xu et al. 2025). In parallel, an *A. argyi* hairy-root system and recombinant assay showed that AYFOMT2 methylates jaceosidin to eupatilin and that transformation increases eupatilin accumulation (Peng et al. 2025). The AYFOMT2 result is a non-terpene branch control, but it demonstrates the kind of tissue transformation and step-level enzyme validation that comparative terpene studies need. Neither study supports transfer of regulator or enzyme function across species by homology alone.

The regulatory network has continued to expand in a direction that is relevant to comparative experiments. AaMYC2-type directly bound E-boxes in the *AaADS* and *AaCYP71AV1* promoters, while co-expression of AaMYC2 with AaMYC2-LIKE increased trichome density and artemisinin more than either factor alone (Khan et al. 2024; Hurrah et al. 2025). AaSHI1 activated the same core pathway promoters in transient assays, AaWRKY6 bound the *AaDBR2* promoter during growth-stage experiments, and AaABI5 connected light and ABA signaling to multiple pathway genes (Yang et al. 2024; Wang et al. 2023; Li et al. 2023). AabZIP1 adds ABA-JA cross-talk through AaMYC2 and also affects cuticular-wax genes under drought (Shu et al. 2022). These studies establish a dense, experimentally tractable *A. annua* network, but their engineered increases in artemisinin are not evidence that the same regulators control volatile terpenes, are conserved in other *Artemisia*, or improve parasite outcomes.

Post-translational and catabolic controls further complicate the genotype-to-metabolite map. AaAPK1, an ABA-responsive SnRK2-family kinase, interacted with and phosphorylated AabZIP1; the phosphorylated regulator increased transactivation of artemisinin-pathway genes, and AaAPK1 overexpression increased artemisinin in the reported experiments (Zhang et al. 2018). A class III peroxidase, Aa547, showed modeled affinity for artemisinin and deoxyartemisinin, localized to peroxisomes, and tracked inversely with artemisinin but directly with lignin in silencing and overexpression experiments (Nair et al. 2019). The kinase result supports a regulatory node and the peroxidase result a candidate turnover branch, but neither permits direct transfer to a second species without phosphorylation, enzyme-product, compartment, and flux measurements.

UV-B provides a related example of coordinated but not necessarily synergistic regulation. A leaf transcriptome identified 10,705 differentially expressed genes and 533 transcription factors after UV-B; reporter assays implicated AaMYB4 in activating *AaADS*, *AaDBR2*, and a flavonoid-pathway gene (Li et al. 2021). Parallel transcriptional regulation can explain correlated accumulation without proving that flavonoids potentiate artemisinin at the parasite. Comparative studies should therefore measure both transcript and chemical interaction terms, including matched parasite exposure, rather than treating co-induction as pharmacological synergy. Production studies add two important causal boundaries. Co-transformation of *HMGR*, *FPS*, and *DBR2* with the trichome regulators *AaHD1* and *AaORA* increased artemisinin up to 3.2-fold in T0 plants and up to 2.2-fold in selected T1 progeny, showing that pathway and compartment engineering can be combined while progeny stability remains variable (Hassani et al. 2023). Separately, endophytes isolated from environmentally distinct plants differed in their microbial conversion of artemisinin: *Colletotrichum* converted the peroxide bridge to an ether, whereas an environment-dependent isolate did not (Maehara et al. 2023). These results support a plant-microbe and engineering layer in the model, but neither proves that endophytes control native plant flux or that engineered/formulated material has improved antiparasitic efficacy.

Protein-level validation remains uneven. A CYP71AV1 expression and purification study established a recombinant heme-protein preparation and preliminary crystallization conditions for the enzyme that oxidizes amorph-4,11-diene toward artemisinic acid (Shi et al. 2018). This is useful structural-method evidence, but a preliminary crystallization condition is not a solved structure, and protein production alone does not establish native activity in another *Artemisia* species. The comparison should retain protein purification, product formation, and pathway flux as separate evidence fields.

Chromatin accessibility provides a complementary tissue-level mechanism. ATAC-seq comparison of glandular trichomes and leaves identified thousands of differentially accessible regions, with greater accessibility near DBR2 and CYP71AV1 in glandular trichomes and enrichment of MYB, bZIP, C2H2, and AP2-like regulatory motifs (Zhou et al. 2021). These data support an epigenomic hypothesis for pathway compartmentalization, but accessibility is associative: causal chromatin perturbation, protein abundance, flux, and cross-species conservation remain to be established.

Regulatory perturbations also connect artemisinin chemistry to broader plant signaling. Salicylic-acid treatment produced an early reactive-oxygen burst and increased artemisinin, artemisinic acid, and dihydroartemisinic acid while changing HMGR and ADS responses more than FDS or CYP71AV1 expression (Pu et al. 2009). RNAi reduction of cinnamate-4-hydroxylase similarly accumulated trans-cinnamic acid and salicylic acid and increased artemisinin while reducing phenylpropanoid outputs (Kumar et al. 2016). These experiments support pathway cross-talk and redox-sensitive terminal chemistry, not a direct parasite mechanism or a field-production prescription.

Germacranolide biosynthesis offers a second comparative framework: FPP can be cyclized by germacrene A synthase, oxidized by germacrene A oxidase, and converted through P450-dependent chemistry and lactonization. The broad presence of a scaffold family does not imply identical product profiles because enzyme specificity, tissue localization, substrate competition, and down-

stream modification differ.


In *A. annua*, a glandular-trichome cDNA library yielded a germacrene A synthase that converted FPP to germacrene A in heterologous assays (Bertea et al. 2006). This creates an experimentally anchored comparator for germacranolide-like pathways and illustrates a general strategy for other species: first establish synthase product, then test downstream oxidases, tissue expression, and matching metabolites. Sequence similarity to germacrene synthase, by itself, is insufficient because closely related TPS proteins can differ in product spectra and because downstream enzymes may redirect a shared scaffold.

A historical biochemical anchor sits downstream of this framework: a purified *A. annua* leaf protein converted arteannuin B to artemisinin with 58% conversion under the reported assay conditions, with a  $K_m$  of 0.5 mM and a pH optimum near 7.0–7.2 (Dhingra and Narasu 2001). This supports a transformation step at the protein-assay level, but the paper does not resolve a modern gene identity, native in-plant flux, subcellular transport, or conservation of the activity in other *Artemisia* species.

Chemical structure work provides an earlier precursor anchor: artemisinic acid was isolated from *A. annua* hexane extract and characterized by X-ray crystallography as a possible biogenetic precursor of artemisinin (Misra et al. 1993). Isolation and structural plausibility are important evidence for pathway substrate space, but they are not equivalent to isotope-traced conversion, enzyme assignment, or malaria activity.

Pathway evidence boundary. The ordered pathway model is a set of evidence-qualified hypotheses. Direct enzyme characterization supports individual steps; a homologous sequence or transcript supports candidate function; a detected metabolite supports presence in the sampled material. None of these alone establishes flux, compartment, or in-vivo production in a second species. Multi-substrate requirements, physiological direction, stereochemistry, and missing transport or cofactor information must remain explicit in the Terpedia pathway records.

Figure 1. Conceptual route from precursor supply to phenotype.



#### 4. Species and chemotype diversity

*A. annua* combines artemisinin-pathway chemistry with variable volatile oils that may include artemisia ketone, camphor, germa-crene D, and 1,8-cineole. *A. absinthium* can show thujone-, camphor-, cineole-, davanone-, or chamazulene-associated profiles depending on provenance and stage. *A. herba-alba*, *A. vulgaris*, and *A. argyi* likewise show regional and tissue-level variation. Multi-species Canadian oils reinforce that composition can differ substantially among taxa and regions (Lopes-Lutz et al. 2008). *A. dracuncululus* is a useful boundary case because estragole may dominate some oils; estragole is not a terpene and should remain in a broader volatile-metabolite record.

Preparation can redistribute the same plant's volatile evidence. In Serbian sweet wormwood, the essential oil was dominated by artemisia ketone and trans-caryophyllene, whereas the hydrosol concentrated the more water-soluble camphor, 1,8-cineole, and artemisia ketone; a synthesis of 61 accessions placed *A. annua* oils into four broad chemotype groups (Aćimović et al. 2022). This is a practical warning for malaria and wormwood comparisons: oil, hydrosol, tea, alcoholic extract, and dried leaf are different exposure matrices, not interchangeable labels for the same terpene dose.

Geography and climate can alter both the relative abundance and the dominant chemical class. Phenological shifts may change monoterpene and sesquiterpene proportions between vegetative and flowering stages. Drying and distillation can remove, transform, or concentrate constituents. These effects explain why percentage tables cannot be pooled without specimen-level metadata



and analytical harmonization.

A literature-derived analysis of Chinese *A. annua* records associated artemisinin content with latitude and climate, with temperature, sunshine duration, and rainfall identified as major ecological covariates and higher reported content toward southern/subtropical regions (Zhang et al. 2011). This strengthens climate and provenance as pre-specified covariates, but the result is an ecological association assembled from heterogeneous literature, not a causal experiment or a universal environmental optimum.

Developmental chemistry is not restricted to wormwood. In davana (*A. pallens*), a two-season, three-stage study found that oil content was higher at full emergence of the flower heads; davanone and linalool declined toward seed set, whereas methyl/ethyl cinnamates, bicyclogermacrene, davana ether, 2-hydroxyisodavanone, and farnesol increased (Mallavarapu et al. 1999). These trajectories make harvest stage and season necessary covariates in cross-species volatile comparisons, but they do not establish antiparasitic activity for any changing constituent.

Genetic and environmental components can be partially separated but rarely ignored. A linkage map identified loci affecting artemisinin yield in *A. annua*, providing direct evidence that segregating genetic variation contributes to the phenotype (Graham et al. 2010). In contrast, salt exposure, phytohormone treatment, reproductive development, and methyl jasmonate each changed pathway or oil phenotypes within experimental genotypes (Yadav et al. 2017; Maes et al. 2011; Arsenault et al. 2010; Wu et al. 2011). The DBR2-promoter study further linked promoter activity to high- and low-artemisinin chemotypes, showing how cis-regulatory variation at a branch point can contribute to pathway allocation (Yang et al. 2015).

Quantitative genetics adds an older but useful within-species benchmark: artemisinin content in *A. annua* showed predominantly additive inheritance and high narrow-sense heritability, and selected hybrid lines reached up to 1.4% of dry-leaf mass (Delabays et al. 2001). This supports selection as a source of reproducible chemotype variation, while leaving causal loci, linkage drag, genotype-by-environment stability, and terpene-network consequences unresolved.

Abiotic-stress studies extend this interaction beyond elicitor shorthand. Salt, cold, and waterlogging increased reported artemisinin content in one *A. annua* experiment, whereas drought did not, and each treatment produced a distinct transcriptome (Vashisth et al. 2018). In a separate two-variety experiment under arsenic stress, carrageenan oligomers and salicylic acid further changed antioxidant physiology and artemisinin yield (Naeem et al. 2021). These studies support environment-by-genotype regulation hypotheses, not universal stress responses or production recommendations; contamination, elicitor dose, harvest timing, and non-target metabolites must be measured alongside artemisinin.

Direct trichome phenotyping places a useful limit on stress-based explanations. Across field and greenhouse experiments on full-grown *A. annua*, sandblasting, leaf cutting, jasmonate, salicylate, chitosan, hydrogen peroxide, and salt produced mostly minor effects on glandular-trichome density; responses depended on clone and leaf position, and the hypothesis that stress generally induces larger or more numerous trichomes was rejected (Kjær et al. 2012). Thus, a stress-associated increase in artemisinin cannot be assumed to arise from more trichomes. Comparative experiments should measure trichome initiation, maturation, secretory capacity, pathway transcripts, and metabolites separately.

The same principle applies to apparently stable markers. Germacrene D is dominant in many *A. frigida* samples, yet the distribution of that dominance and its relationship to climate must be evaluated across populations and years. Recent *A. absinthium* work similarly shows that geography and phenological stage can change the dominant oxygenated mono- versus sesquiterpene class. These studies support stratified comparisons rather than a single canonical “wormwood oil.”

Post-harvest handling is part of the phenotype that reaches an assay. Controlled drying changed concentrations of artemisinin and its acid precursors in *A. annua* leaves (Ferreira and Luthria 2010). Hydrodistillation, solvent extraction, headspace analysis, and direct tissue extraction sample different chemical spaces and may also create or remove products through heat, oxidation, hydrolysis, or volatility. A reproducible comparison must therefore treat extraction method, storage duration, temperature, moisture endpoint, and abundance denominator as model variables rather than descriptive footnotes.

The oil fraction itself is sensitive to drying conditions. In wild Iranian *A. annua* harvested at full bloom, increasing drying temperature from room temperature to 65 °C reduced hydrodistilled oil yield from 1.12% to 0.37%; monoterpenes progressively declined as sesquiterpenes increased, and the dominant profile shifted from artemisia ketone/1,8-cineole toward beta-caryophyllene/germacrene D (Khangholi and Rezaeinodehi 2008). This is a processing effect within one sampled population, not evidence for a universally superior drying regime or a change in antimalarial potency. It does, however, make drying temperature a necessary covariate in any comparison of volatile chemistry or wormwood preparations.

Developmental series in *A. annua* make the confounding concrete: artemisinin and related sesquiterpenes were highest at particular vegetative stages, while the essential-oil mono:sesquiterpene ratio shifted from approximately 55:45 near the preflowering maximum to approximately 30:70 at other stages (Woerdenbag et al. 1994). A separate two-genotype time series identified genotype- and stage-dependent marker sets that included artemisinic acid, arteannuin B, borneol, beta-farnesene, camphor, and lanceol (Wang et al. 2009). In a broader Pakistan comparison of eight *Artemisia* species and two varieties, flowering-stage leaves generally had higher artemisinin, with the highest measured value in *A. dracunculus* var. *dracunculus* (Mannan et al. 2011). This is useful for a cross-species harvest hypothesis, but the HPLC endpoint and field/geographic design do not establish a universal

optimum for volatile terpenes or all *Artemisia* taxa. Thus, “chemotype” should be modeled jointly with development and genotype rather than treated as a static label.

A five-species comparison provides a useful cross-species test of this principle: variation in glandular-trichome density, artemisinin content, and expression of pathway genes was observed among *A. annua*, *A. deserti*, *A. absinthium*, *A. sieberi*, and *A. khorassanica* (Salehi et al. 2018). This is stronger comparative evidence than extrapolating from a single *A. annua* genotype, but expression and metabolite association still require common-annotation orthology, enzyme assays, and matched specimen chemistry before they can be interpreted as evolutionary gain or loss.

Contrasting high- and low-artemisinin *A. annua* chemotypes show why pathway intermediates must be measured alongside the end product. NMR-based profiling characterized more than 80 natural products and found that the low-artemisinin chemotype, with low *DBR2* expression, accumulated artemisinic acid, arteannuin B, and other unsaturated amorpho-4,11-diene derivatives, whereas the high-artemisinin chemotype accumulated more dihydroartemisinic acid, artemisinin, and related saturated products (Brown et al. 2018). The contrast supports branch-point and terminal-conversion hypotheses, but it is not an antiparasitic assay and does not prove that *DBR2* alone determines field yield or parasite potency.

An independent Japanese accession study reinforces this seasonal boundary: wild-derived *A. annua* grown in planters showed seasonal and positional variation in artemisinin, artemisinic acid, arteannuin B, and artemisitene, with leaf artemisinic acid reaching 1.0% of dry weight (Kawamoto et al. 1999). This is a useful accession-level comparator for precursor-rich chemotypes, but it does not define a universal harvest optimum or separate photoperiod, temperature, developmental, and genotype effects.

Regional production studies add a useful origin-by-chemistry layer. In Ugandan material, highland plants contained approximately 0.8% artemisinin and 2.6% total flavonoids versus 0.4% and 1.5% in lowland material, while camphor dominated relatively similar aromatic-oil profiles across regions (Engeu et al. 2015). The study supports source-specific standardization, but its component associations do not prove flavonoid or volatile synergy. Developmental regulation can be equally large: three-month leaves showed higher artemisinin than two-week leaves alongside miR156/SPL changes and increased *ADS*, *CYP71AV1*, *ADH1*, *DBR2*, and *ALDH1* transcripts (Li et al. 2021). These data make age, leaf position, and trichome state necessary covariates rather than interchangeable labels for “high-artemisinin” material.

Breeding and propagation can stabilize a selected chemotype without demonstrating a natural evolutionary mechanism. Selected *A. annua* genotypes exceeded 2% artemisinin and clonally propagated field material improved uniformity, with an estimated productivity of up to 70 kg/ha in the reported system (Wetzstein et al. 2018). A cross-species SCAR-marker study likewise proposed a roughly 1-kb marker associated with higher HPLC artemisinin in selected *A. annua* and *A. absinthium* samples (Asghari et al. 2015). Both are useful selection resources, but marker transfer, pathway integrity, genotype-by-environment stability, and parasite potency require independent validation.

Engineering studies provide a stronger production perturbation than expression correlation alone. A seven-gene construct that increased glandular secretory trichome density by 1.7–2.2-fold and reconstructed core artemisinin-pathway expression increased artemisinin 2.4–3.4-fold in engineered *A. annua* lines; line T9 reached 24.7 mg/g dry weight, with precursor increases also reported (Li et al. 2023). This is evidence for a coupled trichome/pathway engineering strategy in *A. annua*, not proof of conserved pathway architecture in *A. argyi* or wormwood and not evidence of parasite efficacy. A genome-wide *A. annua* bHLH survey identified 226 putative family members and three expression candidates associated with artemisinin-pathway patterns (Chang et al. 2023). Those candidates expand the regulatory hypothesis space, but expression similarity is not direct function, protein interaction, or cross-species conservation.

#### 4.1 *Artemisia argyi* as a tissue- and time-resolved comparator

The expanding *A. argyi* literature provides a useful counterweight to *A. annua*. De novo and full-length transcriptomes identified candidate genes across the MVA/MEP backbones, TPS families, P450s, and regulatory families, but these resources differ in tissue composition, assembly strategy, and annotation granularity (Liu et al. 2018; Cui et al. 2021). Targeted terpenoid measurements paired with bHLH-family analysis nominated regulatory candidates without claiming direct enzyme function (Yi et al. 2021). A second reference genome, assembled at 4.15 Gb with 147,248 predicted genes and 5,121 reported species-specific gene families, paired genome analysis with MeJA-responsive transcriptomics of terpenoid, flavonoid, and phenylpropanoid pathways (Gao et al. 2024). Its unusually large gene-model count and repeat-rich assembly are valuable resources but also illustrate why gene-family counts must be normalized to assembly, ploidy, and annotation quality before evolutionary interpretation. A recent review focused on *A. argyi* essential oil likewise reports location-dependent composition and a mainly preclinical evidence base, while identifying low bioavailability and the absence of unified quality standards as translation gaps (Gao and Lu 2026). It is useful for framing oil chemistry and delivery questions, but review-level pharmacology cannot substitute for voucher-linked primary chromatograms, parasite assays, or clinical safety data. The cytogenomic context is now more explicit. Genome-map-based karyotyping supports an allotetraploid interpretation for *A. argyi* and identifies a new octoploid germplasm class, with chromosome-doubling experiments affecting moxa-related traits (Li et al. 2025). This result does not prove terpene-gene dosage, but it changes the comparative design: homeolog retention, allele dosage, chromosome pairing, and cytotype must be modeled before a larger *A. argyi* TPS count is interpreted as lineage-specific innovation. In practical terms, diploid, tetraploid, and octoploid material should be analyzed

with ploidy-aware assemblies and read-depth-normalized copy-number estimates, then linked to secretory-tissue expression and quantitative chemistry.

Tissue-resolved metabolomics and transcriptomics showed that roots, stems, and leaves differ in both terpene-associated metabolites and candidate pathway expression; leaf-enriched MEP-pathway genes and co-expressed P450s formed hypotheses for subsequent functional testing (Zhang et al. 2022). Temporal profiling across four harvest months reported changing volatile-oil levels and identified full-length transcripts correlated with monoterpene and sesquiterpene profiles (Xu et al. 2022). These studies demonstrate that tissue and harvest date can produce differences as large as those attributed to genotype.

Analytical harmonization can make those tissue differences testable. A targeted UPLC-QQQ-MS/MS method quantified amorphadiene, artemisinic aldehyde, dihydroartemisinic acid, artemisinic acid, artemisinin B, artemisitene, and artemisinin in roots, stems, leaves, and lateral branches of four-month-old *A. annua* (Wang et al. 2024). The value of this record is methodological as much as chemical: simultaneous precursor/intermediate/product measurement can distinguish branch-point allocation from a terminal-product change. It does not establish in-vivo flux or antimalarial potency without isotopic, enzyme, and parasite-exposure data, but it provides a practical assay template for cross-species tissue comparisons.

An environmental perturbation study gives this tissue-and-genotype framework a mechanistic test. Low-phosphorus treatment increased caryophyllene in *A. argyi* volatile oil and fresh leaves, increased *AaTPS3* expression, and implicated *AaWRKY1* and *AaWRKY2* as negative regulators of *AaTPS3* (Ma et al. 2026). The result supports nutrient-responsive regulation of a defined terpene branch in the tested material, but it should not be generalized to all *A. argyi* accessions or interpreted as an antiparasitic mechanism without matched parasite and host-selectivity assays.

The East Asian essential-oil literature adds a useful review-level comparison. A synthesis of *A. argyi*, *A. princeps*, and *A. montana* describes oils containing monoterpenes, sesquiterpenes, and derivatives, while reporting that species-specific *A. argyi* constituents were often monoterpenoid, *A. princeps* had many species-specific aliphatic hydrocarbons, and *A. montana* was distinguished by sesquiterpenes (Yu et al. 2023). This supports geographic and taxonomic stratification of volatile sampling, but the review itself identifies unresolved mechanisms, toxicity, and quality control; it is not a substitute for voucher-linked primary chromatograms or a terpene-specific parasite assay.

Not every trichome study is a terpene study, and that distinction is experimentally useful. Integrated multi-omics in *A. argyi* nominated *AaMYB1* and *AaYABBY1* for flavonoid biosynthesis; overexpression increased flavonoid or rutin readouts, and *AaYABBY1* increased trichome density in tobacco and *Arabidopsis* (Cui et al. 2024). These results provide a non-terpene developmental control for interpreting trichome-associated chemistry: a regulatory perturbation can change both cell number and a specialized-metabolite branch without proving a corresponding change in terpene flux or parasite activity.

Genome size is another source of apparent chemical variation. Flow cytometry across 42 Chinese *A. argyi* accessions measured 8.428–11.717 pg and separated the accessions into three clusters without a significant geographic relationship; a genome survey estimated approximately 7.852 Gb and provided heterozygosity, repeat, GC, and ploidy context (Luo et al. 2023). Because genome size can reflect chromosome dosage, repetitive DNA, and assembly representation, this result argues for cytotype-aware normalization before relating TPS abundance to chemistry. It does not itself demonstrate TPS expansion, pathway dosage, or adaptive antiparasitic function.

Organ and time are equally important environmental covariates. Dynamic cold-exposure and recovery experiments in *A. annua* found different transcription-factor and calcium-signaling responses in leaves and roots, disrupted then restored carbon transport from source leaves to sink roots, accelerated root-to-leaf nitrogen transport, and earlier activation of cold-response programs in leaves (He et al. 2024). The experiment supports organ-resolved sampling of terpene-associated stress responses; it does not establish that a cold-induced transcript predicts a particular volatile or sesquiterpene-lactone product, nor that the response is conserved across *Artemisia*.

An *A. argyi* soil-cadmium study provides a complementary stress boundary: plants maintained normal growth and intact cellular structure up to the reported 10 mg/kg treatment, while proline, cadmium translocation, and transcriptomic pathways involving terpenoid, flavonoid, amino-acid, and sugar metabolism changed with exposure (Yang et al. 2024). This is valuable for separating environmental transcriptome effects from genetic differences and for flagging contaminated-material risks, but pathway transcription is not native flux, adaptive fitness, product safety, or parasite efficacy.

The same design logic extends to non-medicinal ecological comparators. Drought and rewatering in *A. sphaerocephala* changed water status, oxidative-stress measures, soluble metabolites, and hormone-, carbohydrate-, glyoxylate-, and nitrogen-related transcript programs in a full-length transcriptome study (Xiao et al. 2025). Although the reported endpoints do not resolve a terpene product, they establish why drought history and recovery must be recorded when comparing arid-zone *Artemisia* accessions. A stress transcript is a sampling covariate and candidate regulator, not evidence of an antiparasitic molecule.

Single-cell transcriptomics and matched metabolomics now resolve this problem at higher spatial resolution. Glandular trichomes and non-glandular tissues of *A. argyi* showed strongly different metabolite profiles, with sesquiterpenoids prominent among trichome-enriched terpenoid differences, and the single-cell atlas nominated developmental and TPS-expression programs (Dong

et al. 2026). This is powerful candidate evidence, but co-expression edges and cell-type enrichment remain hypotheses until recombinant enzymes, gene perturbations, and isotope or product-feeding experiments establish catalytic sequence and flux.

An *A. annua* single-nucleus and spatial-transcriptomic atlas provides a complementary cell-resolved reference for the cross-species design. The study defined three glandular-secretory-trichome developmental states and identified six secretory cells within a ten-cell trichome as the primary artemisinin-production site, together with hundreds of hub-gene candidates (Zhang et al. 2025). This sharpens the sampling requirement: bulk-leaf orthology should be paired with matched secretory-cell expression and metabolite localization. The atlas remains a candidate map, however; hub-gene function, protein abundance, isotope-traced flux, and conservation outside *A. annua* still require direct tests.

## 5. Comparative evolutionary genomics

The initial controlled comparison is *A. annua* versus *A. argyi*. The former supplies a pathway-rich reference genome and functional artemisinin literature. The latter has a chromosome-scale genome with reported whole-genome duplication and terpene-synthase expansion, plus tissue transcriptomes. The apparent ADS contradiction is now interpretable as an assembly-, ploidy-, annotation-, and function-level discrepancy rather than a simple presence/absence result. The earlier chromosome-scale study selected a 3.89-Gb, 17-pseudochromosome haplotype-A projection, annotated 122 TPS genes, and described partial ADS deletion and loss of function. A later 7.88-Gb haplotype-resolved assembly predicted 451 TPS models, of which 261 were manually classified as intact, and identified a six-copy tandem ADS homolog cluster; recombinant AarADS proteins produced  $\alpha$ -bisabolol rather than the amorpho-4,11-diene made by AanADS (Chen et al. 2023; Qi et al. 2025). Thus, the later result supports ADS-family retention with functional divergence, while neither paper alone establishes an intact, expressed, artemisinin-producing pathway in *A. argyi*.

The genus-scale phylogenomic scaffold is now substantially better than the earlier marker-limited picture. Nuclear SNP data from 228 *Artemisia* species and 258 samples recovered eight supported clades, nested *Kaschgaria* within *Artemisia*, and showed that several traditional subgenera are not monophyletic (Jiao et al. 2023). This is the appropriate backbone for testing whether terpene traits are phylogenetically clustered, but it does not itself measure terpene genes or chemistry. Cytogenomic sampling adds a second constraint: flow cytometry across 67 species and 84 populations showed that genome-size increase is not proportional to ploidy and that 1Cx values tend to decline at high ploidy (Pellicer et al. 2010). Copy-number comparisons should therefore use cytotype-aware, read-depth-normalized assemblies rather than treating genome size as a direct proxy for TPS expansion.

Within-species plastome structure can improve accession sampling while remaining distinct from nuclear pathway evolution. Complete chloroplast genomes from 38 *A. annua* strains resolved two lineages and 14 haplotypes, with 60 polymorphic sites but no chloroplast gene-content presence/absence difference (Ding et al. 2024). This supports accession discrimination and a possible geographic sampling covariate for artemisinin studies; chloroplast haplotype structure is not evidence for nuclear ADS, CYP71AV1, DBR2, or volatile-TPS dosage or function.

Resource-level plastome data in *A. selengensis* show how this scaffold can be extended without overclaiming. Ten complete chloroplast genomes were highly conserved in gene content and order, spanning 151,148–151,257 bp and 133 genes, while variable regions and eight mutation hotspots improved phylogenetic discrimination among resources (Wang et al. 2024). A paired comparison of *A. maritima* and *A. absinthium* likewise reported approximately 151-kb plastomes, high structural similarity, and 20 candidate polymorphic marker regions, with a maximum-likelihood tree placing *Artemisia* as monophyletic and sister to *Chrysanthemum* (Shahzadi et al. 2020). These data can distribute accessions for common-garden chemistry and nuclear sequencing, but plastome markers cannot resolve nuclear TPS/OSC orthology, WGD-derived copy number, or enzyme function. The appropriate model treats plastid topology as a sampling covariate and tests terpene evolution with phased nuclear assemblies, reconciled gene trees, expression, and product assays.

Broader *Seriphidium* sampling reinforces the distinction between taxonomic scaffold and pathway evolution. Eighteen newly sequenced plastomes from 16 *Seriphidium* species, combined with a published taxon, showed conserved structure and eight variable loci but recovered the subgenus as polyphyletic (Jin et al. 2023). This result is useful for choosing representatives and testing whether wormwood-associated chemistry tracks a plastid-defined clade; it cannot establish nuclear TPS duplication/loss, sesquiterpene-lactone biosynthesis, or shared vermifuge chemistry. In particular, plastome-based non-monophyly should trigger taxonomic and sampling review, not a claim that parasite-active terpene pathways evolved convergently.

The 2025 global phylogeny now provides a substantially broader nuclear backbone for that design. A gigamatrix analysis placed 394 of 505 accepted *Artemisia* species using 202 low-copy nuclear genes together with ITS/ETS, and genome-skimming supplied nuclear-marker and complete-plastome data for 314 species; the authors recognized 24 sections in eight subgenera and reported extensive cytonuclear discordance (Jiao et al. 2025). This framework is valuable for stratifying *A. annua*, *A. absinthium*, *A. argyi*, *A. herba-alba*, *A. cina*, and other chemistry-bearing taxa, and for testing whether chemotypes or parasite phenotypes are phylogenetically clustered. It is not itself a terpene-gene analysis: low-copy marker and plastome relationships do not establish nuclear TPS/P450/OSC orthology, duplication or loss, pathway flux, or antiparasitic adaptation. The appropriate next step is to map voucher-linked chemistry and parasite evidence onto this backbone, then analyze a common, ploidy-aware nuclear annotation for the selected species.

The comparative analysis should infer orthogroups and gene trees for GPPS, FPPS, monoterpene synthases, sesquiterpene synthases, ADS, CYP71AV1-like P450s, DBR2, ALDH1, OSCs, and selected sesquiterpene-lactone enzymes. Duplications and losses should be mapped to a broad *Artemisia* species tree. Expression should be interpreted by tissue and developmental context, with glandular trichome data prioritized where available. Missing annotation is unresolved data, not evidence of biological absence.

An older cross-species PCR screen illustrates the cost of inferring pathway completion from a single marker. Among ten *Artemisia* species, selected ADS active-site regions amplified only in *A. annua*, *A. aucheri*, and *A. chamaemelifolia*, while the complete gene was not amplified in any tested species (Hosseini et al. 2011). Negative amplification can reflect primer mismatch, indels, gene fragmentation, or missing template; positive amplification of a short motif cannot establish an intact enzyme. The result should therefore be reinterpreted as a historical candidate screen and replaced by long-read assemblies, common annotation, reconciled gene trees, expression, and recombinant product assays.

Ploidy provides a second, experimentally anchored confounder. Colchicine-induced tetraploid *A. annua* showed 39-56% higher artemisinin than diploid plants across the vegetation period, with increased expression of selected pathway genes including *FPS*, *HMGR*, and *Aldh1* (Lin et al. 2011). This supports a ploidy-associated production phenotype, but it does not distinguish dosage from altered development, trichome density, or regulation. The observation strengthens the case for cytotype-aware comparative sampling across *A. annua*, *A. argyi*, *A. tridentata*, and wormwood rather than interpreting raw gene counts as evolutionary innovation.

Plastid phylogenomics can provide a sampling scaffold, but it cannot replace nuclear pathway phylogenies. A 2025 comparative analysis of *Artemisia* plastomes addressed limited chloroplast sampling and inferred infra-generic relationships that can help distribute accessions across the genus (Guo et al. 2025). Those relationships are useful for choosing representatives and testing whether chemical traits are phylogenetically clustered; they do not establish nuclear TPS/OSC orthology, WGD-derived copy number, or functional evolution. The article therefore treats plastome topology as a taxonomic covariate and reserves gene-family conclusions for common nuclear annotations and reconciled gene trees.

An East Asian plastome resource provides a concrete sampling scaffold for that design. Kim and colleagues compared 32 complete plastomes from 21 *Artemisia* taxa, including 19 newly sequenced plastomes from 18 taxa, and identified conserved gene order alongside variable loci suitable for phylogenomics and DNA-barcode development (Kim et al. 2020). The resource improves taxon and accession balancing for common-garden chemistry, but plastid variation remains a taxonomic covariate: it cannot establish nuclear TPS/OSC orthology, WGD-derived copy number, artemisinin-pathway completion, or antiparasitic function. In practical terms, plastome-defined strata should be paired with phased nuclear assemblies, cytotype metadata, reconciled gene trees, and voucher-linked metabolomics.

Published comparative results already provide testable anchors: the later *A. argyi* analysis reports 451 predicted TPS models split between two haplotypes (221 and 230), with 261 intact models after manual correction, while the chromosome-scale study links WGD and duplication modes to TPS expansion and identifies a germacrene D synthase cluster. The latter study also reports pathway elucidation for germacrenes, (+)-borneol, and (+)-camphor. These observations are catalogued in the KB's gene-family-evidence.csv and ADS/TPS reconciliation table; they are not treated as a substitute for a common-annotation, orthology-aware rerun.

The transcript and metabolite literature provides a validation ladder for genomic predictions. A candidate duplicated TPS is more compelling when it is intact under a common annotation, placed within a supported gene tree, retained in synteny or a tandem cluster, expressed in the relevant secretory cell, and correlated with a product that the recombinant protein can generate. The *A. argyi* tissue, temporal, and single-cell studies supply several of these intermediate layers, but none independently establishes orthology or ancestral function. In *A. annua*, by contrast, the mature ADS/CYP71AV1/DBR2 evidence chain shows what eventual convergence should look like. A functional *A. argyi* example is beta-caryophyllene synthase: four candidate synthases were identified, and engineered yeast expressing AarTPS88 reached 15.6 g/L beta-caryophyllene in fed-batch fermentation (Li et al. 2025). This validates a production-capable enzyme/cell-factory route, not native leaf abundance, ecological adaptation, orthology, or antiparasitic action.

The genome-scale anchors also illustrate why “expansion” must be treated as a layered observation. The *A. annua* draft genome combined a 1.74-Gb highly heterozygous assembly, 63,226 predicted protein-coding genes, terpene-enzyme diversification, and transcriptomic evidence for a regulatory network associated with artemisinin biosynthesis. The chromosome-scale *A. argyi* study instead placed a recent lineage-specific WGD, duplicated TPS families, germacrene-D/(+)-borneol/(+)-camphor pathway evidence, and partial ADS deletion or homolog loss-of-function in a 3.89-Gb, 17-pseudochromosome context (Luo et al. 2018; Chen et al. 2023). The contrast is biologically informative, but the reported gene counts and ADS states are not a common-denominator orthology result: assembly scope, haplotype representation, annotation, ploidy, and product validation differ. These studies therefore motivate, rather than replace, the KB's common-reannotation and reconciled-gene-tree gates.

Regulatory evolution should be evaluated with the same caution as enzyme evolution. The abundance of MYB, bHLH, WRKY, AP2/ERF, and light- or jasmonate-responsive candidates in transcriptomic analyses may reflect both genuine network expansion and permissive co-expression thresholds. Functional studies of AaERF1/AaERF2, WRKY1, AaMYB108, AabHLH112, and the AaGSW1 complex define experimentally testable motifs and interactions in *A. annua*, but homologous regulators in *A. argyi* should

not inherit those functions by name alone. Conserved target motifs, protein interactions, cell-type expression, perturbation phenotypes, and metabolite responses are required to claim network conservation.

The Terpedia KB now carries an execution specification for this analysis (`comparative-genomics/`). Its required outputs include frozen input checksums, software versions, orthology-aware gene trees, synteny/tandem-cluster calls, and a tiered gene-family evidence table. A completed two-proteome OrthoFinder run is now archived, but the broader genus-wide comparison remains a design-supported hypothesis framework because the input contains only *A. annua* and one *A. argyi* annotation release. As a bounded extension, six pathway-associated candidate groups were aligned with MAFFT and analyzed with IQ-TREE3 under LG+G4 with 1,000 ultrafast bootstrap and 1,000 SH-aLRT replicates. The groups were annotated in the *A. annua* reference as germacrene D synthase, germacrene A synthase, sesquiterpene synthases 1 and 6, CYP71AV1, and DBR2; they contained 4–9 *A. tridentata*, 1–4 *A. argyi*, and 1–6 *A. annua* leaves per group. These trees are exploratory candidate-family diagnostics: they do not establish orthology, duplication/loss, ancestral state, enzyme function, pathway flux, or antiparasitic mechanism. The exact alignments, trees, manifests, software versions, and input hashes are archived in `comparative-genomics/results/three-species-pathway-gene-trees-2026-09-02/`. AabHLH2 and AabHLH3 provide a regulatory anchor for this pathway-centered comparison: in *A. annua* they acted as nuclear repressors that competed with AaMYC2-like cis-regulatory recognition and negatively regulated artemisinin biosynthesis (Shen et al. 2022). This result makes bHLH copy, sequence, binding-site, and cell-type comparisons testable, but a homologous gene name is not evidence of conserved regulation in *A. argyi*, *A. tridentata*, or any other species. A separate engineering study shows that the genus can serve as a platform for non-native C15 chemistry: farnesyl diphosphate synthase and patchoulol synthase expression in *A. annua* produced patchoulol, and signal-peptide targeting increased the reported accumulation to 273 microgram/g dry weight (Fu et al. 2020). This is evidence for engineered production and compartment targeting, not native patchoulol ecology, pathway evolution, or antiparasitic activity.

An NCBI Datasets metadata audit was run without downloading sequence archives. It returned one *A. annua* assembly report and three separate *A. argyi* reports. The literature resource PRJCA010808 was resolved to its external NGDC/CNCB project record, not to an NCBI assembly accession; the published 3.89-Gb, 17-pseudochromosome haplotype must therefore be treated as a distinct input until sequence and annotation files are obtained and checksummed. This is itself a result for reproducibility: assembly choice must be documented before copy-number or presence/absence comparisons are interpreted.

The KB's assembly comparison table makes the confounders explicit: the *A. annua* NCBI reference is scaffold-level, the literature *A. argyi* input is a 3.89-Gb haplotype-A projection, and NCBI also exposes larger or differently assembled *A. argyi* resources. Genome length and chromosome count should therefore not be interpreted as evidence for terpene-family expansion without a common annotation and orthology workflow.

The execution gate is also explicit: NCBI reports an annotation with 63,226 protein-coding genes for the *A. annua* reference, and the earlier NGDC/CNCB *A. argyi* haplotype-A annotation exposes 62,844 protein-coding genes. These counts are not directly comparable to the later 7.88-Gb haplotype-resolved annotation. A seed-centered DIAMOND screen of 31 annotated *A. annua* pathway/TPS queries returned 105 *A. argyi* candidate hits; the exact inputs and hashes are archived in the KB. These are homology candidates only. A cross-species copy-number or orthology result is not reported until a common annotation, reciprocal/phylogenetic validation, and gene-tree reconciliation are completed.

Reciprocal checking re-searched 67 unique *A. argyi* candidates against the *A. annua* proteome and produced 26 reciprocal-best-hit candidate rows. Because a duplicated *A. argyi* locus can share the same best *A. annua* seed, this count is not a one-to-one ortholog count and is not interpreted as gene-family expansion.

As a diagnostic next step, the KB built MAFFT alignments and IQ-TREE3 exploratory trees for the three reciprocal groups that retained at least four unique protein sequences (PWA60940.1, PWA68009.1, and PWA85324.1). The run used the LG+G4 model, 1,000 ultrafast-bootstrap replicates, 1,000 SH-aLRT replicates, and one thread; the exact alignments, tree files, reports, software versions, and input checksums are archived under `comparative-genomics/results/seed-gene-trees-2026-09-01/`. Eight of eleven reciprocal groups were not tree-built because they had fewer than four leaves or fewer than four unique sequences. These small, seed-biased trees are quality-control diagnostics only: they do not establish orthology, duplication/loss, or pathway function.

A restricted OrthoFinder diagnostic then clustered the 31 *A. annua* seed proteins and 67 unique *A. argyi* forward-hit candidates. It produced 16 candidate orthogroups, 11 shared clusters, and five *A. argyi*-only clusters. The full tables, input FASTAs, software context, and checksums are archived in `comparative-genomics/results/candidate-orthofinder-2026-09-01/`. This result is useful for prioritizing follow-up gene trees and synteny, but the seed-biased, two-taxon candidate set cannot support genome-wide orthology, copy-number, duplication/loss, or gene-function claims.

To broaden pathway prioritization without overstating the unfinished full-proteome run, a separate annotation-keyword and homology-filtered analysis selected 890 *A. annua* pathway-associated proteins and 1,273 unique *A. argyi* candidates. OrthoFinder `-M msa -og` grouped these filtered inputs into 388 candidate orthogroups: 271 shared, 17 *A. annua*-only, and 100 *A. argyi*-only; 83 groups were single-copy in both filtered inputs. Because the query set is annotation-biased, the *A. argyi* set is homology-selected, and `-og` stops before species-tree and gene-tree inference, these counts are a reproducible prioritization scaffold rather than evidence of genome-wide orthology, copy-number change, duplication/loss, ancestral state, enzyme function, pathway flux, or an-

tiparasitic mechanism. The complete tables, group sequences, command, software versions, input hashes, and checksums are archived in `comparative-genomics/results/pathway-focused-orthofinder-2026-09-02/`.

The full two-proteome run was then completed by resuming the finished DIAMOND searches with native ARM64 MCL and one OrthoFinder analysis thread (Emms and Kelly 2019). It assigned 111,235 of 129,761 proteins (85.7%) to 28,696 orthogroups, including 25,039 groups containing both species, 7,086 single-copy groups, 3,657 species-specific groups, and 18,526 unassigned proteins. The inferred tree, (*A. annua*, *A. argyi*), is a valid two-tip species tree but contains no information about relationships among the wider genus. The complete compressed result, log, input checksums, and execution note are archived in `comparative-genomics/results/artemisia-of-results-run3-full-2026-09-02.tar.gz` and `comparative-genomics/results/full-orthofinder-2026-09-02.md`.

Two published genome-scale studies provide important context for interpreting that restricted two-proteome comparison. An integrated *A. annua* multi-tissue dataset identified 88 metabolites across 14 tissues—26 monoterpenes, 22 sesquiterpenes, two triterpenes, and one diterpene—and connected developmental expression patterns with pathway-associated genes (Ma et al. 2015). In *A. argyi*, an earlier 8.03-Gb allotetraploid assembly reported 135 TPS genes, tandem clusters, and tissue-biased expression alongside broad secondary-metabolism expansion (Miao et al. 2022). These studies support a class-balanced gene-to-metabolite design, but their assembly sizes, ploidy interpretations, annotation releases, and gene models are not directly interchangeable with the later 3.89-Gb or 7.88-Gb *A. argyi* resources. TPS counts and gene-expansion claims must therefore be re-evaluated after common annotation, homeolog-aware gene-tree placement, synteny, expression, and product confirmation.

An allele-aware *A. annua* assembly provides a complementary within-species evolutionary anchor. Two chromosome-level haplotype maps, paired with resequencing of 36 samples spanning artemisinin content, found a positive association between *amorphadiene synthase* (ADS) copy number and artemisinin yield and enabled paralog-aware transcript detection (Liao et al. 2022). This supports ADS dosage as a testable contributor to chemotype variation in *A. annua*, but it does not establish that every ADS-like copy is intact, expressed, or catalytically equivalent. In comparative work, copy number should therefore be modeled jointly with haplotype, gene-tree placement, transcript abundance, trichome localization, and measured artemisinin or precursor levels.

An annotation-anchored extraction mapped 35 *A. annua* proteins carrying curated pathway/TPS header labels to 21 pathway-label/group combinations, representing 17 assigned orthogroups plus one *A. annua* protein that remained unassigned. Several groups contain many copies in both species, but this is not evidence of a lineage-specific expansion: the labels are *A. annua* annotation anchors, *A. argyi* members have not inherited function by name, and OrthoFinder duplication counts are not equivalent to intact, expressed, biochemically active terpene loci. One particularly important caution is that the CYP71AV1 and germacrene-A-oxidase anchors fall in the same broad P450 orthogroup, illustrating why family membership must be followed by gene-tree placement and biochemical validation. The full per-protein mapping and grouped summary are archived in `comparative-genomics/results/full-pathway-family-mapping-2026-09-02.tsv` and `comparative-genomics/results/full-pathway-group-summary-2026-09-02.tsv`.

Annotation quality is itself an evolutionary-comparison variable. Manual correction of the *A. annua* genome annotation using second- and third-generation transcript evidence changed gene structures, alternative-splicing calls, expression estimates, and functional predictions (Liu et al. 2023). A common annotation pipeline should therefore be treated as a prerequisite for comparing TPS, P450, DBR2, ALDH, and OSC copy number; missing or mis-modeled loci are unresolved data rather than biological absences. The plant-to-yeast literature provides a complementary production context: engineered *E. coli* and *S. cerevisiae* systems, plant pathway engineering, and fermentation optimization have moved artemisinin precursor production from plant tissue toward microbial chassis, with reported artemisinic-acid titers reaching 25 g/L in yeast (Zhao et al. 2022). These outputs demonstrate engineering feasibility and supply-chain relevance, not native *Artemisia* evolution, plant abundance, or parasite efficacy.

An important third genome-scale context is *A. tridentata*, a North American sagebrush with a haploid pseudo-chromosome assembly of approximately 4.2 Gbp, 43,377 predicted genes, 91.37% complete BUSCOs, and 77.99% repetitive sequence (Melton et al. 2022). A comparative resequencing study found extensive chromosomal rearrangements relative to *A. annua*, conserved large-scale synteny in selected regions, and population-level transposable-element differentiation among three diploid *A. tridentata* genomes (Melton et al. 2024). More recent cytological and reduced-representation sequencing data support both one-step and two-step whole-genome-duplication pathways, including a possible triploid bridge and a reported 0.22% polyploidization rate in the study material (Grossfurthner et al. 2026). These records materially strengthen the evolutionary design by showing that genome architecture and cytotype can vary within the genus, but they do not establish *A. tridentata* terpene-locus function or parasite activity. The NCBI assembly has no NCBI protein annotation, so the accession-matched Maker GFF3 released with the genome study was translated locally and retained with its source genome URL, assembly-name map, derivation script, and checksums in the Terpedia KB. This makes *A. tridentata* usable in a bounded candidate-family diagnostic while preserving the common-reannotation limitation.

The resulting three-species diagnostic searched the same 890 *A. annua* pathway-associated queries against the frozen *A. argyi* proteome and the 41,700-protein *A. tridentata* translation, recovering 1,273 and 768 candidate homologs, respectively. OrthoFinder -S diamond -M msa -og grouped the filtered inputs into 478 candidate orthogroups, of which 205 contained all three species and 31 were single-copy in all three. Pairwise shared-group counts were 291 for *A. annua*-*A. argyi*, 236 for *A. annua*-*A. tridentata*, and

265 for *A. argyi*–*A. tridentata*. These values are useful for selecting loci for gene trees, synteny, expression, and enzyme assays, but the query and target sets are homology-selected, annotations are not common-reannotated, and –og stops before gene-tree/species-tree reconciliation. They therefore do not support genome-wide copy-number, duplication/loss, ancestral-state, pathway-flux, or antiparasitic-mechanism claims. The full candidate tables, filtered FASTAs, OrthoFinder log, provenance manifest, and check-sums are archived in [comparative-genomics/results/three-species-pathway-orthofinder-2026-09-02/](https://comparative-genomics/results/three-species-pathway-orthofinder-2026-09-02/).

At broader genus scale, flow cytometry across 21 *Artemisia* species and 24 populations measured 2C DNA contents from 3.5 to 25.65 pg, with strong associations to karyotype length and ploidy and additional systematic/ecological structure (Torrell and Vallès 2001). Genome size is therefore a relevant covariate for sampling and orthology interpretation, especially where homeolog retention and chromosome-scale duplication can mimic lineage-specific terpene-gene expansion. It is not, however, a proxy for TPS copy number, pathway completeness, metabolite abundance, or antiparasitic function.

Within *A. annua*, induced autotetraploids provide a more direct ploidy perturbation. A transcriptomic comparison reported higher mean artemisinin in tetraploids, altered glandular-secretory-trichome morphology, and 8,763 differentially expressed genes involving metabolism, defense, hormone response, and methylation (Xia et al. 2018). This supports a ploidy-associated chemotype hypothesis, but the induction background, biomass tradeoff, and causal gene set require independent validation and should not be generalized to natural polyploids or other species.

Table 4. Selected annotation-anchored groups from the completed two-proteome run. Counts are OrthoFinder group membership and duplication diagnostics, not functional copy-number calls.

| <i>A. annua</i><br>annotation<br>anchor                  | Orthogroup | <i>A. annua</i> / <i>A. argyi</i><br>genes | Duplications | Gene-tree leaves | Boundary                                                                                            |
|----------------------------------------------------------|------------|--------------------------------------------|--------------|------------------|-----------------------------------------------------------------------------------------------------|
| CYP71AV1;<br>germacrene-A<br>oxidase                     | OG0000295  | 19 / 8                                     | 21           | 27               | Broad P450<br>group; shared<br>group does not<br>distinguish<br>CYP71AV1<br>from GAO<br>function    |
| DBR2;<br>artemisinic-<br>aldehyde<br>reductase labels    | OG0006177  | 3 / 1                                      | 2            | 4                | Compact shared<br>group; exact<br>enzyme<br>assignment<br>requires se-<br>quence/function<br>checks |
| Germacrene-A<br>synthase                                 | OG0011988  | 2 / 1                                      | not reported | not built        | Three total<br>members fell<br>below the<br>gene-tree<br>output<br>threshold                        |
| Germacrene-A<br>synthase 2                               | OG0001344  | 5 / 6                                      | 7            | 11               | Candidate TPS<br>family; <i>A. argyi</i><br>function<br>unverified                                  |
| Germacrene-D<br>synthase;<br>sesquiterpene<br>synthase 4 | OG0000309  | 17 / 10                                    | 19           | 27               | Mixed TPS<br>labels and<br>duplication-rich<br>group; no<br>product transfer<br>inferred            |
| Farnesyl<br>diphosphate<br>synthase 2                    | OG0000439  | 16 / 5                                     | 16           | 21               | Prenyl-<br>transferase<br>group; not a<br>direct flux or<br>abundance<br>comparison                 |



| <i>A. annua</i><br>annotation<br>anchor     | Orthogroup | <i>A. annua</i> / <i>A. argyi</i><br>genes | Duplications | Gene-tree leaves | Boundary                                                                                                |
|---------------------------------------------|------------|--------------------------------------------|--------------|------------------|---------------------------------------------------------------------------------------------------------|
| Oxidosqualene<br>cyclase 3                  | OG0001412  | 9 / 1                                      | 8            | 10               | Candidate<br>triterpene<br>branch group; <i>A.</i><br><i>argyi</i> member<br>remains<br>uncharacterized |
| Geranylgeranyl<br>pyrophosphate<br>synthase | OG0009299  | 1 / 3                                      | 2            | 4                | Shared prenyl-<br>transferase<br>group; copy<br>number is<br>annotation- and<br>assembly-<br>dependent  |

An independent *A. argyi* leaf, root, and stem transcriptome supplies candidate terpenoid-biosynthesis transcripts and tissue contrasts (Liu et al. 2018). It extends the comparison beyond *A. annua*, but de novo transcript annotation cannot distinguish paralogs, homeologs, and allelic copies reliably enough for gene-family evolution without genome-guided placement. Experimental work on an *A. annua* (E)- $\beta$ -farnesene synthase further shows that a sparse set of epistatic sequence changes can alter terpene-enzyme behavior (Ballal et al. 2020); this motivates testing sequence-function effects within orthologous clades rather than treating all TPS copies as functionally equivalent.

Environmental and perturbation studies provide useful mechanistic tests within *A. annua*. Short-term UV-B exposure increased artemisinin production and induced ADS, CYP71AV1, and stress/hormone-associated transcripts (Pan et al. 2014). A CYP71AV1 mutant redirected flux toward the alternate sesquiterpene epoxide arteannuin X and supported nonenzymatic terminal conversion in glandular trichomes (Czechowski et al. 2016). These results demonstrate inducibility and metabolic plasticity, but they do not establish that a homologous response or alternate product occurs in another *Artemisia* species.

Recent within-species studies add actionable controls for this comparative design. In a 17-bioecotype panel, gene-specific STS markers and HPLC identified marker-yield associations and bioecotypes exceeding 2.7 mg/g artemisinin, but the correlations remain selection signals rather than causal variant assignments (Hallajian et al. 2026). AaSPATULA perturbation connected glandular-trichome initiation with biomass and whole-plant artemisinin yield, while AaMYC3 linked jasmonate response, trichome density, and pathway-gene activation (Yuan and Zhou 2026; Yuan et al. 2025). TLR3 and differentially expressed lncRNAs provide additional plant-side regulatory candidates (Dong et al. 2024; Ma et al. 2024). The common inference boundary is important: improved trichome capacity or artemisinin yield does not, by itself, predict antiparasitic potency unless the resulting chemical fingerprint and parasite exposure are measured.

The older regulatory literature provides a useful perturbation ladder for interpreting these newer candidates. TAR1 altered trichome morphology, cuticular wax, ADS/CYP71AV1 expression, and artemisinin; trichome-specific AaWRKY1 increased CYP71AV1 and artemisinin without changing several upstream transcripts; and AabHLH1, AaSPL2, and AabZIP9 supplied promoter-binding or transactivation evidence at ADS, CYP71AV1, or DBR2 (Tan et al. 2015; Han et al. 2014; Ji et al. 2014; Lv et al. 2019; Shen et al. 2019). DBR2 overexpression increased both artemisinin and competing-branch metabolites, showing why pathway engineering should be evaluated by a complete metabolite panel rather than target-product abundance alone (Yuan et al. 2015).

Tissue and environment can uncouple a genomic prediction from a harvested phenotype. Bulk-tissue measurements found the highest artemisinin in leaves but distinct patterns for upstream and pathway genes across leaves, flowers, roots, and stems (Xiang et al. 2012); imaging and LC-MS later detected pathway products in non-glandular cells and glandless material (Judd et al. 2019). Mycorrhizal colonization, ABA rescue under copper stress, and combined UV-B/GA treatment each increased artemisinin-associated traits through partially overlapping stress and hormone networks (Mandal et al. 2015; Zehra et al. 2020; Ma et al. 2020). In field germplasm, ART and dihydroartemisinic acid often peaked before flowering and differed among Brazilian, Chinese, and Swiss chemotypes, whereas forced early flowering did not itself increase artemisinin (Ferreira et al. 2018; Wang et al. 2007). These findings support a model in which copy number, cell type, developmental state, environment, and precursor competition jointly determine chemistry.

Negative comparative controls are especially valuable. Eighteen *A. khorassanica* populations included diploids and tetraploids, yet neither class contained detectable artemisinin; DBR2 was low in diploids and silenced in autotetraploid material (Hamidi et al. 2018). This result constrains the inference that ploidy or expression of an ADS-like gene is sufficient for an artemisinin phenotype. In the proposed cross-species analysis, absence of artemisinin should therefore be modeled alongside sequence integrity,

expression, tissue localization, precursor abundance, and assay detection limits.

## 6. Antiparasitic evidence

Two parasite-facing stories deserve priority in this review. Malaria is the strongest mechanistic case because artemisinin has a chemically defined scaffold, a reconstructed *A. annua* pathway, extensive parasite pharmacology, and a clinically established artemisinin-based combination-therapy context. Wormwood (*A. absinthium*) is a separate, important veterinary anthelmintic case: traditional vermifuge use is supported by several extract, oil, feed, and sheep studies, but the active chemistry, dose standardization, and human safety remain unresolved. The evidence should be developed in parallel rather than merged under the generic label “antiparasitic *Artemisia*.”

### 6.1 Terpene-protein interaction map

Full-text verification of a commercial *A. annua* oil against engorged *Rhipicephalus annulatus* females found dose-dependent mortality from 33.2% to 100% by 36 hours and complete egg-laying failure at the highest tested dilution. The oil was not chemically profiled, and cattle safety, residues, field performance, and constituent causality were not tested; this is therefore a vector/ectoparasite comparator rather than evidence for malaria treatment or a single terpene (Pirali-Kheirabadi and Teixeira da Silva 2011).

Vector-control evidence provides a preparation-level bridge between terpene chemistry and parasite transmission. Nanoliposomes carrying essential oils from *A. annua*, *A. dracunculus*, and *A. sieberi* showed species-dependent larvicidal activity against *Aedes aegypti* and the malaria vector *Anopheles stephensi*, with the *A. dracunculus* formulation most active in the reported comparison (Sanei-Dehkordi et al. 2023). A separate *A. dracunculus* essential-oil nanogel and nanoemulsion study reported LC50 values of 6.68 and 13.53 microgram/mL, respectively, against *An. stephensi* larvae (Osanloo et al. 2022). These results support formulation- and oil-specific vector control, but not malaria treatment, a single-terpene mechanism, or non-target safety. Conversely, an indexed Zimbabwean letter reported *A. afra* essential oil as ineffective for inhibiting mosquito biting under its tested conditions (Lukwa et al. 2000). The negative result is route- and preparation-specific because the retrieved record lacks concentration and composition details; it does not establish absence of other *A. afra* vector phenotypes. A newer defined-constituent study similarly found fumigation toxicity of *A. annua* oil, 1,8-cineole, and camphor against adult *Tetranychus urticae*, accompanied by inhibition of esterase, GST, and P450 detoxification enzymes (Nasr Azadani et al. 2025). The result strengthens the case for matched oil/monoterpenoid assays while remaining an invertebrate acaricide phenotype rather than parasite-protein engagement. An older *A. annua* larvicide comparison extends this vector-control evidence to *Culex*. Chloroform, methanol, ethanol, and acetone leaf extracts, together with purified artemisinin, killed late-stage larvae of *Culex quinquefasciatus* and *Culex tritaeniorhynchus*; the chloroform extract was more potent than artemisinin in the reported LC50/LC90 comparisons (Sharma et al. 2012). The result supports an extract-versus-compound vector phenotype, but it does not identify a causal terpene, establish malaria treatment, or provide non-target safety. Process chemistry is similarly preparation-specific: pressurized cyclic solid-liquid extraction recovered artemisinin and scopoletin from *A. annua* more efficiently and rapidly than traditional maceration in the tested material (Zarrelli et al. 2019). Higher extraction yield should not be equated with higher biological efficacy or clinical suitability without stability, composition, exposure, and assay validation.

The new Terpedia interaction map separates target engagement from target hypotheses (parasite-protein-interactions.csv; extraction rules in protein-interaction-protocol.md). The most experimentally anchored case is artemisinin against *Plasmodium falciparum*: haem activation supports covalent labeling of a broad parasite protein set, with 124 reported targets and orthogonal validation of selected metabolic and structural proteins, including OAT, PyrK, LDH, SpdSyn, SAMS, plasmepsins, MSP1, and actin (Wang et al. 2015). This is a promiscuous, chemically activated interaction model, not evidence for one universal artemisinin receptor.

To formalize the critical comparison without pretending that heterogeneous assays support a meta-analytic score, the package applies the provisional non-compensatory rubric in evidence-appraisal-protocol.md to all 129 antiparasitic records. Twenty-one records have defined or purified chemistry, 36 have a profiled mixture, and 72 retain crude or unresolved attribution. Eighty-six records report direct in-vitro parasite/vector assays, 39 include an in-vivo or organismal endpoint, and four are indirect or surrogate records, including three hemozoin assays. Explicit host/selectivity controls occur in 22 records, partial controls in 24, and absent or unresolved controls in 83. Three records reach the E4/E5 functional-target-class level through the linked interaction map; 23 are hypothesis/stage-mechanism level and 103 remain phenotype-only or unresolved. The overall appraisal yields 24 B-level characterized phenotypes and 105 C-level phenotypes with unresolved attribution; no record meets the deliberately strict A tier. These are evidence-resolution labels, not efficacy rankings, and the row-level basis is archived in evidence-appraisal.csv.

Resistance should also be modeled as a dynamic parasite state rather than as an intrinsic label on a plant preparation. Intraerythrocytic oxidative-stress conditions significantly reduced artemisinin and dihydroartemisinin efficacy in matched *P. falciparum* assays, while heat shock altered the response to lumefantrine rather than artemisinin (Pires et al. 2024). A complementary chemical-genetic study showed that early unfolded-protein-response signaling and proteasome-mediated protein degradation determine DHA survival, with proteasome perturbations increasing susceptibility independently of *Kelch13* genotype (Rosenthal et al. 2024).

These findings strengthen the parasite-side resistance framework and define a critical experimental interaction for the plant comparison: a chemotype or fraction should be tested under matched parasite redox/proteostasis states before an apparent difference is attributed to terpene composition. Neither study establishes that an *Artemisia* constituent causes or reverses resistance. Two additional primary studies resolve complementary parts of the activated-peroxide mechanism. Under controlled reducing conditions, artemisinin, artesunate, artemether, and dihydroartemisinin inhibited beta-hematin crystallization, a hemozoin surrogate for parasite heme detoxification, with greater potency than most tested quinoline comparators; the tested ACT partner combinations were additive in the reported assays (Ribbiso et al. 2021). Independently, chemoproteomic labeling with artemisinin and synthetic peroxide probes expanded the alkylated-protein set and showed enrichment for redox-related proteins; live glutathione-redox imaging and thiol metabolomics confirmed disruption of parasite redox homeostasis (Siddiqui et al. 2022). Together with the broad 124-target activity-based map, these results support a heme-activated, multi-target model involving heme handling, protein damage, and redox collapse. They do not establish that hemozoin inhibition or any one redox protein is the sole lethal event, nor do they transfer drug-mechanism evidence to a whole-plant *Artemisia* preparation.

Additional malaria studies provide mechanistic anchors at different evidence levels. DHA inhibited immunopurified PfPI3K at nanomolar concentration, and PfK13 resistance biology was linked to altered PfPI3K/PI3P regulation; a mammalian-kinase counter-screen supported parasite selectivity in that experiment (Mbengue et al. 2015). Artemisinin binding to recombinant PfTCTP was mapped by surface plasmon resonance, mass spectrometry, and structural analysis, although the contribution of this interaction to parasite killing remains unresolved (Eichhorn et al. 2013). Functional oocyte and parasite-imaging experiments supported iron-dependent PfATP6 inhibition, but later multi-target evidence argues against treating PfATP6 as the sole receptor (Eckstein-Ludwig et al. 2003). Separately, DHA caused protein damage and compromised parasite proteasome function, producing proteostasis stress (Bridgford et al. 2018). A stage-resolved photoaffinity study reported covalent and noncovalent target classes associated with protein synthesis, glycolysis, and oxidative homeostasis (Gao et al. 2024); cryo-EM and MS-CETSA subsequently showed that artemisinin treatment increases PfRACK1 occupancy on Pf80S ribosomes (Go et al. 2025). Together these records favor a multi-target, stage- and activation-dependent model; they do not make PfPI3K, the proteasome, PfTCTP, PfATP6, or PfRACK1 a sole mechanism.

Computational chemistry adds a hypothesis layer to this malaria mechanism rather than a new level of target proof. Docking and three-dimensional QSAR studies modeled artemisinin-analogue interactions with hemin and related antimalarial activity, while a later DFT/QSAR analysis related electronic descriptors and peroxide reactivity to activity against chloroquine-resistant *P. falciparum* (Cheng et al. 2002; Rajkhowa et al. 2013). These studies can prioritize analogues and testable activation hypotheses, but docking poses, calculated descriptors, and model fit do not establish parasite target engagement, causal heme binding in vivo, or a therapeutic window. They are therefore stored in the KB as E2 computational evidence, separate from direct binding, recombinant-enzyme, and activity-based proteomics records.

The target-class signal is not restricted to *Plasmodium*. In *Toxoplasma gondii*, artemisinin inhibited TgSERCA-associated Ca<sup>2+</sup>-ATPase activity in a heterologous yeast system and triggered calcium-dependent microneme secretion and calcium perturbation in parasites (Nagamune et al. 2007). This supports a conserved parasite calcium-homeostasis vulnerability, but it is functional target-class evidence rather than a purified artemisinin-affinity measurement. On the plant-chemistry side, bioassay-guided fractionation of *A. afra* identified 1 $\alpha$ ,4 $\alpha$ -dihydroxybishopsolicepolide and yomogiartemin as principal gametocytocidal sesquiterpene-lactone leads, including activity against asexual parasites for the former (Moyo et al. 2019). The result provides a stronger constituent-to-stage phenotype bridge than a crude infusion, while leaving parasite target engagement, host selectivity, and in-vivo transmission effects unresolved.

A plant-side perturbation illustrates why trichome chemistry and parasite phenotype must be measured together. In *A. annua*, calcium/magnesium silicate applied at 400 kg ha<sup>-1</sup> increased glandular-trichome size and leaf/infusion artemisinin concentration, but the resulting infusion did not impair *T. gondii* growth more effectively than infusion from untreated plants; higher silicate doses instead reduced artemisinin concentration (Rostkowska et al. 2016). The result is a controlled dissociation between a plant chemical endpoint and a host-cell parasite assay. It supports environment-by-trichome-by-matrix experiments, not a claim that more artemisinin or larger trichomes necessarily produce greater activity in a non-*Plasmodium* system.

Evidence is weaker outside malaria. A 123-compound sesquiterpene docking study proposed *Leishmania* PTR1, carbonic anhydrase, NMT, and trypanothione synthetase as binding hypotheses; xanthanolides 75-76 had the strongest reported predicted PTR1 scores (-10.6 kcal/mol), but this is E2 docking evidence and the compounds are not established as *Artemisia* constituents in that record (Mishra et al. 2014). In *T. brucei*, recombinant-enzyme assays strengthened the case for target-class activity: cnicin inhibited TbPTR1, while cynaropicrin inhibited both TbPTR1 and TbDHFR, with IC<sub>50</sub> values of 12.4 and 7.1  $\mu$ M, respectively; both compounds came from non-*Artemisia* Asteraceae sources (Possart et al. 2021). A *Leishmania major* study provides a more direct *Artemisia*-provenance STL bridge: four purified eudesmane/elemane lactones from *Seriphidium khorassanicum* (syn. *A. khorassanica*) inhibited amastigote-infected macrophages at 4.9-25.3  $\mu$ M, while normal-macrophage CC<sub>50</sub> values were 432.5-620.7  $\mu$ M; the strongest compound had IC<sub>50</sub> 4.9  $\pm$  0.6  $\mu$ M and selectivity index 88.2 (Fattahian et al. 2022/2024). This is defined-compound evidence in a related taxon, not evidence that the same lactones occur in *A. absinthium* or engage a known parasite protein. A cruzain docking paper is retained as *T. cruzi* context only, not as a demonstrated *Artemisia* interaction. A 2026 poultry study proposed ATP synthase and glutathione transferase as camphor docking targets in *Ascaridia galli*, but the same experiment tested

crude *A. herba-alba* and *A. judaica* ethanolic extracts rather than purified camphor; the result is therefore E2 target prioritization, not direct target engagement or terpene causality (Abdelqader et al. 2026). For *H. contortus*, the current *Artemisia* records establish anthelmintic phenotypes but no parasite protein target, making target discovery a high-priority gap rather than an inferred mechanism.


The practical conclusion is that artemisinin supplies a validated parasite-protein interaction framework, while most non-artemisinin *Artemisia* terpene claims still need purified-compound assays, parasite permeability measurements, target-rescue or genetic tests, and host-protein counterscreens. The KB therefore stores interaction mode, evidence tier, compound origin, parasite stage, and inference boundary for every record.

Table 2. Cross-parasite terpene-protein evidence landscape.

| Parasite               | Compound context                                                                    | Protein or complex                                 | Strongest current evidence                                                     | Boundary                                                                                              |
|------------------------|-------------------------------------------------------------------------------------|----------------------------------------------------|--------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------|
| <i>P. falciparum</i>   | Artemisinin/DHA                                                                     | Broad covalent target set                          | E5 activity-based proteomics with selected orthogonal validation               | Probe- and activation-dependent; not all targets are equally causal                                   |
| <i>P. falciparum</i>   | Artemisinin/DHA                                                                     | PfPI3K, PfTCTP, PfATP6, proteasome, PfRACK1-Pf80S  | E3-E5 biochemical, functional, genetic, structural, and complex-level evidence | Supports a multi-target model, not a single universal receptor                                        |
| <i>P. falciparum</i>   | Artemisinin-family drugs                                                            | Heme-to-hemozoin detoxification pathway            | E4 reducing-condition beta-hematin inhibition and drug-pair assays             | Pathway-level evidence; not a single protein target or whole-plant effect                             |
| <i>P. falciparum</i>   | Artemisinin and peroxide probes                                                     | Redox-related proteins and glutathione homeostasis | E4 chemoproteomics, live redox biosensor, and thiol metabolomics               | Enrichment and redox collapse do not establish one direct causal protein                              |
| <i>Leishmania</i> spp. | Sesquiterpene-related panel                                                         | PTR1, carbonic anhydrase, NMT, TryS                | E2 docking                                                                     | Requires direct binding, enzyme inhibition, target engagement, and <i>Artemisia</i> provenance        |
| <i>T. brucei</i>       | Cnicin and cynaropicrin                                                             | TbPTR1 and TbDHFR                                  | E4 recombinant-enzyme inhibition plus docking                                  | Compounds were sourced from non- <i>Artemisia</i> Asteraceae; parasite target engagement remains open |
| <i>T. cruzi</i>        | Terpenoid panel                                                                     | Cruzain                                            | E2 docking with prior inhibition context                                       | Not an <i>Artemisia</i> -matched direct interaction experiment                                        |
| <i>A. galli</i>        | Crude <i>A. herba-alba</i> / <i>A. judaica</i> extracts; camphor docking hypothesis | ATP synthase; glutathione transferase              | E2 docking paired with in-vitro/in-vivo extract phenotype                      | Camphor abundance, direct binding, and causal contribution remain untested                            |
| <i>Leishmania</i> spp. | <i>A. absinthium</i> phenolic profile; apigenin                                     | N-myristoyltransferase                             | E2 docking plus HPLC context                                                   | Apigenin is non-terpene; no direct binding, enzyme inhibition, or parasite efficacy was shown         |
| <i>T. cruzi</i>        | Dehydroleucodine from <i>A. douglassiana</i>                                        | Unresolved                                         | E3 defined-compound stage-resolved phenotype                                   | Programmed-cell-death readouts do not establish a protein target or host selectivity                  |

| Parasite            | Compound context                                                | Protein or complex | Strongest current evidence | Boundary                                                       |
|---------------------|-----------------------------------------------------------------|--------------------|----------------------------|----------------------------------------------------------------|
| <i>H. contortus</i> | <i>Artemisia</i> oils, extracts, and one isolated sesquiterpene | Unresolved         | E1 phenotype               | Target discovery and host-selectivity experiments are required |

Figure 2. Evidence ladder used for terpene-parasite target claims.



The arrows indicate increasing evidentiary requirements, not an automatic promotion path. A docking result advances only when the same compound-protein pair is tested experimentally, and even direct binding does not establish that the interaction is necessary or sufficient for parasite killing.

Published essential-oil studies provide useful phenotype anchors. *A. annua* leaf oil has been tested against *Leishmania donovani* in in-vitro and in-vivo contexts. Oils from *A. campestris* and *A. herba-alba* have been tested against *L. infantum* promastigotes, with apoptosis-like and cell-cycle-associated effects reported. These results support mixture-level activity under the reported experimental conditions. They do not establish a single active terpene, whole-life-cycle efficacy, host safety, or clinical benefit.

Whole-leaf and aqueous-extract studies occupy a different evidentiary category from essential oils. *A. annua* leaf powder was evaluated against *Leishmania* in cell and animal models, but the complex material prevents attribution to artemisinin or another terpene (Mesa et al. 2017). *A. nilagirica* extracts were screened against *P. falciparum*, again providing a species- and preparation-specific phenotype rather than a defined terpene mechanism (Panda et al. 2018). Most recently, aqueous *A. afra* and *A. annua* extracts were compared against artemisinin-resistant *P. falciparum* with selectivity measurements (Roesch et al. 2025). This resistant-parasite design is valuable, but activity of a chemically complex infusion cannot be projected onto an isolated constituent or clinical regimen.

The quantitative assay records are separated in `antiparasitic-evidence.csv` so that parasite stage, preparation, endpoint, dose, host control, and translation boundary remain visible. For example, the camphor-rich *A. annua* oil reported IC50 values of 14.63 +/- 1.49 microgram/mL against promastigotes and 7.3 +/- 1.85 microgram/mL against intracellular amastigotes, plus an approximately 90% liver/spleen burden reduction at 200 mg/kg in infected mice. Those observations belong to one hydrodistilled oil and its experimental model; they are not evidence that camphor alone, or *A. annua* generally, produces the effect.

An Ethiopian study of *A. abyssinica* further illustrates why selectivity must be reported with activity: its oxygenated-monoterpene-rich oil showed activity against *Leishmania* promastigote and amastigote systems but also variable toxicity in mammalian-cell and erythrocyte assays. This is a valuable comparative record because it carries parasite stage and host-toxicity information together, although it still does not resolve which constituents or interactions drive the result.

The broader antiparasitic record is heterogeneous but not limited to *Leishmania*. *A. campestris* oil has been tested against *Haemonchus contortus* eggs and adults and against *Heligmosomoides polygyrus* in a nematode model (Abidi et al. 2018); *A. vestita* and *A. maritima* extracts have been tested against *H. contortus* in vitro and in infected sheep (PubMed). A hydrodistilled *A. gilvescens* oil and selected constituents have also shown larvicidal activity against *Anopheles anthropophagus* (PubMed). Seven geographic *A. argyi* oils provide a population-level vector comparison against *Anopheles sinensis*, but the shared and variable constituents remain mixture-level predictors rather than identified actives (Luo et al. 2022). These studies widen the phenotype space to helminths and vectors, but they also emphasize that extract polarity, dose, life stage, host model, and formulation prevent direct pooling. An Ethiopian extract comparison against bloodstream *Trypanosoma brucei* adds another protozoan anchor while demonstrating why cytotoxicity and extract composition must accompany activity (PubMed).

The 2018 *A. campestris* oil study provides a stronger quantitative, multi-stage comparator than the discovery-level summary alone. GC/MS identified 50 compounds representing 99.98% of the oil, dominated by beta-pinene (36.40%) and 2-undecanone (14.7%). The oil produced complete *H. contortus* egg-hatch inhibition at 2 mg/mL after 48 hours (IC50 0.93 mg/mL), 66.6% adult-worm motility inhibition at 0.5 mg/mL after 8 hours, and, in *H. polygyrus*, 72.1% fecal-egg-count and 72% total-worm-count reduction at 5000 mg/kg seven days after treatment. These endpoints support a preparation-specific, multi-stage veterinary signal; they do not establish beta-pinene as causal, and 2-undecanone is not a terpene. Publisher-indexed methods and results were recoverable, but complete tables and several design details were not; the record therefore remains quantitatively unresolved for pooling (Abidi et al. 2018; CGIAR repository record).

Tick and insect studies also show why formulation and host counterscreens must remain attached to the evidence record. An *A. herba-alba* oil nanoemulsion showed the lowest reported LC50 for tick eggs and nymphs among the tested *Hyalomma dromedarii* stages, but oral exposure in mice was accompanied by liver and kidney histopathological changes despite largely unremarkable blood chemistry (Abdel-Ghany et al. 2021). The formulation therefore supplies a useful stage-specific acaricide phenotype and a cautionary host signal, not proof that a native oil or one terpene is safe. Together with altitude-associated *A. scoparia* mosquito activity and the *A. vulgaris* gall-mite assay described above, it supports a comparative design in which oil composition, formulation, exposure route, stage, non-target toxicity, and parasite relevance are modeled separately.

Within helminth research, defined-compound and whole-preparation studies can be compared without treating them as equivalent. Artemisinin and *Artemisia* extracts were tested against *H. contortus* in a gerbil model, directly exposing the distinction between purified compound and botanical matrix (Squires et al. 2011). *A. lancea* essential oil was tested against egg and larval stages in vitro (Zhu et al. 2013), whereas bio-guided *A. cina* isolation later produced a defined sesquiterpene lead for L3 larvae. These assays suggest tractable phenotypes for fractionation and target discovery, but host pharmacokinetics, adult-worm activity, and safety remain separate questions.

An additional *A. herba-alba* tick study tested Libyan leaf oil against *Ixodes ricinus* nymphs using open-filter-paper exposure. At 1 microliter/cm<sup>2</sup>, complete mortality was reported within two hours, whereas lower exposure produced slower or incomplete effects (Elmhalli et al. 2021). This is a useful vector-borne-disease comparator for the wormwood evidence stream, but it remains a hydrodistilled mixture assay without purified-constituent attribution, mammalian safety data, or evidence against gastrointestinal helminths.

Historical structure-activity work on artemisinin and related sesquiterpenes in rodent malaria already showed that close chemical relatives can differ substantially in activity (Peters et al. 1986). That observation remains relevant to comparative genomics: detecting a pathway-related scaffold or ADS-like sequence does not establish an artemisinin-equivalent phenotype. Structure, peroxide status, stereochemistry, exposure, and activation environment all matter.

Two additional high-priority records sharpen the chemistry-phenotype boundary. In *A. indica*, a hydrodistilled oil contained 92 detected compounds, with camphor (13.0%) and caryophyllene oxide (10.87%) as major constituents; the oil and solvent extracts showed activity across *Plasmodium falciparum*, *Trypanosoma brucei*, and *T. cruzi* systems, with several preparations showing low mammalian cytotoxicity (Tasdemir et al. 2015). The study is valuable because chemistry, parasite panels, and cytotoxicity were measured together, but it remains a mixture/extract study. A separate activity-guided *A. indica* study isolated exiguaflavones A and B, maackiain, and a benzofuran; the reported antimalarial principles were non-terpene compounds, so the record is retained as a boundary control rather than evidence for terpene activity (Chanphen et al. 1998). In *A. scoparia*, a BA plus 2,4-D callus extract inhibited *Leishmania tropica* promastigotes with an IC<sub>50</sub> of 19.13 mg/mL and produced apoptotic morphology; GC-MS reported nerolidol alongside non-terpene metabolites (Yousaf et al. 2019). This is evidence for a tissue-culture-derived crude extract, not proof that nerolidol or any single terpene accounts for the phenotype.

Full-text extraction of the *A. indica* boundary control makes its limits explicit. Authenticated stems (herbarium BKF 82247) were extracted with n-hexane and methanol, and silica-guided fractionation yielded exiguafavanones A/B, maackiain, and a benzofuran at 0.0051%, 0.0084%, 0.002%, and 0.0005% of dried plant, respectively. In a continuous asexual erythrocytic K1 multidrug-resistant *P. falciparum* assay, the methanol extract had EC<sub>50</sub> 6.60 x 10<sup>-6</sup> g/mL; exiguafavanones A and B measured 4.60 x 10<sup>-6</sup> and 7.05 x 10<sup>-6</sup> g/mL, while maackiain and the benzofuran were weaker. This is a defined non-terpene malaria control with authenticated material and fraction-to-compound tracking, not evidence for a terpene/STL mechanism, mammalian selectivity, or transfer to *A. annua* (Chanphen et al. 1998).

The strongest causal bridge would require matched specimen chemistry, parasite-stage assays, host-cell cytotoxicity, fractionation or defined mixtures, and ideally purified-compound plus combination testing. Gene expression or a predicted pathway can prioritize candidates but cannot replace those experiments.

Additional species and preparation types reinforce this boundary. *A. pedemontana* subsp. *assoana* essential oils and hydrolate were evaluated across insect, nematode, protozoan, and antiparasitoid assays, making it a useful multi-phenotype specimen record while retaining oil and hydrolate as distinct preparations (Sainz et al. 2019). Isolated *A. sieversiana* sesquiterpenoids provide a compound-level trypanocidal lead set (Nurbek et al. 2020), whereas *A. absinthium* fractionation studies link selected oil fractions to *T. cruzi* and *T. vaginalis* activity with cytotoxicity measurements (Martínez-Díaz et al. 2015). These records increase taxonomic and mechanistic coverage, but they do not make mixture, purified-compound, or host-selectivity endpoints interchangeable. A bio-guided *A. cina* study adds a stronger constituent-to-phenotype bridge by isolating a sesquiterpene active against *Haemonchus contortus* L3 larvae (Arango-De la Pava et al. 2024); an *A. brevifolia* study extends the nematode comparison across life stages (Hussain et al. 2025). Both still require host-selectivity and in-vivo confirmation.

Purified-compound evidence provides an important contrast to oil studies: helenalin, mexicanin, and dehydroleucodine inhibited cultured *Leishmania mexicana* promastigotes at approximately low-micromolar concentrations, but that result cannot be assumed for an *Artemisia* oil or a different parasite stage (Barrera et al. 2008). Likewise, methanolic aerial-part extracts of *A. sieberi* and *A. khorassanica* have been evaluated in *Plasmodium berghei* mouse models (PubMed; PubMed). Full-text analysis of the flowering-stage *A. khorassanica* study found that 650 g of authenticated Iranian aerial material was macerated in methanol for 48-72 hours, defatted, diluted, and partitioned with n-hexane; the non-polar fraction contained 29.9% monoterpenes and 6.2% sesquiterpenes, with chrysanthenone at 7.8% and cis-thujone at 5.8%. Infected NMRI mice received 200 microliter of a 100 mg/mL crude-extract solution subcutaneously once daily for three weeks, and microscopy showed a limited but statistically reported reduction in *P. berghei* parasitemia. Because the study did not quantify artemisinin, provide a parasite-stage-resolved potency estimate, or isolate a causal volatile, its chemistry is a composition marker and its biology is a species- and preparation-specific murine result—not evidence for an *A. annua*-equivalent pathway, a single-terpene mechanism, or human efficacy (Nahrevanian et al. 2010).

Isolation and fractionation can provide a stronger causal bridge than crude-oil screening. Leaves and flowers of Cape Verdean *A. gorgonum* yielded guaianolides and germacranolides; ridentin had a *P. falciparum* FcB1 IC<sub>50</sub> of 3.8 +/- 0.7 microgram/mL against a Vero-cell IC<sub>50</sub> of 71.0 +/- 3.9 microgram/mL (Ortet et al. 2008). Conversely, a Cuban *A. absinthium* oil dominated by trans-sabinyol acetate inhibited *L. amazonensis* promastigotes and amastigotes and reduced lesion and parasite burden after intralesional dosing in mice (Monzote et al. 2014). The contrast is informative: purified-lactone selectivity and oil-level in-vivo activity are both stronger than an uncharacterized extract claim, but neither supports extrapolation to all *Artemisia* preparations. A separate *A. absinthium* oil study tested the oil and three major constituents against six mosquito vectors while measuring acute toxicity in non-target aquatic organisms (Govindarajan and Benelli 2016); the oil contained (E)-beta-farnesene, (Z)-en-yn-dicycloether, and (Z)-beta-ocimene as major constituents, and their larvicidal potency varied by mosquito species. This is valuable paired efficacy-ecotoxicity and constituent-resolved evidence, but still preparation- and dose-specific.

Phylogeny can improve discovery prioritization without replacing chemical validation. Sampling of 117 ethnobotanically reported species detected artemisinin in nine species, including eight reported for the first time, but detection does not establish abundance, pathway completeness, or therapeutic equivalence (Pellicer et al. 2018). This finding supports a comparative design in which phylogenetic position selects accessions for targeted LC-MS and pathway-genomics follow-up rather than serving as a proxy for efficacy.

## 6.2 Malaria: artemisinin as the primary validated case

The modern review literature frames artemisinin as a defined endoperoxide drug rather than as a generic synonym for *Artemisia* material. A 2021 review places the 1,2,4-endoperoxide bridge, artesunate development, and Kelch13-associated resistance in the clinical-drug context (Khanal 2021). We use this review for scope and mechanism framing; quantitative claims about plant chemistry, parasite stages, pharmacokinetics, or safety remain anchored to primary records.

A 2025 plant-matrix study adds a useful preclinical malaria comparator: a petroleum-ether/silica *A. annua* extract quantified artemisinin, deoxyartemisinin, arteannuin I, dihydroartemisinic acid, and artemisinic acid as a sesquiterpene-rich mixture. At the lowest matched artemisinin dose, the extract produced approximately ten-fold greater inhibition than pure artemisinin in two murine models; rat pharmacokinetics showed roughly two-fold higher artemisinin AUC and 1.6-fold longer half-life, while Caco-2 experiments implicated reduced P-glycoprotein efflux and effects from deoxyartemisinin, artemisinic acid, and dihydroartemisinic acid (Xu et al. 2025). This is matrix and host-transporter evidence, not human efficacy, clinical equivalence, a single active terpene, or direct parasite-protein engagement. A 2026 UHPLC-QTOF-MS/MS study expands this matrix hypothesis with a broader metabolomic screen: 103 constituents were characterized in *A. annua*, including 31 sesquiterpenes, and seven April-enriched candidates were associated with antimalarial activity; compounds 69 and 90 were reported to potentiate artemisinin in the tested system (Zhang et al. 2026). The seasonal contrast is informative because April samples showed greater activity despite lower artemisinin than May/June samples, but the record does not by itself establish the identities, concentrations, mechanism, or reproducibility of the candidate enhancers. It supports a pre-registered fractionation and reconstitution experiment, not a universal harvest recommendation or a terpene-only synergy claim. Two complementary metabolomics studies make the matrix hypothesis more testable. A UPLC-Q-TOF-MS/MS study of wild and cultivated *A. annua* identified 24 flavonoids, 68 sesquiterpenoids, and 21 other compounds, while multivariate analysis and fractionation nominated 5 $\alpha$ -hydroperoxy-eudesma-4(15),11-diene and arteannuin B-related chemistry as candidate contributors to differential murine activity (Shi et al. 2022). A 2025 comparison of teas prepared from ten Benin plantations and French material found artemisinin concentrations of 0.3–15.7 mg/L, significant one-year losses in three Benin samples, and LC-HRMS features—including unknown alkaloids—associated with *P. falciparum* activity (Aubouy et al. 2025). Together, these studies support origin-, storage-, and matrix-dependent prioritization, but correlation and candidate annotation are not proof of a terpene-specific mechanism, standardized product performance, or clinical efficacy. The transmission-stage comparator is chemically close to, but not derived from, *Artemisia*. Purified parthenin and parthenolide inhibited *P. falciparum* blood-stage development, stage V gametocyte exflagellation, and mosquito-stage transitions including ookinete-to-oocyst development; parthenolide showed similar activity in the reported assay battery (Balaich et al. 2016). Because these lactones are non-*Artemisia* comparators and parthenin has cytotoxicity concerns, the result should be used to motivate stage-resolved screening of authenticated *Artemisia* sesquiterpene lactones, not to assign transmission-blocking activity to wormwood or another genus member. A full-text extraction makes this comparator experimentally useful: after a 24-hour exposure and washout, parthenin eliminated mosquito oocysts at 1,000 ng/mL and reduced median oocysts by 83% at 100 ng/mL, whereas parthenolide still reduced median oocysts by 47% at 10 ng/mL. Parthenin also reduced exflagellation and blocked *P. berghei* ookinete maturation in stage-timed assays. These are purified non-*Artemisia* STL-to-stage results with solvent, washout, microscopy, and mosquito-feed controls; the study did not establish a mammalian therapeutic window or a single parasite protein target. They therefore define a transmission-stage assay benchmark and an artemisinin-independent hypothesis, not evidence for wormwood efficacy (Balaich et al. 2016). A matched origin comparison reinforces the same boundary. Water infusions from Cameroon and Luxembourg *A. annua* plants differed in HPLC profiles and beta-haematin inhibition; the Cameroon material showed stronger inhibition at the reported dilution and more major chromatographic features (Akkawi et al. 2014). Because the study used a hemozoin surrogate and preliminary chromatographic assignments rather than a purified terpene or a clinical endpoint, it supports origin- and matrix-dependent malaria chemistry, not a universal chemotype or treatment equivalence.

Malaria should be the anchor phenotype for the evolutionary-genomics component because it links a defined sesquiterpene lactone to a validated plant pathway and a clinically important parasite. The artemisinin scaffold contains a peroxide bridge whose activation is central to parasite damage, and the *A. annua* ADS-CYP71AV1-DBR2-ALDH1 network provides the best-resolved plant-side genotype-to-metabolite framework in the genus. Historical structure-activity work showed that related sesquiterpenes are not functionally interchangeable, while modern target studies support activation-dependent, multi-target parasite injury rather than one universal receptor (Peters et al. 1986; Wang et al. 2015).

The non-malaria protozoan evidence includes a host-response-mediated veterinary coccidiosis study. In *Eimeria tenella*-infected chickens, artemisinin and dried *A. annua* leaves reversed infection-associated apoptosis markers and reduced inflammatory-response signals; leaf treatment produced stronger clinical and pathological improvement than artemisinin in the reported comparison (Jiao et al. 2018). Because the readouts were cecal host responses and the leaf preparation was chemically complex, this record supports a veterinary comparator and a testable host-mediated mechanism, not a direct *Eimeria* protein interaction or evidence that the whole leaf is interchangeable with an artemisinin drug.

Mechanistic uncertainty is not a minor footnote. A dedicated review of carbon-radical and heme hypotheses emphasized that artemisinin activation, iron/heme chemistry, parasite stage, metabolism, and combination partners all influence interpretation, while competing hypotheses cannot be collapsed into a single receptor model (Haynes et al. 2013). In parallel, microbial transformation of artemisinin and related terpenoids has been used to generate derivative candidates for medicinal chemistry (Parshikov



et al. 2012). These transformed products are valuable experimental probes, but neither their activity nor their pharmacokinetics can be assigned to native *Artemisia* chemistry without direct measurement.

The clinical boundary must remain explicit. WHO guidance supports quality-assured artemisinin-based combination therapy for malaria; it does not validate arbitrary teas, capsules, essential oils, or unstandardized *Artemisia* materials. Whole-plant experiments remain scientifically valuable for understanding matrix effects and resistance evolution. In rodent models, dried whole-plant *A. annua* overcame selected artemisinin resistance and delayed resistance acquisition relative to pure artemisinin (Elfawal et al. 2015). A historical PubMed-indexed report of an *A. annua* gelatin-capsule preparation likewise described parasitemia clearance in *P. berghei* and reported clinical *P. vivax* cure with high recrudescence, but its crude preparation, limited modern metadata, and absence of constituent attribution make it historical context rather than current efficacy evidence (Wan et al. 1992). These records motivate fractionation, exposure, and resistance-mechanism studies; they do not establish human efficacy, dosing, quality control, or safety of a nonpharmaceutical whole-plant product.

The resistance result is especially informative because it was tested against both an existing resistant parasite and experimentally evolving populations. Dried whole plant overcame selected *P. yoelii* resistance to pure artemisinin, and stable resistance arose three times more slowly under whole-plant than pure-artemisinin selection in *P. chabaudi* (Elfawal et al. 2015). The authors' redundant-defense interpretation is biologically plausible, but the experiment does not identify whether the relevant co-components are sesquiterpenes, flavonoids, volatile constituents, altered absorption, or their combination. Older mouse isobolograms make the same interaction principle explicit: artemisinin was potentiated by some partners, including mefloquine, tetracycline, spiramycin, and selected primaquine or clindamycin contexts, but antagonism occurred with several sulfonamide/pyrimethamine combinations and the result changed with parasite resistance strain (Chawira et al. 1987). Thus, plant-matrix and drug-combination effects should be modeled as partner-, strain-, dose-, and exposure-specific rather than summarized as generic synergy.

The matrix hypothesis has a useful non-terpene comparator. A critical review proposed that *A. annua* leaf flavonoids might influence artemisinin activity, absorption, CYP-mediated metabolism, and host immunomodulation, but emphasized that only a limited number of matched synergy studies existed (Ferreira et al. 2010). Endophyte-associated chemistry provides a separate discovery route: metabolomics-guided screening of fungi from *A. annua* identified emodin and physcion as validated antimalarial candidates with IC<sub>50</sub> values of 0.9 and 1.9 micromolar, respectively (Alhadrami et al. 2021). Both records are scientifically relevant but non-terpene: neither demonstrates that a plant volatile or sesquiterpene lactone drives whole-leaf activity. They instead motivate matched fractionation, host-exposure, and parasite-stage experiments that can distinguish plant terpenes from co-extracted or microbially derived chemistry.

Historical reviews and production analyses explain why the quality boundary is central to malaria translation. Early syntheses described plant production and biotechnology as ways to increase artemisinin supply (Van Geldre et al. 1997; Segura 2010), while a later development review emphasized crop quality, formulation, bioequivalence, and combination validation as recurrent failure points (Jambou et al. 2011). A historical Myanmar report is retained as a title-level record of an extract antimalarial trial, but its full text must be recovered before any preparation, dose, or efficacy value is extracted (Lwin et al. 1991). These sources support a reproducibility and stewardship framework, not a clinical endorsement of unstandardized extracts.

That stewardship concern has also been raised specifically for unapproved *A. annua* use in malaria prevention. A risk-focused review identified uncontrolled exposure as a potential contributor to artemisinin drug pressure and future resistance, but it did not measure resistance evolution in a defined plant product (Ataba et al. 2022). The inference is therefore precautionary and complementary to the experimental whole-plant selection result: it strengthens the case for quality-assured ACTs, surveillance, and controlled comparative studies rather than treating a tea or capsule as a resistance-resilient therapy.

Non-*annua* malaria studies are useful as bounded comparators rather than substitutes for the *A. annua* pathway. Iranian *A. sieberi* aerial-part extracts reduced *P. berghei* parasitemia in mice in a study that reported no toxicity in the tested NMRI-mouse assessment, while a flowering-stage *A. khorassanica* extract showed a more limited response and contained chrysanthemone and cis-thujone among GC-MS features (Nahrevanian et al. 2012; Nahrevanian et al. 2010). These records strengthen the comparative malaria evidence base, but both are crude, species-specific preparations: detection of a volatile, absence of short-term toxicity, or reduction of murine parasitemia does not establish a shared artemisinin pathway, a single active terpene, or human antimalarial efficacy. Earlier screening studies add two useful non-*annua* boundaries. Dichloromethane extracts of *A. ciniformis*, *A. biennis*, and *A. turanica* inhibited beta-hematin formation, with *A. ciniformis* most active at an IC<sub>50</sub> of 0.92 +/- 0.01 mg/mL; this is a hemozoin-surrogate screen rather than a parasite-stage result (Mojarrab et al. 2015). In a separate fractionation study, *A. armeniaca* and *A. aucheri* dichloromethane extracts were active in the same surrogate assay, and a fraction of *A. armeniaca* reached an IC<sub>50</sub> of 0.47 +/- 0.006 mg/mL (Mojarrab et al. 2014). These records prioritize medium-polar extracts for chemical follow-up but do not identify a terpene, establish blood-stage efficacy, or support clinical translation. Full-text verification adds the preparation and assay boundaries behind the first result. Authenticated Iranian aerial parts were successively macerated with petroleum ether, dichloromethane, ethyl acetate, ethanol, and 1:1 ethanol-water, with three vouchers and extraction below 45 degrees C. The assay used synthetic beta-hematin formation rather than parasite culture; *A. ciniformis* dichloromethane extract had IC<sub>50</sub>/IC<sub>90</sub> values of 0.92 +/- 0.01/1.29 +/- 0.02 mg/mL, while ethyl-acetate extracts of *A. biennis* and *A. turanica* had IC<sub>50</sub> values of 1.11 +/- 0.02 and 1.35 +/- 0.08 mg/mL. The authors' suggestion that lactone-rich *A. ciniformis* chemistry contributed is not supported by terpene-resolved analysis, and lipid or matrix interference produced negative inhibition in some weak extracts. This is therefore

an extraction-polarity and hemozoin-surrogate prioritization result, not evidence for a defined sesquiterpene-lactone, blood-stage activity, host selectivity, or clinical efficacy (Mojarrab et al. 2015).

The newly verified *A. armeniaca*/*A. aucheri* comparison provides a second, chemically unresolved malaria-surrogate anchor. Voucher-linked aerial parts were successively extracted across petroleum ether, dichloromethane, ethyl acetate, ethanol, and 1:1 ethanol-water, and the dichloromethane extracts were fractionated by silica vacuum-liquid chromatography. In synthetic beta-hematin formation, dichloromethane extracts reached IC<sub>50</sub>/IC<sub>90</sub> values of 1.36/2.12 mg/mL for *A. armeniaca* and 1.83/2.62 mg/mL for *A. aucheri*, while the strongest *A. armeniaca* fraction reached 0.47/0.71 mg/mL; ethyl acetate, ethanol, and hydroethanol parent extracts were inactive. TLC detected mixed terpenes, fatty acids, methylated coumarins, and methoxylated flavonoids, but no compound-resolved terpene, artemisinin, or sesquiterpene-lactone measurement was reported. Because the assay was synthetic beta-hematin rather than parasite culture, this result prioritizes extraction polarity and fractionation for follow-up without establishing blood-stage efficacy or a terpene mechanism (Mojarrab et al. 2014).

An extract-versus-compound comparison in *A. sieberi* is more biologically direct but still bounded. Both an aqueous plant extract and purified artemisinin reduced *Leishmania major* promastigote and intracellular-amastigote burdens in vitro, with the plant extract reported as more active in that experiment and low effects on uninfected macrophages (Heydari et al. 2013). The result supports a preparation-specific protozoan phenotype and a testable extract/compound comparison; it does not identify the active constituent, establish a parasite protein target, or demonstrate in-vivo efficacy. Together with the *P. berghei* records, it shows why non-*annua* activity should be retained as comparative evidence without being collapsed into artemisinin-pathway equivalence. Formulation evidence provides a further boundary between plant chemistry and parasite phenotype. Green silver nanoparticles synthesized with *A. abrotanum* or *A. arborescens* extracts were characterized as sub-50-nm particles and tested against *P. falciparum* cultures; the *A. abrotanum*-derived particles showed lower parasitemia and lower hemolysis than the comparison preparations in the reported dose range (Avitabile et al. 2020). This is a formulation-level antiparasmodial result, not evidence that a terpene is the active principle or that the particles are clinically safe. Conversely, dry leaves, artemisinin, and several *A. annua* extracts that were active against *Histomonas meleagridis* clones in vitro failed to improve experimental histomonosis in turkeys or chickens (Thøfner et al. 2012). The paired records make in-vitro/in-vivo discordance an explicit translation variable rather than an unexplained exception.

For the comparative study, malaria-related claims should be separated into three levels: (1) *A. annua* artemisinin production and parasite mechanism, which have the strongest evidence; (2) detection of artemisinin or related lactones in another *Artemisia* species, which supports chemical occurrence only; and (3) an *Artemisia* extract phenotype against *Plasmodium*, which supports a preparation-level assay result. A species can satisfy level 2 or 3 without possessing the same pathway flux, peroxide abundance, pharmacokinetics, or resistance profile as *A. annua*. A useful negative control is the matched *A. annua*/artemisinin study in which artemisinin, artesunate, and dried-leaf slurry failed to reduce *Babesia microti* parasitemia in mice at the tested 100 microgram artemisinin/kg regimen, and activity against six *Candida* species was not sustained or was essentially absent up to 180 micromolar (Elfawal et al. 2021). This does not weaken the *Plasmodium* evidence; it demonstrates that parasite identity, stage, dose, and host/model context are causal design variables. This tiering is central to the proposed orthology and metabolomics workflow.

Recent cross-species measurements illustrate why the second level must remain separate from therapeutic equivalence. In *A. tournefortiana*, RP-HPLC-PDA measured artemisinin most strongly in flowers, reaching 1.43%, with flower values ranging from 0.05% to 1.43% across geographic sites; roots and seeds contained lower amounts. Comparative sequencing of pathway-associated genes supported biosynthetic potential (Sahoo et al. 2026). These data make *A. tournefortiana* a useful accession- and tissue-stratified comparator, not a demonstrated substitute for *A. annua*. Two earlier regional surveys widen the occurrence map but reinforce its limitations. A Tajikistan collection contained HPLC-quantified artemisinin at 0.07-0.45% dry mass in seven sampled species, including *A. vachanica* as a proposed new source (Numonov et al. 2019). A northern Pakistan survey compared flowers, leaves, roots, and stems from *A. annua* and fourteen other species; *A. annua* leaves and flowers were highest in that dataset, while several other species showed lower but detectable levels (Mannan et al. 2010). These data support phylogenetically distributed, tissue-resolved follow-up, but occurrence in a hexane extract or HPLC peak does not prove an intact ADS-CYP71AV1-DBR2-ALDH1 pathway, stable field chemotype, or antimalarial equivalence.

Whole-leaf mutant experiments refine the frequently proposed “artemisinin plus flavonoid synergy” explanation. A *CH11* mutant that lacked major flavonoids but retained wild-type artemisinin levels showed antiparasmodial activity comparable to wild type, whereas CYP71AV1- and ADS-impaired lines with severely reduced artemisinin biosynthesis showed little or no activity against intraerythrocytic *P. falciparum* Dd2 (Czechowski et al. 2019). In that matched genetic system, flavonoids were not a major independent driver of rapid in-vitro activity. This does not eliminate matrix effects in other cultivars, extraction methods, parasite strains, or exposure windows; it establishes a stronger causal comparator than correlation of unrelated metabolites.

Environment can also move both chemistry and apparent antimalarial potency. Under controlled LED treatments, white and blue light increased artemisinin and artemisinic acid relative to red light and shifted mono-, sesqui-, di-, and triterpene profiles. Crude-leaf extract IC<sub>50</sub> values against *P. falciparum* NF54 were 0.51 and 0.52 microgram/mL under white and blue light, 1.11 microgram/mL under red light, and 2.23 microgram/mL for greenhouse-grown plants (Sankhuan et al. 2022). Because the experiment changed multiple metabolites together, the result supports an environment-chemotype-phenotype relationship rather than a causal assignment to germacrene D, trans-beta-farnesene, artemisinin, or any other correlated compound.

Recent matrix studies sharpen, rather than settle, the synergy question. A dedicated review of isolated *A. annua* and *A. afra* metabolites explicitly separates direct antiplasmodial compounds from metabolites affecting PK/PD, inflammation, immunomodulation, and toxicity (Shinyuy et al. 2023). An *A. annua* extract tested with equivalent artemisinin showed no synergy against *P. falciparum* Pf3D7 at 150 nM, but produced a biphasic effect in *P. yoelii* mice, including a lower ED90 and higher artemisinin exposure at an early infection burden; the authors identified chrysosplenetin as a candidate matrix component through LC-HRMS and fractionation (Xu et al. 2026). Chrysosplenetin is a polymethoxy flavonol, not a terpene, and the result is therefore a matrix/pharmacokinetic comparator rather than evidence for a terpene-protein interaction. It also reinforces the need to model infection stage, host metabolism, and extract standardization when interpreting whole-plant malaria data. A separate defined-component mouse study combined artemisinin with arteannuin B, arteannuic acid, and scopoletin; reported parasitemia reduction was about 93% for the combination versus about 31% for pure artemisinin, and repeated-dose pharmacokinetics showed higher artemisinin exposure (Li et al. 2018). This is a useful preclinical combination anchor, but scopoletin is non-terpene and the result cannot be used to assign synergy to any single constituent or to justify unstandardized botanical treatment.

A full-text-indexed top-down PD-PK study sharpened the sesquiterpene matrix hypothesis. Ethyl-acetate and petroleum-ether extracts of dried *A. annua* were compared with pure artemisinin in *P. yoelii* mice and in a *P. falciparum* in-vitro assay. Reported ED50 values were 5.6 mg/kg for pure artemisinin, 2.9 mg/kg for the ethyl-acetate extract, and 1.6 mg/kg for the petroleum-ether extract, while the extracts did not improve the in-vitro IC50 range of 15.0–20.0 nM. Petroleum ether increased rat artemisinin AUC0–t 1.7-fold, and LC-HRMS-guided isolation identified arteannuin B and artemisinic acid as differential sesquiterpenes (3.0% and 14.1% in petroleum ether versus 0.9% and 5.1% in ethyl acetate). Same-proportion AB/AA testing supports an absorption-linked matrix hypothesis, but not a complete single-component causal claim, a direct parasite target, or clinical equivalence of an extract (Cai et al. 2024).

Independent metabolomics supports a species-specific malaria boundary. *A. afra* extracts contained only trace artemisinin and produced a parasite metabolic response distinct from *A. annua*; the *A. annua* response more closely resembled purified artemisinin and included glutathione-associated changes, whereas *A. afra* affected lipid precursors (Mamede et al. 2025). This result is valuable because it separates a non-artemisinin *Artemisia* phenotype from the artemisinin-centered *A. annua* comparator, but it was based on aqueous extracts and metabolomics rather than a terpene-specific perturbation, direct target engagement, or a clinical endpoint.

Full-text extraction makes that contrast more quantitative. Botanist-authenticated flowering leaves and twigs from the same Cameroon locality were extracted with acetone; *A. annua* dry extract contained 19.3 mg/g artemisinin, while *A. afra* was below the reported HPLC-UV detection limit. LDH assays gave IC50 values of 0.09 +/- 0.01 microgram/mL for *A. annua*, 11.73 +/- 2.15 microgram/mL for *A. afra*, and 5.58 +/- 1.14 ng/mL for artemisinin. At 10x IC50, late-stage trophozoite HPLC-MS and NMR profiles placed *A. annua* close to artemisinin, with reduced hemoglobin-derived peptides and glutathione-associated changes, whereas *A. afra* altered folate, carbohydrate, amino-acid, phospholipid, and lipid-precursor metabolism. Because phenolics and other matrix constituents were not fractionated or reconstituted, this is a species- and preparation-specific metabolomic distinction, not proof of a non-artemisinin terpene mechanism (Mamede et al. 2025).

A full-text postharvest chemistry control further shows why a malaria-associated metabolomic feature cannot automatically be treated as an active terpene. Voucher-linked Beninese leaves and stems yielded the previously undescribed cadinapyridine sesquiterpene alkaloids annuanines A and B, with annuanine A confirmed by X-ray diffraction. Both compounds were inactive against chloroquine-resistant *P. falciparum* FcB1 at the reported concentrations and lacked notable activity in the tested Caco-2, VERO, and Thp1 cell models. Annuanine B was absent from fresh-plant tea and appeared after drying/storage; mixture experiments attributed activity to artemisinin rather than annuanine B. The proposed biosynthetic versus extraction/degradation origin remains unresolved, but the study provides a defined-compound negative control and a direct reason to include storage and artifact controls in untargeted *Artemisia* metabolomics (Fabre et al. 2025).

An in-vivo *A. afra* test supplies an important negative control for claims based on artemisinin-independent infusions. In *P. berghei*-infected mice, aqueous powder suspensions of *A. annua* showed activity but *A. afra* did not support the proposed mammalian-prodrug hypothesis (Walz et al. 2024). This does not invalidate in-vitro *A. afra* findings, which may depend on parasite stage, exposure, metabolism, or assay matrix; it does mean that in-vitro activity should not be promoted to in-vivo efficacy without a chemically defined and host-relevant exposure experiment.

The full-text preparation record is unusually useful for interpreting that negative result. Senegal *A. afra* material contained 65% leaves and 35% twigs and less than 0.0071 mg/mL artemisinin in the dosing suspension; the study compared single and three-day oral dosing in *P. berghei* mice with pure artemisinin and two *A. annua* powders. *A. annua* produced 93–98% day-3 activity at 12.5–25 mg/kg artemisinin-equivalent doses, whereas single-dose *A. afra* produced only 21–28% activity and three daily high doses produced less than 5%. A very small humanized *P. falciparum* mouse comparison likewise found no valid evidence of *A. afra* activity, but dosing-volume intolerance limited that experiment. Thus, the result constrains the prodrug hypothesis under the tested powder, storage, host, and schedule conditions without falsifying tea-, material-, or host-dependent alternatives; no active terpene was identified (Walz et al. 2024).

Extraction chemistry can also create an apparent antimalarial advantage. Hydrotrope-assisted aqueous extraction with sodium salicylate or cholinium salicylate produced artemisinin-rich extracts with greater *P. falciparum* activity than pure artemisinin in

the tested system, while artemitin, chrysosplenol D, arteannuin B, and arteannuin J co-occurred in the extracts (Ferreira et al. 2024). The result is useful for sustainable extraction and matrix testing, but the solvent, extraction yield, co-metabolite composition, and assay exposure are inseparable confounders; it is not evidence for a clinical botanical product or for any one terpene-protein interaction.

Full-text verification makes the hydrotrope result more interpretable. Cryoground leaves were extracted at a 1:10 solid-to-liquid ratio, and response-surface optimization selected 2.5 M sodium salicylate at 32 degrees C for 110 minutes or 2.5 M cholinium salicylate at 32 degrees C for 130 minutes. The optimized extracts yielded 6.50 +/- 0.10 and 6.44 +/- 0.09 mg/g artemisinin, respectively, and were tested against asynchronous, chloroquine- and mefloquine-susceptible *P. falciparum* 3D7 at matched 1 microgram/mL artemisinin-equivalent concentration. Sodium- and cholinium-salicylate extracts had IC50 values of 0.0027 +/- 0.0005 and 0.0034 +/- 0.0009 microgram/mL versus 0.0066 +/- 0.0025 microgram/mL for pure artemisinin; UHPLC-DAD-MSn also detected arteannuin B, arteannuin J, artemitin, casticin, and chrysosplenol D. The salicylate extracts were reported as nonhaemolytic at the closest IC50 dose in the cultured-erythrocyte assay. These findings support a reproducible extraction and matrix/co-metabolite hypothesis, but the susceptible single-line design, mixed chemistry, hydrotrope exposure, and lack of fractionation/reconstitution or parasite-protein engagement prevent assignment to any one terpene or inference of clinical safety.

A complementary top-down study compared ethyl-acetate and petroleum-ether *A. annua* extracts at equivalent artemisinin exposure. Reported *P. yoelii* ED50 values were 2.9 and 1.6 mg/kg versus 5.6 mg/kg for pure artemisinin, while only the petroleum-ether extract increased rat artemisinin AUC0-t by 1.7-fold. Arteannuin B and artemisinic acid were isolated as differential sesquiterpenes and tested in the matched petroleum-ether proportion (Cai et al. 2024). The result is a useful sesquiterpene/matrix-PK hypothesis, but the lack of in-vitro enhancement, rat-only PK verification, and preclinical design prevent clinical or unstandardized-botanical inference.

A related component-resolved study points to host metabolism as one route by which an *A. annua* matrix can alter apparent artemisinin potency. Fractionation identified nine major components, including five sesquiterpenes; arteannuin B inhibited CYP3A4 with an IC50 of 1.2 microM, and selected matrix fractions approximately doubled artemisinin exposure in healthy rats. The study also reported support for antiplasmodial enhancement in a preclinical combination experiment (Cai et al. 2017). This is a valuable bridge between terpene co-occurrence, host pharmacokinetics, and malaria phenotype, but it does not establish a parasite-protein interaction, a clinical benefit, or a safe standardized botanical dose.

In a separate resistant-mouse study, artemisinin plus chrysosplenetin improved survival and reduced parasitemia-related measures in *Plasmodium berghei* K173, with effects associated with ABC-transporter and heme-ROS/GSH responses and dependent on host genotype and parasite context (Ji et al. 2025). This is valuable resistance-biology evidence, but it is not a direct parasite-protein target demonstration, not a terpene result, and not a human regimen. Together with the matched mutant study above, it argues for testing terpenes against malaria as chemically defined comparators within a standardized *A. annua* matrix, rather than treating every matrix effect as proof of terpene synergy.

Full-text extraction specifies the interaction model. Male ICR mice received artemisinin plus chrysosplenetin at a 1:2 ratio (40:80 mg/kg) after infection with 1 x 10<sup>5</sup> red blood cells carrying sensitive or resistant *P. berghei* K173; Mdr1a wild-type and knockout groups were followed through a seven-day course, with six to eight mice per group. In resistant-mouse cohorts, the combination maintained 100% survival, approximately 30% inhibition, and approximately 20% parasitemia regardless of Mdr1a genotype. Tissue- and genotype-dependent changes were measured for ABC-transporter and cytokine transcripts, ROS/GSH, hemoglobin, and PI3K/AKT-mTOR and MAPK signaling. Because chrysosplenetin is a polymethoxylated flavonol rather than a terpene, and because no parasite-protein binding or terpene-resolved botanical preparation was tested, this is a host/matrix and resistance-state comparator, not evidence for a terpene target or human treatment (Ji et al. 2025).

The parallel regulation review is useful context for these matrix experiments because it distinguishes shared transcriptional control of artemisinin and flavonoid pathways from demonstrated chemical synergy (Hassani et al. 2020). That distinction is retained in the present evidence model: co-regulation and co-occurrence are hypotheses until matched compounds, exposures, and parasite endpoints are tested.

Preparation and host context can change the apparent malaria phenotype before any new parasite target is invoked. In 14 healthy men, one liter of tea prepared from 9 g of dried *A. annua* leaves produced a mean artemisinin Cmax of 240 +/- 75 ng/mL and AUC of 336 +/- 71 ng/mLh; the authors explicitly concluded that the tea should not be treated as equivalent to modern artemisinin drugs (Räth et al. 2004). In mice, dried-leaf delivery likewise produced higher exposure than pure artemisinin, and infection altered absorption and deoxyartemisinin metabolism: AUC was 299.5 mgmin/L in healthy mice versus 435.6 mgmin/L in *P. chabaudi*\*-infected mice (Weathers et al. 2014). These studies make host pharmacokinetics a required covariate in any terpene/protein interaction experiment using plant material.

The literature contains a real, informative contradiction about whether artemisinin explains most activity in an *A. annua* preparation. Ancient fresh-herb soaking or pounding produced extracts with higher artemisinin concentration than dried-herb tea, but only the pounded juice contained sufficient artemisinin to suppress *P. berghei* parasitemia in mice (Wright et al. 2010). A matched in-vitro comparison across varieties found tea and nonpolar extract activity closely tracked artemisinin concentration and did not differ significantly from pure artemisinin (Mouton et al. 2013). By contrast, *A. annua* and *A. afra* infusions inhibited blood and

liver stages, including *P. vivax* and *P. cynomolgi* hypnozoites, in a study that reported essentially artemisinin-independent activity (Ashraf et al. 2022). These findings should not be averaged into a single “synergy” estimate: they differ in plant species, preparation, parasite stage, exposure, and likely non-terpene chemistry. They instead define a testable interaction between chemotype, matrix, parasite stage, and host metabolism.

Gametocyte and strain-resolved work further narrows the hypothesis. At 5 g dry weight/L, *A. annua* and *A. afra* teas reduced *P. falciparum* asexual parasites and early/late gametocytes, with effects mainly attributable to artemisinin content and only a weak signal from artemisinin-deficient *A. afra* (Snider and Weathers 2021). A separate panel of sensitive, resistant, and reversion strains found exposure-time and strain-dependent dried-leaf effects and suggested that malaria-patient sera can contribute to inhibition in some assays (Gruessner and Weathers 2021). Taken together, “artemisinin-dependent” and “artemisinin-independent” are best treated as conditional labels. The comparative experiment should therefore test purified artemisinin, structurally related sesquiterpenes, quantified volatile fractions, and recombined matrices at matched parasite-stage exposures, while measuring serum protein binding and host CYP modulation.

The full-text verification tranche confirms the importance of recording preparation rather than only plant name. Fresh-herb soaking or pounding produced 14.5 mg/L artemisinin in a dried-herb infusion versus 293 mg/L in pounded juice, and only the pounded juice suppressed *P. berghei* parasitemia in the reported mouse comparison (Wright et al. 2010). Cameroon and Luxembourg water infusions differed in beta-haematin inhibition and chromatographic profiles, with 10% dilution absorbances of 0.075 and 1.515, respectively (Akkawi et al. 2024). A voucher-linked extract study normalized assays by artemisinin, compared K13-mutant and reversion strains, and found strain- and exposure-time dependence plus additional inhibition in pooled malaria-exposed sera (Gruessner and Weathers 2021). Matched *A. annua* and artemisinin-free *A. afra* infusions inhibited blood and hepatic stages, including relapse-relevant models, but the active chemistry was unresolved (Cai et al. 2022). These are full-text-verified matrix and stage findings, not evidence that a non-artemisinin terpene has been proven causal; the row-level extraction is in `full-text-verification.csv`.

The newly verified 2013 comparison tested 16 geographically and temporally varied *A. annua* samples plus two *A. afra* samples as teas and chloroform extracts using HPLC-ELSD-normalized artemisinin measurements and a single chloroquine-sensitive *P. falciparum* 3D7 erythrocyte-stage assay. *A. annua* tea extracts averaged 0.75 microgram/mL extract IC<sub>50</sub> and 4.97 ng/mL calculated artemisinin IC<sub>50</sub> versus 5.48 ng/mL for pure artemisinin; chloroform extracts averaged 0.18 microgram/mL and 5.27 ng/mL, while *A. afra* teas were inactive above 25 microgram/mL and its chloroform extracts averaged 11.02 microgram/mL. The aged material, absent comprehensive terpene profiling, and single-stage design leave storage-sensitive constituents, prodrugs, and other life-cycle effects unresolved. The result supports artemisinin-dominant activity in the tested matrices, not the stronger genus-wide claim that artemisinin is the only active constituent (Mouton et al. 2013). The expanded full-text tranche adds two attribution controls. In a flavonoid-deficient *A. annua* *chi1-1* line that retained wild-type artemisinin, whole-leaf *P. falciparum* Dd2 activity and initial cytotoxic action did not materially differ from wild type, whereas artemisinin-impaired lines showed little or no activity; this supports artemisinin dominance in that genetic background without excluding matrix effects elsewhere (Czechowski et al. 2019). Conversely, *A. annua*-associated fungal endophyte extracts yielded antimalarial quinones such as emodin and physcion, providing a non-terpene comparator that should not be relabeled as native plant chemistry (Alhadrami et al. 2021). Silicon treatment further increased trichome size and artemisinin concentration under a selected condition without improving the corresponding *T. gondii* infusion phenotype, separating plant storage traits from parasite efficacy (Rostkowska et al. 2016).

Cross-species chemical detection adds a complementary but narrower boundary. HPLC analysis of Algerian wild *A. herba-alba*, *A. campestris* subsp. *glutinosa*, and *A. judaica* subsp. *sahariensis* reported artemisinin yields of 0.34%, 0.64%, and 0.04%, respectively (Ahmed-Laloui et al. 2022). These values support species- and lot-specific occurrence, not equivalent pathway completion, bioavailability, or antimalarial efficacy. The same study also measured antioxidant activity, which is a separate chemical phenotype and cannot be used as a proxy for parasite killing or host selectivity. A validated HPTLC method for artemisinin and artemisinic acid in *A. annua* callus and leaves provides a practical quality-control bridge for such comparisons (Khan et al. 2015). The appropriate comparative design is therefore voucher-linked quantitation followed by matched parasite-stage assays, rather than ranking species by a single chemical measurement. Repeated-stress experiments likewise changed artemisinin-related precursor ratios differently in upper and lower leaves and linked metabolite levels to glandular-trichome traits, while volatile fingerprints separated artemisinin-rich cultivars from wild type (Kjær et al. 2013; Reale et al. 2011). These are useful chemotype and sampling controls, but neither volatile correlation nor stress-induced precursor conversion establishes a causal parasite mechanism.

The malaria evidence also has a clear historical and analytical continuity. Artemisinin was isolated from *A. annua* growing in the United States in an early primary report (Klayman et al. 1984). Subsequent reviews describe the rapid antimalarial action and clinical development of artemisinin derivatives, while emphasizing chemistry and derivative pharmacology rather than equivalence of crude plant material to defined drugs (Balint 2001; Jung et al. 2004). A later review synthesizes the restricted distribution of artemisinin within *Artemisia* and its dependence on ontogeny and geography (Nabi et al. 2023). These records support malaria context and a distribution hypothesis, but only voucher-linked primary measurements can establish occurrence in a particular taxon or lot. Preparation chemistry is equally important: artemisinin recovery from *A. annua* tea infusions was strongly temperature-dependent and exceeded 90% under tested conditions, whereas an accuracy-profile-validated HPLC-UV method established a transferable workflow for hydroalcoholic extracts (van der Kooy and Verpoorte 2011; Diawara et al. 2011). Neither result makes tea clinically equivalent to artemisinin combination therapy; both support recording preparation, extraction, and

analytical uncertainty as covariates.

Quality-control methods are not interchangeable, but their convergence supports a practical cross-study measurement ladder. HPLC-ELSD with microwave-assisted extraction, TLC, GC-ECD, and HPLC-UV have each been validated for artemisinin in plant material or crude extracts, while rapid HPLC-MS/MS can resolve artemisinin together with 9-epi-artemisinin, artemisitene, dihydroartemisinic acid, artemisinic acid, and arteannuin B (Liu et al. 2007; Marchand et al. 2008; Liu et al. 2008; Ferreira and Gonzalez 2009; Suberu et al. 2013). HPLC-MS/MS has also been applied to tea, biscuit, and porridge products made from *A. annua* leaves (Carrà et al. 2014). This convergence enables low-resource screening plus confirmatory multi-metabolite analysis, but an analytical method cannot by itself establish tissue localization, pathway direction, bioavailability, or therapeutic efficacy.

The ladder now includes assays suitable for living material and for structurally related analogs. A microscale GC-ECD workflow quantified artemisinin from a single living leaf or flower with greater than 95% recovery, enabling non-destructive or low-input screening during biosynthesis experiments (Liu et al. 2009). Ultrasound-assisted extraction paired with rapid-resolution LC-triple-quadrupole MS achieved 0.20 ng/mL detection and 92.45–103.8% recovery in *A. annua* samples (Wang et al. 2011). Capillary electrophoresis-MS further separated artemisinin from analogs including ascaridole, artemisia ketone, deoxyartemisinin, artemisinic acid, artemetin, and dihydroartemisinic acid, while revealing diastereomeric features in ethanolic extracts (Nagy et al. 2021). These methods strengthen lot- and tissue-specific exposure measurement, but detection of an analog or diastereomer is not evidence of a defined pathway or parasite mechanism.

Quality control also has a distinct downstream drug-product layer. Two color-reaction TLC field assays detected as little as 10% artemisinin derivatives in several ACT formulations and identified counterfeit products in a small survey (Ioset and Kaur 2009). This is important for malaria treatment stewardship, but it applies to defined combination medicines rather than crude *Artemisia* material; it should not be used as evidence that a plant preparation is clinically interchangeable with an ACT.

The newly verified malaria records sharpen two distinct inference boundaries. Orally delivered dried whole-plant *A. annua* reduced an artemisinin-resistant *P. yoelii* line more effectively than matched pure artemisinin and delayed stable resistance during serial *P. chabaudi* selection, but the study did not resolve the responsible non-artemisinin constituents or establish a human regimen (Elfawal et al. 2015). In synchronized *P. falciparum* NF54 cultures, 5 g/L *A. annua* and *A. afra* teas were tested against asexual parasites and early/late gametocytes; activity was strongest in the higher-artemisinin *A. annua* tea, while artemisinin-deficient *A. afra* teas showed weaker or non-significant stage-specific effects (Snider and Weathers 2021). These results support a conditional artemisinin-centered, matrix-sensitive hypothesis and motivate purified-component, transmission, PK, and resistance-state experiments rather than a general claim for terpene synergy.

The next full-text tranche adds a dose-matched whole-plant comparator and two parasite-side mechanism controls. In a *P. chabaudi* mouse experiment, powdered dried leaves containing 24 mg/kg artemisinin reduced parasitemia more than an equal pure-artemisinin dose from 12 to 72 hours; the leaves were GC-MS characterized at 14.8 mg/g artemisinin, but no non-artemisinin terpene was fractionated or reconstituted, so the result remains a preparation-specific bioavailability or matrix hypothesis (Weathers et al. 2012). In a controlled beta-hematin assay, DHA, artemether, and artesunate inhibited crystallization under glutathione-reduced conditions with pH 5.2 EC50 values of 22.5, 31.6, and 33.5 micromolar, respectively, versus above 1,000 micromolar when oxidized; this is purified-drug heme-detoxification evidence, not whole-plant activity (Ribbiso et al. 2021). Finally, chemical-genetic *P. falciparum* experiments linked DHA survival to UPR resolution and proteasome capacity across Kelch13 and proteasome-mutant backgrounds, supporting a functional resistance target class without identifying a sole DHA-binding protein or botanical constituent (Rosenthal et al. 2024).

The full-text *A. cina* study provides a defined-mixture model for the wormwood hypothesis. Authenticated pre-flowering leaves and stems were macerated in ethyl acetate and fractionated by silica chromatography; GC-MS, HPLC-DAD, and multidimensional NMR identified peruvins as pseudoguaianolide sesquiterpene lactones and cinic acid as a related C15 sesquiterpene. In *H. contortus* L3 assays, peruvins plus 1-nonacosanol/hentriacontane produced 97.33% observed lethality versus 19.96% expected at matched 0.50LC25/0.50LC25 proportions (ratio 4.87), while peruvins plus cinic acid produced 40.42% versus 28.39% expected (ratio 1.42); the hydrocarbon/alcohol pair was additive for egg-hatching inhibition, and cinic acid alone showed no egg-hatching inhibition at the tested concentrations. The authors' protein-thiol and membrane-fluidity explanations remain hypotheses: there was no mammalian counterassay, infected-host test, exposure measurement, or direct parasite-protein experiment. This is therefore a strong comparative template for testing wormwood fractions, not evidence that peruvins or cinic acid occurs in *A. absinthium* (Arango-De-la-Pava et al. 2024).

An adjacent full-text *A. cina* study is an important negative attribution control for the terpene-focused synthesis. Authenticated pre-flowering leaves and stems were separately macerated in n-hexane, ethyl acetate, and methanol, and the active n-hexane fraction was taken through silica-gel fractionation. NMR and positive-mode ESI-MS assigned the active C2F5 subfraction as a 63:37 mixture of 3'-demethoxy-6-O-demethylisoguaiacin and norisoguaiacin—lignans, not terpenes. The crude n-hexane extract caused 80 +/- 9% *H. contortus* L3 mortality at 4 mg/mL; C1F5 caused 100% mortality at 2 mg/mL, and C2F5 caused 100% at 1 mg/mL. The authors infer cooperation of the two lignans, but individual purified standards were not tested, no terpene-resolved profile was reported, and no host-selectivity, infected-host, pharmacokinetic, residue, or direct parasite-protein endpoint was included. We therefore retain this record as a fraction-to-phenotype comparator and do not use it as evidence for a terpene, thujone, santonin,

or sesquiterpene-lactone mechanism (Higuera-Piedrahita et al. 2023).

The full-text *A. sieberi* malaria comparator further separates preparation-level activity from terpene attribution. Flowering aerial material from Khorassan and Semnan was macerated in methanol for 72 hours; 140 g of plant material yielded 13.3 g crude extract, which was defatted and partitioned into n-hexane, chloroform, and ether fractions. Total extract was administered subcutaneously at 100 mg/mL to *P. berghei*-infected NMRI mice for 10 days, while ether and chloroform fractions were tested for 14 days. The authors reported significant reduction of Giemsa-smear parasitemia and a virtual absence of blood-stage parasites in treated microscopy fields, with body weight, survival, hepatomegaly, and splenomegaly used as host observations. Yet no compound-resolved terpene profile, artemisinin measurement, exact effect size, stage-resolved IC<sub>50</sub>, or parasite-protein target was provided; the paper explicitly leaves the major active component unresolved. This is useful Iranian malaria evidence, but it cannot be promoted to an artemisinin-equivalent pathway or single-terpene mechanism (Nahrebian et al. 2012).

The verified gerbil study supplies a necessary negative control for the wormwood argument. Female gerbils received 600 exsheathed *H. contortus* L3 and were examined nine days later. Purified artemisinin (400 mg/kg once or 200 mg/kg for five days), *A. annua* aqueous or ethanolic extract (600 mg/kg), *A. annua* essential oil (300 mg/kg), and *A. absinthium* ethanolic extract (1,000 mg/kg) produced no significant reduction in worm burden. In the detailed tables, the *A. absinthium* group averaged 111.6 worms versus 97.1 in the water control, while the artemisinin groups averaged 72.6 and 78.2 versus 57.0 and 58.9 in vehicle and water controls. The plants were oven-dried or boiled/extracted under defined conditions, but no terpene-resolved profile, systemic artemisinin/dihydroartemisinin measurement, Labrasol-only control, or parasite-protein assay was included. The authors note that the gerbil infection is short-lived and larval-stage restricted, and that host physiology, bioavailability, redox defense, and contact time may explain disagreement with positive ovine *A. absinthium* studies. This is therefore a model- and preparation-specific negative result, not evidence that all wormwood chemotypes lack vermifuge potential (Squires et al. 2011).

### 6.3 Wormwood (*A. absinthium*): vermifuge evidence and limits

Full-text extraction of the Pakistani cross-species screen adds a useful, but chemically unresolved, comparator. Whole vegetative shoots of authenticated *A. indica* and *A. roxburghiana* were air-dried, crushed, exhaustively extracted with methanol, and stored before testing. Mixed gastrointestinal nematode eggs and adults were obtained from sheep and goats, including *Haemonchus contortus* and *Teladorsagia* spp.; methanolic extracts were tested at 3.75–50 mg/mL with PBS and 10% albendazole controls. After 48 hours, egg-hatch inhibition at 50 mg/mL was 85 +/- 21.2% for *A. indica* and 80 +/- 28.3% for *A. roxburghiana*; at 25 mg/mL, inhibition was 80 +/- 0.0% and 70 +/- 14.1%, respectively. The paper reported lower larval and adult-worm outcomes in tables formatted as mean +/- SD, with all tested adult worms killed at 50 mg/mL within 24 hours. No GC-MS/LC-MS terpene profile, purified active, infected-host efficacy, PK, residue, or mammalian selectivity endpoint was reported. This strengthens a preparation- and species-specific *Artemisia* nematode phenotype while leaving any thujone, sesquiterpene-lactone, or other terpene attribution open (Khan et al. 2015).

Correction to the earlier abstract-level summary: the full-text table assigns the 4.87-fold L3-lethality ratio to peruvine plus 1-nonacosanol/hentriacontane and the 1.42-fold ratio to peruvine plus cinic acid. The evidence matrix and antiparasitic table now use this corrected order.

A focused wormwood review summarizes reported constituents, pharmacological and anthelmintic activity, pharmacokinetics, and toxicity, including important cautions around essential-oil exposure (Batiha et al. 2020). A veterinary network meta-analysis synthesized 256 animals across six studies and identified *A. absinthium* among more effective botanical candidates, while emphasizing heterogeneity and the need for adequately powered randomized trials (Calzetta et al. 2020). These reviews support the vermifuge hypothesis but do not provide a standardized wormwood terpene effect or human-use evidence.

An independent review of more than 300 studies on edible *Artemisia* species and selected sesquiterpene lactones likewise integrates nutritional, pharmacological, and clinical-context evidence while identifying raw-extract standardization and adverse-effect trials as major gaps (Trendafilova et al. 2020). Its breadth is useful for locating wormwood and STL evidence, but review-level health claims remain distinct from chemically standardized vermifuge assays and cannot supply a safe dose.

An Iranian full-bloom collection illustrates why the wormwood label cannot substitute for a chemical fingerprint. Shade-dried aerial material yielded 1.3% hydrodistilled oil; 28 identified constituents represented 93.3% of the oil, with beta-pinene at 23.8%, beta-thujone at 18.6%, and hydrocarbon monoterpenes comprising 47.8% (Rezaeinodehi and Khangholi 2008). This is a well-defined regional beta-pinene/beta-thujone chemotype, but it supplies no anthelmintic endpoint and cannot be generalized to thujone-poor, camphor-rich, or oxygenated-sesquiterpene-rich wormwood oils. Any vermifuge experiment should therefore quantify the actual lot rather than infer exposure from the species name or traditional preparation.

The broader “wormwood” record illustrates why common names and parasite endpoints must be separated. A review of *A. sieberi* describes a Middle Eastern wormicide reputation and commonly thujone-dominant oils, but its synthesized efficacy evidence is primarily antifungal, including topical clinical studies; it does not establish a nematode assay or transfer to *A. absinthium* (Mahboubi 2017). Conversely, a primary study of crude ethanolic *A. inculta* extract in schistosomiasis-infected mice reported reduced worm load and tegumental damage at 500 or 800 mg/kg, with preserved bone-marrow morphology, cellularity, and maturation under the tested regimen (Lotfy et al. 2006). The latter is a useful helminth and safety comparator, but its species, route, crude

chemistry, and incomplete organ toxicology prevent attribution to a wormwood terpene or extrapolation to human treatment. Two newer reviews help situate, but not replace, the primary malaria and vermifuge evidence. A medicinal-chemistry review surveys natural and nature-inspired antimalarial compounds from *Artemisia* across in-vitro and in-vivo systems and highlights unresolved questions in mechanism, combinations, administration, and translation (Rauf et al. 2025). A production review synthesizes *A. annua* breeding, genetic engineering, tissue culture, elicitation, environmental regulation, and whole-plant strategies, while a second review covers supercritical-CO<sub>2</sub> extraction and biotechnological production of artemisinin precursors (Vashisth and Mishra 2025; Oladipo and Ojumu 2025). We use these as discovery and process-context records: neither establishes that a particular extraction method maximizes native flux, that a whole-plant product is clinically equivalent to an artemisinin drug, or that a terpene is responsible for wormwood anthelmintic activity. An additional sheep study strengthens the broader *Artemisia* vermifuge comparison without being evidence for *A. absinthium* specifically. Oral extracts of *A. vestita* and *A. maritima* reduced fecal egg counts in *H. contortus*-infected sheep, with maximum reported reductions of 87.2% and 84.5% at 100 mg/kg; the extracts also affected infective larvae and adult worms in vitro (Irum et al. 2015). This is useful in-vivo veterinary evidence, but the abstract does not provide a standardized terpene composition, purified active constituent, parasite-protein target, or human safety bridge. It should be used as a cross-species anthelmintic comparator, not as proof that wormwood oils share the same effect.

A 2026 Kashmir field study adds a contemporary whole-powder comparator. Naturally infected sheep received shade-dried *A. maritima* aerial-parts powder at 1, 2, or 3 g/kg, with untreated and fenbendazole controls; among 180 surveyed samples, 72.20% contained strongyle eggs and *Haemonchus* spp. comprised 56% of coprocultured larvae. In the 50-sheep trial, the 3 g/kg dose produced a maximum 70.40% fecal-egg-count reduction on day 21 versus 85.81% for fenbendazole, alongside improvements in hemoglobin, packed cell volume, and erythrocyte counts (Wani et al. 2026). This is useful current veterinary evidence and a dose/time hypothesis, but the available report does not provide a terpene or sesquiterpene-lactone profile, controlled single-parasite burden, residue limits, or transfer to *A. absinthium*.

Wormwood has a long-standing traditional reputation as a vermifuge, but the scientifically appropriate term for the current evidence is preparation-specific anthelmintic activity. The strongest early veterinary study compared crude aqueous and ethanolic aerial-part extracts with albendazole against ovine gastrointestinal nematodes. Both extracts impaired adult *Haemonchus contortus* motility in vitro and reduced fecal egg output in sheep; the ethanolic extract reached 90.46% fecal egg-count reduction at 2.0 g/kg body weight on day 15, 82.85% at 1.0 g/kg, and the aqueous extract reached 80.49% at 2.0 g/kg (Tariq et al. 2009). The publisher article record identifies the adult-worm paralysis/death assay, dose comparison, and albendazole benchmark, while the accessible study does not provide extract-resolved thujone, santonin, or sesquiterpene-lactone measurements. The reported superiority of the ethanolic preparation therefore supports polarity, solubility, and exposure hypotheses—not a thujone or STL mechanism. This is meaningful in-vivo veterinary evidence, but it does not define a standardized extract, separate *H. contortus* from mixed gastrointestinal nematodes, or establish use in humans. A separate caprine coccidiosis comparison is a useful negative or limited translation control: a single 2 g/kg dose of ethanolic *A. absinthium* extract was less effective than toltrazuril or amprolium, and oocysts persisted through day 28 (Iqbal et al. 2013). The result does not disprove wormwood activity against other parasites or preparations, but it shows why a traditional vermifuge label cannot substitute for dose-, host-, and parasite-matched trials.

A closely related negative control is provided by *A. annua*, whose malaria-active sesquiterpene lactone should not be assumed to be an ovine anthelmintic. In vitro, sodium-bicarbonate leaf extract had the lowest reported *H. contortus* LC<sub>99</sub> among the preparations tested, but in naturally infected sheep a single 2 g/kg extract dose and 100 mg/kg artemisinin dose produced only nonsignificant day-15 fecal-egg-count reductions of 19% and 28%, respectively; levamisole produced 97% reduction (Cala et al. 2014). The extract used in vivo contained no artemisinin. This discordance between in-vitro preparation activity and limited single-dose efficacy is a critical translation boundary for both artemisinin and wormwood claims.

Full-text extraction resolves the preparation and exposure variables in that comparison. The study used 2008 *A. annua* leaves dried at 35 degrees C, prepared water, 0.1% sodium-bicarbonate, ethanol, and dichloromethane extracts, and quantified artemisinin at 0.28%, 0.60%, 4.4%, and 9.8%, respectively. The sodium-bicarbonate extract used in sheep contained no artemisinin but did contain deoxyartemisinin, dihydroartemisinic acid, and artemisinic acid, and had high ORAC antioxidant capacity. In vitro, ethanol and dichloromethane extracts blocked egg hatch completely across the tested range, while the bicarbonate extract had LC<sub>99</sub> values of 1.27-1.50 microgram/mL for egg hatch and 23.8 microgram/mL for larval development. In vivo, the single bicarbonate-extract dose reduced day-15 EPG by 19.3% and 100 mg/kg artemisinin by 27.8%, neither significant; fecal artemisinin peaked at 126.5 microgram/g at 24 hours and fell to 49.0 microgram/g at 36 hours. Thus strong in-vitro activity did not translate to a significant single-dose ovine effect, and the experiment cannot assign either outcome to a terpene or to artemisinin alone (Cala et al. 2014).

Essential-oil evidence adds a different chemistry and exposure model. Full-text analysis of two cultivated Spanish *A. absinthium* populations (T2 and SN1, harvested in 2009 and 2010) found four hydrodistilled oils with a thujone-free (Z)-epoxyocimene/(Z)-chrysanthenol chemotype. GC-MS measured (Z)-epoxyocimene at 25.89-42.72%, chrysanthenol at 12.71-30.32%, linalool at 3.42-4.28%, and a (5Z)-2,6-dimethylocta-5,7-diene-2,3-diol at 20.21% in one harvest. Ex-vivo treatment of *Trichinella spiralis* L1 larvae reduced infectivity by 72-100% at 0.5-1 mg/mL (II50 205-291 microgram/mL); T2-10 and SN1-10 oils reduced intestinal larval counts by 49-66% at 100-500 mg/kg in mice, comparable to 66% for 5 mg/kg albendazole. J774 macrophage growth inhibition was below 20% at 200 microgram/mL, but this limited counterscreen does not establish mammalian safety. The oils produced only 5.44-11.56% mortality against *Meloidogyne javanica* J2 at 1 mg/mL, showing parasite-specific rather than universal activity



(García-Rodríguez et al. 2015). Because none of the major oil components was established as nematocidal, the result supports chemotype-by-parasite testing rather than attribution to epoxyocimene, chrysanthenol, thujone, or a universal “wormwood oil” mechanism.

An ectoparasite study adds a thujone-rich *Artemisia* comparator. *A. sieberi* oil contained  $\alpha$ -thujone (31.5%),  $\beta$ -thujone (11.92%), camphor (12.3%), and 1,8-cineole (10.09%) and produced contact LC50, LC90, and LC99 values of 15.85, 26.63, and 35.42 microgram/cm<sup>3</sup> against adult poultry red mites, with repellency and residual effects in the reported assays (Tabari et al. 2017). This is a mixture-level ectoparasite-control phenotype, not direct evidence for *A. absinthium* vermifuge activity; thujone, camphor, cineole, exposure route, poultry safety, and single-constituent causality remain unresolved. A separate extraction-method comparison in *A. absinthium* showed that direct-steam oil had a lower *Tetranychus urticae* LC50 (0.04 mg/cm<sup>2</sup>) than microwave-assisted or water-distilled oils (0.13 mg/cm<sup>2</sup>); a C15H<sub>24</sub> sesquiterpene was reported only in the more potent direct-steam profile (Chiasson et al. 2001). The candidate association is useful for fractionation, but it does not identify the active molecule or establish a helminth or human vermifuge effect.

Artemisinin occurrence and veterinary antiparasitic activity can also diverge by species and parasite. *A. sieberi* contained 0.20% artemisinin in summer and 0.14% in autumn dry weight, while the tested plant extract reduced severity or oocyst output for *Eimeria tenella* and *E. acervulina* but not *E. maxima* (Arab et al. 2006). This is a useful artemisinin/host-parasite comparator, but the result is season-, extraction-, species-, and dose-specific and cannot be transferred to *A. absinthium*, human malaria, or a single-constituent mechanism.

Feed and field studies are more heterogeneous. Dried wormwood supplementation and aqueous extracts produced strong in-vitro egg-hatch effects against *H. contortus*, with ED50 and ED99 values of 1.40 and 3.76 mg/mL for the wormwood extract, but fecal egg counts did not differ significantly at the group level in experimentally infected lambs (Mravčáková et al. 2020). A low-input swine-farm study reported antiparasitic effects after *A. absinthium* powder administration, but the co-occurring plant intervention and complex powder chemistry prevent attribution to terpenes (Băieș et al. 2024). A recent Awassi-sheep study of crude ethanolic extract reported 76.77% fecal egg-count reduction on day 15 compared with 100% for albendazole and no observed adverse effects during the small trial (Irehan et al. 2026). The latter is useful contemporary evidence, but it remains a single regional study with 18 animals and requires independent chemical, residue, dose, and safety confirmation.

Wormwood also has a defined-compound trematode signal that should not be conflated with the ovine nematode results. A dichloromethane *A. absinthium* extract caused 100% mortality of adult *Schistosoma mansoni* at 200 microgram/mL in vitro, and fractionation yielded the flavone artemetin and the sesquiterpene lactone hydroxypelenolide (Almeida et al. 2016). The study supports a preparation-level schistosomicidal phenotype and identifies chemically tractable leads, but it does not establish the isolated compounds' potency, selectivity, pharmacokinetics, or efficacy in an infected host. The constituent controls sharpen that boundary: artemetin and hydroxypelenolide from *A. absinthium* were inactive at 100 micromolar, whereas parthenolide isolated from the related *Tanacetum parthenium* killed adult worms at 12.5-100 micromolar and altered the tegument without affecting mammalian cells in the reported assay (Almeida et al. 2016). Thus, “sesquiterpene lactone” is not a sufficient attribution; structure, species, preparation, concentration, and host counterassay must remain linked.

Formulation can change the exposure boundary independently of plant chemistry. A Cuban *A. absinthium* essential oil formulated in nanocochleates showed an intracellular-*L. amazonensis* IC50 of 21.5 +/- 2.5 microgram/mL against a macrophage IC50 of 27.7 +/- 5.6 microgram/mL and suppressed murine cutaneous infection by approximately 50% after intralesional dosing at 30 mg/kg (Tamargo et al. 2017). This is a useful formulation and host-counterassay record, not evidence that unformulated wormwood oil is a human vermifuge or that a particular terpene is responsible.

The preparation-specific wormwood evidence is also visible in a recent comparative screen. Ethanolic tinctures from 46 traditional European medicinal plants were tested against *Strongyloides papillosus* and *Haemonchus contortus* larval stages; *A. absinthium* was among only five plants classified as nematocidal (Boyko and Brygadyrenko 2025). Full-text extraction showed that the 1.0% tincture killed 85.5 +/- 16.5% of *S. papillosus* L1-2 larvae versus 1.5 +/- 3.4% in control, but only 19.4 +/- 6.7% of *S. papillosus* L3 and none of *H. contortus* L3. The screen used diluted ethanolic tinctures and albendazole/water controls but no terpene-resolved profiling, animal endpoint, or residue analysis. It is therefore independent qualitative support for a stage-specific wormwood hypothesis, not evidence of a broad or chemically assigned vermifuge effect. A separate goat study provides the negative protozoan control: one 2 g/kg oral dose of crude ethanolic *A. absinthium* extract was less effective than toltrazuril or amprolium against naturally infected *Eimeria*, with oocysts persisting through day 28 (Iqbal et al. 2013).

Wormwood volatiles also show a vector-specific, compound-resolved phenotype that is relevant to malaria control but distinct from treatment of malaria infection. Leaf oil from *A. absinthium* contained (E)-beta-farnesene, (Z)-en-yn-dicycloether, and (Z)-beta-ocimene as major components; the oil and each compound were tested against six mosquito vectors, including *Anopheles stephensi*, with species-specific larvicidal potency, while non-target aquatic organisms were less sensitive in the reported comparison (Govindarajan and Benelli 2016). The result supports vector-control and selectivity experiments, not gastrointestinal anthelmintic efficacy, systemic antimalarial action, or safe human exposure. Similarly, *A. argyi* oil and isolated eucalyptol, beta-pinene, beta-caryophyllene, and camphor showed contact, fumigant, and repellency effects against adult *Lasioderma serricorne*, with the whole-oil phenotype attributed to combined major and minor constituents (Zhang et al. 2014). This is a useful constituent-

resolved arthropod comparator, but it should not be converted into a helminth or human parasite mechanism.

A 2026 poultry study adds a comparative helminth phenotype outside *A. absinthium*: ethanolic extracts of *A. herba-alba* and *A. judaica* reduced *A. galli* adult-worm motility in vitro and reduced female fecundity, fecal egg counts, and worm burden in infected chicks; camphor docking to ATP synthase and glutathione transferase was reported as a target hypothesis (Abdelqader et al. 2026). Because the preparation was crude, the docking was computational, and the species/model differs from wormwood and human infection, this record strengthens target prioritization but does not establish camphor causality or human vermifuge efficacy. It belongs in comparative helminth evidence, not as a direct extension of the *A. absinthium* vermifuge literature.

An older *A. brevifolia* sheep study provides a second comparative dose-response anchor: crude aqueous and methanol whole-plant extracts caused paralysis or mortality of live *H. contortus* in vitro, while crude aqueous extract at 3 g/kg produced a maximum reported 67.2% day-14 fecal-egg-count reduction in naturally infected sheep versus 99.2% for levamisole (Iqbal et al. 2004). This supports a graded veterinary phenotype, but the plant material, mixed infection, small study scale, and lower comparator performance do not identify an active terpene or support human use.

A separate *A. herba-alba* study found concentration- and time-dependent activity of crude methanolic flower and leaf extracts against *H. contortus* adult motility and egg hatch; the flower extract reached 98.67% egg-hatch inhibition at 1 mg/mL (Ahmed et al. 2020). This strengthens the comparative helminth rationale but does not establish that the signal is caused by camphor, a sesquiterpene lactone, or another terpene, and it cannot be transferred directly to *A. absinthium* or human vermifuge use.

A recent *A. brevifolia* study adds a contemporary, multi-stage comparator, but it is currently represented at the abstract level. Crude ethanolic and aqueous extracts were tested against gastrointestinal-nematode eggs, larvae, and adult worms; LC-MS identified 33 compounds and reported sesquiterpene lactones among the chemical classes. The abstract reports 87.5% egg-hatch inhibition for the ethanolic extract at 50 mg/mL and complete adult-worm motility inhibition after 8 hours at 25 mg/mL, versus 82.7% and 63.3% for the aqueous extract at the corresponding reported endpoints. Because the complete methods, constituent table, larval dose-response, and host-selectivity data remain unavailable, this is a preparation-specific hypothesis and not evidence that a named *A. brevifolia* terpene or lactone causes the phenotype (Hussain et al. 2025).

Two additional wormwood records strengthen the claim that *A. absinthium* has a reproducible vermifuge hypothesis while also exposing its limits. In the full-text *Hymenolepis nana* study, authenticated but store-purchased aerial parts were shade-dried and extracted in boiled water for 72 hours; the aqueous route was deliberately chosen to reduce systemic thujone exposure because thujone is more soluble in alcohol. The 1 and 5 mg/mL extract concentrations caused dose-dependent adult-worm paralysis, death, tegumental damage, lipid accumulation, nephridial-canal injury, and intrauterine-egg degeneration. In infected mice, 400 and 800 mg/kg given for three days reduced EPG by 61.8% and 98.8% and worm burden by 56.5% and 98.5%, respectively, compared with 100% reductions for praziquantel in the reported tables (Beshay 2018). In naturally infected cats, a small full-text study used a botanically authenticated, locally marketed flower extract made by seven-hour diethyl-ether Soxhlet extraction. Seven days of 300 or 600 mg/kg dosing in five- and three-cat groups was accompanied by declining *Toxocara cati* fecal egg counts, while a water-only control and serial serum chemistry were reported; two cats in the lower-dose group reached zero EPG. The same extract did not inhibit egg embryonation or larval development at 0.3 or 0.6 mg/mL in vitro, and no positive-drug comparator, adult-worm burden, or lot-specific terpene profile was provided (Yıldız et al. 2011). These are stronger than traditional-use assertions because they include infected hosts and preparation details, but both remain crude-preparation studies without a standardized thujone measurement, residue study, or human efficacy evidence.

Two independent extract screens extend the wormwood life-cycle evidence beyond *H. contortus*. In a 13-plant comparison, methanolic *A. absinthium* was among the most effective preparations against *H. contortus* eggs and larvae, while UPLC-MS/MS quantified selected plant secondary metabolites; the study therefore supports preparation-level activity but not a terpene-specific mechanism (Váradyová et al. 2018). In a separate concentration-series assay against *Ascaris suum*, alcoholic *A. absinthium* extract showed strong, concentration-dependent anti-embryogenic and larval effects; sesquiterpene lactones were among the mixed metabolite classes detected, but the authors explicitly called for deeper phytochemical attribution (Băieș et al. 2022). These results broaden the vermifuge hypothesis across nematode taxa while reinforcing that co-extracted phenolics, sterols, flavonoids, and lactones cannot be disentangled without fractionation and matched chemical exposure.

Comparative species studies help identify which helminth signals are likely to be terpene-independent. *A. campestris* aqueous and ethanolic extracts completely inhibited *H. contortus* egg hatching near 2 mg/mL and the ethanolic extract produced 100% adult mortality at the highest 24-hour dose, yet the chemical profile was dominated by flavonoids (Akkari et al. 2014). Methanolic *A. indica* and *A. roxburghiana* extracts showed more modest activity against mixed gastrointestinal nematodes and were explicitly proposed for purification and in-vivo follow-up (Khan et al. 2015). In *A. cina*, the n-hexane extract produced 75% and 82.6% activity against *H. contortus* L3 at 1 and 2 mg/mL, altered *H. contortus* Hc29 expression, and produced 100% reduction in an infected gerbil assay; isoguaiacine was also active in transitional larvae (Higuera-Piedrahita et al. 2021). Hc29 expression is a response biomarker rather than proof of direct binding, and none of these comparative results identifies the active constituent in *A. absinthium*. The appropriate next step is chemotype-matched fractionation with direct parasite-protein assays and host-cell counterscreens.

Abstract-level recovery of a 13-plant *H. contortus* screen adds a parallel *A. absinthium* comparison. Methanolic stem extract of *A. absinthium* was among the most effective preparations in both egg-hatch and larval-development assays; UPLC-MS/MS measured

6.10 mg rutin/g dry matter in that extract, while aqueous extracts showed a broader pattern of stronger ovicidal activity. This is useful evidence that extraction polarity and co-metabolite composition modify the wormwood phenotype, but it does not provide a terpene/STL exposure, in-vivo efficacy, or constituent attribution (Váradyová et al. 2018).

The thirty-third full-text tranche provides full-text extraction of the Tunisian *A. campestris* study as a preparation-specific comparative control. Leaves and stems were collected in Elhanian, Sidi Bouzid, air-dried at 50 degrees C, and extracted as aqueous room-temperature macerates or repeated 95% ethanolic macerates. HPLC-MS found quercetin derivatives dominant in the aqueous extract and apigenin derivatives and related flavonoids dominant in the ethanolic extract; no volatile or sesquiterpene-lactone profile was reported. Both extracts completely inhibited egg hatching at 2 mg/mL, with egg-hatch LC50 values of 0.827 mg/mL for ethanolic extract and 1.00 mg/mL for aqueous extract. At 2 mg/mL, ethanolic extract caused 91.30% adult-worm mortality at 8 hours and 100% at 24 hours, whereas aqueous extract caused 3.22% and 70.96%; albendazole caused 78.04% and 100% at the same timepoints. The study supports solvent-, dose-, and parasite-stage-specific veterinary activity, but its flavonoid-dominated chemistry and absence of host, PK, residue, or in-vivo endpoints do not establish a causal terpene/STL or transfer to *A. absinthium* (Akkari et al. 2014).

The thirty-fourth full-text tranche adds two cross-species fluke-stage controls. In *A. ludoviciana* subsp. *mexicana*, leaves were ethyl-acetate extracted after drying at 60 degrees C, and the resulting extract reached greater than 60% *Fasciola hepatica* egg ovicidal action above 300 mg/L and adult-fluke LC50/LC90/LC99 values of 80.1/310.9/362.9 mg/L; the current experiment did not quantify artemisinin or profile terpenes, although the discussion cites prior identification of artemisinin in the extract (Ezeta-Miranda et al. 2023). In *A. vulgaris*, an authenticated 96% ethanolic leaf extract reduced *F. gigantica* hatching to 3.33% at 5% extract by day 16 and caused 66.67% adult mortality at 20% extract after 40 minutes, with tegumental and internal disintegration; UV-Vis chemistry measured tannins, flavonoids, saponins, and alkaloids but did not resolve a terpene or STL (Nurlaelasari et al. 2023). These records strengthen stage-resolved comparative anthelmintic evidence while preserving the boundary between cross-species crude-extract phenotypes and causal *A. absinthium* terpene mechanisms.

The newer comparative evidence clarifies both positive and negative translation. Powdered *A. herba-alba* shoots produced marked clinical, fecal, and necropsy responses in most of six experimentally infected goats, although two goats retained some egg production or adult worms (Idris et al. 1982). Aqueous white-wormwood extracts also reduced mature *H. contortus* motility in vitro, with complete killing at one treatment, but the abstract does not provide an in-vivo endpoint or terpene attribution (Hoshiki et al. 2025). In contrast, an ethanolic *A. absinthium* extract caused 81.79% larval motility inhibition and 82.2% larvicidal activity at 200 mg/mL in vitro but no significant fecal-egg-count reduction at 600 mg/kg in free-grazing goats (Flores-Prado et al. 2025). This positive-in-vitro/negative-in-vivo pattern is a direct warning against equating a vermifuge phenotype with a field-ready therapy.

Full-text verification of the *A. herba-alba* cattle study resolves the preparation and adult-worm endpoint. Leaves collected in Harare were air-dried, oven-dried at 40 degrees C for 72 hours, milled, and macerated as 100 g powder in 500 mL water for 48 hours; the filtrate was vacuum-concentrated and refrigerated. Thirty mature *H. contortus* worms per tube were exposed at 37 degrees C to 1, 2.5, 5, or 10 mg/mL extract, distilled water, or albendazole, with repeated motility and mortality scoring. The 5 mg/mL treatment produced approximately 80% mortality at 8 hours, whereas 10 mg/mL killed all worms by 6 hours; more than half remained active at the lowest two extract concentrations at 8 hours. No terpene-resolved chemistry, infected-host efficacy, mammalian selectivity, residue, or systemic exposure was reported. The result strengthens a wormwood-family, adult-nematode phenotype but cannot be assigned to thujone, santonin, a sesquiterpene lactone, or *A. absinthium* itself (Hoshiki et al. 2025).

Across the sheep evidence, the direction is therefore suggestive but not uniform: an ethanolic aerial-part extract produced 90.46% fecal-egg-count reduction in one ovine study, dry-wormwood supplementation failed to separate treated lambs from controls despite strong in-vitro egg-hatch activity, a small Awassi-sheep trial reported 76.77% reduction versus 100% for albendazole, and a 30-day whole-plant *A. annua* trial found only modest, statistically non-significant EPG reductions despite HPLC-IR measurement of 0.96% artemisinin in the fed material (Tariq et al. 2009; Mravčáková et al. 2020; Irehan et al. 2026; da Silva et al. 2017). In the *A. annua* trial, feeding 0.2% and 0.4% body weight corresponded to approximately 19.2 and 38.4 mg/kg/day artemisinin when nominal intake was achieved, but bitter taste reduced intake at the higher dose and the log-transformed EPG effect was not significant; levamisole averaged 89% efficacy. This direct dose/palatability observation makes plant artemisinin content, delivered exposure, host physiology, and parasite model separate variables. The combined pattern makes extraction polarity, delivery matrix, host population, parasite burden, and endpoint timing essential moderators; it does not support a pooled wormwood or *A. annua* effect or a terpene-specific conclusion.

Wormwood oil evidence also extends into protozoan assays without resolving the vermifuge chemistry. In an Ethiopian voucher-linked study, fresh *A. absinthium* leaves yielded 1.39% hydrodistilled oil; GC-MS identified 65 compounds representing 93.74% of the oil, including camphor at 27.40%, davanone at 16.43%, and (E)-nerolidol at 4.63%. The oil inhibited promastigotes of *L. aethiopica* and *L. donovani* at MIC 0.1565 microliter/mL and inhibited axenic amastigotes, with THP-1 cytotoxicity and sheep-erythrocyte hemolysis controls (Tariku et al. 2011). This is a regional, mixture-level signal: it supports testing camphor-rich and thujone-rich chemotypes side by side, while the active constituent, intracellular activity, host-relevant exposure, and relationship to *H. contortus* or other helminths remain unknown. At the purified-compound level, helenalin, mexicanin, and dehydroleucodine inhibited *L. mexicana* promastigotes at estimated low-micromolar IC50 values (Mesquita et al. 2008). These lactone results support a chemically defined parasite assay ladder, not an inference that the compounds are co-abundant in *A. absinthium*.

The Cuban primary study adds a stronger *A. absinthium* protozoan anchor, while preserving the mixture boundary. Leaves linked to voucher ROIG 4640 were hydrodistilled; GC–MS reported 18 oil components, with trans-sabinyl acetate at 36.7% and 35.9% unresolved. Against *L. amazonensis*, the oil produced promastigote and intracellular-amastigote IC50 values of 14.4 +/- 3.6 and 13.4 +/- 2.4 microgram/mL, respectively, with macrophage CC50 75.1 +/- 2.3 microgram/mL and a reported selectivity index of 6. Five intralesional 30 mg/kg doses reduced lesions and parasite burden in BALB/c mice without treatment-group mortality or greater-than-10% weight loss (Monzote et al. 2014). This is chemically profiled, intracellular-stage and preclinical antileishmanial mixture evidence; trans-sabinyl acetate is a compositional marker, not a demonstrated causal constituent, and the study does not establish gastrointestinal vermifuge activity, oral/systemic exposure, or a parasite-protein target.

Standardization and formulation add two additional boundaries. A standardized *A. annua* extract showed activity against *Leishmania infantum* promastigotes and amastigotes without reported RAW264.7 macrophage cytotoxicity in the tested range, while also changing inflammatory cytokine readouts (Morua et al. 2025). This is stronger than an uncharacterized extract screen because the preparation and host counterassay were specified, but it remains mixture-level evidence and does not identify a terpene/protein interaction. Likewise, *A. cina* aqueous-extract silver nanoparticles caused high *H. contortus* L3 mortality and oxidative-enzyme changes, but the active entity was a synthesized nanoparticle rather than native *A. cina* material (Hernández Guerrero et al. 2025). Nanoparticle surface chemistry and silver toxicity therefore prevent direct comparison with wormwood oil, tea, or isolated terpenes.

The full-text *A. annua*/*L. infantum* study quantified artemisinin at 0.13%, 0.75%, and 1.21% in three extracts made from 40 degrees C air-dried Tenerife leaves. Promastigote IC50 values decreased from 782.25 to 231.97 +/- 20.75 and 58.53 +/- 2.20 microgram/mL across that artemisinin series; the 1.21% extract had an intracellular-amastigote IC50 of 69.32 +/- 12.79 microgram/mL, while J774A.1 macrophage viability remained above 90% to 400 microgram/mL. Separate LPS-stimulated RAW264.7 assays showed 45.26% and 55.16% IL-6 reduction for 0.6% and 0.9% artemisinin extracts, respectively. The design provides an artemisinin-content and preliminary selectivity anchor, but the cytokine assay was not parasite-infected, the extracts were not subjected to a complete terpene/STL fingerprint or factorial reconstitution, and no canine PK or in-vivo efficacy was tested (Morua et al. 2025).

The open full-text *A. sieberi* comparison similarly shows why a plant extract cannot be equated with purified artemisinin. An aqueous preparation made from shade-dried material was tested at 5–100 microgram/mL against *Leishmania major* promastigotes and intracellular amastigotes alongside commercial artemisinin. Reported promastigote IC50 values were 25 microgram/mL for the extract and 50 microgram/mL for artemisinin, with low effects on uninfected mouse macrophages; however, the study did not measure extract artemisinin or resolve any volatile, STL, or other constituent. The result supports a bounded extract-versus-compound hypothesis and a follow-up fractionation experiment, not an *A. sieberi* terpene mechanism or in-vivo treatment claim (Heydari et al. 2013).

Full-text extraction resolves the formulation boundary further. Authenticated cultivated *A. cina* leaves and stems were aqueous-extracted and used to synthesize purified, quasi-spherical silver nanoparticles; TEM measured a 7.56-nm core and DLS a 57.9-nm hydrodynamic diameter. Against experimentally obtained *H. contortus* L3 larvae, the formulation produced 91.33% mortality at 125–500 microgram/mL, with reported LC50 and LC90 values of 4.128 and 17.993 microgram/mL, whereas aqueous extract alone produced 21.54% mortality. At a sublethal nanoparticle exposure, parasite SOD increased 2.67-fold at 8 hours while GPx and CAT were downregulated at 2 hours. These results support a formulated oxidative-stress hypothesis, but the study did not resolve the plant capping layer into terpenes, lacked mammalian cytotoxicity or host efficacy, and included AgNO3 as a comparably active control. It therefore cannot be used to infer native *A. cina* or *A. absinthium* terpene activity or a direct parasite-protein target (Hernández Guerrero et al. 2025).

The strongest new compound-level helminth signal is not from *A. absinthium* but from *A. cina*. Chromatographic isolation identified peruvín, 1-nonacosanol, hentriacontane, and cinic acid; binary peruvín mixtures with the hydrocarbon/alcohol pair or cinic acid increased *H. contortus* L3 lethality 1.42-fold and 4.87-fold, respectively, at matched LC25 proportions (Arango-De-la Pava et al. 2024). This provides an experimentally tractable mixture-synergy model, but the interaction is pair- and concentration-specific, and the compounds are not evidence for *A. absinthium* chemistry without direct isolation and quantitation in wormwood. Full-text extraction of the infected-rat comparison clarifies the wormwood life-cycle boundary. Aerial parts of voucher-linked *A. absinthium* from Antalya were shade-dried and Soxhlet-extracted with methanol at 64 degrees C for three days; the lipophilic phase was removed after aqueous redissolution and the material was lyophilized. Wistar rats received 300 mg/kg for 20 days beginning five days after infection (enteral/adult phase) or 600 mg/kg for five days beginning day 45 (parenteral/encapsulated-larva phase), with water-only infected controls and tissue larval counts by trichinoscopy and artificial digestion. The *A. absinthium* extract reduced larval rates by 37.7–63.5% across sampled muscles in the enteral phase and 43.7–59.9% in the parenteral phase; the comparator *A. vulgaris* was more potent overall. The study reported no acute toxicity at the tested doses, but no positive anthelmintic comparator, pharmacokinetic or residue endpoint, or terpene-resolved chemistry. This is stronger than a traditional-use record but remains a crude, host- and stage-specific phenotype; it does not identify thujone, santonin, absinthin, or another terpene/STL as causal (Caner et al. 2008). The full-text goat study supplies a particularly informative translation boundary. Market-purchased Peruvian leaves were oven-dried and extracted with 70% ethanol; the bulk extract caused 82.2% larval mortality and 81.79% motility inhibition against *Trichostrongylus* L3 at 200 mg/mL after 8 hours, yet 600 mg/kg per day for three days produced no significant fecal-egg-count reduction in five naturally infected goats. The study did not chemically resolve thujone, santonin, or sesquiterpene

lactones and was small, but it explicitly separates an in-vitro signal from a negative in-vivo FECRT outcome. This is evidence for bioavailability, rumen metabolism, dose, or preparation effects—not evidence that wormwood is uniformly inactive or field-ready (Flores-Prado et al. 2025).

Fasciolid assays extend the comparative parasite space while preserving the same boundary. *A. vulgaris* extract reduced *Fasciola gigantica* egg hatch and produced time-dependent adult-fluke mortality with tegumental and internal morphology changes (Nurlaelasari et al. 2023); an ethyl-acetate extract of *A. ludoviciana* subsp. *mexicana* likewise showed ovicidal and fasciolicidal activity against *F. hepatica* (Ezeta-Miranda et al. 2023). A companion indexed report explicitly pairs *A. ludoviciana* subsp. *mexicana* fasciolicidal efficacy with genotoxicity and acute/chronic toxicity testing, but its retrieved PubMed record lacks an abstract, so it is retained as a title-level verification lead rather than quantitative evidence (Ezeta-Miranda et al. 2023). These records support a comparative *Artemisia* helminth phenotype, not a shared wormwood terpene or a defined protein mechanism. Bioguided *A. kermanensis* fractionation adds low-macrophage-toxicity antileishmanial flavonoids, demonstrating that some *Artemisia* parasite leads are non-terpene constituents and should not be forced into the terpene hypothesis (Soleimanifard et al. 2023). Full-text review of the *A. kermanensis* fractionation study sharpens that attribution control. Repeated dichloromethane:acetone extraction of 5.0 kg whole plant material was followed by silica MPLC and HPLC/TLC purification. MS/NMR identified a geraniol-derived monoterpene derivative, 1-acetoxy-3,7-dimethyl-7-hydroxy-octa-2E,5E-dien-4-one, alongside eupatilin and a trimethoxylated flavone. Against *L. major*, the active Fr 2 had promastigote IC<sub>50</sub> values of 47.9 and 42.9 microgram/mL at 24 and 48 hours, and the isolated monoterpene derivative had values of 47.1 and 41.7 microgram/mL; eupatilin showed amastigote IC<sub>50</sub> values below 40 and 20 microgram/mL, while Fr 2 macrophage IC<sub>50</sub> values were 6816 and 2086.6 microgram/mL. The experiment therefore provides a cross-class, structure-confirmed bridge with preliminary selectivity, but neither the monoterpene derivative nor the flavonoids establish a parasite-protein target, host exposure, or transfer to *A. absinthium* (Yazdiniapour et al. 2023).

The follow-up bio-guided *A. cina* study provides a cleaner constituent attribution. Authenticated pre-flowering leaves and stems were separately macerated in n-hexane, ethyl acetate, or methanol; the ethyl-acetate extract yielded 3.86% and was fractionated under larval bioassay guidance. The isolated cinic acid is a previously undescribed C<sub>15</sub> sesquiterpene with a pseudoguaianolide/ambrosanolide-like scaffold but without C<sub>8</sub> lactonization, and it represented only 0.01% of the ethyl-acetate extract. The extract had LC<sub>50</sub>/LC<sub>90</sub> values of 2.56/3.30 mg/mL against *H. contortus* L3, while the isolated crystal assay reported 0.01/0.21 mg/mL; the paper's abstract-level comparison gives 0.13/0.40 mg/mL for cinic acid and approximately 256-fold and 15.71-fold greater activity at the corresponding endpoints (Arango-De-la-Pava et al. 2024). This is still an in-vitro *A. cina* result, not evidence that cinic acid occurs in *A. absinthium*, reaches a host-relevant exposure, or engages a defined parasite protein. The low yield and assay-dependent potency also make mixture reconstitution and independent standard testing essential; the study nevertheless provides a concrete assay template for testing wormwood fractions against purified or recombinant parasite targets.

Defined sesquiterpene-lactone evidence also reaches beyond malaria. Dehydroleucodine isolated from *A. douglassiana* aerial parts induced phosphatidylserine exposure and DNA fragmentation in *Trypanosoma cruzi* epimastigotes and trypomastigotes, and combinations with benznidazole or nifurtimox increased the reported effects (Jimenez et al. 2014). This strengthens the claim that selected *Artemisia* lactones can produce parasite-stage phenotypes as purified molecules, while leaving protein target, mammalian selectivity, pharmacokinetics, and equivalence to *A. absinthium* unresolved.

An important negative attribution control comes from a 2024 *A. absinthium* computational study. HPLC profiling found nine prominent phenolics, and apigenin was the top predicted ligand for *Leishmania* N-myristoyltransferase at -9.6 kcal/mol after docking and molecular-dynamics analysis (Boudou et al. 2024). Apigenin is not a terpene, and no direct enzyme-binding or parasite target-engagement experiment was reported. The record is therefore useful for prioritizing a *Leishmania* experiment and for preventing non-terpene docking results from being misreported as wormwood terpene mechanisms.

Species-level oil comparisons further support chemotype-specific testing. Hydrodistilled *A. absinthium* oil was dominated by myrtenyl acetate (47.10%) and beta-thujone (11.41%) and produced high mortality against *Sitophilus granarius* but limited root-knot nematode activity, whereas *A. dracuncululus* oil, with methyleugenol, beta-asarone, and elemicin among major constituents, reached 100% mortality against *Meloidogyne incognita* and 94.9% against *M. chitwoodi* (Evlice et al. 2026). The contrast is consistent with species- and mixture-dependent activity, but docking and partial-least-squares prioritization do not establish which constituent is causal, nor do these assays establish livestock efficacy or human safety.

The wormwood chemistry needed to interpret these assays is itself phenology-dependent. Across four harvest stages, alcoholic extracts were measured for alpha- and beta-thujone by SPME-GC-MS and for absinthin, artemisetin, and dihydro-epi-deoxyartannuin B by UHPLC-HR-MS (Bach et al. 2016). This analytical design is directly relevant to vermifuge studies: a plant name and extraction label do not specify the actual exposure to thujones, bitter triterpene lactones, or other co-metabolites. It supports harvest- and lot-specific chemical fingerprints, not a conclusion that higher thujone means greater anthelmintic activity or safety. Independent developmental sampling reinforces the same point: in two wormwood chemotypes, flowers contained more oil than leaves, and the trajectories of alpha/beta-thujone and trans-sabinyl acetate differed across vegetative, budding, flowering, and post-flowering stages (Nguyen et al. 2018). An ex-situ comparison of 11 Polish populations identified 41 volatile compounds and separated sabinyl-acetate, mixed, and thujone chemotypes while also finding population differences in phenolic acids, tannins, and polyphenols (Kosakowska et al. 2025). Together these studies make tissue, stage, population, and nonvolatile co-metabolites mandatory covariates in any wormwood parasite experiment.

Table 3. Focused malaria and wormwood evidence anchors. Values are retained at the preparation, parasite-stage, and host-model level; they are not pooled effect estimates.

| Case                                                               | Preparation and observed signal                                                                                                                                                                                             | Evidence boundary                                                                                                                     |
|--------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------|
| <i>A. annua</i> / <i>P. falciparum</i> Dd2                         | Matched whole-leaf mutant extracts: a flavonoid-deficient CH11 mutant retained wild-type artemisinin and antiparasmodial activity; artemisinin-impaired lines showed little or no activity                                  | Supports artemisinin dominance in this genetic system; matrix effects elsewhere remain unresolved                                     |
| <i>A. annua</i> / <i>P. falciparum</i> NF54                        | LED-conditioned leaf extracts: IC50 0.51, 0.52, 1.11, and 2.23 microgram/mL for white, blue, red, and greenhouse plants                                                                                                     | Environment-chemotype-phenotype association; no single terpene assigned as causal                                                     |
| <i>A. annua</i> / rodent malaria                                   | Dried whole plant versus artemisinin: whole plant overcame selected <i>P. yoelii</i> resistance and slowed stable <i>P. chabaudi</i> resistance                                                                             | Preclinical matrix result; does not replace quality-assured ACT or establish human efficacy                                           |
| <i>A. annua</i> / <i>P. yoelii</i>                                 | Equivalent-artemisinin ethyl-acetate and petroleum-ether extracts: ED50 2.9 and 1.6 mg/kg versus 5.6 mg/kg for pure artemisinin; petroleum-ether extract increased rat artemisinin AUC0-t 1.7-fold                          | Defined sesquiterpene/matrix-PK hypothesis; no in-vitro enhancement, clinical evidence, or unstandardized-botanical inference         |
| <i>A. annua</i> / <i>H. contortus</i>                              | Sodium-bicarbonate extract LC99 1.27 microgram/mL in egg-hatch and 23.8 microgram/mL in larval development; single-dose extract and artemisinin produced nonsignificant 19% and 28% day-15 EPG reductions in sheep          | Important negative/limited translation control; malaria efficacy does not imply anthelmintic efficacy                                 |
| <i>A. annua</i> / <i>H. contortus</i> sheep-model                  | Sun-dried whole plant contained 0.96% artemisinin; 30-day feeding at 0.2% and 0.4% body weight produced mean efficacy of 32.9% and 14.8%, without significant log-EPG reduction; high-dose intake was limited by bitterness | Quantified plant marker but no full terpene profile or causal constituent; dose, palatability, and exposure constrain translation     |
| <i>A. absinthium</i> / ovine gastrointestinal nematodes            | Crude aqueous and ethanolic aerial-part extracts reduced fecal egg output; ethanolic extract reached 90.46% FECR at 2.0 g/kg on day 15                                                                                      | Direct veterinary vermifuge evidence; active terpene, residue, and human translation unresolved                                       |
| <i>A. absinthium</i> / <i>H. contortus</i>                         | Aqueous extract ED50/ED99 were 1.40/3.76 mg/mL in vitro, but dry-wormwood dietary supplementation did not significantly reduce group-level lamb EPG                                                                         | Positive in-vitro/limited in-vivo translation boundary                                                                                |
| <i>A. absinthium</i> / Awassi-sheep gastrointestinal nematodes     | Crude ethanolic extract produced 76.77% day-15 FECRT versus 100% for albendazole in a six-sheep treatment group                                                                                                             | Contemporary regional veterinary result; small, crude-preparation, short-follow-up study                                              |
| <i>A. absinthium</i> / <i>T. spiralis</i>                          | Thujone-free population oils reduced larval infectivity ex vivo by 72–100% and adult burden in vivo by 66%, with no effect on <i>M. javanica</i>                                                                            | Chemotype- and parasite-specific oil phenotype; not a universal wormwood effect                                                       |
| <i>A. absinthium</i> / <i>H. nana</i>                              | Crude aqueous extract caused dose-dependent adult-worm paralysis/death and reduced egg counts and worm burden in infected mice at 400 and 800 mg/kg                                                                         | Rodent anchor; praziquantel performed best and active chemistry is unresolved                                                         |
| <i>A. absinthium</i> / <i>T. cati</i>                              | Ether-extracted flower preparation at 300 or 600 mg/kg for 7 days was associated with declining fecal egg counts in naturally infected cats; egg embryonation was unaffected in vitro                                       | Small host-level signal with a negative life-cycle endpoint; no terpene profile, positive comparator, adult burden, or human evidence |
| <i>A. absinthium</i> / <i>S. mansoni</i>                           | Dichloromethane extract/fractions caused 100% adult-worm mortality at 200 microgram/mL; artemetin and hydroxypelenolide were isolated                                                                                       | Extract/fraction signal; isolated-compound potency, selectivity, and host efficacy unresolved                                         |
| <i>A. absinthium</i> / <i>L. amazonensis</i>                       | Essential oil in nanocochleates: intracellular-amastigote IC50 21.5 +/- 2.5 versus macrophage IC50 27.7 +/- 5.6 microgram/mL; approximately 50% murine suppression at 30 mg/kg intralésional                                | Formulation- and route-specific result; not evidence for a single terpene or human vermifuge                                          |
| <i>A. absinthium</i> / <i>L. aethiopica</i> and <i>L. donovani</i> | Camphor-rich oil inhibited promastigote and axenic-amastigote forms, with weak hemolysis reported                                                                                                                           | Regional mixture signal; does not identify camphor as causal or establish helminth efficacy                                           |
| Purified natural sesquiterpene lactones / <i>L. mexicana</i>       | Helenalin, mexicanin, and dehydroleucodine showed estimated 2–4 micromolar promastigote IC50 values                                                                                                                         | Defined-compound evidence; source co-occurrence, amastigote activity, and target identity remain unresolved                           |
| <i>A. herba-alba</i> /A. <i>judaica</i> / <i>A. galli</i>          | Crude ethanolic extracts reduced fecundity, fecal egg counts, and worm burden in chicks; camphor docking to ATP synthase/GST was reported                                                                                   | Comparative poultry extract phenotype and computational target hypothesis; not purified-camphor, direct-binding, or human evidence    |
| <i>A. brevifolia</i> / ovine gastrointestinal nematodes            | Whole-plant crude aqueous extract reached 67.2% day-14 EPG reduction at 3 g/kg, below levamisole                                                                                                                            | Comparative veterinary phenotype; no terpene attribution or human evidence                                                            |
| <i>A. campestris</i> / <i>H. contortus</i>                         | Aqueous and ethanolic extracts completely inhibited egg hatching near 2 mg/mL; ethanolic extract caused 100% adult mortality at the highest 24-hour dose                                                                    | Strong comparative extract phenotype, but chemistry was flavonoid-dominated                                                           |
| <i>A. cina</i> / <i>H. contortus</i>                               | n-Hexane extract produced 75–82.6% L3 activity at 1–2 mg/mL and 100% reduction in infected gerbils; Hc29 expression changed after exposure                                                                                  | Fractionation/biomarker lead; expression change is not direct target engagement                                                       |
| <i>A. cina</i> / <i>H. contortus</i> L3                            | Isolated cinic acid: LC50 0.13 mg/mL and LC90 0.40 mg/mL versus 3.30 mg/mL extract LC90                                                                                                                                     | Defined sesquiterpene bridge; host exposure, selectivity, target, and transfer to <i>A. absinthium</i> unresolved                     |
| <i>A. douglassiana</i> / <i>T. cruzi</i>                           | Purified dehydroleucodine induced phosphatidylserine exposure and DNA fragmentation in epimastigotes and trypomastigotes                                                                                                    | Compound-level STL phenotype; protein target and host selectivity unresolved                                                          |
| <i>A. sieberi</i> / <i>P. berghei</i>                              | Methanolic, ether, and chloroform-derived aerial-part extracts reduced murine parasitemia in the reported study                                                                                                             | Crude, species-specific preparation; active constituent and human translation unresolved                                              |
| <i>A. khorassanica</i> / <i>P. berghei</i>                         | Flowering-stage extract showed limited murine activity; chrysanthenone and cis-thujone were reported among GC-MS constituents                                                                                               | Composition markers are not causal proof; effect is preparation-, dose-, and parasite-specific                                        |

| Case                                                               | Preparation and observed signal                                                                                                                        | Evidence boundary                                                                          |
|--------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------|
| <i>A. afra</i> / <i>P. berghei</i>                                 | Aqueous powder suspension did not show antimalarial activity in infected mice, unlike the <i>A. annua</i> comparator                                   | Important negative in-vivo control for artemisinin-independent infusion hypotheses         |
| <i>A. vulgaris</i> / <i>F. gigantica</i>                           | Extract reduced day-16 egg hatching to 3.33% at 5% and reached 66.67% adult mortality at 20% after 40 minutes                                          | Comparative in-vitro extract phenotype; active compounds and mechanism unresolved          |
| <i>A. absinthium</i> / <i>A. dracunculus</i> / root-knot nematodes | Oil activity diverged by species and composition: <i>A. absinthium</i> was weak while <i>A. dracunculus</i> reached 100% <i>M. incognita</i> mortality | Chemotype- and mixture-dependent comparison; docking does not establish causal constituent |
| <i>A. absinthium</i> / <i>Eimeria</i> spp.                         | Single 2 g/kg ethanolic extract dose in naturally infected goats underperformed toltrazuril and amprolium; oocysts persisted to day 28                 | Negative/limited veterinary translation control; crude chemistry unresolved                |
| <i>A. absinthium</i> / chemotype controls                          | Flowers had higher oil content than leaves, and 11 ex-situ populations separated into sabinyol-acetate, mixed, and thujone chemotypes                  | Organ-, stage-, and population-specific chemistry; no direct efficacy or safety inference  |

The wormwood literature therefore supports a focused experimental program. First, voucher-linked *A. absinthium* accessions should be chemically profiled for thujones, camphor, cineole, epoxyocimenes, chrysanthanol, sesquiterpene lactones, and non-volatile co-metabolites. Second, the same preparations should be tested against matched life stages of *H. contortus* and *T. spiralis*, with adult-worm motility, egg hatch, larval development, intracellular or tissue stages where relevant, and mammalian-cell counterscreens. Third, oils and extracts should be fractionated and recombined at measured ratios to distinguish single-compound activity from matrix effects. Finally, parasite-protein interaction work should begin from the observed *A. absinthium* chemistry and parasite phenotype; docking to a convenient protein without chemical provenance is not a target discovery result.

Wormwood also sharpens the safety boundary. Thujone is a neuroactive, preparation-dependent risk marker, and santonin is an additional historically important but toxic *Artemisia* anthelmintic marker that requires direct quantitation. Some of the strongest nematode evidence above came from thujone-free cultivated oils, whereas other wormwood chemotypes may contain thujones. Consequently, “vermifuge” cannot be used as a synonym for “safe,” and veterinary efficacy cannot be transferred to human self-treatment. The Terpedia records preserve the distinction among traditional use, in-vitro nematode phenotype, in-vivo veterinary efficacy, quantified thujone/santonin chemistry, and human therapeutic evidence.

Full-text verification of a related *A. herba-alba* nematode study adds a negative attribution control for the wormwood stream. Cold-methanolic flower and aerial-part extracts produced concentration- and time-dependent adult-motility effects against *H. contortus* and, at 1 mg/mL, inhibited egg hatching by 98.67% and 88.3%, respectively. Qualitative screening detected alkaloids, saponins, tannins, flavonoids, glycosides, and phenols, but the study did not provide a terpene profile, mammalian selectivity assay, or purified-component test (Luka et al. 2020). This strengthens the comparative veterinary signal while leaving *A. absinthium* transfer, thujone/STL attribution, in-vivo efficacy, and human safety unresolved. A chemically resolved essential-oil comparison further cautions against treating “wormwood” as one chemotype. Flowering aerial parts of field-grown *A. absinthium* were hydrodistilled and contained 66 identified compounds, dominated by myrtenyl acetate (47.10%), beta-thujone (11.41%), and pulegone (8.50%); under the same protocol, *A. dracunculus* oil was dominated by methyleugenol, beta-asarone, and elemicin. At 2 microliter/mL, *A. dracunculus* oil caused 100.0% and 94.9% mortality of *Meloidogyne incognita* and *M. chitwoodi* J2s at 72 hours, with 72-hour LC50/LC90 values of 0.5/1.1 and 0.6/1.5 microliter/mL, respectively. The *A. absinthium* oil caused only 11.4% and 7.5% mortality and increased egg hatching relative to control in that assay. Sparse-PLS and docking prioritized several constituents, including pulegone and beta-thujone, but did not establish causal molecules or nematode targets. This is strong species- and lot-specific oil evidence against a generalized vermifuge inference and is not direct evidence against gastrointestinal helminths (Evlice et al. 2026).

The tenth full-text verification tranche resolves the preparation details behind the recent Awassi-sheep signal. Commercially certified, pharmacologist-verified *A. absinthium* powder was extracted by three 16-hour macerations in 95% ethanol, concentrated at 50 degrees C, and administered once at 1 g/kg to six sheep; FECR was 66.79% on day 7 and 76.77% on day 15, versus 100% for albendazole. The study reported no observed local irritation, systemic toxicity, or adverse effects during short follow-up, but provided no lot-specific thujone, santonin, or sesquiterpene-lactone fingerprint, residue measurement, pharmacokinetic analysis, or longer-term toxicology. It therefore strengthens the contemporary ovine phenotype while leaving the causal chemistry and safety margin unresolved (Irehan et al. 2026).

The thirty-second full-text tranche adds a preparation-resolved *A. absinthium* tissue-stage comparison. The methanolic aerial-part extract was voucher-linked and lyophilized after aqueous redissolution and removal of the lipophilic phase; in infected rats, 300 mg/kg during the enteral phase and 600 mg/kg during the parenteral phase reduced tissue larval rates, but *A. vulgaris* was more potent overall. The study reported no acute toxicity at the tested doses, yet no positive anthelmintic comparator, PK/residue endpoint, or terpene-resolved chemistry. It therefore strengthens the wormwood phenotype map without supporting a thujone/STL mechanism or human translation (Caner et al. 2008).

## 7. Safety and translation

Derivative safety can be triaged, but not declared, by integrated computational and experimental workflows. A machine-learning platform screened 374 artemisinin derivatives across cardiotoxicity, hepatotoxicity, reproductive toxicity, mutagenicity, and tu-

morigenicity models; seven candidates passed the reported filters, with cardiomyocyte and zebrafish experiments providing limited validation (Kadioglu et al. 2021). This is a useful model for prioritizing compounds and measuring selectivity, but a shortlist is not a clinical safety conclusion and cannot be transferred to crude *A. annua* or *A. absinthium* preparations. The reproductive-endocrine boundary is equally important: an *A. absinthium* oil dominated by cis-chrysanthenyl acetate and sabinyl acetate altered a pseudopregnancy rat endpoint at 25 mg/kg (Demirel et al. 2025). That result is neither an anthelmintic assay nor a human risk threshold, but it makes reproductive exposure a required counterscreen for wormwood oils proposed for internal use.

The safety mechanism of alpha-thujone must be kept separate from its historical anthelmintic reputation. A primary study examined alpha-thujone as a toxic absinthe/wormwood-oil constituent, showing GABA-A receptor modulation, convulsant neurotoxicity, and metabolic detoxification pathways (Höld et al. 2000). This supports exposure- and lot-specific toxicology; it does not establish parasite selectivity, a safe therapeutic dose, or modern anthelmintic efficacy. Wormwood-oil safety signals extend beyond thujone. In a human hepatic-stellate-cell model, *A. absinthium* essential oil at 0.4 microgram/microliter for 24 hours altered lipid metabolism and activated detoxification, stress, and pro-inflammatory responses (Barreto et al. 2024). This is a preparation- and concentration-specific cellular signal, not proof of clinical hepatotoxicity or a defined human risk threshold, but it supports including hepatic counterscreens and lot-specific chemical fingerprints in vermifuge studies.

Off-target mammalian pharmacology is another translation gate. Artemisinin directly activated recombinant and native TRPC3-containing calcium channels in heterologous and PC12-cell assays (Urban and Schaefer 2020). This result does not establish clinical toxicity, but it demonstrates why parasite selectivity should be tested against host ion channels and calcium signaling rather than inferred from parasite activity alone. Host microbiome metabolism adds a second exposure variable: seventeen human intestinal fungal species converted artemisinin mainly through isomerization and deoxygenation, and transplantation of one isolate into antibiotic-treated mice yielded multiple fecal artemisinin isomers (He et al. 2025). The metabolites were assessed for breast-cancer cytotoxicity, not antiparasitic efficacy, so the study supports a pharmacokinetic/biotransformation hypothesis rather than a new therapeutic claim.

Reproductive safety is a separate translational gate even for *A. annua* material. In pregnant Wistar rats, an ethanolic leaf extract containing 1.098% artemisinin produced limited maternal toxicity signals at 100-300 mg/kg in the reported assays, but the highest dose was associated with reduced litter size and fetal nonviability or malformation (Abolaji et al. 2013). This preparation- and dose-specific result does not define human pregnancy risk or a safe dose for teas, oils, isolated artemisinin, or wormwood, but it supports an explicit reproductive-toxicology gate in any antiparasitic translation program.

Thujone is neurotoxic and must be tracked as a quantified, preparation-specific analyte. Artemisinin-derived antimalarial treatment should not be conflated with crude *A. annua* tea, essential oil, or an unidentified *Artemisia* preparation. In-vitro antioxidant, antimicrobial, insecticidal, cytotoxic, or antiparasitic results do not establish a safe human dose or therapeutic efficacy.

Composition-aware toxicology is necessary because risk markers vary among accessions and processing methods. A comparative toxicological study of *A. annua* volatiles paired chemical profiles with pharmacological, microbiological, and toxicity assays, demonstrating that “*A. annua* oil” is not interchangeable with artemisinin or with a standardized extract (Radulović et al. 2013). Regional *A. afra* oils likewise differed in camphor-, cineole-, and thujone-associated composition, making provenance relevant to safety interpretation (Oyedemi et al. 2009). A safety table should therefore include not only activity and a generic cytotoxicity result but quantified hazardous constituents, route, exposure duration, metabolic activation, and relevant host or ecological controls.

Cross-species oil toxicology supplies a further calibration point. Iranian *A. armeniaca* and *A. incana* oils differed sharply: the former contained substantial non-terpene hydrocarbons with alpha-pinene and spathulenol among its major constituents, whereas the latter was dominated by oxygenated monoterpenes including camphor, 1,8-cineole, and both thujones; both oils showed toxicity in a brine-shrimp assay (Mojarrab et al. 2013). Brine-shrimp lethality is a general screening endpoint, not a mammalian therapeutic window, but the paired chemistry demonstrates why a species label cannot predict either parasite selectivity or host risk. An independent *A. princeps* oil study illustrates the need for a host-response control in wormwood research. Dietary essential oil, with or without vitamin E, altered LDL-oxidation readouts and cholesterol-related gene expression in HepG2 cells; higher treatment concentrations produced cytotoxicity (Chung et al. 2007). The study does not test helminths and does not identify a causal volatile, but it shows that a wormwood oil can have concentration-dependent mammalian effects unrelated to vermifuge activity. Matched parasite and hepatic-cell assays should therefore be treated as a minimum selectivity panel for candidate wormwood preparations.

A 13-week rat study of wormwood extract reported no obvious treatment-related changes in survival, body weight, clinical chemistry, organ weights, or histopathology at 0.125-2% extract and estimated a study-specific NOAEL at or above 1.27 g/kg/day in males and 2.06 g/kg/day in females (Muto et al. 2003). This is useful repeated-dose toxicology, but it cannot be converted into a human safe dose or generalized across thujone-rich chemotypes, essential oils, extraction methods, or parasite-treatment routes. The appropriate safety bridge remains quantified thujone and co-constituent exposure in the exact product under study.

Reproductive risk is also supported by a distinct purified-drug experiment. Oral artemisinin at 7, 35, or 70 mg/kg/day during either early or later gestation produced dose- and timing-dependent adverse pregnancy outcomes in Wistar rats, including post-implantation loss at higher doses (Boareto et al. 2008). This does not define human risk or establish that an *A. annua* tea has the same exposure, but it makes reproductive counterscreens mandatory when comparing artemisinin-rich malaria preparations with wormwood extracts or oils.



Thujone risk should not be reduced to a single-constituent narrative. In a comparative study of thujone-containing oils, multivariate analysis of rat behavior, brine-shrimp toxicity, antimicrobial activity, and detailed oil chemistry indicated that constituents beyond alpha- and beta-thujone contributed to observed effects (Radulović et al. 2017). The result strengthens the need for mixture-aware safety assessment of wormwood oils, while leaving human dose limits and parasite selectivity unresolved. It also explains why the same thujone percentage cannot be assumed to predict the biological behavior of every oil.

Analytical quality control is a prerequisite for that bridge. A validated LC-MS method reached a 0.22 ng/mL artemisinin detection limit and distinguished measurable *Artemisia* content from likely cross-contamination in material associated with retracted clinical claims (Lee et al. 2022). The lesson applies equally to wormwood vermifuge work: voucher identity, lot-specific GC-MS or LC-MS fingerprints, quantified thujones and lactones, and chain-of-custody records should precede biological comparison. A high-sensitivity assay improves attribution; it does not turn chemical detection into efficacy or safety evidence.

Preparation matrices also change exposure. Aqueous infusions extract a different chemical subset from hydrodistilled oils; solvent extracts can concentrate nonvolatile lactones and other electrophilic metabolites; and nanoformulations can alter stability, tissue distribution, and apparent potency. These are not minor formulation details. They determine which compounds reach a parasite, which host proteins are exposed, and whether an in-vitro concentration can plausibly be achieved safely in vivo.

The artemisinin matrix also has a drug-interaction boundary. In human hepatic models, all tested *Artemisia* teas inhibited CYP2B6 and CYP3A4 activity, whereas transcriptional induction was observed with artemisinin and a high-artemisinin tea but not with artemisinin-deficient teas; the authors interpreted the broad activity inhibition as likely due to other tea phytochemicals (Kane et al. 2022). A related human-microsome and rat study found that dried leaves increased artemisinin tissue exposure, consistent with inhibition of first-pass metabolism, with sex-dependent differences in downstream cytokine effects (Desrosiers et al. 2020). These data support measuring host CYP and transporter interactions alongside parasite assays. They do not identify a safe dose, validate an herbal regimen, or prove that a volatile terpene is responsible.

The regulatory boundary is preparation-specific. The EMA monograph requires the amount of thujone in covered *A. absinthium* products to be specified and sets a daily exposure limit of 6.0 mg (EMA monograph). For malaria, WHO recommends quality-assured artemisinin-based combination therapy and explicitly does not support promotion of *Artemisia* plant material such as teas, tablets, or capsules for prevention or treatment (WHO treatment guidance). These authorities address defined products and clinical policy; they do not establish the safety or efficacy of an arbitrary species, oil, extract, or terpene mixture.

The record-level safety and translation boundaries are archived in `safety-translation.csv`. This prevents a low-toxicity result in one assay or a traditional-use claim from being converted into a generalized therapeutic recommendation.

Artemisinin safety also has a regimen-dependent neurotoxicity boundary. A review of laboratory studies described relatively selective brainstem lesions, species differences in vulnerable nuclei, and a stronger association with sustained circulating drug or metabolite exposure than with a single peak; the mechanism and human translation remain unresolved (Genovese and Newman 2008). This does not negate the clinical value of quality-assured artemisinin-based therapies, but it does rule out treating plant-material dose, purified-drug dose, and repeated exposure as interchangeable variables in a terpene/protein or whole-plant translation experiment.

Two additional exposure records sharpen this boundary. Artemisinin and artesunate produced dose-dependent DNA damage in cultured human HepG2 cells at the tested concentrations, without significant changes in CASP3 or SOD1 expression (Aquino et al. 2013). Separately, repeated ingestion of an *A. vulgaris* infusion was associated with confirmed serum thujone concentrations of 27.7 and 24.1 microgram/mL on successive days in a poisoning case (Di Lorenzo et al. 2018). These records are not universal dose thresholds or evidence against clinically indicated artemisinin therapy; they show why route, duration, species, product chemistry, and exposure biomarkers must be retained when comparing malaria compounds with wormwood preparations.

A human-lymphocyte study provides an independent in-vitro genotoxicity signal. Artemisinin and artemether increased comet-assay DNA damage and micronucleus frequency at the tested concentrations, while artemisinin also increased necrotic cell death after 48 hours (Cardoso et al. 2019). The experiment does not establish clinical risk, therapeutic exposure, or whole-plant toxicity, but it strengthens the requirement for orthogonal genotoxicity, reproductive, and pharmacokinetic testing before translating artemisinin-rich or wormwood-derived candidates.

## 8. Testable hypotheses and research agenda

The new production and analytical records add two explicit tests to this agenda. Nutrient-by-symbiont effects should be modeled as interactions among soil, phosphate, fungal inoculation, genotype, and cultivation system rather than as a universal artemisinin increase. Likewise, chiral oil composition should be tested as a possible moderator of parasite phenotype and host toxicity, with enantiomer-resolved standards and matched exposure rather than achiral peak percentages alone.

We propose six hypotheses: (1) WGD- and lineage-specific TPS retention predicts chemical diversity only when copy number is paired with expression and metabolomics; (2) antiparasitic chemistry associated with artemisinin-like products tracks pathway completion more closely than total TPS count; (3) essential-oil activity is often a chemotype-specific mixture property; (4)

secretory-tissue expression predicts accumulated compounds better than bulk leaf expression; (5) santonin/thujone abundance and nonvolatile STL profiles predict wormwood anthelmintic phenotypes only after preparation-specific fractionation and host-selectivity testing; and (6) cytotype- and diploidization-aware sampling will explain a measurable fraction of apparent *Artemisia* terpene-family expansion that is otherwise confounded by homeolog retention and annotation choice. Discriminating tests combine frozen assemblies, standardized annotation, orthology-aware gene trees, trichome transcriptomics, targeted metabolomics, enzyme assays, and fractionated parasite assays.

### 8.1 Minimum reporting standard for a future meta-analysis

Each observation should be stored as a specimen-level record with a stable taxon identifier, voucher or accession, tissue, developmental stage, locality and collection date, preparation, extraction yield, analytical platform, retention-index or spectral confirmation, compound identity confidence, abundance units, and assay metadata. A quantitative synthesis should not pool normalized peak percentages with absolute concentrations, purified compounds with oils, or promastigote-only endpoints with intracellular or in-vivo endpoints. This design makes heterogeneity a modeled variable rather than an unexplained nuisance.

### 8.2 Evidence-audit status of this review

The article's auditable evidence is distributed across seven linked layers. `sources.json` registers 538 sources; `evidence-matrix.csv` contains 553 source-linked records spanning chemistry, pathway biology, genomes, chemotypes, reviews, and parasite assays; `antiparasitic-evidence.csv` retains preparation, parasite-stage, dose, endpoint, and host-control fields; `parasite-protein-interact` separates 19 direct, functional, computational, and unresolved-target records; `evidence-appraisal.csv` applies a provisional, non-compensatory appraisal to all 129 antiparasitic records; `full-text-verification.csv` records 81 full-text sources and 96 linked evidence rows with preparation-specific extraction and unresolved eligibility gates; and `claim-audit.csv` maps 402 central article claims to source identifiers and permitted inference boundaries. The package also archives 1,408 bounded title/abstract screening decisions—1,387 within the priority queue and 21 historical or out-of-queue decisions—and 179 compound/specimen extraction records. These counts describe the current versioned evidence state, not a completed systematic-review denominator.

The audit supports an extensive, source-traceable synthesis but does not justify calling the review exhaustive. All 1,387 priority-queue records now have bounded title/abstract decisions; 81 sources have full-text verification, while 29 antiparasitic queue sources, quantitative harmonization, broad multi-species species-tree sampling, and common reannotation remain open. Claims based on the 553-record matrix are source-linked and scope-qualified; completion of title/abstract screening or a bounded full-text tranche does not itself make every retained record an included study. The exact package version, validation output, and SHA-256 manifest are stored in the Terpedia KB snapshot.

## 9. Conclusions

*Artemisia* terpene biology is best understood as a structured interaction between evolutionary history, gene-family innovation, tissue compartmentalization, ecological context, and analytical method. A publishable comparative review must therefore preserve specimen-level provenance and separate observed chemistry, direct enzyme evidence, annotation-supported hypotheses, mixture-level phenotypes, and unestablished claims. The Terpedia-linked dataset and protocol provide the basis for expanding this draft into a fully screened, quantitatively harmonized review.

### Data and code availability

All source tables, figures, and code are versioned in this package. Large comparative-genomics inputs and outputs, accessions, retrieval metadata, manifests, and checksums are archived in the linked Terpedia KB study.