


Homework Assignment #10

10.1

[47] PROBLEM 4.53

Scattering from a hydrogen atom ...

(a) Classical model of a hydrogen atom



Newton's second law

$$m a_r = F_r$$


$$-m \frac{v^2}{r} = -\frac{K e^2}{r^2}$$

$$\frac{1}{2} m v^2 = \frac{K e^2}{2r} ; U = -\frac{K e^2}{r}$$


Thus $T = -\frac{1}{2} U$ and $E = T + U = \frac{1}{2} U$.

(b) The scattering process

Before the collision



After the collision



In general, $E = \frac{1}{2} m v_1^2 - \frac{K e^2}{r_1} + \frac{1}{2} m v_2^2 - \frac{K e^2}{r_2} + \frac{e^2}{|\vec{r}_1 - \vec{r}_2|}$

(c) Conservation of energy

Energy before the collision $E = T_2 + \frac{K e^2}{2r}$


" after the collision $E = T_1 + \frac{K e^2}{2r'}$

Energy is conserved, so

$$T_1 = T_2 + \frac{K e^2}{2} \left(\frac{1}{r} - \frac{1}{r'} \right).$$

"Frequency" for an underdamped oscillator...

Consider a damped oscillator with $\beta < \omega_0$ (underdamped);



We can write the solution $x(t) = A e^{-\beta t} \cos(\omega_1 t - \delta)$ where δ is arbitrary. W.L.O.G. set $\delta = 0$.

$$x = A e^{-\beta t} \cos(\omega_1 t) \quad \text{and} \quad v = \dot{x} = A \left[-\beta e^{-\beta t} \cos(\omega_1 t) - \omega_1 e^{-\beta t} \sin(\omega_1 t) \right].$$

(a) Let τ_1 = the time between maxima.

$$\begin{aligned} \text{(a) The maxima of } x(t) \text{ occur} & \quad -\omega_1 e^{-\beta t} \sin \\ \text{when } v = 0; \text{ that is, } & \tan \omega_1 t = -\beta/\omega_1. \\ \tan(\omega_1(t_1 + \tau)) = \tan \omega_1 t & \Rightarrow \omega_1 \tau = 2\pi \\ \text{Thus } \tau = 2\pi/\omega_1. & \end{aligned}$$

(b) Let $\tau_1' = 2 \times$ the time between zeros.

(c) Suppose $\beta = \omega_0/2$; then $\omega_1 = (\omega_0^2 - \beta^2)^{1/2} = \omega_0 \sqrt{3}/2$

(b) The zeros of $x(t)$ occur when $\cos(\omega_1 t) = 0$.

$$\omega_1 t_n = \pi/2 + n\pi$$

$$\text{Define } \tau = t_{n+2} - t_n = \frac{1}{\omega_1} \left[\frac{\pi}{2} + (n+2)\pi - \frac{\pi}{2} - n\pi \right]$$

$$\text{Thus } \tau = 2\pi/\omega_1.$$

$$\text{(c) Suppose } \beta = \frac{\omega_0}{2}. \text{ Then } \omega_1 = \sqrt{\omega_0^2 - \beta^2} = \frac{\sqrt{3}}{2} \omega_0.$$


(This problem involves some computer calculations.)

(a) $x(t) = C_1 \exp(p_1 t) + C_2 \exp(p_2 t)$ where $p_{\{1,2\}} = -\beta \pm \sqrt{\beta^2 - \omega_0^2}$

Solve for C_1 and C_2 from $x(0) = x_0$ and $x'(0) = v_0$

$$C_1 = (p_2 x_0 + v_0) / (p_2 - p_1) \quad \text{and} \quad C_2 = (p_1 x_0 + v_0) / (p_1 - p_2)$$

(b) Sketches of the graphs of $x(t)$ for (i) $v_0 = 0$ and for (ii) $x_0 = 0$:



(c)

Let $\beta \rightarrow 0$. Then $p_1 \rightarrow p_1 \rightarrow i \omega_0$ and $p_2 \rightarrow -i \omega_0$.

Then $x(t) = C_1 \exp(i \omega_0 t) + C_2 \exp(-i \omega_0 t)$

which is an undamped oscillation.

[50] PROBLEM 5.37

A driven underdamped oscillator ...

10.4

(This problem is a computer problem, based on Example 5.3.)


Consider a driven damped oscillator, with these parameter values

$$\begin{aligned} \omega &= 2\pi & \text{and} & & \omega_0 &= \pi/2 \\ \beta &= 0.2 \omega_0 & \text{and} & & f_0 &= 1000. \end{aligned}$$

Note that $\beta < \omega_0$, so this is an underdamped oscillator.

Initial conditions: $x_0 = 0$ and $v_0 = 0$.

Plot $x(t)$



Compare with Example 5.3 and explain the similarities and difference:

/1/ Problem 5.37 has transient oscillations with a longer period; $\omega_0 = \pi/2$ compared to $\omega_0 = 10 \pi$.

/2/ Problem 5.37 has a longer decay time; $\beta = 0.1 \pi$ compared to $\beta = 0.5 \pi$.

/3/ Problem 5.37 and Example 5.3 have the same frequency of steady state motion; $\omega = 2\pi$ in both cases.

Another interpretation of the quality factor Q of a resonance ...

Consider a driven oscillator with $\omega = \omega_0$. The steady state solution is

$x_p(t) = A \cos(\omega t - \delta)$ where $A = f_0 / (2\beta \omega_0)$ and $\delta = \pi / 2$.

(A) The total energy is

$$E = \frac{1}{2} m \dot{x}^2 + \frac{1}{2} k x^2 = \frac{1}{2} m A^2 \omega^2 \sin^2(\omega t - \delta) + \frac{1}{2} m \omega_0^2 A^2 \cos^2(\omega t - \delta)$$

$$= \frac{1}{2} m \omega^2 A^2 \quad (\text{because } \omega = \omega_0)$$

(B) The energy dissipated in one period is

$$\Delta E = \int F_{\text{damping}} v dt = \int_0^{2\pi/\omega} m \cdot 2\beta \cdot v^2 dt = 2m\beta A^2 \omega^2 \left[\int \sin^2(u) du \right] \omega$$

$$= 2\pi m \beta \omega A^2$$

(C) Now calculate $E / \Delta E$. The result is $\omega / (4\pi\beta) = Q / (2\pi)$.

Thus $Q = 2\pi E / \Delta E$.

Oscillator driven by rectangular pulses ...

(This problem is a computer problem, based on Example 5.5)

- Parameter values

$$\tau_0 = 1; \quad \omega_0 = 2\pi; \quad \beta = 0.1; \quad ;$$

$$\Delta\tau = 0.25; \quad f_{\max} = 1; \quad ;$$

$$\tau = 1.0, 1.5, 2.0 \text{ and } 2.5; \quad \omega = 2\pi/\tau.$$

- Fourier coefficients for $f(t) = \sum_{n=0}^{\infty} a_n \cos(n\omega t)$

$$a_0 = \frac{f_{\max} \Delta\tau}{\tau} = \frac{0.25}{\tau}$$

$$a_n = \frac{2f_{\max}}{\pi n} \sin\left(\frac{\pi n \Delta\tau}{\tau}\right) = \frac{2}{\pi n} \sin\left(\frac{\pi n}{4\tau}\right) \text{ for } n \geq 1$$

- Forced coefficients for $x(t) = \sum_{n=0}^{\infty} A_n \cos(n\omega t - \delta_n)$

$$A_n = \frac{a_n}{\sqrt{(\omega_0^2 - n^2\omega^2)^2 + 4\beta^2 n^2 \omega^2}} = \frac{a_n/2\pi}{\sqrt{2\pi(1 - n^2/\tau^2)^2 + 0.04 \frac{n^2}{\tau^2}}}$$

$$\delta_n = \arctan \left[\frac{2\beta n \omega}{(2\pi)^2 - n^2 \omega^2} \right]$$

- Computer program and graphs next page.

Oscillator driven by rectangular pulses ...

(This problem is a computer problem, based on Example 5.5)

Problem 5.52


In[152]:= Remove["Global`*"]

```

In[173]:= Do[
  tau = 1.00000001 + 0.5 * (j - 1); omega = 2 * Pi / tau; beta = 0.1;
  a[0] = 0.25 / tau;
  Do[a[n] = 2 / (Pi * n) * Sin[Pi * n / 4 / tau], {n, 1, 10}];
  Do[A[n] = a[n] / (2 * Pi) / Sqrt[2 * Pi * (1 - n^2 / tau^2)^2 + 0.04 * n^2 / tau^2],
    {n, 0, 11}];
  Do[delta[n] = ArcTan[(2 * beta * n * omega) / (4 * Pi^2 - n^2 * omega^2)],
    {n, 0, 11}];
  x[t_] := Sum[A[n] * Cos[n * omega * t - delta[n]], {n, 0, 10}];
  lbl = StringJoin["τ = ", ToString[tau], " τ₀"];
  pl[j] = Plot[x[t], {t, 0, 7}, PlotRange -> {{0, 7}, {-0.4, 0.4}},
    PlotLabel -> lbl],
  {j, 1, 4}]

```

In[175]:= Show[GraphicsGrid[{{pl[1], pl[2]}, {pl[3], pl[4]}}], ImageSize -> Large]



Homework Assignment 10

Name _____

due in class Friday, November 11

Cover sheet : Staple this page in front of your solutions, with answers where indicated.

[47] Problem 4.53 **

What is $T_1 - T_2$? $= Ke^2/2 (1/r - 1/r')$

2 points

[48] Problem 5.25 **

What is the time between maxima? Also, what is ω_1 for part (c)?

$\tau = 2\pi / \omega_1$ $\omega_1 = \sqrt{3}/2 \omega_0$

2 points

[49] Problem 5.30 **

Hand in the computer program and computer plots. Check plots

2 points

[50] Problem 5.37 **

Hand in the computer program and computer plot.

Explain the similarities and differences compared to Example 5.3.

2 points

/1/ The decay time is longer; β is smaller.

/2/ The period of transient oscillations is longer; ω_0 is smaller.

/3/ The period of steady state oscillations is the same; ω is the same.

[50x] Problem 5.44 **

Express Q in terms of the parameters (m, ω, β). $Q = \omega / 2\beta$

2 points

[50xx] Problem 5.52 *** [Computer]

Hand in the computer program and computer plot.

Compare your results with those of the example.

The amplitudes are larger than in Figure 5.25.

3 points
