


• **FEDERAL UNIVERSITY OF AGRICULTURE ABEOKUTA**

PHS 101 PAST QUESTIONS



DEPARTMENT:

1. Use dimensional analysis to determine which of the following equations is certainly wrong.
 I. $x = vt$ II. $F = mv/a$ III. $F = m/a$ IV. $h = \sqrt{2g}$ V. $v = (2gh)^{1/2}$
 A. I only B. I and II C. H and II D. III, IV and V E. II only
2. A road journey takes 4 hours and 10 min at 80 km/h, including a half hour break for lunch. How much time would be saved by travelling at 100 km/h instead?
 A. 48 min B. 54 min C. 60 min D. 36 min E. 24 min
3. The position of a particle is given by $x = 5 \sin(120t)$ m where t is measured in seconds. Find the instantaneous velocity at 0.5 s.
 A. 5 m/s B. 4 m/s C. 520 m/s D. 300 m/s E. 600 m/s
4. A particle has an acceleration of $a = 7i + 2j \text{ ms}^{-2}$ for a period of 5 s. After 5 s the velocity is $v = 5i - 2k \text{ m/s}$. What was the initial velocity?
 A. $40i - 12k$ B. $17i - 9j$ C. $40i - 10j - 2k$ D. $17i - 7j - 2k$ E. $17i - 7j - 2k$
5. An Olympic athlete completes a long jump of 8.3 m with an initial speed of 9 m/s. What was the maximum increase in height of his waistline?
 A. 0.99 m B. 1.19 m C. 1.10 m D. 2.38 m E. 2.20 m
6. A ball thrown vertically upwards returns to its starting in 4 s. Find its initial speed.
 A. 0 m/s B. 10 m/s C. 20 m/s D. 20 m/s E. 80 m/s
7. How much force does it take to give a 20000 kg locomotive an acceleration of 1 ms^{-2} on a level track with a coefficient of rolling friction of 0.03?
 A. 6000 N B. 30 kN C. 24 kN D. 12 kN E. 36 kN
8. A 60 kg parachutist and her 7 kg parachute fall at a constant 6 m/s. Find the force on the woman due to the chute. A. 657 N B. 519 N C. 69 N D. 588 N E. 47 N
9. An 80 kg person pushes a 20 kg crate over a rough surface. Take $\mu_s = 0.8$ for the person and $\mu_k = 0.4$ for the crate. What is the maximum possible acceleration of the crate?
 A. 25.3 ms^{-2} B. 27.4 ms^{-2} C. 9.8 ms^{-2} D. 9.2 ms^{-2} E. 19.2 ms^{-2}
10. A resultant force of 20 N gives a body of mass m an acceleration of 8.0 ms^{-2} , and a body of mass m' an acceleration of 24 ms^{-2} . What acceleration will this force cause the two masses to acquire if fastened together? A. 12.0 ms^{-2} B. 6.0 ms^{-2} C. 32.0 ms^{-2} D. 16.0 ms^{-2} E. 3.0 ms^{-2}
11. The ice point and the triple point differ by what amount?
 A. 273 K B. 0.01 K C. 0 K D. 32 K E. 5/9 K

D'OLA #SUCCESS

2007/2008 SESSION FIRST EXAMINATIONS
PHS 101 GENERAL PHYSICS I

INSTRUCTION: Answer ALL questions.

TIME ALLOWED: 30 Minutes. Where necessary, take $g = 9.8 \text{ m/s}^2$.

NAME: _____

MACHINE NO: _____

DEPARTMENT: _____

1. Use dimensional analysis to determine which of the following equations is certainly wrong: I. $\lambda = v/f$ II. $F = mv/a$ III. $F = mv/g$ IV. $h = v^2/2$ V. $x = (2gl)^{1/2}$ A. I only B. I and II C. I and III D. III, IV and V E. If only

2. A road journey takes 4 hours and 30 min, at 80 km/h, including a half hour break for lunch. How much time would be saved by travelling at 100 km/h instead? A. 48 min B. 54 min C. 60 min D. 36 min E. 24 min

3. Train A has a length of 1 km and travels at 50 m/s. Train B has a length of 0.5 km and starts just as the rear of train A passes the front of train B. Train B has an acceleration of 3 m/s^2 and a maximum speed of 40 m/s. Use this information to answer questions 3 and 4.

3. When does B pass A, that is, when does the rear of B pass the front of A? A. 180 s B. 150 s C. 210 s D. 230 s E. 167 s

4. How far has A travelled in this time? A. 1.8 km B. 10.5 km C. 18 km D. 23 km E. 25 km

5. The position of a particle is given by $x = 5 \sin(120t) \text{ m}$ where t is time in seconds. Find the instantaneous velocity at 0.5 s.

- A. -4 m/s B. 4 m/s C. 520 m/s D. 600 m/s E. 600 m/s

6. A particle has an acceleration of $a = 2t + 3 \text{ m/s}^2$ for a period of 3 s. At the end of this time the velocity is $v_2 = 51 + 2t \text{ m/s}$. What was the initial velocity?

- A. $40i - 12j$ B. $17i - 9j$ C. $40i - 10j - 2k$ D. $17i - 7j - 2k$ E. $17i + 7j - 2k$

7. An Olympic athlete completes a long jump of 8.3 m with an initial speed of 9.7 m/s . What was the maximum increase in height of his waist-line?

- A. 0.99 m B. 1.19 m C. 1.10 m D. 2.38 m E. 2.20 m

8. A doughnut-shaped space station has an outer rim of radius of 1 km. What angular period should it rotate for a person at the rim to experience an acceleration of $g/5$? A. 140 s B. 140.5 s C. 4.44 s D. 4.49 s E. 1.5 s

9. A 21 g particle is subject to two forces that produce a resultant acceleration of $a = 4i - 2j \text{ m/s}^2$. If $F_1 = 3i + 2j$ N, find F_2 . A. $9i - 8j$ B. $7i - 4j$ C. $7i - 4j - 3k$ D. $3i - 5j - 3k$ E. $3i - 5j - 3k$

10. A 60 kg parachutist has a 7 kg parachute falling at a constant rate. Find the force on the woman due to the chute.

- A. 657 N B. 519 N C. 60 N D. 58 N E. 412 N

11. Two blocks with masses $m_A = 0.3 \text{ kg}$ and $m_B = 0.3 \text{ kg}$ hang in equilibrium as shown in figure 1. Find the tensions in the rope, T_1 and T_2 .

- A. 4.9 N, 2.9 N B. 3.9 N, 2.9 N C. 3.9 N, 2.9 N D. 3.9 N, 2.9 N E. 3.9 N, 2.9 N

D'OLA #SUCCESS

10 - 11

A projectile is launched from a point on level ground with velocity 30 m/s at an angle α above the

horizontal. The angle of position α is the vertical height reached a maximum?

(A) 20° (B) 45° (C) 90° (D) 180°

The angle of projection α is the horizontal range a maximum?

(A) 45° (B) 15° (C) 90° (D) 10°

12 - 13

A car at 25 m/s undergoes uniform deceleration when the brakes are applied, slowing down to

2.5 m/s

What is the deceleration in ms^{-2} ?

(A) -2.0 (B) 5.2 (C) -5.2 (D) 7.2

How far does it travel during this period?

(A) 9.0 m (B) 45 m (C) 4.5 m (D) 18.8 m

How far further does it travel before stopping if the deceleration remains constant?

(A) 2.5 m (B) 10.4 m (C) 9 m (D) 15 m

14 - 15

A ball rolls down from rest from a 1.4 m height

How long does it take to reach the ground?

(A) 1.6 s (B) 2.85 s (C) 2.85 s (D) 1.6

What is the velocity just before striking the ground?

(A) 16.6 m/s (B) 17.8 m/s (C) 17.8 m/s (D) 0.87 m/s

16 - 17

A man wishes to go to a particular point 1200 m east and 2100 m north of his present position.

What is the difference in a straight line to the point in question?

(A) 3300 m (B) 900 m (C) 900 m (D) 3320 m (E) 4260 m

At what angle should he proceed?

(A) 66.3° (B) 60.9° (C) 66.9° (D) 67°

18 - 20

A car of mass 1500 kg travels in a circular path of radius 20 m on a horizontal road surface at a speed of 10 m/s .

What is the frictional force necessary to prevent slipping?

(A) 10^4 N (B) $1.472 \times 10^4 \text{ N}$ (C) $1.284 \times 10^4 \text{ N}$ (D) $1.372 \times 10^4 \text{ N}$ (E) 15.343 N

What is the normal force between the road and the automobile?

(A) 10^4 N (B) $1.472 \times 10^4 \text{ N}$ (C) $1.284 \times 10^4 \text{ N}$ (D) $1.372 \times 10^4 \text{ N}$ (E) 15.343 N

What is the coefficient of friction between the tires and the road? Is it required to supply this force?

(A) 0.40 (B) 0.60 (C) 0.80 (D) 0.90

A thermometer which was not accurately calibrated indicates -0.5°C at the lower fixed point and 100°C at the upper fixed point. What temperature does the thermometer register when the actual temperature is 60°C ?

D'OLA #SUCCESS

NAME: _____

MATRIC NO: _____

DEPARTMENT: _____

PHYSICS

V = at
10x3

1. What is the dimension of G (gravitational constant) in this relation, $F = G \frac{Mm}{r^2}$ where other symbols have their usual meaning. A. $M^{-1}L^2T^{-2}$ B. ML^3T^{-2} C. $M^2L^2T^{-2}$ D. M^2LT^{-2}
2. What displacement must be added to a 50 cm displacement in the +x-direction to give a resultant displacement of 85 cm at 25° ? A. 53 cm at 45° B. 53 cm at 44° C. 45 cm at 50° D. 45 cm at 53°
3. A body falls freely from rest. Find the distance it falls in 3.0 s. A. 44 m B. 67 m C. 45 m D. 43 m
4. A child holding a wagon from rolling straight back down a driveway that is inclined at 20° to the horizontal. If the wagon weighs 150 N, with what force must the child pull on the handle if the handle is parallel to the incline? A. 48 N B. 51 N C. 49 N D. 60 N
5. A force acts on 2-kg mass and gives it an acceleration of 3 ms^{-2} . What acceleration is produced by the same force when acting on a mass of 4 kg? A. 2 ms^{-2} B. 3 ms^{-2} C. 4 ms^{-2} D. 5 ms^{-2}
6. Compute the work done against gravity by a pump that discharges 600 liters of fuel oil into a tank 20 m above the pump's intake. One cubic centimeter of fuel oil has a mass of 0.82 g. A. 98.4 kJ B. 98 kJ C. 9800 kJ D. 94.6 kJ
7. Compute the power output of a machine that lifts a 300 kg crate through a height of 20.0 m in a time of 60.0 s. A. 163 W B. 1.63 kW C. 162.3 kW D. 126 W
8. An engine expends 40.0 hp in propelling a car along a level track at a constant speed of 15.0 ms^{-1} . How large is the total retarding force acting on the car. A. 1.99 kN B. 129 N C. 199 N D. 289 kN
9. A 40 000 kg freight car is coasting at a speed of 5.0 ms^{-1} along a straight track when it strikes a 30 000 kg stationary freight car and couples to it. What will be their combined speed after impact? A. 3.0 ms^{-1} B. 2.78 ms^{-1} C. 5.0 ms^{-1} D. 2.9 ms^{-1}
10. During collision, which of the following expression represents an inelastic case? A. $e < 1$ B. $e > 1$ C. $e = 1$ D. $e = 0$
11. At very low temperature, the viscosity of certain liquids drops to virtually zero in a phenomenon called, A. Superconductivity B. Superfluidity C. Superelasticity D. Resonance

D'OLA #SUCCESS

13. At very low temperature, the viscosity of certain liquids drops to virtually zero in a phenomenon called ~~A. Superconductivity~~ ☒ ~~B. Superfluidity~~ ☒ C. Superthermal capacity
D. Resonance E. All of the above
14. "It is impossible to construct a continuously operating machine that does work with energy from external source" This is a statement of ~~A. First law of thermodynamics~~
B. Second law of thermodynamics C. Zeroth law of thermodynamics
D. All of the above E. Joule's law
14. A process whereby ΔU , ΔQ and W in the first law of thermodynamics are non zero is referred to as ~~A. Isothermal~~ B. Isochoric C. Adiabatic ☒ D. Isobaric E. Isometric
15. The mechanical equivalent of heat is expressed as
~~A. $mc\Delta\theta$~~ B. $t\Delta x$ ☒ C. W/Q D. $1/t$ E. $1/T$

D'OLA #SUCCESS

NAME: _____
MATIC NO: _____
DEPARTMENT: _____

E: 29-08-2001

TIME ALLOWED: 96 Minutes

INSTRUCTIONS: ANSWER ALL QUESTIONS AND SUBMIT THE DETACHED ANSWER SHEET.

Questions 1-2

To keep an object moving in a circle at constant speed, a force is required given as $F = \frac{mv^2}{r}$ where m = mass, v = velocity and r = radius of its circular path.

- Determine the numerical values of a, b and c
 A. 1, 2 and 3 B. 1, 2 and -3 C. -1, -2 and 3 D. 1, 2 and -1
 E. 3, 2 and -1.

Which of the following represent the correct expression for the centripetal force?
 A. $F = \frac{Kmv^2}{r}$ B. $F = \frac{Kmv^2}{r^2}$ C. $F = \frac{Kmv^2}{r^3}$ D. $F = \frac{Kmv^2}{2r}$ E. $F = \frac{Kmv^2}{r}$

Which of the following is the correct expression for Kepler's third law?

- A. $T^2 = 4\pi^2/GM$ B. $T = 4\pi^2/GM$ C. $T^2 = 2\pi^2/GM$ D. $T = \pi^2/GM$
 E. None of the above.

D'OLA #SUCCESS

For Questions 4-5

A 2.0kg block is attached to a spring for which $K = 20 \text{ N/m}$. It is held at an extension of 5.0cm and then released at $t = 0$.

4. Find the displacement as a function of time.

- A. $X = 0.5 \sin(10t + \pi/2) \text{ m}$ B. $X = 0.05 \sin(10t + \pi/2) \text{ m}$ C. $X = 5.0 \sin(10t + \pi/2) \text{ m}$
 D. $X = 5.0 \sin(10t - \pi/2) \text{ m}$ E. $X = 0.5 \sin(10t - \pi/2) \text{ m}$

5. Find the velocity and acceleration when $X = +A/2$.

- A. 4.3 m/s and -2.5 m/s^2 B. 0.43 m/s and 2.5 m/s^2
 C. 0.43 m/s and -2.5 m/s^2 D. 2.5 m/s and 0.43 m/s^2 E. None of the above.

6. When the net work done on an object is equal to the change in its kinetic energy, we have:

- A. Conservation of linear momentum B. Conservation of energy
 C. Work-energy theorem D. Mechanical energy E. None of the above

7. A car moving with a velocity of 18 km/h accelerates at 1.5 m/s^2 . The velocity and position of the car at this time are:

- A. 1.25 m/s and 20 m B. 2.0 m/s and 12.5 m C. 20 m/s and 12.5 m
 D. 20 m/s and 12.5 m E. 12.5 m/s and 2.0 m

8. Two cars, 7 km away from each other, move towards each other. The first car has a constant velocity of 54 km/h . At what time and where will they meet?

- A. 200 sec and 300 m B. 200 sec and 30 m C. 300 sec and 200 m
 D. 20.0 sec and 300 m E. 200 sec and 3.00 m

TIME ALLOWED: 30 Minutes

Where necessary take $g = 10 \text{ ms}^{-2}$

NAME

MATRIC NO

DEPARTMENT

D'OLA #SUCCESS

1. The speed v of a wave on a string depends on the tension F in the string and the mass per unit length $\frac{m}{l}$ of the string. If it is known that $[F] = [MLT^{-2}]$ and the constants a and b in the equation for the speed of a wave on a string $v = \text{constant } F^a (\frac{m}{l})^b$

A. 1, 2 D. 1, $\frac{1}{2}$ C. $\frac{1}{2}$, $\frac{1}{2}$ D. 2, $\frac{1}{2}$ E. $\frac{1}{2}$, $\frac{1}{2}$

or Q2-Q3

A car is accelerating uniformly as it passes two checkpoints that are 30m apart. The time taken between checkpoints is 4.0s and the car's speed at the first checkpoint is 5.0ms⁻¹.

Find the car's acceleration A. 1.5ms⁻² B. 7.5ms⁻² C. 60ms⁻² D. 3.0ms⁻² E. 4.5ms⁻² Ans 1-25

Find its speed at the second checkpoint. A. 5ms⁻¹ B. 10ms⁻¹ C. 15ms⁻¹ D. 7.5ms⁻¹ E. 6.0ms⁻¹

2. A plane starts from rest and accelerates along the ground before takeoff. It moves 600m in 12s. Find the distance moved during the twelfth second.

A. 10m B. 600m C. 26m D. 720m E. 360m

A ball thrown vertically upwards returns to its starting in 4s. Find its initial speed.

A. 4ms⁻¹ B. 16ms⁻¹ C. 2ms⁻¹ D. 20ms⁻¹ E. 10ms⁻¹

3. How much force does it take to give a 20000kg locomotive an acceleration of 1 ms⁻² on a level track with a coefficient of rolling friction of 0.03?

A. 6000N B. 30kN C. 24kN D. 12kN E. 6N

A resultant force of 20N gives a body of mass m an acceleration of 2.0ms⁻², and a body of mass $2m$ an acceleration of 24ms⁻². What acceleration will this force cause the two masses to acquire if fastened together?

A. 12.0ms⁻² B. 6.0ms⁻² C. 12.0ms⁻² D. 16.0ms⁻² E. 3.0ms⁻²

A 7kg object is subjected to two forces $F_1 = 20i + 30j$ N and $F_2 = 3i - 50j$ N. Find the acceleration of the object.

A. $\frac{20}{7}i + \frac{12}{7}j$ B. $\frac{12}{7}i + \frac{20}{7}j$ C. $\frac{12}{7}i - \frac{20}{7}j$ D. $\frac{20}{7}i + \frac{30}{7}j$ E. $\frac{8}{7}i - \frac{30}{7}j$

In Fig. 1, $FP = 20$ N, $m_1 = m_2 = 2$ kg, and the acceleration is 0.50ms⁻², what will be the tension in the connecting cord if the frictional forces on the two blocks are equal? How large is the frictional force on either block?

A. 10N, 3N B. 14N, 3N C. 8N, 6N D. 10N, 5.5N E. 14N, 6N

$$471 = \frac{20}{7}$$

$$281 = 20j$$

15 9 - 10
A projectile is launched from a point on level ground with velocity 30 m/s at an angle α above the horizontal.

At what angle of position α is the vertical height reached a maximum?

(A) 20° (C) 45° (D) 90° (E) 180°

At what angle of projection α is the horizontal range a maximum?

(A) 45° (C) 15° (D) 90° (E) 10°

15 11 - 13

A car moving at 25 m/s undergoes uniform deceleration when the brakes are applied, slowing down to rest.

It takes 4 s to stop.

What is the deceleration in ms^{-2} ?

(A) -2.0 (C) 5.2 (D) -5.2 (E) 7.2

How far does it travel during this period?

(A) 9.0 m (C) 45 m (D) 4.5 m (E) 18.8 m

How far does it travel before stopping if the deceleration remains constant?

(A) 2.5 m (C) 10.4 m (D) 9 m (E) 15 m

15 14 - 15

A ball rolls down from rest from a 1.4 m height.

How long does it take to reach the ground?

(A) 1.07 s (C) -2.85 s (D) 2.85 s (E) 1.6 s

What is the velocity just before striking the ground?

(A) 16.6 m/s (C) -17.8 m/s (D) 17.8 m/s (E) 0.87 m/s

15 16 - 17

A person wants to go to a particular point 1200 m east and 2100 m north of his present position.

What is the difference in a straight line to the point in question?

(A) 3300 m (C) 900 m (D) 3320 m (E) 4260 m

At what angle should he proceed?

(A) 66.3° (C) 60.9° (D) 66.9° (E) 67°

15 18 - 20

A car of mass 1500 kg travels in a circular path of radius 20 m on a horizontal road surface at a constant speed of 10 m/s .

What is the frictional force necessary to prevent slipping?

(A) 10^4 N (B) $1.472 \times 10^4 \text{ N}$ (C) $1.284 \times 10^4 \text{ N}$ (D) $1.372 \times 10^4 \text{ N}$ (E) 15.543 N

What is the normal force between the road and the automobile?

(A) 10^4 N (B) $1.472 \times 10^4 \text{ N}$ (C) $1.284 \times 10^4 \text{ N}$ (D) $1.372 \times 10^4 \text{ N}$ (E) 15.53 N

What is the coefficient of friction between the tires and the road? It is required to supply this force?

(A) 0.40 (C) 0.60 (D) 0.80 (E) 0.90

15 21 - 22
A thermometer was not accurately calibrated and reads -0.5°C at the lower fixed point and 100°C at the upper fixed point. What temperature does the thermometer register when the actual temperature is 60°C ?

D'OLA #SUCCESS

PHS 101

University of Agriculture, Abeokuta.

Department of Physics

2003/2004 Continuous Assessment Test

PHS 101-GENERAL PHYSICS I

Time allowed: 30mins

Shade clearly across only one correct letter option in the spaces provided at the end. All the following physical quantities have derived units except?

- * 1. A. Acceleration. B. Density. C. Speed. D. Current. E. Pressure

2. To keep an object moving in a circle at a constant speed requires a force called?

- A. Tangential force. B. Frictional force. C. Centrifugal force. D. Gravitational force. E. Centripetal force

3. A uniform ladder 8.0m long weighs 220N, rests on rough ground and is propped against a vertical rough wall at an angle θ to the horizontal. If $\mu = 0.4$ for the ground and the wall surfaces, find the value of θ when slip is about to take place

- A. 46.9° B. 38.7° C. 34.6° D. 43.6° E. 68.7°

4. A train accelerates uniformly from rest to reach 54km/h in 200seconds, after which the speed remains constant for 300seconds. At the end of this time the train decelerates to rest in 160seconds. Find the total distance traveled.

- A. 15.0km B. 7.125km C. 71.25km D. 712.5km E. 150.0km

For Questions 5 and 6

A bullet is fired vertically upwards with an initial velocity of 98m/s from the top of a building 100m high. ($g = 9.8\text{ms}^{-2}$)

5. Find the maximum height reached above the ground.

- A. 49.0m B. 4900.0m C. 490.0m D. 98.0m E. 980.0m

6. Find the total time before reaching the ground as well as the velocity on landing.

- A. 10sec. And 107.5m/s B. 20.97sec. and 10.75m/s C. 20.97sec. and 107.5m/s D. 10sec. And 10.75m/s E. 10sec. And 20.97m/s

7. A body moving with S.H.M has a velocity of 3m/s when 375mm from its midposition and an acceleration of 1m/s² when 250mm from the midposition. Calculate the periodic time and the amplitude.

- A. 3.142sec and 1.55m B. 1.75sec and 1.55m C. 1.57sec and 1.55m D. 3.142sec and 1.55m E. None of the above

8. A body oscillates along a straight line with S.H.M. The frequency is 0.1Hz and the amplitude 300mm. Find the position of maximum displacement.

- A. 205.2mm B. 281.1mm C. 299.3mm D. 21.7mm E. 21.9mm


9. If at the triple point (273.16K) of water, the resistance of the element in a platinum resistance thermometer is 8.75 Ω , what is the resistance at room temperature (301.82K) if the latter is measured on the scale of the resistance thermometer?

- A. 7.92 Ω B. 9.424 Ω C. 9.124 Ω D. 9.63 Ω E. 7.424 Ω

10. A thermometer which was not accurately calibrated indicates 47°C at the lower fixed point and 106°C at the upper fixed point. What is the true temperature when the thermometer registers 63.4°C?

- A. 166.7°C B. 39.0°C C. 16.4°C D. 1667.0°C E. 16.7°C

D'OLA #SUCCESS



NAME: Ask me again

MATRIC NO:

DEPARTMENT:

1. Use dimensional analysis to determine which of the following equations is certainly wrong
 I. $\lambda = vt$ II. $F = mv/a$ III. $F = mv/t$ IV. $h = v^2/2g$ V. $v = (2gh)^{1/2}$
 A. I only B. I and II C. II and III D. III, IV and V E. II only

2. A road journey takes 4 hours and 30 min at 80 km/h, including a half hour break for lunch. How much time would be saved by travelling at 100 km/h instead?
 A. 48 min B. 54 min C. 60 min D. 36 min E. 24 min

3. The position of a particle is given by $s = 5 \sin(120t)$ m where t is measured in seconds. Find the instantaneous velocity at 0.5s.
 A. 5 m/s B. 4 m/s C. 520 m/s D. 300 m/s E. 600 m/s

4. A particle has an acceleration of $a = 7i + 2j$ ms⁻² for a period of 5s. After it a time the velocity is $v = 5i - 2k$ m/s. What was the initial velocity?
 A. $40i - 12k$ B. $17i - 9j$ C. $40i - 10j - 2k$ D. $17i - 7j - 2k$ E. $7i + j - 2k$

5. An Olympic athlete completes a long jump of 8.3m with an initial speed of 9 ms⁻¹. What was the maximum increase in height of his waistline?
 A. 0.99m B. 1.19m C. 1.10m D. 2.38m E. 2.20m

6. A ball thrown vertically upwards returns to its starting in 4s. Find its initial speed.
 A. 0 ms⁻¹ B. 16 ms⁻¹ C. 20 ms⁻¹ D. 20 ms⁻¹ E. 80 ms⁻¹

7. How much force does it take to give a 20000 kg locomotive an acceleration of 1 ms⁻² on a level track with a coefficient of rolling friction of 0.03?
 A. 6000 N B. 30 kN C. 24 kN D. 12 kN E. 36 kN

8. A 60 kg parachutist and her 7 kg parachute fall at a constant 6 m/s. Find the force on the woman due to the chute. A. 657 N B. 519 N C. 69 N D. 585 N E. 412 N

9. An 80 kg person pushes a 20 kg crate over a rough surface. Take $\mu_s = 0.8$ for the person and $\mu_k = 0.4$ for the crate. What is the maximum possible acceleration of the crate?
 A. 25.3 ms⁻² B. 27.4 ms⁻² C. 9.8 ms⁻² D. 3.92 ms⁻² E. 39.2 ms⁻²

10. A resultant force of 20 N gives a body of mass m an acceleration of 8.0 ms⁻², and a body of mass m' an acceleration of 24 ms⁻². What acceleration will this force cause the two masses to acquire if fastened together? A. 12.0 ms⁻² B. 6.0 ms⁻² C. 32.0 ms⁻² D. 16.0 ms⁻² E. 8.0 ms⁻²

11. The ice point and the triple point differ by what amount?
 A. 373 K B. 0.01 K C. 0 K D. 32 K E. 5/9 K

D'OLA #SUCCESS

12. At very low temperature, the viscosity of certain liquid drops to virtually zero in a phenomenon called ☒ A. Superconductivity ☒ B. Superfluidity ☐ C. Supercritical fluidity ☐ D. Resonance ☐ E. All of the above
13. "It is impossible to construct a continuously operating machine that does work with energy from external source" This is a statement of ☒ A. First law of thermodynamics ☐ B. Second law of thermodynamics ☐ C. Zeroth law of thermodynamics ☐ D. All of the above ☐ E. Joule's law
14. A process whereby ΔU , ΔQ and W in the first law of thermodynamics are non zero is referred to as ☐ A. Isothermal ☐ B. Isochoric ☐ C. Adiabatic ☒ D. Isobaric ☐ E. Isomeric
15. The mechanical equivalent of heat is expressed as ☒ A. $mc\Delta\theta$ ☐ B. $\rho\Delta x$ ☒ C. W/Q ☐ D. $1/v$ ☐ E. $1/f$

D'OLA #SUCCESS

NAME: _____
 MATRIC NO: _____
 DEPARTMENT: _____

Office Use

12. The ice point and the triple point differ by what amount? A. 273 K B. 0.01 K
C. 0 K D. 32 K
13. It is impossible to construct a continuously operating machine that does work with energy from internal source. This is a statement of A. First law of thermodynamics B. second law of thermodynamics C. zeroth law of thermodynamics D. all of the above
14. The process that occur so rapidly that there is no transfer of heat between the system and the environment are known as A. Cyclic processes B. Adiabatic processes
C. Isochoric processes D. Isothermal processes
15. Which of the following is not correct of a cyclic process?
A. $W > 0$ and $Q < 0$ for counter clockwise cycle B. $W < 0$ and $Q > 0$ for a clockwise cycle
C. $Q + W = 0$ D. None of the above

D'OLA #SUCCESS

NAME:

MATRIC NO: DEPARTMENT:

	A	B	C	D
1.				
2.				
3.				
4.				
5.				
6.				
7.				
8.				
9.				
10.				
11.				
12.				
13.				
14.				
15.				

or Questions 9-11

A stone is thrown vertically upward with an initial velocity of 12 m/s from the top of a building. It takes 3.4 sec for the stone to hit the ground.

Find the height of the building.

- A. 170.0 m B. 1.70 m C. 17.0 m D. 1700 m E. None of the above

Find the maximum height reached.

- A. 4.2 m B. 2.42 m C. 242.0 m D. 0.242 m E. 240.0 m

Find the velocity of the stone when it reaches the ground.

- A. 22.0 m/s B. 0.22 m/s C. 22.0 m/s D. 2.2 m/s E. None of the above

Find the gravitational force the sun exerts on the earth given that the mass of the sun is $M_s = 1.99 \times 10^{30} \text{ kg}$ and mass of the earth is $M_e = 5.98 \times 10^{24} \text{ kg}$ and the mean distance between the sun and the earth is $1.50 \times 10^{11} \text{ m}$.

- A. $3.53 \times 10^{12} \text{ N}$ B. $3.53 \times 10^{19} \text{ N}$ C. $3.53 \times 10^{23} \text{ N}$ D. $0.353 \times 10^{23} \text{ N}$ E. $0.353 \times 10^{19} \text{ N}$

Which of the following is/are true of frictional force?

- A. Friction always opposes the motion.
B. Friction is always parallel to the surface.
C. The maximum value of static friction is $f_{\text{max}} = \mu_s N$.
D. If the object is sliding $f_k = \mu_k N$.
E. All of the above.

D'OLA #SUCCESS

or questions 14-15

A block is released from rest on a 53° inclined plane. The coefficient of kinetic friction between the block and the surface is 0.7 .

The acceleration of the block is:

- A. 5.5 m/s^2 B. 38.0 m/s^2 C. 3.8 m/s^2 D. 55 m/s^2 E. 0.38 m/s^2

The velocity of the block after traveling 4 m is:

- A. 3.8 m/s B. 3.5 m/s C. 0.38 m/s D. 55 m/s E. None of the above

A 0.5 kg block is hung from a spring and brought to equilibrium, the spring stretches 0.4 m . The spring constant of the spring is:

- A. 250 N/m B. 2.50 N/m C. 25 N/m D. 0.25 N/m E. 0.025 N/m

A 0.2 m long rod is rotating about an axis through its end with a constant angular velocity of 12 rev/s . Find its angular velocity, the velocity of its center and the velocity of its tip.

- A. 12.6 rad/s , 5.0 m/s and 10 m/s B. 12.6 rad/s , 10 m/s and 5.0 m/s
C. 10 rad/s , 12.6 m/s and 10.0 m/s D. 10 rad/s , 5.0 m/s and 12.6 m/s
E. 20 rad/s , 5.0 m/s and 10.0 m/s

A block is thrown along a horizontal surface with an initial velocity of 9 m/s . The coefficient of kinetic friction between the block and the surface is 0.6 . How far will the block move before it stops?

- A. 7.5 m B. 0.675 m C. 0.75 m D. 0.75 m E. 13.5 m

When a 0.5kg block is hung from a spring and brought to rest, the spring stretches 0.2m. Find the spring constant of the spring.

- A. 250N/m B. 25N/m C. 2.5N/m D. 10.0N/m E. 0.1N/m

How much power is required to push a 100kg box along a horizontal surface at a constant speed of 5m/s if the coefficient of kinetic friction between the box and the surface is 0.3?

- A. 300W B. 30W C. 1500W D. 150W E. 3000W

Which of the following expression represents Hooke's law?

- A. $F = kx$ B. $F = -1/2Kx^2$ C. $F = \mu N$ D. $F = Kx^2$ E. $F = Kx$

Questions 22-24

Volume of liquid passing per second, V , through a pipe when the flow is steady, depends on coefficient of viscosity η of the liquid, radius r of the pipe and the pressure gradient P/L cause the flow.

Which of the following is the correct dimensional expression?

- A. $L^3T^{-1} = (MLT^{-1})^2 L^2 (ML^{-1}T^{-2})^2$
 B. $L^3T^{-1} = (MLT^{-1})^2 L^2 (ML^{-1}T^{-2})$
 C. $L^3T^{-1} = (ML^{-1}T^{-1})^2 L^2 (ML^{-1}T^{-2})$
 D. $L^3T^{-1} = (ML^{-1}T^{-1})^2 L^2 (ML^{-1}T^{-2})^2$
 E. None of the above.

What are the values of x , y , z ?

- A. $x=-1$, $y=4$ and $z=1$ B. $x=-2$, $y=3$ and $z=0$ C. $x=1$, $y=-4$ and $z=1$
 D. $x=-1$, $y=4$ and $z=-1$ E. $x=-2$, $y=-3$ and $z=1$

Which of the following is the correct expression for the volume V ?

- A. $V = KPr^3/\eta l$ B. $V = KPr^4/\eta l$ C. $V = KPr^2/\eta l$
 D. $V = KPr^4/\eta l^2$ E. $V = KPr^3/\eta l^2$

D'OLA #SUCCESS

Questions 25-26:

A block of mass 100g is attached to the end of a spring whose force constant is 100N/m. The block slides on a horizontal surface for which $\mu_k = 0.1$, and is extended 5cm and then released.

Find the work done by the spring up to the point at which it is compressed by 3cm.

- A. 0.032J B. 0.078J C. 0.024J D. 0.002J E. 0.24J

Find the net work done on the block up to this point.

- A. 0.032J B. 0.0078J C. 0.024J D. 0.002J E. 0.24J

A block of mass 100g is dropped 2cm along a horizontal surface by a force acting at an angle of 33° to the horizontal. The initial speed is 3m/s and $\mu_k = 1/3$. Find the change in kinetic energy and the final speed of the block.

- A. 2J and 5m/s B. 32J and 0.5m/s C. 32J and 5m/s
 D. 20J and 5m/s E. None of the above.

- A. 63.4°C B. 36.4°C C. 46.3°C D. 43.6°C E. 34.6°C
22. The resistance R_{θ} of platinum wire at temperature $\theta^{\circ}\text{C}$, measured on the gas scale is given by $R_{\theta} = R_0(1 + a\theta + b\theta^2)$, where $a = 3.9 \times 10^{-3}$, $b = -5.6 \times 10^{-7}$, and $R_0 = 9.8\Omega$. What is the resistance at 100°C ?
 A. 46.13Ω B. 34.16Ω C. 13.47Ω D. 74.13Ω E. 31.47Ω
23. At what temperature do the Celsius and Fahrenheit scales have the same numerical value?
~~A. -40°C~~ B. -12°C C. -30°C D. -60°C E. 0°C
24. A container of gas has a volume of 0.10m^3 at a pressure of $2.0 \times 10^5 \text{Nm}^{-2}$ and a temperature of 27°C . Find the new pressure if the gas is heated at constant volume to 87°C .
 A. $4.2 \times 10^5 \text{Nm}^{-2}$ ~~B. $2.4 \times 10^5 \text{Nm}^{-2}$~~ C. $1.2 \times 10^5 \text{Nm}^{-2}$
 D. $2.1 \times 10^5 \text{Nm}^{-2}$ E. $5.0 \times 10^4 \text{Nm}^{-2}$
25. In an experiment to verify Newton's law of cooling, it was observed from the cooling curve that at 60°C the rate of cooling of the body was 1.0Ks^{-1} and at 53.4°C the rate of cooling was 0.8Ks^{-1} . Calculate the room temperature at the time of the experiment.
 A. 27°C B. 56.4°C C. 30°C D. 25°C E. 15°C

D'OLA #SUCCESS

For Questions 26-29

The first law of thermodynamics may be expressed by the relation $\Delta Q = \Delta U + \Delta W$, where Q is the quantity of heat given to a system of internal energy U and W is the external work done. Assume the system is an ideal gas enclosed in a cylinder with a piston.

1. $\Delta U = \Delta Q - \Delta W$
2. $\Delta W = V\Delta P$, where V is the volume of the gas and P is the pressure
3. $\Delta U = -\Delta W$ - adiabatic
4. $\Delta Q = -\Delta W$ - isothermal
5. $\Delta Q = \Delta U$ - isochoric

26. Which of the following cases 1 - 5 above represent(s) an ISOTHERMAL change?
 A. 1 only B. 2 only ~~C. 3 only~~ D. 4 only E. 5 only
27. Which of the cases represent(s) an ISOBARIC change?
~~A. 1 only~~ B. 2 only C. 3 only D. 4 only E. 5 only
28. Which of the cases represent(s) an ISOCHORIC change?
 A. 1 only B. 2 only C. 3 only D. 4 only E. 5 only
29. Which of the cases represent(s) an ADIABATIC change?
 A. 1 only B. 2 only C. 3 only D. 4 only E. 5 only

For Questions 30-33