


Problem Set #2
CHEM101A: General College Chemistry

Donald Aingworth IV

August 29, 2025

1 Topic A Problem 12

What mass of Fe_2O_3 would react with 20.00 g of Zn? The chemical equation for this reaction is:




1.1 Solution

The simple stoichiometry is the way to go here.

$$20.00\text{g} \times \frac{1 \text{ mol Zn}}{65.38\text{g}} \times \frac{1 \text{ Fe}_2\text{O}_3}{3 \text{ Zn}} \times \frac{159.7 \text{ g Fe}_2\text{O}_3}{1 \text{ mol Fe}_2\text{O}_3} = \boxed{16.28 \text{ g Fe}_2\text{O}_3} \quad (2)$$

2 Topic A Problem 13

x moles of C_4H_{10} reacts with oxygen according to the following equation:



- a) How many moles of water are formed?
- b) How many moles of oxygen are consumed?

2.1 Solution (a)

The ratio of C_4H_{10} used to H_2O created in this reaction is 1:5. With x moles of C_4H_{10} , that would give us $\boxed{5x \text{ mol H}_2\text{O}}$.

2.2 Solution (b)

The ratio of C_4H_{10} used to O_2 consumed in this reaction is 2:13. With x moles of C_4H_{10} , that would give us $\boxed{\frac{13}{2}x \text{ mol O}_2}$.

3 Topic A Problem 14

10.00 g of N_2 is mixed with 33.61 g of F_2 , and the elements react according to the following equation:



- a) Which element is the limiting reactant?
- b) What is the theoretical yield of NF_3 ?
- c) If the reaction goes to completion, how many grams of the excess reactant will remain?
- d) Set up an ICE table for this reaction.

3.1 Solution (a)

First, we calculate the theoretical yields for each for the reactants.

$$m_{\text{N}_2} = 10.00 \text{ g} \times \frac{1 \text{ mol N}_2}{28.02 \text{ g N}_2} \times \frac{2 \text{ NF}_3}{1 \text{ N}_2} \times \frac{71.01 \text{ g NF}_3}{1 \text{ mol NF}_3} = 50.69 \text{ g NF}_3 \quad (5)$$

$$m_{\text{F}_2} = 33.61 \text{ g} \times \frac{1 \text{ mol F}_2}{38.00 \text{ g F}_2} \times \frac{2 \text{ NF}_3}{3 \text{ F}_2} \times \frac{71.01 \text{ g NF}_3}{1 \text{ mol NF}_3} = 41.87 \text{ g NF}_3 \quad (6)$$

With a lower final mass, $\boxed{\text{F}_2}$ is the limiting reactant.

3.2 Solution (b)

The theoretical yield was found in part (a). $\boxed{41.87 \text{ g NF}_3}$

3.3 Solution (c)

Use a similar strategy to part (a).

$$33.61 \text{ g} \times \frac{1 \text{ mol F}_2}{38.00 \text{ g F}_2} \times \frac{1 \text{ N}_2}{3 \text{ F}_2} \times \frac{28.02 \text{ g N}_2}{1 \text{ mol N}_2} = 8.261 \text{ g NF}_3 \quad (7)$$

Subtract this from the available mass of N_2 to get the final N_2 .

$$10.00 \text{ g N}_2 - 8.261 \text{ g N}_2 = \boxed{1.74 \text{ g N}_2} \quad (8)$$

3.4 Solution (d)

I used tabular for this table. Please excuse any poor or improper formatting.

mol	N_2	$+ 3 \text{F}_2$	$\longrightarrow 2 \text{NF}_3$
I	0.3569	0.8844	0
C	-0.2948	-0.8844	0.5896
E	0.0621	0	0.5896

For those interested in how I went about getting these values, I can explain. I started with the initial mass of F_2 , which has been previously established to be the limiting reactant, and converted that to moles. I did (roughly) the same thing for the known quantity of N_2 initially. We also start with no NF_3 . Assuming the percentage yield to be 100%, every mole of F_2 would be used, so the Change row for F_2 would be the negative of the initial quantity of F_2 . Multiply that by the ratio of N_2 to F_2 ($\frac{1}{3}$) to get the Change row of N_2 . The same can be done for NF_2 , just taking the negative thereof and with a ratio of $\frac{2}{3}$ instead of $\frac{1}{3}$. With all of this, we only have to add the initial and the change together (respecting the positive or negative signs) to get the values for the End row.

4 Topic A Problem 15

- a) If 58.26 g of iodine reacts with excess aluminum, what is the theoretical yield of aluminum iodide? The reaction is $2 \text{Al} + 3 \text{I}_2 \longrightarrow 2 \text{AlI}_3$.
- b) If 56.11 g of aluminum iodide is actually formed in the reaction in part a, what is the percent yield of aluminum iodide?

4.1 Solution (a)

Watch me use the power of Stiochiometry Magic.

$$58.26 \text{ g} \times \frac{1 \text{ mol I}_2}{253.8 \text{ g I}_2} \times \frac{2 \text{ AlI}_3}{3 \text{ I}_2} \times \frac{407.68 \text{ g AlI}_3}{1 \text{ mol AlI}_3} = \boxed{62.39 \text{ g AlI}_3} \quad (9)$$


4.2 Solution (b)

Here we use the formula for the pecent yield.

$$\text{PY} = \frac{\text{AY}}{\text{TY}} \times 100\% = \frac{56.11 \text{ g}}{62.39 \text{ g}} \times 100\% = 0.8994 \times 100\% = \boxed{89.94\%} \quad (10)$$

5 Topic A Problem 16

A chemist mixes 16.00 g of HCl with 10.00 g of Mg and obtains an 81.3% yield of MgCl_2 . What mass of MgCl_2 did the chemist obtain? The chemical reaction is:



5.1 Solution

First calculate the theoretical yield of MgCl_2 in the cases of HCl and Mg being the limiting reactants.

$$MM(\text{MgCl}_2) = 24.31 \text{ g/mol} + 2 * 35.45 \text{ g/mol} = 95.21 \text{ g/mol} \quad (12)$$

$$MM(\text{HCl}) = 1.008 \text{ g/mol} + 35.45 \text{ g/mol} = 36.458 \text{ g/mol} \quad (13)$$

$$m_{\text{Mg}} = 10.00 \text{ g} \times \frac{1 \text{ mol Mg}}{24.31 \text{ g Mg}} \times \frac{1 \text{ MgCl}_2}{1 \text{ Mg}} \times \frac{95.21 \text{ g MgCl}_2}{1 \text{ mol MgCl}_2} \quad (14)$$

$$= 39.16 \text{ g MgCl}_2 \quad (15)$$

$$m_{\text{HCl}} = 16.00 \text{ g} \times \frac{1 \text{ mol HCl}}{36.458 \text{ g HCl}} \times \frac{1 \text{ MgCl}_2}{2 \text{ HCl}} \times \frac{95.21 \text{ g MgCl}_2}{1 \text{ mol MgCl}_2} \quad (16)$$

$$= 20.89 \text{ g MgCl}_2 \quad (17)$$

The latter is lower, so the HCl would be the limiting reactant and 20.89 g MgCl_2 would be the theoretical yield. Multiplying this by the (decimal version of) the percentage yield to get the actual yield.


$$20.89 \text{ g MgCl}_2 * 0.813 = \boxed{17.0 \text{ g MgCl}_2} \quad (18)$$

6 Topic A Problem 17

How many milliliters of liquid Br_2 (density = 3.1 g/mL) will react with 6.143 g of Cr, if the product of this reaction is CrBr_3 ?

6.1 Solution

First write a chemical equation for this and balance it.




The rest of the path is paved with the magic of Stoichiometry.

$$6.143 \text{ g Cr} \times \frac{1 \text{ mol Cr}}{52.00 \text{ g Cr}} \times \frac{3 \text{ Br}_2}{2 \text{ Cr}} \times \frac{159.8 \text{ g Br}_2}{1 \text{ mol Br}_2} \times \frac{1 \text{ mL}}{3.1 \text{ g}} = \boxed{9.1 \text{ mL Br}_2} \quad (19)$$

7 Topic A Problem 18

Ethane (C_2H_6) reacts with oxygen according to the following chemical equation:



- a) If you mix 5 moles of C_2H_6 with 13 moles of O_2 , how many moles of each substance will you end up with, assuming the reaction goes to completion? Include an ICE table in your answer.
- b) If you mix 81.43 g of C_2H_6 with 194.60 g of O_2 , how many grams of each substance will you end up with, assuming the reaction goes to completion? Include an ICE table in your answer. (Note: your ICE table should be in terms of moles.)
- c) A chemist mixes 3.414 moles of O_2 with an unknown number of moles of C_2H_6 . The chemist obtains 1.657 moles of O_2 . How many moles of C_2H_6 must have been present originally, assuming the reaction went to completion? Include an ICE table in your answer.

7.1 Solution (a)

mol	$2 \text{C}_2\text{H}_6$	+	7O_2	\longrightarrow	4CO_2	+	$6 \text{H}_2\text{O}$
I	5		13		0		0
C	$-\frac{26}{7}$		-13		$\frac{52}{7}$		$\frac{78}{7}$
E	$\frac{9}{7}$		0		$\frac{52}{7}$		$\frac{78}{7}$

You will end up with $\boxed{1.286 \text{ mol C}_2\text{H}_6}$, $\boxed{0 \text{ mol O}_2}$, $\boxed{7.429 \text{ mol CO}_2}$, and $\boxed{11.143 \text{ mol H}_2\text{O}}$.

7.2 Solution (b)

mol	$2 \text{C}_2\text{H}_6$	+	7O_2	\longrightarrow	4CO_2	+	$6 \text{H}_2\text{O}$
I	2.708		6.081		0		0
C	-1.737		-6.081		3.475		5.212
E	0.971		0		3.475		5.212

Final table

Calculations

Convert grams to moles for oxygen and ethane.

$$n(\text{O}_2) = \frac{m}{MM} = \frac{194.60 \text{ g O}_2}{32.00 \text{ g/mol}} = 6.08125 \text{ mol O}_2 \quad (21)$$

$$n(\text{C}_2\text{H}_6) = \frac{m}{MM} = \frac{81.43 \text{ g C}_2\text{H}_6}{30.068 \text{ g/mol}} = 2.708194759 \text{ mol C}_2\text{H}_6 \quad (22)$$

Next, we check which would result in the most product, for the coefficient c of each reactant.

$$ML = \frac{n}{c} \quad (23)$$

$$ML(\text{O}_2) = \frac{6.08125 \text{ mol O}_2}{7 \text{ O}_2} = 0.86875 \text{ mol} \quad (24)$$

$$ML(\text{C}_2\text{H}_6) = \frac{2.708194759 \text{ mol C}_2\text{H}_6}{2 \text{ C}_2\text{H}_6} = 1.357 \text{ mol} \quad (25)$$

This makes the O_2 the limiting reactant. We can just fill out the ICE table's 'C' (change) values from here by using ratios of coefficients of reactants and products. I won't go write all my calculations here, but they tend to have a simple formula.

$$n_2 = n_1 * \frac{c_2}{c_1} \quad (26)$$

This leads into the filling out of the bottom row (end), with a simple formula, $E = I + C$. From here, all we need to do is convert moles to grams.

$$MM(\text{C}_2\text{H}_6) = 2 * 12.01 \text{ g/mol} + 6 * 1.008 \text{ g/mol} = 30.068 \text{ g/mol} \quad (27)$$

$$MM(\text{O}_2) = 2 * 16.00 \text{ g/mol} = 32.00 \text{ g/mol} \quad (28)$$

$$MM(\text{CO}_2) = 12.01 + 2 * 16.00 \text{ g/mol} = 44.01 \text{ g/mol} \quad (29)$$

$$MM(\text{H}_2\text{O}) = 2 * 1.008 \text{ g/mol} + 16.00 \text{ g/mol} = 18.016 \text{ g/mol} \quad (30)$$

$$m = MM * n \quad (31)$$

$$m(\text{C}_2\text{H}_6) = 30.068 \text{ g/mol C}_2\text{H}_6 * 0.971 \text{ mol} = \boxed{29.20 \text{ g C}_2\text{H}_6} \quad (32)$$

$$m(\text{O}_2) = 32.00 \text{ g/mol O}_2 * 0 \text{ mol} = \boxed{0 \text{ g O}_2} \quad (33)$$

$$m(\text{CO}_2) = 44.01 \text{ g/mol CO}_2 * 3.475 \text{ mol} = \boxed{152.9 \text{ g CO}_2} \quad (34)$$

$$m(\text{H}_2\text{O}) = 18.016 \text{ g/mol H}_2\text{O} * 5.212 \text{ mol} = \boxed{93.90 \text{ g H}_2\text{O}} \quad (35)$$

7.3 Solution (c)

We'll just put together an ICE table and fill it out until we have enough information to get the answer. We don't even need to fill it out completely.

mol	2 C ₂ H ₆	+	7 O ₂	→	4 CO ₂	+	6 H ₂ O
I	0.502		3.414		0		0
C	-0.502		-1.757				
E	0		1.657				

I have a few steps I made for this.


1. We can start by filling out our knowns. In the interest of candor, we don't even know if the initial amount of CO_2 and H_2O is 0, that's just an assumption I made.

2. Since the O_2 has some left over at the end, it is definitely not the limiting reactant. That makes C_2H_6 the limiting reactant, so none of it should be left by the end and we can fill that out.
3. The formula of $\text{C} = \text{E} - \text{I}$ can be applied for Oxygen, giving us our change in O_2 .
 - If you ever took Physics or Calculus (maybe even Algebra), you'll recognize this as equivalent to $\Delta x = x_f - x_i$.
4. We can use the coefficient ratio mentioned in part (b), this time $\frac{2}{7}$, for the change in C_2H_6 .
5. The formula $\text{C} = \text{E} - \text{I}$ can be turned into $\text{I} = \text{E} - \text{C}$ to get initial quantity of C_2H_6 .

With everything filled out, we have the final answer of $0.502 \text{ mol C}_2\text{H}_6$.

8 Topic A Problem 19

Ammonia reacts with oxygen according to the following chemical equation:



Suppose you mix x moles of NH_3 with y moles of O_2 .

- If NH_3 is the limiting reactant, how many moles of each substance will you end up with, assuming the reaction goes to completion? Include an ICE table in your answer.
- If O_2 is the limiting reactant, how many moles of each substance will you end up with, assuming the reaction goes to completion? Include an ICE table in your answer.
- If you end up with $0.4y$ moles of O_2 , what must the relationship be between x and y , assuming the reaction goes to completion?

8.1 Solution (a)

mol	4NH_3	+	3O_2	\longrightarrow	2N_2	+	$6\text{H}_2\text{O}$
I	x		y		0		0
C	$-x$		$-\frac{3}{4}x$		$\frac{1}{2}x$		$\frac{3}{2}x$
E	0		$y - \frac{3}{4}x$		$\frac{1}{2}x$		$\frac{3}{2}x$

My answers are found in the End (E) section of the ICE table. I did everything in my head.

8.2 Solution (b)

mol	4NH_3	+	3O_2	\longrightarrow	2N_2	+	$6\text{H}_2\text{O}$
I	x		y		0		0
C	$-\frac{4}{3}y$		$-y$		$\frac{2}{3}y$		$2y$
E	$x - \frac{4}{3}y$		0		$\frac{2}{3}y$		$2y$

My answers are found in the End (E) section of the ICE table. I did everything in my head.

8.3 Solution (c)

mol	4NH_3	+	3O_2	\longrightarrow	2N_2	+	$6\text{H}_2\text{O}$
I	x		y				
C	$-x$		$-0.6y$				
E	0		$0.4y$				

Out of respect for the unknowns, I have left the boxes unnecessary to calculate for completion of this problem blank, all of which lie in the products.

- The things I did initially fill out are the final amount of O_2 and the initial amounts of O_2 and NH_3 .

2. The fact that there is O_2 remaining suggests that the NH_3 is the limiting reactant, so it would be 0 at the end.
3. $\text{E} = \text{C} + \text{I} \rightarrow \text{C} = \text{E} - \text{I}$ gives us our change values. This in turn gives us an equation for their relationship, bearing in mind the ratio of their coefficients.

$$x * \frac{3}{4} = \frac{3}{5}y \quad (37)$$

$$\boxed{x = \frac{4}{5}y = 0.8y} \quad (38)$$

9 Topic A Problem 20

You have x grams of $\text{Na}_2\text{Cr}_2\text{O}_7$. How many grams of CrCl_3 will be formed if the $\text{Na}_2\text{Cr}_2\text{O}_7$ undergoes the reaction below? Express your answer in terms of x .



9.1 Solution

First, we find the molar masses of each involved chemical that we need, which are only $\text{Na}_2\text{Cr}_2\text{O}_7$ and CrCl_3 .

$$MM(\text{Na}_2\text{Cr}_2\text{O}_7) = 2 * 22.99 \text{ g/mol} + 2 * 52.00 \text{ g/mol} + 7 * 16.00 \text{ g/mol} \quad (40)$$

$$= 261.98 \text{ g/mol} \quad (41)$$

$$MM(\text{CrCl}_3) = 52.00 \text{ g/mol} + 3 * 35.45 \text{ g/mol} = 158.35 \text{ g/mol} \quad (42)$$


From here, the magic of Stoichiometry will guide us.

$$m(\text{CrCl}_3) = x \times \frac{1 \text{ mol}}{261.98 \text{ mol}} \times \frac{2 \text{ CrCl}_3}{1 \text{ Na}_2\text{Cr}_2\text{O}_7} \times \frac{158.35 \text{ g}}{1 \text{ mol}} \quad (43)$$

$$= \boxed{1.209x \text{ g CrCl}_3} \quad (44)$$

10 Topic A Problem 21

A metal sample weighing 1.410 g contains a mixture of copper and aluminum. When excess HCl is added to this sample, the aluminum reacts as follows:



849 mL of H_2 (density 0.08264 g/L) is produced. Calculate the mass percentage of each element in the original sample. Note that copper does not react with HCl.

10.1 Solution

The way we start this is by finding the amount (mass) of Al that reacts. This is done through the magic of stoichiometry.

$$m(\text{Al}) = 0.849 \text{ L} \times \frac{0.08264 \text{ g}}{1 \text{ L}} \times \frac{1 \text{ mol}}{2.016 \text{ g}} \times \frac{2 \text{ Al}}{3 \text{ H}_2} \times \frac{26.98 \text{ g}}{1 \text{ mol}} \quad (46)$$

$$= 0.626 \text{ g Al} \quad (47)$$

From here, we use the formula for mass percentage of $MP = \frac{m}{m_\Sigma}$ for each chemical's mass. The value of m_Σ is the total mass, in this case $m_{1.410}$.

$$MP(\text{Al}) = \frac{0.626}{1.410} \times 100\% = \boxed{44.4\% \text{ Al}} \quad (48)$$

$$MP(\text{Cu}) = \left(1 - \frac{0.626}{1.410}\right) \times 100\% = \boxed{55.6\% \text{ Cu}} \quad (49)$$

11 Topic B Problem 1

Answer each of the following questions about making solutions.

a) If you dissolve 4.18 g of solid $\text{Mg}(\text{NO}_3)_2$ in enough water to make 150 mL of solution, what will be the molarity of the resulting solution?

b) If you need to make 100 mL of 1.08 M CaCl_2 , what mass of solid CaCl_2 will you need?

c) You have 25.0 g of solid KCl , and you use all of it to make a 0.500 M KCl solution. What volume of solution did you make?

11.1 Solution (a)

First convert grams to moles.

$$MM(\text{Mg}(\text{NO}_3)_2) = 24.31 + 2 * 14.01 + 6 * 16.00 = 148.33 \text{ g/mol} \quad (50)$$

$$n(\text{Mg}(\text{NO}_3)_2) = \frac{4.18 \text{ gram Mg}(\text{NO}_3)_2}{148.33 \text{ g/mol}} = 0.0282 \text{ mol} \quad (51)$$

From here, we just find the molarity.

$$[\text{Mg}(\text{NO}_3)_2] = \frac{n}{V} = \frac{0.0282 \text{ mol}}{0.150 \text{ L}} = \boxed{0.188 \text{ M Mg}(\text{NO}_3)_2} \quad (52)$$

11.2 Solution (b)

First find the number of moles of CaCl_2 .

$$n = V * [\text{CaCl}_2] = 0.100 \text{ L} * 1.08 \text{ M CaCl}_2 = 0.108 \text{ mol CaCl}_2 \quad (53)$$

From here, use the molar mass to find the total mass.

$$MM(\text{CaCl}_2) = 40.08 + 2 * 35.45 = 110.98 \text{ g/mol} \quad (54)$$

$$m = n * MM = 0.108 \text{ mol CaCl}_2 * 110.98 \text{ g/mol} = \boxed{12.0 \text{ M CaCl}_2} \quad (55)$$

11.3 Solution (c)

First convert grams to moles.

$$MM(\text{KCl}) = 39.10 + 35.45 = 74.55 \text{ g/mol} \quad (56)$$

$$n = \frac{25.0 \text{ g KCl}}{74.55 \text{ g/mol}} = 0.335 \text{ mol KCl} \quad (57)$$

From here, we can find the volume.

$$V = \frac{n}{[\text{KCl}]} = \frac{0.335 \text{ mol KCl}}{0.500 \text{ M KCl}} = \boxed{0.671 \text{ L}} \quad (58)$$

12 Topic B Problem 2

Answer the following questions about dilutions.

- a) If you add 100 mL of water to 10 mL of 0.605 M HCl, what will be the molarity of the resulting solution?
- b) You have 200 mL of 1.50 M HNO₃. If you wish to dilute this solution to a final concentration of 0.300 M, what volume of water should you add?
- c) You need to make 1.50 liters of 0.400 M NaOH by diluting a 2.00 M NaOH solution. What volume of the 2.00 M NaOH should you use, and what volume of water should you add to it?

12.1 Solution (a)

Use the relationship between concentration and volume.

$$C_1 V_1 = C_2 V_2 \quad (59)$$

$$C_2 = C_1 \frac{V_1}{V_2} = 0.605 * \frac{100}{110} = \boxed{0.550 \text{ M HCl}} \quad (60)$$

12.2 Solution (b)

First the relationship between concentration and volume.

$$C_1 V_1 = C_2 V_2 \quad (61)$$

$$V_2 = V_1 \frac{C_1}{C_2} = 200 * \frac{1.50}{0.300} = 200 * 5 = 1000 \text{ mL} \quad (62)$$

Now take the difference between final and initial volume to find the added volume.

$$\Delta V = V_2 - V_1 = 1000 - 200 = \boxed{800 \text{ mL}} \quad (63)$$

12.3 Solution (c)

Here we can use the same relationship between concentrations and volumes. We can start with making a ratio for the volume that would turn 1 liter of 2.00 M NaOH to 0.400 M NaOH.

$$C_1 V_1 = C_2 V_2 \quad (64)$$

$$V_2 = V_1 \frac{C_1}{C_2} = 1 * \frac{2.0}{0.4} = 5 \text{ L} \quad (65)$$

This means that every liter of 2.00 M NaOH would require 4 liters of water to form 0.400 M NaOH. For each chemical (pure water and 2.00 M NaOH), we can multiply the final volume by its necessary ratio ($\frac{4}{5}$ and $\frac{1}{5}$, respectively).

$$V_{\text{water}} = 1.5 \text{ L} * \frac{4}{5} = 1.2 \text{ L} \quad (66)$$

$$V_{\text{NaOH}} = 1.5 \text{ L} * \frac{1}{5} = 0.3 \text{ L} \quad (67)$$

For the water, we need $\boxed{1.2\text{ L}}$. For the NaOH, we need $\boxed{0.3\text{ L}}$.

13 Topic B Problem 3

All of the compounds below dissolve in water. Which of them are strong electrolytes, which are weak electrolytes, and which are nonelectrolytes?

- | | | | |
|-------------------------------|----------------------------|--------------------------------------|-------------------------------|
| a) NaCl | d) MgCrO_4 | g) $\text{C}_2\text{H}_5\text{OH}$ | j) H_2SO_4 |
| b) $\text{Mg}(\text{NO}_3)_2$ | e) H_3PO_4 | h) $\text{HC}_3\text{H}_5\text{O}_3$ | k) NH_4Br |
| c) HClO_2 | f) AgF | i) CH_3CN | l) $(\text{CH}_3)_2\text{CO}$ |

13.1 Solution

- | | | |
|-----------------------|-----------------------|-----------------------|
| a) Strong electrolyte | e) Weak electrolyte | i) Non-electrolyte |
| b) Strong electrolyte | f) Strong electrolyte | j) Strong electrolyte |
| c) Weak electrolyte | g) Non-electrolyte | k) Strong electrolyte |
| d) Strong electrolyte | h) Weak electrolyte | l) Non-electrolyte |

I *really* need to work on this.

14 Topic B Problem 4

What ions (if any) are present in each of the following solutions, and what is the molar concentration of each ion?

a) 0.1 M NaBr

c) 0.2 M FeCl₃

b) 0.04 M KNO₃

d) 1.5 M (NH₄)₂SO₄

14.1 Solution

a/ 0.1 M Na⁺, 0.1 M Br⁻

b/ 0.04 M K⁺, 0.04 M NO₃⁻

c/ 0.2 M Fe³⁺, 0.6 M Cl⁻

d/ 3.0 M NH₄⁺, 1.5 M SO₄²⁻

15 Topic B Problem 5

How many moles of each ion are present in 175 mL of 0.147 M $\text{Fe}_2(\text{SO}_4)_3$?

15.1 Solution

Start with the iron (III) Fe^{3+} . There are 2 Fe^{3+} ions per $\text{Fe}_2(\text{SO}_4)_3$ atom, so we can find the molarity of the iron (III) atom. By multiplying that by the volume, we will find the number of moles of iron ions.

$$n = 2 \frac{\text{Fe}^{3+}}{\text{Fe}_2(\text{SO}_4)_3} * 0.147 \text{ M Fe}_2(\text{SO}_4)_3 * 0.175 \text{ liter} = \boxed{0.0515 \text{ mol Fe}^{3+}} \quad (68)$$

Following that, there are three SO_4^{2-} ions per $\text{Fe}_2(\text{SO}_4)_3$ atom, so we can find the molarity of the sulfate. Multiply the molarity by the volume to get the number of moles.

$$n = 3 \frac{\text{SO}_4^{2-}}{\text{Fe}_2(\text{SO}_4)_3} * 0.147 \text{ M Fe}_2(\text{SO}_4)_3 * 0.175 \text{ liter} = \boxed{0.0772 \text{ mol SO}_4^{2-}} \quad (69)$$

16 Topic B Problem 6

Which of the following are acceptable ways to make one liter of 1 M NaCl?

- a) Put 1 liter of water into a container, then add 1 mole of NaCl and stir until the NaCl dissolves.
- b) Put 1 mole of NaCl into a container, then add 1 liter of water and stir until the NaCl dissolves.
- c) Put 1 mole of NaCl into a container, then add water with stirring until the total volume reaches 1 liter.

16.1 Solution

The one mole of NaCl will inevitably have some volume, which will not be lost when it dissolves. Both (a) and (b) result in a solution whose volume is the sum of the volume of the NaCl and the water it is in, with exactly one mole of NaCl. The molarity of the result will be $\frac{1 \text{ mol}}{1+x \text{ L}}$ NaCl, for some volume of the NaCl x . It doesn't matter if the resultant molarity is greater or smaller than $\frac{1 \text{ mol}}{1 \text{ L}}$ NaCl, but it is undeniable that they are not equal, leaving the two strategies as unusable. Strategy (c) works, though. You measure the moles from the start, and you know exactly how much water there is.

17 Topic B Problem 7

Janet dissolves 6.50 g of solid potassium phosphate in enough water to make 100.0 mL of solution. Farid then adds enough water to the solution to reduce the concentration of potassium ions to 0.250 M. How much water did Farid add?

17.1 Solution

What's in a name? The name "Potassium Phosphate" suggests that the chemical has the formula K_3PO_4 . We can convert grams to moles and use that to find the concentration, then use $C_1V_1 = C_2V_2$ to find the final amount of water, then subtract the initial volume to find the volume added.

$$MM(\text{K}_3\text{PO}_4) = 3 * 39.10 + 30.97 + 4 * 16.00 = 212.27 \text{ g/mol} \quad (70)$$

$$[\text{K}_3\text{PO}_4] = 6.50 \text{ g} \times \frac{1 \text{ mol}}{212.27 \text{ g}} \times \frac{1}{0.100 \text{ L}} \text{ K}_3\text{PO}_4 = 0.306 \text{ mol/L} \quad (71)$$

$$C_1V_1 = C_2V_2 \quad (72)$$

$$V_2 = V_1 \frac{C_1}{C_2} = 0.1 \times \frac{0.306}{0.250} = 0.12249 \text{ L} \quad (73)$$

$$\Delta V = V_2 - V_1 = 0.122495 - 0.100 = \boxed{0.0225 \text{ L}} \quad (74)$$

18 Topic B Problem 8

Gerardo dissolves 8.213 g of solid $\text{Mg}(\text{NO}_3)_2$ in enough water to make 200.0 mL of solution. Marciela then adds enough solid $\text{Al}(\text{NO}_3)_3$ to increase the concentration of nitrate ions to 0.900 M. Assuming that the solution volume does not change significantly, what mass of $\text{Al}(\text{NO}_3)_3$ did Marciela add?

18.1 Solution

First calculate the molarity of NO_3 .

$$MM(\text{Mg}(\text{NO}_3)_2) = 24.31 + 2 * 14.01 + 6 * 16.00 \quad (75)$$

$$= 148.33 \text{ g/mol } \text{Mg}(\text{NO}_3)_2 \quad (76)$$

$$MM(\text{Al}(\text{NO}_3)_3) = 26.98 + 3 * 14.01 + 9 * 16.00 \quad (77)$$

$$= 213.01 \text{ g/mol } \text{Al}(\text{NO}_3)_3 \quad (78)$$

$$[\text{Mg}(\text{NO}_3)_2] = 8.213 \times \frac{1 \text{ mol}}{148.33 \text{ g}} \times \frac{1}{0.2000 \text{ L}} \quad (79)$$

$$= 0.2768 \text{ M } \text{Mg}(\text{NO}_3)_2 \quad (80)$$

$$[\text{NO}_3]_i = 2 * 0.2768 \text{ M} = 0.5537 \text{ M } \text{NO}_3 \quad (81)$$

This means that the added molarity is $0.3463 \text{ M } \text{NO}_3$, which makes the molarity of $\text{Al}(\text{NO}_3)_3$ to be $0.3463 \text{ M } \text{NO}_3 \times \frac{1}{3} = 0.1154 \text{ M } \text{Al}(\text{NO}_3)_3$. Multiply this by the volume to get the number of moles of $\text{Al}(\text{NO}_3)_3$ added. This can be used with the molar mass to find the mass of $\text{Al}(\text{NO}_3)_3$ used.

$$m(\text{Al}(\text{NO}_3)_3) = MM(\text{Al}(\text{NO}_3)_3) * [\text{Al}(\text{NO}_3)_3] * V \quad (82)$$

$$= 213.01 \text{ g/mol } \text{Al}(\text{NO}_3)_3 \times 0.1154 \text{ M } \text{Al}(\text{NO}_3)_3 \times 0.2000 \text{ L} \quad (83)$$

$$= 4.918 \text{ g} \quad (84)$$

This means that Marciela added 4.918 g.

19 Topic B Problem 9

Chantelle dissolves 2.35 g of NaCl, 3.12 g of CaCl₂, and 1.88 g of FeCl₃ in enough water to make 175 mL of solution. What is the molarity of chloride ions in this solution?

19.1 Solution

Find the number of moles of Cl from each chemical and sum them together.

$$MM(\text{NaCl}) = 22.99 + 35.45 = 58.44 \text{ g/mol} \quad (85)$$

$$MM(\text{CaCl}_2) = 40.08 + 2 * 35.45 = 110.98 \text{ g/mol} \quad (86)$$

$$MM(\text{FeCl}_3) = 55.85 + 3 * 35.45 = 162.2 \text{ g/mol} \quad (87)$$

$$\begin{aligned} n(\text{Cl}^-) &= \sum \frac{c_i m_i}{MM} = \frac{2.35}{58.44} + \frac{2 * 3.12}{110.98} + \frac{3 * 1.88}{162.2} \\ &= 0.0402 + 0.0562 + 0.0348 = 0.1312 \text{ mole Cl}^- \end{aligned} \quad (88)$$

$$= 0.0402 + 0.0562 + 0.0348 = 0.1312 \text{ mole Cl}^- \quad (89)$$

This can be divided by the volume to get the molarity.

$$[\text{Cl}^-] = \frac{n}{V} = \frac{0.1312 \text{ mole Cl}^-}{0.175 \text{ L}} = \boxed{0.750 \text{ M Cl}^-} \quad (90)$$

20 Topic B Problem 10

Wenzhou prepares 200 mL of a solution of SnCl_4 in which the concentration of chloride ions is 0.240 M.

a) What is the molarity of the SnCl_4 solution (i.e. what should the bottle be labeled)?

b) What mass of SnCl_4 did Wenzhou use?

20.1 Solution (a)

The concentration of SnCl_4 is a quarter of the concentration of Cl^- , since the ratio of SnCl_4 to Cl^- is 4:1.

$$0.240 \text{ M Cl}^- \times \frac{1 \text{ SnCl}_4}{4 \text{ Cl}^-} = \boxed{0.060 \text{ M SnCl}_4} \quad (91)$$

20.2 Solution (b)

Multiply molarity by volume, then multiply it by molar mass.

$$MM(\text{SnCl}_4) = 118.7 + 4 * 35.45 = 260.5 \text{ g/mol} \quad (92)$$

$$m(\text{SnCl}_4) = [\text{SnCl}_4] * V * MM(\text{SnCl}_4) \quad (93)$$

$$= 0.060 \text{ M SnCl}_4 \times 0.200 \text{ L} \times 260.5 \text{ g/mol} \quad (94)$$

$$= \boxed{3.13 \text{ g}} \quad (95)$$

21 Topic B Problem 11

A beaker holds x liters of 0.2 M AlBr_3 . Give answers to each part below in terms of x .

- a) How many moles of aluminum ions are in this solution?
- b) How many moles of bromide ions are in this solution?
- c) How much water must you add if want to dilute the original solution to a concentration of 0.02 M?

21.1 Solution (a)

The molarity of the aluminum is the molarity of the aluminum bromide. Multiply this by the number of liters to get the number of moles of ions.

$$n = 0.2 \text{ M Al}^{3+} \times x \text{ L} = \boxed{0.2x \text{ mol Al}^{3+}} \quad (96)$$

21.2 Solution (b)

The molarity of the bromine is thrice the molarity of the aluminum bromide. Multiply this by the number of liters to get the number of moles of ions.

$$n = 0.6 \text{ M Br}^- \times x \text{ L} = \boxed{0.6x \text{ mol Br}^-} \quad (97)$$

21.3 Solution (c)

First use $C_1V_1 = C_2V_2$ (I should really find out what that really is), then use $\Delta V = V_f - V_i$.

$$C_1V_1 = C_2V_2 \quad (98)$$

$$V_2 = V_1 \frac{C_1}{C_2} = x * \frac{0.2}{0.02} = 10x \text{ L} \quad (99)$$

$$\Delta V = V_f - V_i = 10x - x = \boxed{9x} \quad (100)$$

22 Topic B Problem 12

Using the solubility rules, determine which of the following compounds are insoluble in water. There is a solubility rules handout available in Canvas.

- | | | |
|--------------------------------------|---------------------|---------------------------------|
| a) $\text{K}_2\text{Cr}_2\text{O}_7$ | d) ZnBr_2 | g) $\text{Ba}_3(\text{PO}_4)_2$ |
| b) $\text{Mn}(\text{NO}_3)_2$ | e) MgSO_4 | |
| c) FeS | f) NaHCO_3 | |

22.1 Solution

Time for the final boss of this week.

- a/ Soluble (contains K^+)
- b/ Soluble (contains NO_3^-)
- c/ Insoluble (S^{2-} is insoluble with Fe^{2+})
- d/ Soluble (contains Br^- , soluble with Zn^{2+})
- e/ Soluble (contains SO_4^{2-} , soluble with Mg^{2+})
- f/ Soluble (contains HCO_3^- , which contains CO_3^{2-} , soluble with Na^+)
- g/ Insoluble (contains PO_4^{3-} , insoluble with Ba^{2+})

Contents

1	Topic A Problem 12	2
1.1	Solution	2
2	Topic A Problem 13	3
2.1	Solution (a)	3
2.2	Solution (b)	3
3	Topic A Problem 14	4
3.1	Solution (a)	4
3.2	Solution (b)	4
3.3	Solution (c)	4
3.4	Solution (d)	4
4	Topic A Problem 15	6
4.1	Solution (a)	6
4.2	Solution (b)	6
5	Topic A Problem 16	7
5.1	Solution	7
6	Topic A Problem 17	8
6.1	Solution	8
7	Topic A Problem 18	9
7.1	Solution (a)	9
7.2	Solution (b)	9
7.3	Solution (c)	10
8	Topic A Problem 19	12
8.1	Solution (a)	12
8.2	Solution (b)	12
8.3	Solution (c)	12
9	Topic A Problem 20	14
9.1	Solution	14
10	Topic A Problem 21	15
10.1	Solution	15
11	Topic B Problem 1	16
11.1	Solution (a)	16
11.2	Solution (b)	16
11.3	Solution (c)	16

12 Topic B Problem 2	17
12.1 Solution (a)	17
12.2 Solution (b)	17
12.3 Solution (c)	17
13 Topic B Problem 3	19
13.1 Solution	19
14 Topic B Problem 4	20
14.1 Solution	20
15 Topic B Problem 5	21
15.1 Solution	21
16 Topic B Problem 6	22
16.1 Solution	22
17 Topic B Problem 7	23
17.1 Solution	23
18 Topic B Problem 8	24
18.1 Solution	24
19 Topic B Problem 9	25
19.1 Solution	25
20 Topic B Problem 10	26
20.1 Solution (a)	26
20.2 Solution (b)	26
21 Topic B Problem 11	27
21.1 Solution (a)	27
21.2 Solution (b)	27
21.3 Solution (c)	27
22 Topic B Problem 12	28
22.1 Solution	28