

Problem Set #10

CHEM101A: General College Chemistry

Donald Aingworth

October 24, 2025

23 Topic E Problem 23

What are the possible values of m_ℓ and m_s for a 4f electron?

23.1 Solution

The value of ℓ is 3. This means that the values of m_ℓ are $-3, -2, -1, 0, 1, 2, 3$. The electron can have either positive or negative spin, so the possible values of

m_s are $+\frac{1}{2}, -\frac{1}{2}$.

24 Topic E Problem 24


Explain why it is impossible for an orbital to have $n = 3$ and $\ell = 3$. (Hint: think about what these numbers are telling you about nodes.)

24.1 Solution

The value of n is 0 and it is 1 more than the number of nodes. ℓ describes the number of angular nodes. If the value of n is 3, there are $n - 1 = 2$ total nodes. If the value of ℓ is 3, there are 3 angular nodes. The prior two statements contradict each other.

25 Topic E Problem 25

The two graphs below show the electron density and radial probability for an atomic orbital. (Both graphs show the same orbital.) Use the graphs to answer questions a through e below.




- Which graph is the electron density plot?
- What is the most probable distance between the electron and the nucleus for this orbital? (You will need to estimate it from one of the graphs.)
- How many radial nodes does this orbital have? How can you tell?
- Does this orbital have any angular nodes? How can you tell?
- If $n = 4$ for this orbital, what orbital is it?

25.1 Solution

- The electron density plot is the one on the left (Ψ^2).
- The most probable distance from the center would be about 1250 pm.
- The electron density plot goes to zero at two points, so there would be two radial nodes.
- Since the graph of the electron density is zero at the origin, this means that there are radial nodes.
- 4p

26 Topic E Problem 26

The radial probability plot below is for a p orbital. What type of p orbital is it (2p, 3p, 4p, etc.)? Explain your reasoning.




26.1 Solution

It's a p orbital, so $\ell = 1$ and there is one angular node. There are four points at which the radial probability goes to zero, meaning four (4) radial nodes. Adding the number of radial and angular nodes together, we get $4 + 1 = 5$. Add one to the number of total nodes to get the value of n to get $n = 1 + 5 = 6$. The answer is 6p.

27 Topic E Problem 27

One of the electron density graphs below is for a 2p orbital and one is for a 3d orbital.

- Which one is which? Explain your answer.
- Explain why both of these graphs show just one “hump” (i.e. there is no place where the graph goes to zero).
- Explain why both of these graphs start at the origin.
- Give two examples of orbitals whose electron density plots would not start at the origin, and explain your answer.



27.1 Solution

- The graph on the left is the 3d orbital graph. The 2p graph would be generally further in due to having a smaller orbital radius.
- Both the 2p and 3d orbitals have their highest possible number of angular orbitals. For the 2p, $\ell = 1$ and $n = 2$ allows for only 1 total nodes, meaning no radial orbitals. For the 3d, $\ell = 2$ and $n = 3$ allows for only 2 total nodes, meaning no radial orbitals.
- Both have angular nodes as previously stated, so both start at zero at the origin.
- 1s and 5s

28 Topic E Problem 28

A partial energy diagram for lithium (Li) is shown below. Answer questions a through d, using the information on this diagram and your understanding of emission spectra. This is a review problem.

ionized state	_____	0 kJ/mol
4p	_____	-83 kJ/mol
3p	_____	-150 kJ/mol
2p	_____	-342 kJ/mol
2s	_____	-520 kJ/mol

- Calculate ΔE for the $2p \rightarrow 2s$ transition in lithium, in kJ/mol.
- Calculate the wavelength of light emitted during the $2p \rightarrow 2s$ transition, in nm.
- When the outer electron undergoes a $5p \rightarrow 2s$ transition, the atom emits 256nm light. Calculate the energy of the 5p orbital, in kJ/mol.

28.1 Solution (a)

Calculate the difference.

$$\Delta E = E_f - E_i = -520 \text{ kJ/mol} - (-342 \text{ kJ/mol}) = \boxed{-178 \text{ kJ/mol}} \quad (1)$$

28.2 Solution (b)

Take the absolute value and divide by Avogadro's number to get the energy of the emitted photon.

$$E = \frac{|\Delta E|}{N_A} = \frac{178 \text{ kJ/mol}}{6.022 \times 10^{23} \text{ mol}^{-1}} = 2.956 \times 10^{-19} \text{ J} \quad (2)$$

Convert this to the wavelength.

$$\lambda = \frac{hc}{E} = \frac{1.9864748 \times 10^{-25} \text{ J m}}{2.956 \times 10^{-19} \text{ J}} = \boxed{672 \text{ nm}} \quad (3)$$

28.3 Solution (c)

First convert wavelength to energy.

$$\Delta E = \frac{hc}{\lambda} * N_A = \frac{1.9864748 \times 10^{-25} \text{ J m}}{256 \times 10^{-9} \text{ m}} * 6.022 \times 10^{23} \text{ mol}^{-1} \quad (4)$$

$$= 476.3 \text{ kJ/mol} \quad (5)$$

Add this to the 2s energy to get the 5p energy.

$$E_i = E_f + \Delta E = -520 + 476.3 = \boxed{-53 \text{ kJ/mol}} \quad (6)$$

29 Topic E Problem 29

Write ground-state electron configurations for the following atoms and ions. You may use inert gas abbreviations (for example, $[\text{Ne}]3s^1$ instead of $1s^22s^22p^63s^1$).

- | | | |
|------------------|---------------------|---------------------|
| a) Rb | d) S^{2-} | g) Co |
| b) Rb^+ | e) Cd | h) Co^{2+} |
| c) S | f) Cd^{2+} | i) Co^{3+} |

29.1 Solution

- | | |
|--|---|
| a/ $\text{Rb} \rightarrow [\text{Kr}] 5s^1$ | f/ $\text{Cd}^{2+} \rightarrow [\text{Kr}] 4d^{10}$ |
| b/ $\text{Rb}^+ \rightarrow [\text{Kr}]$ | g/ $\text{Co} \rightarrow [\text{Ar}] 4s^23d^7$ |
| c/ $\text{S} \rightarrow [\text{Ne}] 3s^23p^4$ | h/ $\text{Co}^{2+} \rightarrow [\text{Ar}] 3d^7$ |
| d/ $\text{S}^{2-} \rightarrow [\text{Ar}]$ | i/ $\text{Co}^{3+} \rightarrow [\text{Ar}] 3d^6$ |
| e/ $\text{Cd} \rightarrow [\text{Kr}] 5s^24d^{10}$ | |

30 Topic E Problem 30

- a) Which has the higher energy in a hydrogen atom, the 3s orbital or the $3p_x$ orbital?
- b) Which has the higher energy in a phosphorus atom, the 3s orbital or the $3p_x$ orbital?
- c) Which has the higher energy in a hydrogen atom, the 4s or the $3d_{xy}$ orbital?
- d) Which has the higher energy in a Mn atom, the 4s or the $3d_{xy}$ orbital?
- e) Which has the higher energy in a Mn^{2+} ion, the 4s or the $3d_{xy}$ orbital?

30.1 Solution

- | | | |
|---------------------|--------------|-------|
| a/ Identical energy | c/ 4s | e/ 4s |
| b/ $3p_x$ | d/ $3d_{xy}$ | |

31 Topic E Problem 31

Which of the following configurations are ground states, which are excited states, and which are impossible configurations for an uncharged lithium atom?


- | | | |
|---------------|---------------|----------------|
| a) $1s^3$ | c) $1s^22s^1$ | e) $1s^287f^1$ |
| b) $1s^21p^1$ | d) $1s^22p^1$ | |

31.1 Solution

- | | | |
|---------------|------------|------------|
| a/ Impossible | c/ Ground | e/ Excited |
| b/ Impossible | d/ Excited | |

32 Topic E Problem 32

One possible electron configuration for an oxygen atom is $[\text{He}]2s^22p^4$. Which of the following orbital energy diagrams represent the ground state, which represent excited states, and which represent impossible arrangements for the 2p electrons in an uncharged oxygen atom?



32.1 Solution

a/ Excited

d/ Impossible

b/ Excited

e/ Ground

c/ Ground

f/ Impossible

33 Topic E Problem 33

Draw orbital energy diagrams for the 3d and 4s orbitals in the ground states of the following atoms. Do not show any other orbitals (but include arrows for the electrons).

a) Mn

b) Ni

c) Mn^{2+}

34 Topic E Problem 34

Which ground-state atoms in period 4 (elements 19 through 36) have...

- a) no unpaired electrons?
- b) two unpaired electrons?

34.1 Solution

a/ Ca, Zn, Kr

b/ Ti, Ni, Ge, Se

35 Topic E Problem 35

Draw an orbital energy diagram for the following configurations.

1. An atom that has the configuration $1s^2 2s^2 2p^4$ and is diamagnetic.
2. An atom that has the configuration $1s^2 2s^2 2p^4$ and is paramagnetic.

36 Topic E Problem 36

Which of the following configurations must be paramagnetic, which could be paramagnetic, and which cannot possibly be paramagnetic (i.e. they must be diamagnetic)?

- | | |
|------------------------|--|
| a) [Ne]3s | c) [Ne]3s ² 3p |
| b) [Ne]3s ² | d) [Ne]3s ² 3p ² |

36.1 Solution

- | | |
|-----------------|--------------------------|
| a) Paramagnetic | c) Paramagnetic |
| b) Diamagnetic | d) Could be paramagnetic |

37 Topic E Problem 37

- a) How many electrons have $n = 4$ in a ground-state atom of technetium (Tc)?
- b) How many electrons have $\ell = 1$ in a ground-state atom of arsenic (As)?
- c) How many electrons have $m_\ell = 1$ in a ground-state atom of krypton (Kr)?
- d) How many electrons have $m_s = -\frac{1}{2}$ in a ground-state atom of radium (Ra)?
- e) What is the maximum number of electrons that could have $m_\ell = 2$ in a ground-state atom of iron?
- f) What is the minimum number of electrons that could have $m_s = \frac{1}{2}$ in a ground-state atom of oxygen?

37.1 Solution

- | | | |
|-------|-------|------|
| a) 13 | c) 8 | e) 2 |
| b) 15 | d) 44 | f) 3 |

38 Topic E Problem 38

Explain each of the following observations. Explanations such as “Ca is larger than Mg because atoms get larger as you down a column of the periodic table” are not acceptable; you must tell me why this trend occurs.

- a) The atomic radius of Na is larger than the atomic radius of Mg.
- b) The atomic radius of K is larger than the atomic radius of Na.
- c) The ionic radius of S^{2-} is larger than the ionic radius of Cl^- .
- d) The ionic radius of Zr^{3+} is larger than the ionic radius of Zr^{4+} .

39 Topic E Problem 39

Arrange the elements Al, Ga, Ne, and S in order of increasing ionization energy (i.e. from lowest to highest). You should not need to look up the ionization energies to answer this question.

39.1 Solution

Atoms with a larger radius (i.e. closer to the metals) have lower ionization energies. The same applies for atoms with a larger number of core electrons.

$$\boxed{\text{Ga} < \text{Al} < \text{S} < \text{Ne}} \quad (7)$$

40 Topic E Problem 40

The list below shows the ionization energies for elements 36 through 40, in kJ/mol:

Element 36: 1351	Element 38: 549	Element 40: 640
Element 37: 403	Element 39: 600	

- Explain why the ionization energies increase as you go from element 37 to element 40.
- Explain why the ionization energy drops dramatically as you go from element 36 to element 37.
- Would you expect the ionization energy of element 35 to be lower than 1351 kJ/mol, or higher than 1351 kJ/mol?

40.1 Solution

For the hell of it, I'm going to answer this in the voice of Heisenberg, the well-known chemist.

- A higher element number without a completed shell means the charge in and attraction from the nucleus is higher. This results in a higher first ionization energy because the valence electron is pulled on harder by the nucleus of the atom.
- Element 36 happens to be a noble gas. Noble gases have full shells. This means that the electrons have maximum pull by the nucleus and the outermost shell has minimum push by core electrons. This results in overall lower energy required for first ionization of the noble gas.
- I expect it to have lower first ionization energy due to a lower charge from the nucleus and an equal number of core electrons. This would mean that the nucleus pulls less on the electrons.

41 Topic E Problem 41

An element in period 3 (elements 10 through 18) has the following ionization energies. Identify the element. Note: IE 1 is the energy required to remove the first electron, IE 2 is the energy required to remove the second electron, etc.

IE 1 = 787 kJ/mol	IE 3 = 3232 kJ/mol	IE 5 = 16,091 kJ/mol
IE 2 = 1577 kJ/mol	IE 4 = 4356 kJ/mol	IE 6 = 19,805 kJ/mol

41.1 Solution

Note that there is a giant jump between IE 4 and IE 5. We can assume that that would be the point where it starts breaking into a lower shell. There is only one possible column break that would take place in this range, being from Sodium to Neon. Assuming neon's electron configuration to be the appearance after IE 4, we can count the previous shifts.

- IE 4 initial appearance: Na
- IE 3 initial appearance: Mg
- IE 2 initial appearance: Al
- IE 1 initial appearance: Si

This solidifies out final answer to be Si.

42 Topic E Problem 42

The ionization energy of chlorine is 1251 kJ/mol. Based on this value, which of the following conclusions is reasonable? Select the correct statement, and fill in the blank with the correct orbital name.

- The energy of the ____ orbital(s) in chlorine is 1251 kJ/mol.
- The energy of the ____ orbital(s) in chlorine is -1251 kJ/mol.

42.1 Solution

The energy of the 3p orbital(s) in chlorine is -1251 kJ/mol.

Contents

23 Topic E Problem 23	2
23.1 Solution	2
24 Topic E Problem 24	2
24.1 Solution	2
25 Topic E Problem 25	3
25.1 Solution	3
26 Topic E Problem 26	4
26.1 Solution	4
27 Topic E Problem 27	5
27.1 Solution	5
28 Topic E Problem 28	6
28.1 Solution (a)	6
28.2 Solution (b)	6
28.3 Solution (c)	7
29 Topic E Problem 29	8
29.1 Solution	8
30 Topic E Problem 30	9
30.1 Solution	9
31 Topic E Problem 31	10
31.1 Solution	10
32 Topic E Problem 32	11
32.1 Solution	11
33 Topic E Problem 33	12
34 Topic E Problem 34	13
34.1 Solution	13
35 Topic E Problem 35	14
36 Topic E Problem 36	15
36.1 Solution	15
37 Topic E Problem 37	15
37.1 Solution	15
38 Topic E Problem 38	16

39 Topic E Problem 39	17
39.1 Solution	17
40 Topic E Problem 40	17
40.1 Solution	17
41 Topic E Problem 41	18
41.1 Solution	18
42 Topic E Problem 42	18
42.1 Solution	18