

Problem Set #6

CHEM101A: General College Chemistry

Donald Aingworth IV

September 26, 2025

1 Topic D Problem 8

Consider the following two facts about the reaction of HCl with NaOH

Fact #1: ΔE is a negative number for this reaction, which means that the energy of the chemicals decreases.

Fact #2: When HCl and NaOH are mixed, the mixture becomes hotter, which means that the energy of the chemicals increases.

Explain how both of these statements can be true at the same time.

1.1 Solution

2 Topic D Problem 9

When aqueous HCl reacts with solid NaHCO₃, the mixture becomes colder.

1. Does the kinetic energy of the chemicals increase, decrease, or remain the same during the reaction?
2. Does the potential energy of the chemical increase, decrease, or remain the same during the reaction?
3. What is the sign of ΔE for this reaction?

2.1 Solution

3 Topic D Problem 10

For the reaction $2 \text{Al(s)} + 6 \text{H}^+(\text{aq}) \rightarrow 2 \text{Al}^{3+}(\text{aq}) + 3 \text{H}_2(\text{g})$, $\Delta E = -1057 \text{ kJ}$

a) How much energy is given off when 2.822 g of Al reacts with hydrogen ions, assuming the hydrogen ions are in excess? b) How much energy is given off when 1.413 g of Al is added to 23.0 mL of a solution that contains 5.89 M H^+ ? c) If you want to obtain 50.0 kJ of energy from this reaction, how many grams of aluminum and how many mL of 3.00 M HCl must you use? d) If this reaction produces 13.3 L of $\text{H}_2(\text{g})$ at 25.0°C and 1.014 atm, how much energy does it give off?

3.1 Solution

4 Topic D Problem 11

When 4.00 g of Br₂ reacts with excess Al to form AlBr₃, 8.80 kJ of energy is given off. Calculate ΔE for the reaction $2 \text{ Al(s)} + 3 \text{ Br}_2\text{(s)} \rightarrow 2 \text{ AlBr}_3\text{(s)}$.

4.1 Solution

5 Topic D Problem 12

A chemist burns 0.5177 g of liquid hexane (C_6H_{14}) in a bomb calorimeter that has a heat capacity of $4287 \text{ J}/^\circ\text{C}$. The temperature of the calorimeter rises from 21.382°C to 27.170°C during the reaction. Calculate ΔE for the following reaction: $2 \text{C}_6\text{H}_{14}(\text{l}) + 19 \text{O}_2(\text{g}) \rightarrow 12 \text{CO}_2(\text{g}) + 14 \text{H}_2\text{O}(\text{l})$

5.1 Solution

6 Topic D Problem 13

Explain why some reactions give off different amounts of heat when they are carried out at constant pressure versus constant volume.

6.1 Solution

7 Topic D Problem 14

A reaction has $\Delta E = -50 \text{ kJ}$. a) If this reaction occurs, will the reaction mixture become hotter, or colder? b) Is this an exothermic reaction, or is it an endothermic reaction? c) After this reaction has occurred, will the potential energy of the chemical mixture be higher than, lower than, or the same as it was before the reaction? d) After this reaction has occurred, will the kinetic energy of the universe be higher than, lower than, or the same as it was before the reaction (assuming no other reaction has occurred)?

7.1 Solution

8 Topic D Problem 15

A reaction has $\Delta E = 10 \text{ kJ}$, and it converts a solid into a gas. a) If this reaction occurs in a closed, rigid container, will the amount of heat that is absorbed be larger than 10 kJ, smaller than 10 kJ, or exactly 10 kJ? Justify your answer. b) If this reaction occurs in an open container, will the amount of heat that is absorbed be larger than 10 kJ, smaller than 10 kJ, or exactly 10 kJ? Justify your answer. c) Will PV work be done in either part a or part b? If so, which part(s)? d) What is the sign of the PV work when PV work occurs?

8.1 Solution

9 Topic D Problem 16

A reaction has $\Delta E = -10$ kJ, and it converts a gas into a liquid. a) If this reaction occurs in a closed, rigid container, will the amount of heat that is given off be larger than 10 kJ, smaller than 10 kJ, or exactly 10 kJ? Justify your answer. b) If this reaction occurs in an open container, will the amount of heat that is given off be larger than 10 kJ, smaller than 10 kJ, or exactly 10 kJ? Justify your answer. c) Will PV work be done in either part a or part b? If so, which part(s)? d) What is the sign of the PV work when PV work occurs?

9.1 Solution

10 Topic D Problem 17

A reaction has $\Delta E = -10$ kJ, and it converts a liquid into a gas. a) If this reaction occurs in a closed, rigid container, will the amount of heat that is given off be larger than 10 kJ, smaller than 10 kJ, or exactly 10 kJ? Justify your answer. b) If this reaction occurs in an open container, will the amount of heat that is given off be larger than 10 kJ, smaller than 10 kJ, or exactly 10 kJ? Justify your answer. c) Will PV work be done in either part a or part b? If so, which part(s)? d) What is the sign of the PV work when PV work occurs?

10.1 Solution

11 Topic D Problem 18

a) What is the PV work when 3.00 moles of liquid water boils at 100°C at constant pressure? Give your answer in kJ, and include the correct sign. b) What is the PV work when 3.00 moles of steam condenses at 100°C at constant pressure? Give your answer in kJ, and include the correct sign.

11.1 Solution

12 Topic D Problem 19

Consider the following reaction: $4 \text{ C}_3\text{H}_9\text{N}(\text{l}) + 21 \text{ O}_2(\text{g}) \rightarrow 12 \text{ CO}_2(\text{g}) + 18 \text{ H}_2\text{O}(\text{l}) + 2 \text{ N}_2(\text{g})$ $\Delta E = -9667 \text{ kJ}$ The following questions (parts a through j) refer to this reaction. a) Calculate ΔH for this reaction at 25°C . b) How much heat will be given off if 8.250 g of $\text{C}_3\text{H}_9\text{N}$ reacts with excess O_2 in a closed, rigid container at 25°C ? c) How much heat will be given off if 8.250g of $\text{C}_3\text{H}_9\text{N}$ reacts with excess O_2 in an open container at 25°C ? d) Calculate the PV work in part b. Include the correct sign. e) Calculate the PV work in part c. Include the correct sign. f) If you want to obtain 200.0 kJ of heat by reacting $\text{C}_3\text{H}_9\text{N}$ with O_2 in an open container, what mass of $\text{C}_3\text{H}_9\text{N}$ must you use? g) How much heat is produced when 2.199 g of liquid $\text{C}_3\text{H}_9\text{N}$ reacts with 5.738 g of gaseous O_2 in an open container? h) If you burn enough liquid $\text{C}_3\text{H}_9\text{N}$ to produce 841.2 kJ of heat in an open container, what volume of gaseous N_2 will you form at 25.0°C and 752 torr? i) If this reaction is carried out in a closed, rigid container and produces 31.74 g of liquid H_2O , how much heat does it produce?

12.1 Solution

13 Topic D Problem 20

When 2.810 g of solid Al reacts with excess gaseous F₂ in an open container at 25°C, 157.3 kJ of heat is produced. a) Calculate ΔH and ΔE for the following reaction at 25°C: $2 \text{ Al(s)} + 3 \text{ F}_2\text{(g)} \rightarrow 2 \text{ AlF}_3\text{(s)}$ b) How much heat will be produced if 8.493 g of solid Al reacts with 16.610 g of gaseous F₂ in a closed, rigid container at 25°C? Use your answer from part a to solve this problem.

13.1 Solution

14 Topic D Problem 21

A chemist burns 1.628 g of liquid isopropyl alcohol ($\text{C}_3\text{H}_8\text{O}$) in a bomb calorimeter that has a heat capacity of $3927 \text{ J}/^\circ\text{C}$. The temperature of the calorimeter rises from 19.085°C to 31.683°C during the reaction. Calculate ΔH and ΔE for the following reaction at 25°C : $2 \text{ C}_3\text{H}_8\text{O}(\text{l}) + 9 \text{ O}_2(\text{g}) \rightarrow 6 \text{ CO}_2(\text{g}) + 8 \text{ H}_2\text{O}(\text{l})$

14.1 Solution

15 Topic D Problem 22

For the reaction $2 \text{C(s)} + \text{O}_2\text{(g)} \rightarrow 2 \text{CO(g)}$, $\Delta E = -219 \text{ kJ}$. When will ΔH also be -219 kJ ? a) When the reaction is carried out in an open container. b) When the reaction is carried out in a closed, rigid container. c) ΔH will always equal ΔE . d) ΔH will never equal ΔE .

15.1 Solution