

Homework #2

Math 104: Intro to Analysis

Summer 2026

UC Berkeley

Donald Aingworth IV

July 10, 2026

Contents

0	Various Theorems	2
0.1	Monotone Decrease I	2
0.2	Monotone Decrease II	2
0.3	Inverese Triangle Inequality	2
1	Problem 1	3
1.1	Solution (a)	3
2	Problem 2	5
3	Problem 3	6
3.1	Solution (a)	6
3.2	Solution (b)	6
3.3	Solution (c)	7
3.4	Solution (d)	8
4	Problem 4	9
4.1	Solution (a)	9
4.2	Solution (b)	9
5	Problem 5	11
5.1	Solution (a)	11
5.2	Solution (b)	11
6	Problem 6	12
6.1	Solution (a)	12
6.2	Solution (b)	12

0 Various Theorems

0.1 Monotone Decrease I

$a_n = \frac{1}{n(\log n)(\log \log n)^p}$ for $n \in \mathbb{N}$ is monotonically decreasing if $p > 0$.

Proof. Let $p > 0$. We know $n > 0$ since $n \in \mathbb{N}$. This also means $n \geq 1$. Logarithms are monotone, as mentioned in Rudin (p.63). Therefore, $\log n$ and by extension $\log \log n$ and $(\log \log n)^p$ are monotonically increasing. Therefore, $n(\log n)(\log \log n)^p$ is monotonically increasing. On the other hand, this also means that $\frac{1}{n(\log n)(\log \log n)^p}$ is monotonically decreasing. OEΔ

0.2 Monotone Decrease II

$b_n = \frac{1}{n(\log 2)(\log(n) + \log(\log 2))^p}$ for $n \in \mathbb{N}$ is monotonically decreasing if $p > 0$.

Proof. This uses a lot of what was used in Monotone Decrease I. n is monotone increasing. $\log(n)$ is monotone increasing and $\log(\log(2))$ is constant, so $\log(n) + \log(\log(2))$ and by extension $(\log(n) + \log(\log(2)))^p$ if $p > 0$ are monotone increasing. $\log(2)$ is constant, so $n(\log 2)(\log(n) + \log(\log 2))^p$ is monotone increasing if $p > 0$. Therefore, $\frac{1}{n(\log 2)(\log(n) + \log(\log 2))^p}$ is monotone decreasing if $p > 0$. OEΔ

0.3 Inverse Triangle Inequality

For $a, b, c \in (X, d)$, $|d(p, q) - d(q, r)| \leq d(p, r)$.

Proof. From the triangle inequality, we have two equations.

$$d(p, q) \leq d(p, r) + d(q, r) \tag{1}$$

$$d(q, r) \leq d(p, r) + d(p, q) \tag{2}$$

We can subtract $d(q, r)$ and $d(p, q)$ from them respectively.

$$d(p, q) - d(q, r) \leq d(p, r) \tag{3}$$

$$d(q, r) - d(p, q) \leq d(p, r) \tag{4}$$

Reverse the latter.

$$d(p, q) - d(q, r) \geq -d(p, r) \tag{5}$$

By the definition of the absolute value, we can say that $|d(p, q) - d(q, r)| \leq d(p, r)$. OEΔ

1 Problem 1

For which values of $p \in \mathbb{R}$ are the following series convergent or divergent? Prove your answers rigorously, using only tests we explicitly covered in class. *Hint:* for both of these series you should use something with the name “Cauchy” in it.

$$(a) \sum_{n=3}^{\infty} \frac{1}{n(\log n)(\log \log n)^p}$$

$$(b) \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n^p}.$$

1.1 Solution (a)

Proof. Since $2 < e$, $\log(2) < \log(e) = 1$. Furthermore, through the use of a calculator, $\log 2 = 0.6931$ and $\log \log 2 = -0.3665$.

Let $p \leq 0$. This means $a_n = \frac{(\log \log n)^q}{n(\log n)}$ for some $q \geq 0$ such that $q = -p$. $\log n$ is monotonically increasing, so by extension so is $\log \log n$. Therefore, since $\log n$ is monotonically increasing and unbounded, so is $\log \log n$ and so is $(\log \log n)^q$. Therefore, for some $N_0 \in \mathbb{N}$, for all $n \geq N_0$, $\log \log n \geq 1$ and $(\log \log n)^q \geq 1$. Therefore if $n \geq N_0$, $\frac{(\log \log n)^q}{n \log n} \geq \frac{1}{n \log n}$. Recall that if $a_n \geq d_n \geq 0$ for $n \geq N_0$ and if $\sum d_n$ diverges, then $\sum a_n$ diverges (Rudin, p.60). In Rudin (p.62), it is proven that $\sum_{n=2}^{\infty} \frac{1}{n \log n}$ diverges. Therefore, $\sum_{n=3}^{\infty} \frac{(\log \log n)^q}{n \log n}$ diverges.

Going forward, we will therefore only operate on cases where $p > 0$. We established that if $p > 0$, there exists $N_0 \in \mathbb{N}$ such that for all $n \geq N_0$, $\frac{1}{n(\log n)(\log \log n)^p} > 0^1$. Furthermore, it is established earlier in this document that $\frac{1}{n(\log n)(\log \log n)^p}$ is monotonically decreasing.

We will use the Cauchy Condensation Test, which says that for monotonically decreasing, positive a_n , $\sum_{n=1}^{\infty} a_n$ converges if and only if $\sum_{n=1}^{\infty} 2^n a_{2^n}$ converges. Consider it in this case.

$$a_n = \frac{1}{n(\log n)(\log \log n)^p} \quad (6)$$

$$b_n = 2^n a_{2^n} = \frac{2^n}{2^n(\log 2^n)(\log \log 2^n)^p} = \frac{1}{n(\log 2)(\log(n \log 2))^p} = \frac{1}{n(\log 2)(\log(n) + \log(\log 2))^p} \quad (7)$$

Now, we can use how $\frac{1}{n \log 2 (\log(n) + \log(\log 2))^p}$ is positive and monotonically decreasing to use the Cauchy Condensation Test again.

$$b_n = \frac{1}{n(\log 2)(\log(n) + \log(\log 2))^p} \quad (8)$$

$$c_n = 2^n b_{2^n} = \frac{2^n}{2^n(\log 2)(\log(2^n) + \log(\log 2))^p} = \frac{1}{(\log 2)(n \log 2 + \log(\log 2))^p} \quad (9)$$

Here, we have two cases: either $p > 1$ or $p \leq 1$.

First consider the case where $p \leq 1$, where $n \in \mathbb{N}$ and $n \geq 3$. Now, we know that $\log(2) < 1$, so $n \log(2) < n$. Therefore, since $\log \log 2 < 0$, $n \log(2) + \log \log(2) < n$ and by extension $(n \log(2) + \log \log(2))^p < n$. Therefore, $\frac{1}{(n \log(2) + \log \log(2))^p} > \frac{1}{n}$. If we sum this for $n \geq 3$, we find that $\sum_{n=3}^{\infty} \frac{1}{(n \log(2) + \log \log(2))^p} > \sum_{n=3}^{\infty} \frac{1}{n}$. Since $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges to ∞ , we can equally say that $\sum_{n=3}^{\infty} \frac{1}{n}$ diverges to ∞ . By

¹Might need further proof, but see prior paragraph.

However, since n is monotonically increasing and unbounded, for every $m \in \mathbb{N}$, there is an N_1 such that if $n > N_1$, $n \log(2) + \log \log(2) \geq m$.

$$\frac{1}{n^{p-1}} > \frac{1}{(n \log 2 + \log \log 2)^p} \quad (10)$$

$$(n \log 2 + \log \log 2)^p > n^{p-1} \quad (11)$$

$$n \log(2) + \log \log 2 > n \text{ if } n \geq 3 \quad (12)$$

$$(n \log(2) + \log \log 2)^p > n^p > n^{p-1} \text{ if } n \geq 3 \quad (13)$$

OEΔ

2 Problem 2

A series $\sum_{n=1}^{\infty} a_n$ is said to be *absolutely convergent* if $\sum_{n=1}^{\infty} |a_n|$ is convergent. Prove that every absolutely convergent series is convergent. *Hint:* use the Cauchy criterion.

Proof. Suppose that $\sum_{n=1}^{\infty} a_n$ is absolutely convergent. This means that $\sum_{n=1}^{\infty} |a_n|$ is convergent and by extension Cauchy. By the Cauchy Criterion, this means that for all $\varepsilon > 0$, there exists some $N \in \mathbb{N}$ such that if $m \geq n \geq N$, $|\sum_{k=n}^m a_k| \leq \varepsilon$.

Recall the triangle inequality for the usual metric: if $a, b, c \in \mathbb{R}$, $|a - c| \leq |a - b| + |b - c|$. Since $|a - c| = |a - b + b - c|$, $|a - b + b - c| \leq |a - b| + |b - c|$. Suppose that for some $a_k = a - b$ and $a_{k+1} = b - c$. Therefore, $|a_k + a_{k+1}| \leq |a_k| + |a_{k+1}|$. Since $m - n$ is finite, we can extend this process to every $k \in [n, m - 1] \cap \mathbb{N}$ to conclude that $|\sum_{k=n}^m a_k| \leq \sum_{k=n}^m |a_k|$. By the absolute value, $|\sum_{k=n}^m a_k| \geq 0$, meaning $\sum_{k=n}^m |a_k| \geq 0$ and therefore $\sum_{k=n}^m |a_k| = |\sum_{k=n}^m |a_k||$. This means that $|\sum_{k=n}^m a_k| \leq |\sum_{k=n}^m |a_k||$.

Transistively, we can say from our earlier convergence statement that for all $\varepsilon > 0$, there exists some $N \in \mathbb{N}$ such that if $m \geq n \geq N$, $|\sum_{k=n}^m a_k| \leq \varepsilon$. Therefore, by the Cauchy Criterion, $\sum_{n=1}^{\infty} a_n$ is convergent. Ergo, if a series is absolutely convergent, it is convergent. OEΔ

3 Problem 3

Let X, Y be sets and $f : X \rightarrow Y$ a function.

- (a) Show that, for $E, F \subset Y$ we have $f^{-1}(E \cup F) = f^{-1}(E) \cup f^{-1}(F)$ and $f^{-1}(E \cap F) = f^{-1}(E) \cap f^{-1}(F)$.
- (b) Show that, for $E, F \subset X$, we have $f(E \cup F) = f(E) \cup f(F)$. Is it also always true that $f(E \cap F) = f(E) \cap f(F)$? Prove or give a counterexample.
- (c) Let $E \subset X$. Show that $E \subset f^{-1}(f(E))$. Show that equality holds for all E if and only if f is injective.
- (d) Let $E \subset Y$. Show that $f(f^{-1}(E)) \subset E$. Show that equality holds for all E if and only if f is surjective.

3.1 Solution (a)

First we prove that $f^{-1}(E \cup F) = f^{-1}(E) \cup f^{-1}(F)$.

Proof. Begin with $f^{-1}(E \cup F)$. If $x \in f^{-1}(E \cup F)$, then $f(x) \in E \cup F$. Therefore, $f(x) \in E$ or $f(x) \in F$. If $f(x) \in E$, then $x \in f^{-1}(E)$. If $f(x) \in F$, then $x \in f^{-1}(F)$. Therefore, since either $f(x) \in E$ or $f(x) \in F$, $x \in f^{-1}(E)$ or $x \in f^{-1}(F)$. Therefore, $x \in f^{-1}(E) \cup f^{-1}(F)$. This means $f^{-1}(E \cup F) \subset f^{-1}(E) \cup f^{-1}(F)$.

For the reverse direction, begin with $x \in f^{-1}(E) \cup f^{-1}(F)$. This means that either $x \in f^{-1}(E)$ or $x \in f^{-1}(F)$. Without loss of generality, suppose that $x \in f^{-1}(E)$. This means that, $f(x) \in E$. Quite logically, this means that $f(x) \in E$ or $f(x) \in F$. Therefore, $f(x) \in E \cup F$. Therefore, $x \in f^{-1}(E \cup F)$. Concluding, $f^{-1}(E) \cup f^{-1}(F) \subset f^{-1}(E \cup F)$.

Since $f^{-1}(E \cup F) \subset f^{-1}(E) \cup f^{-1}(F)$ and $f^{-1}(E) \cup f^{-1}(F) \subset f^{-1}(E \cup F)$, $f^{-1}(E \cup F) = f^{-1}(E) \cup f^{-1}(F)$. OEΔ

Next, we prove that $f^{-1}(E \cap F) = f^{-1}(E) \cap f^{-1}(F)$.

Proof. Begin with $f^{-1}(E \cap F)$. If $x \in f^{-1}(E \cap F)$, then $f(x) \in E \cap F$. Therefore, $f(x) \in E$ and $f(x) \in F$. Since $f(x) \in E$, $x \in f^{-1}(E)$. Since $f(x) \in F$, $x \in f^{-1}(F)$. Therefore, $x \in f^{-1}(E) \cap f^{-1}(F)$. Therefore, $f^{-1}(E \cap F) \subset f^{-1}(E) \cap f^{-1}(F)$.

Suppose $x \in f^{-1}(E) \cap f^{-1}(F)$. Then, $x \in f^{-1}(E)$ and $x \in f^{-1}(F)$. Since $x \in f^{-1}(E)$, $f(x) \in E$. Since $x \in f^{-1}(F)$, $f(x) \in F$. Therefore, $f(x) \in E \cap F$. Therefore, $x \in f^{-1}(E \cap F)$. Therefore, $f^{-1}(E) \cap f^{-1}(F) \subset f^{-1}(E \cap F)$.

Since $f^{-1}(E \cap F) \subset f^{-1}(E) \cap f^{-1}(F)$ and $f^{-1}(E) \cap f^{-1}(F) \subset f^{-1}(E \cap F)$, $f^{-1}(E \cap F) = f^{-1}(E) \cap f^{-1}(F)$. OEΔ

3.2 Solution (b)

First, we prove that $f(E \cup F) = f(E) \cup f(F)$.

Proof. Suppose that $x \in X$ and $f(x) \in f(E \cup F)$. This means that $x \in E \cup F$. Therefore, either $x \in E$ or $x \in F$. Without loss of generality, suppose that $x \in E$. Therefore, $f(x) \in f(E)$. Using the same steps, we can show that $f(x) \in f(F)$. Therefore, $f(x) \in f(E) \cup f(F)$. This means that $f(E \cup F) \subset f(E) \cup f(F)$.

Suppose that $x \in X$ and $f(x) \in f(E) \cup f(F)$. This means that either $f(x) \in f(E)$ or $f(x) \in f(F)$. Therefore either $x \in E$ or $x \in F$. This means that $x \in E \cup F$. Therefore, $f(x) \in f(E \cup F)$. This means that $f(E) \cup f(F) \subset f(E \cup F)$.

Since $f(E \cup F) \subset f(E) \cup f(F)$ and $f(E) \cup f(F) \subset f(E \cup F)$, $f(E \cup F) = f(E) \cup f(F)$. OEΔ

Now we disprove that $f(E \cap F) = f(E) \cap f(F)$.

Proof by Counterexample. Take the case of $E = (0, \infty)$ and $F = (-\infty, 0)$ where $f : \mathbb{R} \rightarrow \mathbb{R}$ and $f(x) = |x|$. Taking the intersection, $E \cap F = \emptyset$. Therefore, $f(E \cap F) = \emptyset$.

Now, applying the function to E and F , we find $f(E) = (0, \infty)$ and $f(F) = (0, \infty)$. Taking the intersection of the two, we find $f(E) \cap f(F) = (0, \infty)$.

Since $(0, \infty) \neq \emptyset$, $f(E \cap F) \neq f(E) \cap f(F)$. OEΔ

3.3 Solution (c)

First, we prove that $E \subset f^{-1}(f(E))$.

Proof. Let $x \in E$. Therefore, $f(x) \in f(E)$. Since $f(x) \in f(E)$ and $f^{-1}(f(E)) = \{x \in X \mid f(x) \in f(E)\}$, $x \in f^{-1}(f(E))$. Therefore, $E \subset f^{-1}(f(E))$. OEΔ

Next we prove that if f is injective, then $E = f^{-1}(f(E))$.

Proof. Injective means that if $a, b \in E$, $f(a) = f(b)$ implies that $a = b$. Suppose that f is injective. Recall the definitions that $f(E) = \{f(x) \mid x \in E\}$ and $f^{-1}(f(E)) = \{x \mid f(x) \in f(E)\}$.

Take any element in $x \in f^{-1}(f(E))$. By the definition of $f^{-1}(f(E))$, this means that $f(x) \in f(E)$. By the definition of $f(E)$, there exists $y \in E$ such that $f(x) = f(y)$. Since f is injective, this means that $x = y$. Therefore, since $y \in E$ and $x = y$, $x \in E$. This means that if $x \in f^{-1}(f(E))$, $x \in E$. Therefore, $f^{-1}(f(E)) \subset E$.

We proved earlier that $E \subset f^{-1}(f(E))$. Therefore, $f^{-1}(f(E)) \subset E$ and $E \subset f^{-1}(f(E))$. Combining them, that means $E = f^{-1}(f(E))$. OEΔ

Third, we prove that if $E = f^{-1}(f(E))$, then f is injective.

Proof. Recall the definitions that $f(E) = \{f(x) \mid x \in E\}$ and $f^{-1}(f(E)) = \{x \mid f(x) \in f(E)\}$.

Suppose that $E = f^{-1}(f(E))$ and $x, y \in X$. Let $x \in E$ and $y \notin E$, so $x \neq y$. Therefore, if $x \in f^{-1}(f(E))$ and $y \notin f^{-1}(f(E))$. This also means that $f(x) \in f(E)$. Additionally, $f(y) \notin f(E)$, because if $f(y) \in f(E)$, then by definition $y \in f^{-1}(f(E))$, which is a contradiction. Therefore, $x \neq y$ implies $f(x) \neq f(y)$. By contrapositive, this means that $f(x) = f(y)$ implies $x = y$. Ergo, f is injective. OEΔ

Last, we consolidate.

Proof. We know that if f is injective, then $E = f^{-1}(f(E))$. We also know that if $E = f^{-1}(f(E))$, then f is injective. Therefore, f is injective if and only if $E = f^{-1}(f(E))$. OEΔ

3.4 Solution (d)

First we prove that $f(f^{-1}(E)) \subset E$.

Proof. Take $y \in f(f^{-1}(E))$. Therefore, there exists some element $x \in f^{-1}(E)$ such that $f(x) = y$. Since $x \in f^{-1}(E)$, then $f(x) \in E$. Since $y = f(x)$, $y \in E$ for any $y \in f(f^{-1}(E))$. Ergo, $f(f^{-1}(E)) \subset E$. OEΔ

Second, we prove that if f is surjective, then $f(f^{-1}(E)) = E$.

Proof. Take any $x \in E$. Since f is surjective, for every element $z \in E$, there is an element $y \in X$ such that $f(y) = z$. Therefore, there is an element $y \in X$ such that $f(y) = x$. By the definition of $f^{-1}(E)$, this means that $y \in f^{-1}(E)$. Furthermore, by the definition of $f(f^{-1}(E))$, since $y \in f^{-1}(E)$, $f(y) \in f(f^{-1}(E))$. Since $f(y) = x$, $x \in f(f^{-1}(E))$. Therefore, $x \in E$ implies $x \in f(f^{-1}(E))$. Ergo, $E \subset f(f^{-1}(E))$.

We have established that $E \subset f(f^{-1}(E))$. We have established that $f(f^{-1}(E)) \subset E$. Ergo, $f(f^{-1}(E)) = E$. OEΔ

Third, we prove that if $f(f^{-1}(E)) = E$, then f is surjective.

Proof. Suppose that for all E , $f(f^{-1}(E)) = E$. This means that if $y \in E$ then $y \in f(f^{-1}(E))$. Since $y \in f(f^{-1}(E))$, there exists $x \in f^{-1}(E)$ where $f(x) = y$. Since $f^{-1}(E) \subset X$, $x \in X$. Therefore, for any $y \in E$, there exists $x \in X$ such that $f(x) = y$. Ergo, f is surjective within E for all $E \subset Y$.

Suppose that $E = Y$. That would mean that f is surjective in E and as such surjective in Y . Therefore, f is surjective over Y . OEΔ

Last, we consolidate.

Proof. We know that if f is surjective, then $E = f(f^{-1}(E))$. We also know that if $E = f(f^{-1}(E))$, then f is surjective. Therefore, f is surjective if and only if $E = f(f^{-1}(E))$. OEΔ

4 Problem 4

Let (X, d_X) and (Y, d_Y) be metric spaces.

- (a) A function $f : X \rightarrow Y$ (not necessarily continuous) is said to be *locally bounded* if, for each $p \in X$, there is an $r > 0$ so that $f(B_r(p))$ is a bounded set in Y . Show that every locally bounded function on a *compact* metric space is bounded (i.e. $f(X)$ is a bounded set in Y).
- (b) A function $f : X \rightarrow Y$ is said to be *locally constant* if, for each $p \in X$, there is an $r > 0$ so that f is constant on $B_r(p)$. Show that every locally constant function on a *connected* metric space is constant. *Hint:* show that $f^{-1}(\{y\})$ is clopen in X for any $y \in Y$

4.1 Solution (a)

Proof. Let $f : X \rightarrow Y$ be locally bounded and let (X, d_X) be compact. Since (X, d_X) is compact, every open cover of X has a finite subcover. By the definition of a locally bounded function, for each $p \in X$, there is an $r_p > 0$ so that $f(B_{r_p}(p))$ is bounded in Y . Create a collection out of every $B_{r_p}(p)$ and call it P . Since every open ball contains its center and open balls are open, the collection is an open cover of X . Since X is compact, there is a cover of X made up of a finite number of elements in P , which we can call F .

Create a new set called G defined as $G = \{f(S)\}_{S \in F}$. Since every element in F is an open ball, applying f to every open ball returns a bounded set in Y , and F is finite, G is a finite collection of bounded sets in Y . Since the union of a finite number of bounded sets is bounded², the union of all elements in G , $H = \bigcup_{S \in G} S$, is a bounded set. Furthermore, since H is a collection of bounded sets in Y , $H \subset Y$.

Now, we prove that if $f(x) \in f(X)$, then $f(x) \in H$. Take any element $f(x) \in f(X)$. By the definition of $f(X)$, $x \in X$. Since F is a cover of X , there exists $T \in F$ such that $x \in T$. Therefore, $f(x) \in f(T)$. Since $H = \bigcup_{S \in G} S = \bigcup_{S \in F} f(S)$, $f(T) \subset H$. $f(x) \in f(T)$ and $f(T) \subset H$ means $f(x) \in H$. Therefore, $f(x) \in f(X)$ implies $f(x) \in H$. Therefore, $f(X) \subset H$.

Lastly, a few facts to put everything together. Since H is bounded and $f(X) \subset H$, $f(X)$ is bounded. Since $H \subset Y$ and $f(X) \subset H$, $f(X) \subset Y$. Therefore, $f(X)$ is a bounded set in Y . Ergo, every locally bounded function on a compact metric space is bounded. OEΔ

4.2 Solution (b)

Proof by Contradiction. Let $f : X \rightarrow Y$ be locally constant and let (X, d_X) and (Y, d_Y) be metric spaces. Furthermore, let (X, d_X) be connected, meaning the only clopen sets in (X, d_X) are $\emptyset$ and X .

Suppose for contradiction that f is not constant. This means that there exists a set $V \subset Y$ with more than one element where $V = f(X)$. For all $v \in V$, let X_v be the set of elements $p \in X$ where $f(p) = v$. Since f is locally constant, for each $p \in X$, there is an $r > 0$ such that for all $q \in B_r(p)$, $f(p) = f(q)$. If $p \in X_v$, then $q \in X_v$ and by extension $B_r(p) \subset X_v$. Therefore, for all $v \in V$, X_v is open.

Since f is a well-defined function, for every $p \in X$, there exists $f(p) \in Y$. Therefore, for every $p \in X$, there exists $v \in V$ where $f(p) = v$. Therefore, for every $p \in X$, there exists $v \in V$ such that $p \in X_v$. Therefore, $X = \bigcup_{v \in V} X_v$ and for any $w \in V$, $X_w^c = X \setminus X_w = \bigcup_{v \in V \setminus \{w\}} X_v$.

Take any $w \in V$. Since $V = f(X)$, $w \in f(X)$, meaning there is some element $x \in X$ where $f(x) = w$. Therefore, $X_w \neq \emptyset$. Furthermore, f is not constant, so $X_w \neq X$. We already proved

²Prove or cite a source.

that for every $v \in V$, X_v is open. Therefore, since the union of open sets is open, $\bigcup_{v \in V \setminus \{w\}} X_v$ is also open. This means that X_w^c is open. Since a set is closed if and only if its complement is open, X_w is closed. Therefore, X_w is both open and closed, or clopen. This contradicts the fact that X is connected, since the only clopen subsets of connected set X are $\emptyset$ and X . Ergo, $f(X)$ is constant. OEΔ

5 Problem 5

Using the $\epsilon - \delta$ definition of continuity directly, prove that the following functions are continuous.

(a) $f : \mathbb{R} \rightarrow \mathbb{R}; \quad f(x) = 104x.$

(b) $f : \mathbb{R} \rightarrow \mathbb{R}; \quad f(x) = x^2.$

5.1 Solution (a)

Proof. For a function to be continuous, it must be continuous at every point in its domain. Recall the $\epsilon - \delta$ definition of continuous at a point: for all $\epsilon > 0$, there exists some $\delta > 0$ such that $d_X(x, p) < \delta$ implies $d_Y(f(x), f(p)) < \epsilon$. In the real numbers and the usual metric, this means that $|x - p| < \delta$ implies that $|f(x) - f(p)| < \epsilon$.

Let $\epsilon > 0$ and $\delta = \frac{\epsilon}{104} > 0$. Suppose that $|x - p| < \delta$. This means that $|x - p| < \frac{\epsilon}{104}$. Multiplying everything by 104, we find that $\epsilon > 104|x - p| = |104x - 104p|$. Since $f(x) = 104x$, $|f(x) - f(p)| < \epsilon$. Putting things together, this means that for any $\epsilon > 0$, there exists some $\delta > 0$ such that $|x - p| < \delta$ implies $|f(x) - f(p)| < \epsilon$. Ergo, $f(x) = 104x$ is continuous. OEΔ

5.2 Solution (b)

Proof. Let $\epsilon > 0$. Take any $x, p \in \mathbb{R}$. By the definition of the absolute value, $|x + p| \geq 0$. There are two conditions from here, that either $x = p$ or $x \neq p$.

Suppose $x = p$. This means that $|x - p| = 0$, so for any $\delta > 0$, $\delta > |x - p|$. Furthermore, since $|f(x) - f(p)| = |x^2 - p^2| = |x - p||x + p|$, $|x^2 - p^2| = |x - p||x + p| = 0|x + p| = 0$. Therefore, $\epsilon > |f(x) - f(p)|$, so if $x = p$, the $\epsilon - \delta$ definition is satisfied.

Suppose $x \neq p$, so $|x + p| \neq 0$, so $|x + p| > 0$ by the definition of the absolute value. This means $|x + p| > 0$. Let $\delta = \frac{\epsilon}{|x + p|}$ and suppose that $|x - p| < \delta$. Therefore, $|x - p| < \frac{\epsilon}{|x + p|}$. Multiply both sides by $|x + p|$ to find that $|x - p||x + p| < \epsilon$. Therefore, $|(x - p)(x + p)| < \epsilon$. Therefore, $|x^2 - p^2| < \epsilon$. This means $|f(x) - f(p)| < \epsilon$. Therefore, the $\epsilon - \delta$ definition is satisfied.

The $\epsilon - \delta$ definition is satisfied in all cases. Ergo, $f(x) = x^2$ is continuous. OEΔ

6 Problem 6

Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be defined by $f(x) = \begin{cases} x, & \text{if } x \in \mathbb{Q} \\ 0, & \text{if } x \notin \mathbb{Q} \end{cases}$. Using the $\epsilon - \delta$ definition of continuity,

- (a) prove that f is not continuous at any point $x \neq 0$;
- (b) prove that f is continuous at $x = 0$.

6.1 Solution (a)

This is an altered version of Thomae's Function. We consider this in two cases: $x \in \mathbb{Q} \setminus \{0\}$ and $x \notin \mathbb{Q}$.

First we prove that f is not continuous at x if $x \in \mathbb{Q} \setminus \{0\}$.

Proof. Take $\epsilon = \frac{|f(x)|}{2}$. Take any $\delta > 0$ such that $|x - p| < \delta$. Since the irrational numbers are dense in the real numbers, between any two real numbers, there is an irrational number. This means that there is an irrational number p where $x - \delta < p < x + \delta$. This means that $-\delta < p - x < \delta$, and by extension $|p - x| = |x - p| < \delta$.

Now take $|f(x) - f(p)|$. Since $x \in \mathbb{Q}$ and $p \notin \mathbb{Q}$. Therefore, $|f(x) - f(p)| = |x - 0| = |x|$. Therefore, since $\epsilon = \frac{|f(x)|}{2} = \frac{|x|}{2}$, $|x| = |f(x) - f(p)| \geq \epsilon$. Ergo, by the $\epsilon - \delta$ definition, f is not continuous at $x \in \mathbb{Q} \setminus \{0\}$. OEΔ

Next, we prove that f is not continuous at x if $x \in \mathbb{Q}$.

Proof. Take any $\delta > 0$ such that $|x - p| < \delta$. Since the rational numbers are dense in the real numbers, there is a non-zero rational number p where $x - \delta < p < x + \delta$. Therefore, $|x - p| < \delta$.

Now, look at the epsilon-side. Since p is rational but x is irrational, $|f(x) - f(p)| = |p - 0| = |p|$. If $\epsilon > |p| > 0$, then $\epsilon > |f(x) - f(p)|$. However, if $0 < \epsilon \leq |p|$, then $\epsilon \leq |f(x) - f(p)|$. Ergo, by the $\epsilon - \delta$ definition of continuous, f is not continuous at x if $x \in \mathbb{Q}$. OEΔ

If necessary, we consolidate this.

Proof. We know that f is not continuous at x if $x \in \mathbb{Q}$. We know f is not continuous at x if $x \in \mathbb{Q} \setminus \{0\}$. Therefore, f is not continuous at x if $x \in \mathbb{R} \setminus \{0\}$. Ergo, f is not continuous at any point $x \neq 0$. OEΔ

6.2 Solution (b)

Proof. Let $\epsilon > 0$. Suppose that $\delta = \epsilon$. Assume that $|x - p| < \delta$. This means that $|p - x| < \delta$. Since $x = 0$, $|p| < \delta$. Since $\delta = \epsilon$, this also means $|p| < \epsilon$.

Now, we apply this to the function. Suppose that $p \notin \mathbb{Q}$. This means that $|x - p| = |0 - 0| = |0| = 0$. By definition, we know that this works, since $\epsilon > 0$. Suppose that $p \in \mathbb{Q}$. This means that $|x - p| = |p - 0| = |p|$. Earlier in the proof, we established that $|p| < \epsilon$, so it holds.

Ergo, f is continuous at $x = 0$. OEΔ