

Chapter 34 End-of-Chapter Problems

Halliday & Resnick, 10th Edition

Donald Aingworth IV

Hit me where it Matters

1 Problem 1

You look through a camera toward an image of a hummingbird in a plane mirror. The camera is 4.30 m in front of the mirror. The bird is at camera level, 5.00 m to your right and 3.30 m from the mirror. What is the distance between the camera and the apparent position of the bird's image in the mirror?

1.1 Solution

In a plane mirror, the object will appear as far from the mirror as it is legitimately. Add the distance from the mirror of the camera to the bird's distance to the mirror.

$$\Delta x = 4.30 \text{ m} + 3.30 \text{ m} = 7.60 \text{ m} \quad (1)$$

Here use the Pythagorean theorem to find the distance.

$$s = \sqrt{\Delta x^2 + \Delta y^2} = \sqrt{7.60^2 + 5.00^2} = \sqrt{57.76 + 25.00} = \sqrt{82.76} = \boxed{9.097 \text{ m}} \quad (2)$$

2 Problem 3

In Fig. 34-32, an isotropic point source of light S is positioned at distance d from a viewing screen A and the light intensity I_P at point P (level with S) is measured. Then a plane mirror M is placed behind S at distance d . By how much is I_P multiplied by the presence of the mirror?




Figure 34-32

2.1 Solution

The equation for the intensity of light.

$$I = \frac{P_s}{4\pi r^2} \quad (3)$$

This gives us an equation for I_P .

$$I_P = \frac{P_s}{4\pi d^2} \quad (4)$$

Using this, we can create an equation for the intensity at point P from the light that reflected from the mirror (call it I_M).

$$I_M = \frac{P_s}{4\pi(3d)^2} \quad (5)$$

We can use these to calculate the total intensity and relate that to I_P .

$$I_{net} = I_P + I_M = \frac{P_s}{4\pi d^2} + \frac{P_s}{4\pi(3d)^2} \quad (6)$$

$$= \frac{P_s}{4\pi d^2} \left(1 + \frac{1}{3^2}\right) = I_P \left(\frac{10}{9}\right) \quad (7)$$

$$\frac{I_{net}}{I_P} = \boxed{\frac{10}{9} = 1.11} \quad (8)$$

3 Problem 7

A concave shaving mirror has a radius of curvature of 35.0 cm. It is positioned so that the (upright) image of a man's face is 2.50 times the size of the face. How far is the mirror from the face?

3.1 Solution

The upright image size difference tells us the magnification and solve for the position of the image (i).

$$m = -\frac{i}{p} = 2.50 \quad (9)$$

$$i = -2.50p \quad (10)$$

Usng this and the focal point, we can find a value of p .

$$\frac{1}{p} + \frac{1}{i} = \frac{2}{r} \quad (11)$$

$$\frac{1}{p} - \frac{1}{2.50p} = \frac{2}{r} \quad (12)$$

$$\frac{1}{p} \left(1 - \frac{2}{5}\right) = \frac{2}{r} \quad (13)$$

$$p = \frac{r}{2} \times \frac{3}{5} = \frac{3r}{10} = \frac{3}{10} \times 35.0 \text{ cm} = \boxed{10.5 \text{ cm}} \quad (14)$$

4 Problem 9-16

Object O stands on the central axis of a spherical mirror. For this situation, each problem in Table 34-3 gives object distance p_s (centimeters), the type of mirror, and then the distance (centimeters, without proper sign) between the focal point and the mirror. Find (a) the radius of curvature r (including sign), (b) the image distance i , and (c) the lateral magnification m . Also, determine whether the image is (d) real (R) or virtual (V), (e) inverted (I) from object O or non-inverted (NI), and (f) on the same side of the mirror as O or on the opposite side.

	p	Mirror	r	i	m	R/V	I/NI	Side
9	+18	Concave, 12	24	36	-2	Real	Inverted	Same
10	+15	Concave, 10	20	30	-2	Real	Inverted	Same
11	+8	Convex, 10	-20	-4.4	0.556	Virtual	Non-inverted	Other side
12	+24	Concave, 36	72	-72	3	Virtual	Non-inverted	Other side
13	+12	Concave, 18	36	-36	3	Virtual	Non-inverted	Other side
14	+22	Convex, 35	-70	-13.5	1.59	Virtual	Inverted	Other side
15	+10	Convex, 8	-16	-4.4	0.4	Virtual	Non-inverted	Other side
16	+17	Convex, 14	-28	-7.68	0.45	Virtual	Non-inverted	Other side

4.1 Equations used

$$\frac{1}{p} + \frac{1}{i} = \frac{1}{f} \rightarrow i = \left(\frac{1}{f} - \frac{1}{p} \right) \quad (15)$$

$$m = -\frac{i}{p} \quad (16)$$

Real means $i > 0$. Inverted means $m < 0$.

5 Problem 17-29

Object O stands on the central axis of a spherical or plane mirror. For this situation, each problem in Table 34-4 refers to (a) the type of mirror, (b) the focal distance f , (c) the radius of curvature r , (d) the object distance p , (e) the image distance i , and (f) the lateral magnification m . (All distances are in centimeters.) It also refers to whether (g) the image is real (R) or virtual (V), (h) inverted (I) or non-inverted (NI) from O, and (i) on the same side of the mirror as object O or on the opposite side. Fill in the missing information. Where only a sign is missing, answer with the sign.

Table 34-4 Problems 17 through 29: More Mirrors. See the setup for these problems.

	(a) Type	(b) f	(c) r	(d) p	(e) i	(f) m	(g) R/V	(h) I/NI	(i) Side
17	Concave	20		+10					
18				+24		0.50		I	
19			-40		-10				
20				+40		-0.70			
21		+20		+30					
22		20				+0.10			
23		30				+0.20			
24				+60		-0.50			
25				+30		0.40		I	
26		20		+60					
27		-30			-15				
28				+10		+1.0			
29	Convex		40		4.0				

5.1 Solution

	(a) Type	(b) f	(c) r	(d) p	(e) i	(f) m	(g) R/V	(h) I/NI	(i) Side
17	Concave	+20	+40	+10	-20	2	Virtual	Non-inverted	Other side
19	Convex	-20	-40	20	-10	0.5	Virtual	Non-inverted	Other side
21	Concave	+20	+40	+30	+60	-2	Real	Inverted	Same side
23	Convex	-30	-60	+120	-24	+0.2	Virtual	Non-inverted	Other side
25	Concave	+8.6	+17	+30	-12	-0.40	Virtual	Inverted	Other side
27	Convex	-30	-60	+30	-15	+0.5	Virtual	Non-inverted	Other side
29	Convex	-20	-40	+5	-4	+0.8	Virtual	Non-inverted	Other side

6 Problem 32-38

An object O stands on the central axis of a spherical refracting surface. For this situation, each problem in Table 34-5 refers to the index of refraction n_1 where the object is located, (a) the index of refraction n_2 on the other side of the refracting surface, (b) the object distance p , (c) the radius of curvature r of the surface, and (d) the image distance i . (All distances are in centimeters.) Fill in the missing information, including whether the image is (e) real (R) or virtual (V) and (f) on the same side of the surface as object O or on the opposite side.

Table 34-5 Problems 32 through 38: Spherical Refracting Surfaces. See the setup for these problems.

	n_1	(a) n_2	(b) p	(c) r	(d) i	(e) R/V	(f) Side
32	1.0	1.5	+10	+30			
33	1.0	1.5	+10		-13		
34	1.5		+100	-30	+600		
35	1.5	1.0	+70	+30			
36	1.5	1.0		-30	-7.5		
37	1.5	1.0	+10		-6.0		
38	1.0	1.5		+30	+600		

6.1 Solution

7 Problem 39

In Fig. 34-38, a beam of parallel light rays from a laser is incident on a solid transparent sphere of index of refraction n . (a) If a point image is produced at the back of the sphere, what is the index of refraction of the sphere? (b) What index of refraction, if any, will produce a point image at the center of the sphere?




Figure 34-38 Problem 39.

7.1 Solution (a)

Assume that the index of refraction outside the sphere to be equal to 1. The source would have to be infinitely far away to produce parallel lines, so we can say that $p = \infty$ and $\lim_{p \rightarrow \infty} \frac{n_1}{p} = 0$. We have an equation that includes known information and our sought-out index of refraction of the sphere.

$$\frac{n_1}{p} + \frac{n_2}{i} = \frac{n_2}{i} = \frac{n_2 - 1}{r} = \frac{n_2}{r} - \frac{1}{r} \quad (17)$$

The image hitting the other side at only one point means the image distance will be twice the radius.

$$\frac{n_2}{2r} = \frac{n_2}{r} - \frac{1}{r} \quad (18)$$

$$n_2 - \frac{n_2}{2} = \frac{n_2}{2} = 1 \quad (19)$$

$$n_2 = \boxed{2} \quad (20)$$

7.2 Solution (b)

This is similar, so we can do the same thing as we did for part (a). This time we just set $i = r$.

$$\frac{n_2}{r} = \frac{n_2}{r} - \frac{1}{r} \quad (21)$$

$$0 = -\frac{1}{r} \rightarrow 0 = -1 \quad (22)$$

This makes it impossible.

8 Problem 41

A lens is made of glass having an index of refraction of 1.5. One side of the lens is flat, and the other is convex with a radius of curvature of 20 cm. (a) Find the focal length of the lens. (b) If an object is placed 40 cm in front of the lens, where is the image?

8.1 Solution (a)

Use an equation for the focal length.

$$\frac{1}{f} = (n - 1) \left(\frac{1}{r_1} - \frac{1}{r_2} \right) \quad (23)$$

The value of r_1 will be ∞ , while the value of r_2 will be -20 cm. We know that $n = 1.5$.

$$\frac{1}{f} = (1.5 - 1) \left(0 - \frac{1}{-20 \text{ cm}} \right) = \frac{1}{2} \times \frac{1}{20 \text{ cm}} = \frac{1}{40 \text{ cm}} \quad (24)$$

$$f = \boxed{40 \text{ cm}} \quad (25)$$

8.2 Solution (b)

Since it is at the focal point, the image will be infinitely far away.

9 Problem 45

You produce an image of the Sun on a screen, using a thin lens whose focal length is 20.0 cm. What is the diameter of the image? (See Appendix C for needed data on the Sun.)

9.1 Solution

The sun is about 1.5×10^{11} m away. Its radius is about 6.96×10^8 m, leaving it with a diameter of 1.392×10^9 m. Our focal length is 0.200 m. We can use this to find the image distance.

$$\frac{1}{p} + \frac{1}{i} = \frac{1}{f} \quad (26)$$

$$i = \left(\frac{1}{f} - \frac{1}{p} \right)^{-1} = \left(\frac{1}{0.200 \text{ m}} - \frac{1}{1.5 \times 10^{11} \text{ m}} \right)^{-1} \quad (27)$$

$$\approx \left(\frac{1}{0.200} \right)^{-1} = 0.200 \text{ m} \quad (28)$$

This can be used to find the magnification.

$$m = -\frac{i}{p} = -\frac{0.200 \text{ m}}{1.5 \times 10^{11} \text{ m}} = -1.33 \times 10^{-12} \quad (29)$$

Multiply the diameter by the magnification to find the image diameter.

$$d' = |md| = |-1.33 \times 10^{-12} \times 1.392 \times 10^9 \text{ m}| \quad (30)$$

$$= \left| -\frac{29}{15625} \text{ m} \right| = \boxed{1.856 \times 10^{-3} \text{ m}} \quad (31)$$

10 Problem 49

An illuminated slide is held 44 cm from a screen. How far from the slide must a lens of focal length 11 cm be placed (between the slide and the screen) to form an image of the slide's picture on the screen?

10.1 Solution

Use the thin lens equation.

$$\frac{1}{p} + \frac{1}{i} = \frac{1}{f} \quad (32)$$

There is also an equation that would allow for the lens to be between the slide and the screen.

$$p + i = 44 \text{ cm} \quad (33)$$

Combine these and solve for p .

$$i = 44 \text{ cm} - p \quad (34)$$

$$\frac{1}{11 \text{ cm}} = \frac{1}{p} + \frac{1}{44 \text{ cm} - p} = \frac{44 \text{ cm} - p + p}{p(44 \text{ cm} - p)} \quad (35)$$

$$44p - p^2 = 484 \quad (36)$$

$$0 = p^2 - 44p + 484 \quad (37)$$

$$p = \frac{44 \pm \sqrt{44^2 - 4 \times 484}}{2} = \boxed{22 \text{ cm}} \quad (38)$$

11 Problem 53

11.1 Solution

12 Problem 57

12.1 Solution

13 Problem 63

13.1 Solution

14 Problem 69

14.1 Solution

15 Problem 73

15.1 Solution

16 Problem 75

16.1 Solution

17 Problem 81

17.1 Solution

18 Problem 83

18.1 Solution

19 Problem 85

19.1 Solution

20 Problem 87

20.1 Solution

21 Problem 99

21.1 Solution

22 Problem 105

22.1 Solution

23 Problem 109

23.1 Solution

Contents

1 Problem 1	2
1.1 Solution	2
2 Problem 3	3
2.1 Solution	3
3 Problem 7	4
3.1 Solution	4
4 Problem 9-16	5
4.1 Equations used	5
5 Problem 17-29	6
5.1 Solution	6
6 Problem 32-38	7
6.1 Solution	7
7 Problem 39	8
7.1 Solution (a)	8
7.2 Solution (b)	8
8 Problem 41	9
8.1 Solution (a)	9
8.2 Solution (b)	9
9 Problem 45	10
9.1 Solution	10
10 Problem 49	11
10.1 Solution	11
11 Problem 53	12
11.1 Solution	12
12 Problem 57	13
12.1 Solution	13

13 Problem 63	14
13.1 Solution	14
14 Problem 69	15
14.1 Solution	15
15 Problem 73	16
15.1 Solution	16
16 Problem 75	17
16.1 Solution	17
17 Problem 81	18
17.1 Solution	18
18 Problem 83	19
18.1 Solution	19
19 Problem 85	20
19.1 Solution	20
20 Problem 87	21
20.1 Solution	21
21 Problem 99	22
21.1 Solution	22
22 Problem 105	23
22.1 Solution	23
23 Problem 109	24
23.1 Solution	24