

# Chapter 33 End-of-Chapter Problems

Halliday & Resnick, 10th Edition

Donald Aingworth IV

Hit me where it Matters

# 1 Problem 1

A certain helium-neon laser emits red light in a narrow band of wavelengths centered at 632.8 nm and with a “wavelength width” (such as on the scale of Fig. 33-1) of 0.0100 nm. What is the corresponding “frequency width” for the emission?

## 1.1 Solution

Use the traditional formula for the wavelength. Here, the speed of the wave is the speed of light. We can treat this like an error and raw value issue.

$$v = \lambda f \rightarrow f = \frac{c}{\lambda} \quad (1)$$

$$\frac{\delta f}{f} = \frac{\delta \lambda}{\lambda} \quad (2)$$

$$\delta f = f * \frac{\delta \lambda}{\lambda} = c * \frac{\delta \lambda}{\lambda^2} \quad (3)$$

$$= 2.998 \times 10^8 \text{ m/s} * \frac{0.0100 \text{ nm}}{(632.8 \text{ nm})^2} \quad (4)$$

$$= \boxed{7.49 \text{ GHz}} \quad (5)$$

## 2 Problem 5

What inductance must be connected to a 17 pF capacitor in an oscillator capable of generating 550 nm (i.e., visible) electromagnetic waves? Comment on your answer.

### 2.1 Solution

For an  $LC$  circuit, the angular frequency is  $\frac{1}{\sqrt{LC}}$ .

$$\omega = \frac{1}{\sqrt{LC}} \quad (6)$$

The linear frequency is calculatable from this.

$$f = \frac{\omega}{2\pi} = \frac{1}{2\pi\sqrt{LC}} \quad (7)$$

This can be related to the wave speed (the speed of light in the case of an EM wave).

$$v = \lambda f \quad (8)$$

$$c = \lambda f = \frac{\lambda}{2\pi\sqrt{LC}} \quad (9)$$

This can be solved for the inductance ( $L$ ) and found a solution for.

$$c = \frac{\lambda}{2\pi\sqrt{L}\sqrt{C}} \quad (10)$$

$$\sqrt{L} = \frac{\lambda}{2\pi c\sqrt{C}} \quad (11)$$

$$L = \frac{\lambda^2}{(2\pi)^2 c^2 C} \quad (12)$$

$$= \frac{(550 \text{ nm})^2}{4\pi^2 (2.998 \times 10^8 \text{ m/s})^2 * 17 \text{ pF}} \quad (13)$$

$$= \boxed{5.015 \times 10^{-21} \text{ H}} \quad (14)$$

### 3 Problem 7

What is the intensity of a traveling plane electromagnetic wave if  $B_m$  is  $1.0 \times 10^{-4} \text{ T}$ ?

#### 3.1 Solution

Start by calculating the electric wave magnitude.

$$c = \frac{E_m}{B_m} \quad (15)$$

$$E_m = c B_m = 2.998 \times 10^8 \text{ m/s} * 1.0 \times 10^{-4} \text{ T} = 2.998 \times 10^4 \text{ N/C} \quad (16)$$

The intensity can be calculated from this. Bear in mind that  $E_{\text{rms}} = \frac{E_m}{\sqrt{2}}$ .

$$I = \frac{1}{c\mu_0} E_{\text{rms}}^2 = \frac{E_m^2}{2c\mu_0} = \frac{cB_m^2}{2\mu_0} \quad (17)$$

$$= \frac{2.998 \times 10^8 \text{ m/s} * 10^{-8} \text{ T}^2}{2 * 1.257 \times 10^{-6} \text{ H/m}} \quad (18)$$

$$= \boxed{1.193 \times 10^6 \text{ W/m}} \quad (19)$$

### 4 Problem 9

Some neodymium-glass lasers can provide 100 TW of power in 1.0 ns pulses at a wavelength of 0.26  $\mu\text{m}$ . How much energy is contained in a single pulse?

#### 4.1 Solution

Multiply the power of the laser by the frequency of the pulses to get the energy of a single pulse.

$$E = P * f = 100 \times 10^{12} \text{ W} * 1.0 \times 10^{-9} \text{ s} = \boxed{100 \text{ kJ}} \quad (20)$$

## 5 Problem 11

A plane electromagnetic wave traveling in the positive direction of an  $x$  axis in vacuum has components  $E_x = E_y = 0$  and  $E_z$  has the below value.

$$E_z = (2.0 \text{ V/m}) \cos [(\pi \times 10^{15} \text{ s}^{-1})(t - x/c)] \quad (21)$$

(a) What is the amplitude of the magnetic field component? (b) Parallel to which axis does the magnetic field oscillate? (c) When the electric field component is in the positive direction of the  $z$  axis at a certain point P, what is the direction of the magnetic field component there?

### 5.1 Solution (a)

The amplitude would be proportional to the electric field component and the speed of light.

$$B_m = \frac{E_m}{c} = \frac{2.0 \text{ V/m}}{2.998 \times 10^8 \text{ m/s}} = \boxed{6.67 \times 10^{-9} \text{ T}} \quad (22)$$

### 5.2 Solution (b)

It would also depend on the  $x$ -value so it would not be parallel to the  $x$  axis. It can't be parallel to the  $z$  axis because the electric field already is. The only axis left is the  $y$  axis.

### 5.3 Solution (c)

The the magnetic wave along its axis is negatively proportional to the magnitude of the electric wave on its axis. As such, if the electric wave is in the positive direction of the  $z$  axis, then the magnetic wave will be in the negative  $y$ -direction.

## 6 Problem 13

Sunlight just outside Earth's atmosphere has an intensity of  $1.40 \text{ kW/m}^2$ . Calculate (a)  $E_m$  and (b)  $B_m$  for sunlight there, assuming it to be a plane wave.

### 6.1 Solution (a)

The magnitude of the electric field is calculatable from the intensity. Start with the RMS electric field and the intensity.

$$I = \frac{1}{c\mu_0} E_{rms}^2 \quad (23)$$

The relationship with the magnitude of the electric field and the RMS electric field ( $E_{rms}^2 = \frac{E_m^2}{2}$ ) can be applied here.

$$I = \frac{E_m^2}{2c\mu_0} \quad (24)$$

This can be solved for  $E_m$  to find its value.

$$E_m = \sqrt{2Ic\mu_0} \quad (25)$$

$$= \sqrt{2 * 1.40 \times 10^3 \text{ W/m}^2 * 2.998 \times 10^8 \text{ m/s} * 1.257 \times 10^{-6} \text{ H/m}} \quad (26)$$

$$= \sqrt{1.055 \times 10^6 \text{ N}^2/\text{C}^2} = \boxed{1027 \text{ N/C}} \quad (27)$$

### 6.2 Solution (b)

Use the fraction for electromagnetic waves related to the speed of light.

$$c = \frac{E_m}{B_m} \quad (28)$$

Solve for and find the magnitude of the magnetic wave.

$$B_m = \frac{E_m}{c} = \frac{1027 \text{ N/C}}{2.998 \times 10^8 \text{ m/s}} \quad (29)$$

$$= \boxed{3.426 \times 10^{-6} \text{ T}} \quad (30)$$

## 7 Problem 19

High-power lasers are used to compress a plasma (a gas of charged particles) by radiation pressure. A laser generating radiation pulses with peak power  $1.5 \times 10^3$  MW is focused onto  $1.0 \text{ mm}^2$  of high-electron-density plasma. Find the pressure exerted on the plasma if the plasma reflects all the light beams directly back along their paths.

### 7.1 Solution

## 8 Problem 21

What is the radiation pressure 1.5 m away from a 500 W lightbulb? Assume that the surface on which the pressure is exerted faces the bulb and is perfectly absorbing and that the bulb radiates uniformly in all directions.

### 8.1 Solution



## 9 Problem 29

A small spaceship with a mass of only  $1.5 \times 10^3$  kg (including an astronaut) is drifting in outer space with negligible gravitational forces acting on it. If the astronaut turns on a 10 kW laser beam, what speed will the ship attain in 1.0 day because of the momentum carried away by the beam?

### 9.1 Solution

## 10 Problem 33

In Fig. 33-40, initially unpolarized light is sent into a system of three polarizing sheets whose polarizing directions make angles of  $\theta_1 = 40^\circ$ ,  $\theta_2 = 20^\circ$ , and  $\theta_3 = 40^\circ$  with the direction of the y axis. What percentage of the light's initial intensity is transmitted by the system? (Hint: Be careful with the angles.)

### 10.1 Solution

## 11 Problem 37

We want to rotate the direction of polarization of a beam of polarized light through  $90^\circ$  by sending the beam through one or more polarizing sheets. (a) What is the minimum number of sheets required? (b) What is the minimum number of sheets required if the transmitted intensity is to be more than 60% of the original intensity?

### 11.1 Solution

## 12 Problem 41

A beam of polarized light is sent into a system of two polarizing sheets. Relative to the polarization direction of that incident light, the polarizing directions of the sheets are at angles  $\theta$  for the first sheet and  $90^\circ$  for the second sheet. If 0.10 of the incident intensity is transmitted by the two sheets, what is  $\theta$ ?

### 12.1 Solution

## 13 Problem 43

A beam of partially polarized light can be considered to be a mixture of polarized and unpolarized light. Suppose we send such a beam through a polarizing filter and then rotate the filter through  $360^\circ$  while keeping it perpendicular to the beam. If the transmitted intensity varies by a factor of 5.0 during the rotation, what fraction of the intensity of the original beam is associated with the beam's polarized light?

### 13.1 Solution

## 14 Problem 45

When the rectangular metal tank in Fig. 33-46 is filled to the top with an unknown liquid, observer O, with eyes level with the top of the tank, can just see corner E. A ray that refracts toward O at the top surface of the liquid is shown. If  $D = 85.0$  cm and  $L = 1.10$  m, what is the index of refraction of the liquid?

### 14.1 Solution

## 15 Problem 49

Figure 33-49 shows light reflecting from two perpendicular reflecting surfaces A and B. Find the angle between the incoming ray  $i$  and the outgoing ray  $r'$ .

### 15.1 Solution

## 16 Problem 51

In Fig. 33-51, light is incident at angle  $\theta_1 = 40.1^\circ$  on a boundary between two transparent materials. Some of the light travels down through the next three layers of transparent materials, while some of it reflects upward and then escapes into the air. If  $n_1 = 1.30$ ,  $n_2 = 1.40$ ,  $n_3 = 1.32$ , and  $n_4 = 1.45$ , what is the value of (a)  $\theta_5$  in the air and (b)  $\theta_4$  in the bottom material?

### 16.1 Solution



## 17 Problem 55

In Fig. 33-55, a 2.00-m-long vertical pole extends from the bottom of a swimming pool to a point 50.0 cm above the water. Sunlight is incident at angle  $\theta = 55.0^\circ$ . What is the length of the shadow of the pole on the level bottom of the pool?

### 17.1 Solution

## 18 Problem 59

In Fig. 33-57, a ray of light is perpendicular to the face  $ab$  of a glass prism ( $n = 1.52$ ). Find the largest value for the angle  $\phi$  so that the ray is totally reflected at face  $ac$  if the prism is immersed (a) in air and (b) in water.

### 18.1 Solution

## 19 Problem 65

Figure 33-61 depicts a simplistic optical fiber: a plastic core ( $n_1 = 1.58$ ) is surrounded by a plastic sheath ( $n_2 = 1.53$ ). A light ray is incident on one end of the fiber at angle  $\theta$ . The ray is to undergo total internal reflection at point A, where it encounters the core-sheath boundary. (Thus there is no loss of light through that boundary.) What is the maximum value of  $\theta$  that allows total internal reflection at A?

### 19.1 Solution

## 20 Problem 69

Light that is traveling in water (with an index of refraction of 1.33) is incident on a plate of glass (with index of refraction 1.53). At what angle of incidence does the reflected light end up fully polarized?

### 20.1 Solution

## 21 Problem 71

### 21.1 Solution

## 22 Problem 75

### 22.1 Solution

## **23 Problem 83**

### **23.1 Solution**

## 24 Problem 87

### 24.1 Solution



## 25 Problem 89

### 25.1 Solution

## 26 Problem 91

### 26.1 Solution

## 27 Problem 105

### 27.1 Solution

# Contents

<b>1</b>	<b>Problem 1</b>	<b>2</b>
1.1	Solution . . . . .	2
<b>2</b>	<b>Problem 5</b>	<b>3</b>
2.1	Solution . . . . .	3
<b>3</b>	<b>Problem 7</b>	<b>4</b>
3.1	Solution . . . . .	4
<b>4</b>	<b>Problem 9</b>	<b>4</b>
4.1	Solution . . . . .	4
<b>5</b>	<b>Problem 11</b>	<b>5</b>
5.1	Solution (a) . . . . .	5
5.2	Solution (b) . . . . .	5
5.3	Solution (c) . . . . .	5
<b>6</b>	<b>Problem 13</b>	<b>6</b>
6.1	Solution (a) . . . . .	6
6.2	Solution (b) . . . . .	6
<b>7</b>	<b>Problem 19</b>	<b>7</b>
7.1	Solution . . . . .	7
<b>8</b>	<b>Problem 21</b>	<b>8</b>
8.1	Solution . . . . .	8
<b>9</b>	<b>Problem 29</b>	<b>9</b>
9.1	Solution . . . . .	9
<b>10</b>	<b>Problem 33</b>	<b>10</b>
10.1	Solution . . . . .	10
<b>11</b>	<b>Problem 37</b>	<b>11</b>
11.1	Solution . . . . .	11
<b>12</b>	<b>Problem 41</b>	<b>12</b>
12.1	Solution . . . . .	12

<b>13 Problem 43</b>	<b>13</b>
13.1 Solution . . . . .	13
<b>14 Problem 45</b>	<b>14</b>
14.1 Solution . . . . .	14
<b>15 Problem 49</b>	<b>15</b>
15.1 Solution . . . . .	15
<b>16 Problem 51</b>	<b>16</b>
16.1 Solution . . . . .	16
<b>17 Problem 55</b>	<b>17</b>
17.1 Solution . . . . .	17
<b>18 Problem 59</b>	<b>18</b>
18.1 Solution . . . . .	18
<b>19 Problem 65</b>	<b>19</b>
19.1 Solution . . . . .	19
<b>20 Problem 69</b>	<b>20</b>
20.1 Solution . . . . .	20
<b>21 Problem 71</b>	<b>21</b>
21.1 Solution . . . . .	21
<b>22 Problem 75</b>	<b>22</b>
22.1 Solution . . . . .	22
<b>23 Problem 83</b>	<b>23</b>
23.1 Solution . . . . .	23
<b>24 Problem 87</b>	<b>24</b>
24.1 Solution . . . . .	24
<b>25 Problem 89</b>	<b>25</b>
25.1 Solution . . . . .	25

<b>26 Problem 91</b>	<b>26</b>
26.1 Solution . . . . .	26
<b>27 Problem 105</b>	<b>27</b>
27.1 Solution . . . . .	27