

Worksheet #6

PHYS 4C: Waves and Thermodynamics

Donald Aingworth IV

September 22, 2025

Problem

One of the fundamental principles of Classical Mechanics is that if the positions and velocities of all particles in an isolated system are known at some initial time and the internal interactions are completely understood (i.e., all force laws are known), then it should be possible (at least in theory) to determine the motion of the entire system for all subsequent time.

Suppose we have an infinite string with given tension T and mass/length μ and it is known at time $t = 0$ that $y(x, 0) = Y(x)$ and that all parts of the string are completely at rest. Determine $y(x, t)$ for the string (do not assume that the wave must travel in one direction or the other). Does your result make sense?

The following procedure is recommended.

- Start with the general expression $y(x, t) = f(x - vt) + g(x + vt)$.
- Derive an expression for $\frac{\partial y}{\partial t}$.
- Set $t = 0$ and construct two equations involving the two unknown functions $f(x)$ and $g(x)$.
- Solve those equations for $f(x)$ and $g(x)$ separately.
- Substitute into the expression for $y(x, t)$.

Solution

Suppose that the string's wave has an equation involving a combination of two different solutions.

$$y(x, t) = f(x - vt) + g(x + vt) \quad (1)$$

We can in turn formulate an equation for the velocity of any point at a given time t by taking the partial derivative with respect to t .

$$\frac{\partial y}{\partial t}(x, t) = v(g(x + vt) - f(x - vt)) \quad (2)$$

Setting $t = 0$, we can form two equations, one from the position and one from the velocity of each particle. Bear in mind that it is given that at time $t = 0$, all parts of the string are at rest.

$$y(x, 0) = Y(x) = f(x) + g(x) \quad (3)$$

$$\frac{\partial y}{\partial t}(x, 0) = 0 = v(g(x) - f(x)) \quad (4)$$

We can solve the second equation and find a very simple value for $g(x)$.

$$0 = v(g(x) - f(x)) \quad (5)$$

$$0 = g(x) - f(x) \quad (6)$$

$$g(x) = f(x) \quad (7)$$

This can be substituted into the first equation.

$$Y(x) = f(x) + g(x) = 2f(x) = 2g(x) \quad (8)$$