

Chapter 16 End-of-Chapter Problems

Halliday & Resnick, 10th Edition

Donald Aingworth IV

Hit me where it Matters

1 Problem 1

If a wave $y(x, t) = (6.0\text{mm}) \sin(kx + (600 \text{ rad/s})t + \phi)$ travels along a string, how much time does any given point on the string take to move between displacements $y = +2.0 \text{ mm}$ and $y = -2.0 \text{ mm}$?

1.1 Solution

Notice that the amplitude of the string is 6.0mm. Our positive and negative final amplitudes of y are 2 and -2 millimeters respectively. Note that these two values have a magnitude of one third of the amplitude. There are four points on the unit circle where the sine of the angle is $\frac{1}{3}$. We can assume that the problem is asking the minimum time, since there are two ways that a point can travel between 2mm and -2mm. These two points would be for $kx + (600 \text{ rad/s})t + \phi = \theta$, starting when $\theta = \arcsin\left(\frac{1}{3}\right)$ and ending when $\theta = \pi - \arcsin\left(\frac{1}{3}\right)$. We can find the difference between these two.

$$\Delta\theta = \theta_f - \theta_i = \pi - \arcsin\left(\frac{1}{3}\right) - \arcsin\left(\frac{1}{3}\right) \quad (1)$$

$$= 2.462 \text{ rad} \quad (2)$$

As far as we consider it, the only thing in $kx + (600 \text{ rad/s})t + \phi$ changing is t . Since we're generalizing, the other things besides $(600 \text{ rad/s})t$ can be considered constants and initial conditions, so we can look at t .

$$\Delta\theta = (kx + (600 \text{ rad/s})t_f + \phi) - (kx + (600 \text{ rad/s})t_i + \phi) \quad (3)$$

$$= (600 \text{ rad/s})(t_f - t_i) \quad (4)$$

$$\Delta t = \frac{\Delta\theta}{(600 \text{ rad/s})} = \frac{2.462 \text{ rad}}{(600 \text{ rad/s})} = \boxed{0.0041 \text{ s}} \quad (5)$$

Note: if you were to find the minimum time between -2mm and 2mm, that would result in 1.1ms, but that is not the question posed here.

2 Problem 3

A wave has an angular frequency of 110 rad/s and a wavelength of 1.80 m. Calculate (a) the angular wave number and (b) the speed of the wave.

2.1 Solution (a)

We know the wavelength of the wave, which we can use to find the wave number.

$$k = \frac{2\pi}{\lambda} = \frac{2\pi}{1.80} = \frac{\pi}{0.9} = \boxed{3.49 \text{ rad/m}} \quad (6)$$

2.2 Solution (b)

We can use our angular wave number we found and the angular velocity to find the wave velocity.

$$v = \frac{\omega}{k} = \frac{110 \text{ rad/s}}{\pi \text{ rad}/0.9 \text{ m}} = \frac{110 * 0.9}{\pi} \text{ m/s} = \boxed{31.51 \text{ m/s}} \quad (7)$$

3 Problem 5

A sinusoidal wave travels along a string. The time for a particular point to move from maximum displacement to zero is 0.170 s. What are the (a) period and (b) frequency? (c) The wavelength is 1.40 m; what is the wave speed?

3.1 Solution (a)

The time to return from maximum displacement in one direction would be a quarter of the time to move from maximum displacement in the same direction (one period).

$$T = 4 \Delta t = 4 * 0.170 \text{ s} = \boxed{0.680 \text{ s}} \quad (8)$$

3.2 Solution (b)

The frequency is the reciprocal of the period.

$$f = \frac{1}{T} = \frac{1}{0.680 \text{ s}} = \boxed{1.47 \text{ Hz}} \quad (9)$$

3.3 Solution (c)

The velocity is equal to the wavelength divided by the period.

$$v = \frac{\lambda}{T} = \frac{1.40 \text{ m}}{0.680 \text{ s}} = \boxed{2.06 \text{ m/s}} \quad (10)$$

4 Problem 9

A sinusoidal wave moving along a string is shown twice in Fig. 16-33, as crest A travels in the positive direction of an x axis by distance $d = 6.0$ cm in 4.0 ms. The tick marks along the axis are separated by 10 cm; height $H = 6.00$ mm. The equation for the wave is in the form $y(x, t) = y_m \sin(kx \pm \omega t)$, so what are (a) y_m , (b) k , (c) ω , and (d) the correct choice of sign in front of ω ?




Figure 16-33 Problem 9.

4.1 Solution

The velocity of the wave would be equivalent to the distance covered divided by the time taken.

$$v = \frac{\Delta x}{\Delta t} = \frac{0.06 \text{ m}}{0.004 \text{ s}} = 15 \text{ m/s} \quad (11)$$

The amplitude would also be equal to half the distance between the top and bottom peaks (H).

$$A = \frac{H}{2} = \frac{6 \text{ mm}}{2} = 3 \text{ mm} = y_m \quad (12)$$

Note that the wave goes through one period over the course of four tick marks of a length of 10 cm each, equivalent to 40 cm total. This would be equivalent to the wavelength, which can be used to find the frequency. This frequency can be used to find the angular velocity.

$$v = \lambda f \rightarrow f = \frac{v}{\lambda} = \frac{15 \text{ m/s}}{0.4 \text{ m}} = 37.5 \text{ Hz} \quad (13)$$

$$f = \frac{\omega}{2\pi} \rightarrow \omega = 2\pi f = 75\pi \text{ rad/s} \quad (14)$$

Knowing our velocity and angular velocity, we can use this to find the angular wave number.

$$v = \frac{\omega}{k} \rightarrow k = \frac{\omega}{v} = \frac{75\pi \text{ rad/s}}{15 \text{ m/s}} = 5\pi \text{ rad/m} \quad (15)$$

We have our values needed to fill out the sinusoidal wave. Since the wave travels in a positive direction, the sign here would be negative.

$$\boxed{y(x, t) = (3 \text{ mm}) \sin((5\pi \text{ rad/m})x - (75\pi \text{ rad/s})t)} \quad (16)$$

5 Problem 15

A stretched string has a mass per unit length of 5.00 g/cm and a tension of 10.0 N. A sinusoidal wave on this string has an amplitude of 0.12 mm and a frequency of 100 Hz and is traveling in the negative direction of an x axis. If the wave equation is of the form $y(x, t) = y_m \sin(kx \pm \omega t)$, what are (a) y_m , (b) k , (c) ω , and (d) the correct choice of sign in front of ω ?

5.1 Solution

We have the value of the amplitude, which would be equal to y_m .

$$y_m = A = 0.12 \text{ mm} \quad (17)$$

Our frequency can give us the value of ω .

$$f = \frac{\omega}{2\pi} \quad (18)$$

$$\omega = 2\pi f = 2\pi * 100 \text{ Hz} = 200\pi \text{ rad/s} \quad (19)$$

For the wave number k , we can start by finding the speed of the wave. This should be done under consideration of a stretched string.

$$v = \sqrt{\frac{\tau}{\mu}} = \sqrt{\frac{10.0 \text{ N}}{5.00 \text{ g/cm}}} = \sqrt{\frac{100.0 \text{ N}}{5.00 \text{ kg/m}}} = \sqrt{20 \text{ m}^2/\text{s}^2} \quad (20)$$

This in turn can be used in conjunction with the angular velocity of the wave.

$$v = \frac{\omega}{k} \quad (21)$$

$$k = \frac{\omega}{v} = \frac{200\pi \text{ rad/s}}{\sqrt{20 \text{ m}^2/\text{s}^2}} = 140.5 \text{ m}^{-1} \quad (22)$$

Since the wave is traveling in the negative direction, the sign would be positive.

$$\boxed{y(x, t) = (1.2 \times 10^{-4} \text{ m}) \sin((140.5 \text{ m}^{-1})x + (200\pi \text{ rad/s})t)} \quad (23)$$

6 Problem 19

What is the speed of a transverse wave in a rope of length 2.00 m and mass 60.0 g under a tension of 500 N?

6.1 Solution

First find the linear density of the rope.

$$\mu = \frac{m}{L} = \frac{0.060 \text{ kg}}{2.00 \text{ m}} = 0.03 \text{ kg/m} \quad (24)$$

Then, use the formula for the speed of the wave.

$$v = \sqrt{\frac{\tau}{\mu}} = \sqrt{\frac{500 \text{ N}}{0.03 \text{ kg/m}}} = \boxed{129.1 \text{ m/s}} \quad (25)$$

7 Problem 21

A 100 g wire is held under a tension of 250 N with one end at $x = 0$ and the other at $x = 10.0$ m. At time $t = 0$, pulse 1 is sent along the wire from the end at $x = 10.0$ m. At time $t = 30.0$ ms, pulse 2 is sent along the wire from the end at $x = 0$. At what position x do the pulses begin to meet?

7.1 Solution

Since the wire is a wire, the speed can be calculated by first finding the linear density and then the wave velocity.

$$\mu = \frac{m}{L} = \frac{0.1 \text{ kg}}{10.0 \text{ m}} = 0.01 \text{ kg/m} \quad (26)$$

$$v = \sqrt{\frac{\tau}{\mu}} = \sqrt{\frac{250 \text{ N}}{0.01 \text{ kg/m}}} = \sqrt{25000 \text{ m}^2/\text{s}^2} = 50 \sqrt{10} \text{ m/s} \quad (27)$$

This will be the speed of both waves, which we can use to find the location of the first wave when the second wave starts up.

$$\Delta x = v * \Delta t = 50 \sqrt{10} \text{ m/s} \times 30 \times 10^{-3} \text{ s} = 1.5 \sqrt{10} \text{ m} \quad (28)$$

$$x_i = L - \Delta x = 10.0 \text{ m} - 1.5 \sqrt{10} \text{ m} = \frac{20 - 3\sqrt{10}}{2} \text{ m} \quad (29)$$

Since the waves will be traveling at the same speed in opposite directions, they will meet at the average point of their starting points.

$$x_{\text{avg}} = \frac{x_i}{2} = \frac{20 - 3\sqrt{10}}{4} \text{ m} = \boxed{2.628 \text{ m}} \quad (30)$$

8 Problem 25

A uniform rope of mass m and length L hangs from a ceiling. (a) Show that the speed of a transverse wave on the rope is a function of y , the distance from the lower end, and is given by $v = \sqrt{gy}$. (b) Show that the time a transverse wave takes to travel the length of the rope is given by $t = 2\sqrt{L/g}$.

8.1 Solution (a)

There are two deterministic factors for the velocity of the wave: the tension force and the density of the wire. There are equations for each of these and they can be put into the velocity equation. The tension force would be equal to the downward force on the wire, which would only be the gravitational force.

$$\tau = F_g = mg * \frac{y}{L} \quad (31)$$

$$\mu = \frac{m}{L} \quad (32)$$

$$v = \sqrt{\frac{mg}{m/L} * \frac{y}{L}} = \sqrt{gy} \quad (33)$$

9 Problem 30

Use the wave equation to find the speed of a wave given in terms of the general function $h(x, t)$:

$$y(x, t) = (4.00\text{mm})h[(30\text{m}^{-1})x + (6.0\text{s}^{-1})t] \quad (34)$$

9.1 Solution

10 Problem 31

Two identical traveling waves, moving in the same direction, are out of phase by $\pi/2$ rad. What is the amplitude of the resultant wave in terms of the common amplitude y_m of the two combining waves?

10.1 Solution

11 Problem 35

Two sinusoidal waves of the same frequency travel in the same direction along a string. If $y_{m1} = 3.0\text{cm}$, $y_{m2} = 4.0\text{cm}$, $\psi_1 = 0$, and $\psi_2 = \pi/2$ rad, what is the amplitude of the resultant wave?

11.1 Solution

12 Problem 37

These two waves travel along the same string:

$$y_1(x, t) = (4.60\text{mm})\sin(2\pi x - 400\pi t) \quad (35)$$

$$y_2(x, t) = (5.60\text{mm})\sin(2\pi x - 400\pi t + 0.80\pi \text{ rad}). \quad (36)$$

What are (a) the amplitude and (b) the phase angle (relative to wave 1) of the resultant wave? (c) If a third wave of amplitude 5.00 mm is also to be sent along the string in the same direction as the first two waves, what should be its phase angle in order to maximize the amplitude of the new resultant wave?

12.1 Solution

13 Problem 41

A string fixed at both ends is 8.40 m long and has a mass of 0.120 kg. It is subjected to a tension of 96.0 N and set oscillating. (a) What is the speed of the waves on the string? (b) What is the longest possible wavelength for a standing wave? (c) Give the frequency of that wave.

13.1 Solution

14 Problem 43

What are (a) the lowest frequency, (b) the second lowest frequency, and (c) the third lowest frequency for standing waves on a wire that is 10.0 m long,

has a mass of 100 g, and is stretched under a tension of 250 N?

14.1 Solution

15 Problem 45

A string that is stretched between fixed supports separated by 75.0 cm has resonant frequencies of 420 and 315 Hz, with no intermediate resonant frequencies. What are (a) the lowest resonant frequency and (b) the wave speed?

15.1 Solution

16 Problem 53

A string oscillates according to the equation

$$y = (0.50\text{cm}) \sin \left[\left(\frac{\pi}{3} \text{cm}^{-1} \right) x \right] \cos \left[(40\pi \text{s}^{-1}) t \right] \quad (37)$$

What are the (a) amplitude and (b) speed of the two waves (identical except for direction of travel) whose superposition gives this oscillation? (c) What is the distance between nodes? (d) What is the transverse speed of a particle of the string at the position $x = 1.5 \text{ cm}$ when $t = \frac{9}{8} \text{ s}$?

16.1 Solution

17 Problem 61

17.1 Solution

18 Problem 65

18.1 Solution

19 Problem 73

19.1 Solution

20 Problem 75

20.1 Solution

21 Problem 83

21.1 Solution

Contents

1 Problem 1	2
1.1 Solution	2
2 Problem 3	3
2.1 Solution (a)	3
2.2 Solution (b)	3
3 Problem 5	4
3.1 Solution (a)	4
3.2 Solution (b)	4
3.3 Solution (c)	4
4 Problem 9	5
4.1 Solution	5
5 Problem 15	6
5.1 Solution	6
6 Problem 19	7
6.1 Solution	7
7 Problem 21	8
7.1 Solution	8
8 Problem 25	9
8.1 Solution (a)	9
9 Problem 30	9
9.1 Solution	9
10 Problem 31	9
10.1 Solution	10
11 Problem 35	10
11.1 Solution	10
12 Problem 37	10
12.1 Solution	10

13 Problem 41	10
13.1 Solution	10
14 Problem 43	10
14.1 Solution	11
15 Problem 45	11
15.1 Solution	11
16 Problem 53	11
16.1 Solution	11
17 Problem 61	12
17.1 Solution	12
18 Problem 65	13
18.1 Solution	13
19 Problem 73	14
19.1 Solution	14
20 Problem 75	15
20.1 Solution	15
21 Problem 83	16
21.1 Solution	16