

Chapter 33 End-of-Chapter Problems

Halliday & Resnick, 10th Edition

Donald Aingworth IV

Hit me where it Matters

1 Problem 1

A certain helium-neon laser emits red light in a narrow band of wavelengths centered at 632.8 nm and with a “wavelength width” (such as on the scale of Fig. 33-1) of 0.0100 nm. What is the corresponding “frequency width” for the emission?

1.1 Solution

Use the traditional formula for the wavelength. Here, the speed of the wave is the speed of light. We can treat this like an error and raw value issue.

$$v = \lambda f \rightarrow f = \frac{c}{\lambda} \quad (1)$$

$$\frac{\delta f}{f} = \frac{\delta \lambda}{\lambda} \quad (2)$$

$$\delta f = f * \frac{\delta \lambda}{\lambda} = c * \frac{\delta \lambda}{\lambda^2} \quad (3)$$

$$= 2.998 \times 10^8 \text{ m/s} * \frac{0.0100 \text{ nm}}{(632.8 \text{ nm})^2} \quad (4)$$

$$= \boxed{7.49 \text{ GHz}} \quad (5)$$

2 Problem 5

What inductance must be connected to a 17 pF capacitor in an oscillator capable of generating 550 nm (i.e., visible) electromagnetic waves? Comment on your answer.

2.1 Solution

For an LC circuit, the angular frequency is $\frac{1}{\sqrt{LC}}$.

$$\omega = \frac{1}{\sqrt{LC}} \quad (6)$$

The linear frequency is calculatable from this.

$$f = \frac{\omega}{2\pi} = \frac{1}{2\pi\sqrt{LC}} \quad (7)$$

This can be related to the wave speed (the speed of light in the case of an EM wave).

$$v = \lambda f \quad (8)$$

$$c = \lambda f = \frac{\lambda}{2\pi\sqrt{LC}} \quad (9)$$

This can be solved for the inductance (L) and found a solution for.

$$c = \frac{\lambda}{2\pi\sqrt{L}\sqrt{C}} \quad (10)$$

$$\sqrt{L} = \frac{\lambda}{2\pi c\sqrt{C}} \quad (11)$$

$$L = \frac{\lambda^2}{(2\pi)^2 c^2 C} \quad (12)$$

$$= \frac{(550 \text{ nm})^2}{4\pi^2 (2.998 \times 10^8 \text{ m/s})^2 * 17 \text{ pF}} \quad (13)$$

$$= \boxed{5.015 \times 10^{-21} \text{ H}} \quad (14)$$

3 Problem 7

What is the intensity of a traveling plane electromagnetic wave if B_m is $1.0 \times 10^{-4} \text{ T}$?

3.1 Solution

Start by calculating the electric wave magnitude.

$$c = \frac{E_m}{B_m} \quad (15)$$

$$E_m = c B_m = 2.998 \times 10^8 \text{ m/s} * 1.0 \times 10^{-4} \text{ T} = 2.998 \times 10^4 \text{ N/C} \quad (16)$$

The intensity can be calculated from this. Bear in mind that $E_{\text{rms}} = \frac{E_m}{\sqrt{2}}$.

$$I = \frac{1}{c\mu_0} E_{\text{rms}}^2 = \frac{E_m^2}{2c\mu_0} = \frac{cB_m^2}{2\mu_0} \quad (17)$$

$$= \frac{2.998 \times 10^8 \text{ m/s} * 10^{-8} \text{ T}^2}{2 * 1.257 \times 10^{-6} \text{ H/m}} \quad (18)$$

$$= \boxed{1.193 \times 10^6 \text{ W/m}} \quad (19)$$

4 Problem 9

Some neodymium-glass lasers can provide 100 TW of power in 1.0 ns pulses at a wavelength of 0.26 μm . How much energy is contained in a single pulse?

4.1 Solution

Multiply the power of the laser by the frequency of the pulses to get the energy of a single pulse.

$$E = P * f = 100 \times 10^{12} \text{ W} * 1.0 \times 10^{-9} \text{ s} = \boxed{100 \text{ kJ}} \quad (20)$$

5 Problem 11

A plane electromagnetic wave traveling in the positive direction of an x axis in vacuum has components $E_x = E_y = 0$ and E_z has the below value.

$$E_z = (2.0 \text{ V/m}) \cos [(\pi \times 10^{15} \text{ s}^{-1})(t - x/c)] \quad (21)$$

(a) What is the amplitude of the magnetic field component? (b) Parallel to which axis does the magnetic field oscillate? (c) When the electric field component is in the positive direction of the z axis at a certain point P, what is the direction of the magnetic field component there?

5.1 Solution (a)

The amplitude would be proportional to the electric field component and the speed of light.

$$B_m = \frac{E_m}{c} = \frac{2.0 \text{ V/m}}{2.998 \times 10^8 \text{ m/s}} = \boxed{6.67 \times 10^{-9} \text{ T}} \quad (22)$$

5.2 Solution (b)

It would also depend on the x -value so it would not be parallel to the x axis. It can't be parallel to the z axis because the electric field already is. The only axis left is the y axis.

5.3 Solution (c)

The the magnetic wave along its axis is negatively proportional to the magnitude of the electric wave on its axis. As such, if the electric wave is in the positive direction of the z axis, then the magnetic wave will be in the $\boxed{\text{negative } y\text{-direction}}$.

6 Problem 13

Sunlight just outside Earth's atmosphere has an intensity of 1.40 kW/m^2 . Calculate (a) E_m and (b) B_m for sunlight there, assuming it to be a plane wave.

6.1 Solution (a)

The magnitude of the electric field is calculatable from the intensity. Start with the RMS electric field and the intensity.

$$I = \frac{1}{c\mu_0} E_{rms}^2 \quad (23)$$

The relationship with the magnitude of the electric field and the RMS electric field ($E_{rms}^2 = \frac{E_m^2}{2}$) can be applied here.

$$I = \frac{E_m^2}{2c\mu_0} \quad (24)$$

This can be solved for E_m to find its value.

$$E_m = \sqrt{2Ic\mu_0} \quad (25)$$

$$= \sqrt{2 * 1.40 \times 10^3 \text{ W/m}^2 * 2.998 \times 10^8 \text{ m/s} * 1.257 \times 10^{-6} \text{ H/m}} \quad (26)$$

$$= \sqrt{1.055 \times 10^6 \text{ N}^2/\text{C}^2} = \boxed{1027 \text{ N/C}} \quad (27)$$

6.2 Solution (b)

Use the fraction for electromagnetic waves related to the speed of light.

$$c = \frac{E_m}{B_m} \quad (28)$$

Solve for and find the magnitude of the magnetic wave.

$$B_m = \frac{E_m}{c} = \frac{1027 \text{ N/C}}{2.998 \times 10^8 \text{ m/s}} \quad (29)$$

$$= \boxed{3.426 \times 10^{-6} \text{ T}} \quad (30)$$

7 Problem 19

High-power lasers are used to compress a plasma (a gas of charged particles) by radiation pressure. A laser generating radiation pulses with peak power 1.5×10^3 MW is focused onto 1.0 mm^2 of high-electron-density plasma. Find the pressure exerted on the plasma if the plasma reflects all the light beams directly back along their paths.

7.1 Solution

The power would be equivalent to the intensity multiplied by the area impacted.

$$P = IA \leftrightarrow I = \frac{P}{A} \quad (31)$$

This in turn can be used in the equation for pressure (and force) from radiation. Bear in mind that since the plasma reflects all light rays, we would use the corresponding equation.

$$p_r = \frac{2I}{c} = \frac{2P}{cA} = \frac{2 * 1.5 \times 10^9 \text{ W}}{(2.998 \times 10^8 \text{ m/s})(1.0 \times 10^{-6} \text{ m}^2)} \quad (32)$$

$$= \boxed{1.0 \times 10^7 \text{ Pa}} \quad (33)$$

8 Problem 21

What is the radiation pressure 1.5 m away from a 500 W lightbulb? Assume that the surface on which the pressure is exerted faces the bulb and is perfectly absorbing and that the bulb radiates uniformly in all directions.

8.1 Solution

Radiation pressure is equal to the intensity divided by the speed of light.

$$p_r = \frac{I}{c} \quad (34)$$

The intensity is equal to the power divided by the area affected.

$$I = \frac{P}{A} = \frac{P}{4\pi r^2} \quad (35)$$

This can be applied to find the radiation pressure.

$$p_r = \frac{I}{c} = \frac{\frac{P}{4\pi r^2}}{c} = \frac{P}{4\pi r^2 c} \quad (36)$$

$$= \frac{500 \text{ W}}{4\pi * (1.5 \text{ m})^2 * 2.998 \times 10^8 \text{ m/s}} \quad (37)$$

$$= \boxed{5.899 \times 10^{-8} \text{ Pa}} \quad (38)$$

9 Problem 29

A small spaceship with a mass of only 1.5×10^3 kg (including an astronaut) is drifting in outer space with negligible gravitational forces acting on it. If the astronaut turns on a 10 kW laser beam, what speed will the ship attain in 1.0 day because of the momentum carried away by the beam?

9.1 Solution

To begin, there are 86400 seconds in a day. Multiply this by the 10 kW, we wind up with 8.64×10^8 J lost or gained from the laser in one day. Divide this energy by the speed of light to get the momentum added, which we then divide by the mass to get the speed increase.

$$\Delta p = \frac{\Delta U}{c} = \frac{8.64 \times 10^8 \text{ J}}{2.998 \times 10^8 \text{ m/s}} = 2.882 \text{ N} \cdot \text{s} \quad (39)$$

$$\Delta v = \frac{\Delta p}{m} = 0.00192 \text{ m/s} \quad (40)$$

10 Problem 33

In Fig. 33-40, initially unpolarized light is sent into a system of three polarizing sheets whose polarizing directions make angles of $\theta_1 = 40^\circ$, $\theta_2 = 20^\circ$, and $\theta_3 = 40^\circ$ with the direction of the y axis. What percentage of the light's initial intensity is transmitted by the system? (Hint: Be careful with the angles.)

10.1 Solution

11 Problem 37

We want to rotate the direction of polarization of a beam of polarized light through 90° by sending the beam through one or more polarizing sheets. (a) What is the minimum number of sheets required? (b) What is the minimum number of sheets required if the transmitted intensity is to be more than 60% of the original intensity?

11.1 Solution

12 Problem 41

A beam of polarized light is sent into a system of two polarizing sheets. Relative to the polarization direction of that incident light, the polarizing directions of the sheets are at angles θ for the first sheet and 90° for the second sheet. If 0.10 of the incident intensity is transmitted by the two sheets, what is θ ?

12.1 Solution

13 Problem 43

A beam of partially polarized light can be considered to be a mixture of polarized and unpolarized light. Suppose we send such a beam through a polarizing filter and then rotate the filter through 360° while keeping it perpendicular to the beam. If the transmitted intensity varies by a factor of 5.0 during the rotation, what fraction of the intensity of the original beam is associated with the beam's polarized light?

13.1 Solution

14 Problem 45

When the rectangular metal tank in Fig. 33-46 is filled to the top with an unknown liquid, observer O, with eyes level with the top of the tank, can just see corner E. A ray that refracts toward O at the top surface of the liquid is shown. If $D = 85.0$ cm and $L = 1.10$ m, what is the index of refraction of the liquid?

14.1 Solution

15 Problem 49

Figure 33-49 shows light reflecting from two perpendicular reflecting surfaces A and B. Find the angle between the incoming ray i and the outgoing ray r' .

15.1 Solution

16 Problem 51

In Fig. 33-51, light is incident at angle $\theta_1 = 40.1^\circ$ on a boundary between two transparent materials. Some of the light travels down through the next three layers of transparent materials, while some of it reflects upward and then escapes into the air. If $n_1 = 1.30$, $n_2 = 1.40$, $n_3 = 1.32$, and $n_4 = 1.45$, what is the value of (a) θ_5 in the air and (b) θ_4 in the bottom material?

16.1 Solution

17 Problem 55

In Fig. 33-55, a 2.00-m-long vertical pole extends from the bottom of a swimming pool to a point 50.0 cm above the water. Sunlight is incident at angle $\theta = 55.0^\circ$. What is the length of the shadow of the pole on the level bottom of the pool?

17.1 Solution

18 Problem 59

In Fig. 33-57, a ray of light is perpendicular to the face ab of a glass prism ($n = 1.52$). Find the largest value for the angle ϕ so that the ray is totally reflected at face ac if the prism is immersed (a) in air and (b) in water.

18.1 Solution

19 Problem 65

Figure 33-61 depicts a simplistic optical fiber: a plastic core ($n_1 = 1.58$) is surrounded by a plastic sheath ($n_2 = 1.53$). A light ray is incident on one end of the fiber at angle θ . The ray is to undergo total internal reflection at point A, where it encounters the core-sheath boundary. (Thus there is no loss of light through that boundary.) What is the maximum value of θ that allows total internal reflection at A?

19.1 Solution

20 Problem 69

Light that is traveling in water (with an index of refraction of 1.33) is incident on a plate of glass (with index of refraction 1.53). At what angle of incidence does the reflected light end up fully polarized?

20.1 Solution

21 Problem 71

21.1 Solution

22 Problem 75

22.1 Solution

23 Problem 83

23.1 Solution

24 Problem 87

24.1 Solution

25 Problem 89

25.1 Solution

26 Problem 91

26.1 Solution

27 Problem 105

27.1 Solution

Contents

1	Problem 1	2
1.1	Solution	2
2	Problem 5	3
2.1	Solution	3
3	Problem 7	4
3.1	Solution	4
4	Problem 9	4
4.1	Solution	4
5	Problem 11	5
5.1	Solution (a)	5
5.2	Solution (b)	5
5.3	Solution (c)	5
6	Problem 13	6
6.1	Solution (a)	6
6.2	Solution (b)	6
7	Problem 19	7
7.1	Solution	7
8	Problem 21	8
8.1	Solution	8
9	Problem 29	9
9.1	Solution	9
10	Problem 33	10
10.1	Solution	10
11	Problem 37	11
11.1	Solution	11
12	Problem 41	12
12.1	Solution	12

13 Problem 43	13
13.1 Solution	13
14 Problem 45	14
14.1 Solution	14
15 Problem 49	15
15.1 Solution	15
16 Problem 51	16
16.1 Solution	16
17 Problem 55	17
17.1 Solution	17
18 Problem 59	18
18.1 Solution	18
19 Problem 65	19
19.1 Solution	19
20 Problem 69	20
20.1 Solution	20
21 Problem 71	21
21.1 Solution	21
22 Problem 75	22
22.1 Solution	22
23 Problem 83	23
23.1 Solution	23
24 Problem 87	24
24.1 Solution	24
25 Problem 89	25
25.1 Solution	25

26 Problem 91	26
26.1 Solution	26
27 Problem 105	27
27.1 Solution	27