

# 1 Problem 1

Find the mass in kilograms of  $7.50 \times 10^{24}$  atoms of arsenic, which has a molar mass of 74.9g/mol.

## 1.1 Solution

Convert atoms to moles.

$$\frac{7.50 \times 10^{24} \text{ atoms}}{6.02 \times 10^{23} \text{ atoms/mol}} = 12.458 \text{ mol} \quad (1)$$

Using the molar mass, convert moles to grams.

$$12.458 \text{ mol} * 74.9 \text{ g/mol} = 933.140 \text{ g} = \boxed{0.933 \text{ kg}} \quad (2)$$

## 2 Problem 3

Oxygen gas having a volume of  $1000\text{cm}^3$  at  $40.0^\circ\text{C}$  and  $1.01 \times 10^5\text{Pa}$  expands until its volume is  $1500\text{cm}^3$  and its pressure is  $1.06 \times 10^5\text{Pa}$ . Find (a) the number of moles of oxygen present and (b) the final temperature of the sample.

### 2.1 Solution (a)

We can use the ideal gas law for this. We apply it to the first case, converting the  $40.0^\circ\text{C}$  to Kelvin.

$$40^\circ\text{C} = 313.15\text{K} \quad (3)$$

$$1000\text{cm}^3 = 1000 \times 10^{-6}\text{m}^3 = 1 \times 10^{-3}\text{m}^3 \quad (4)$$

$$pV = nRT \quad (5)$$

$$n = \frac{pV}{RT} = \frac{1.01 \times 10^5\text{Pa} * 10^{-3}\text{cm}^3}{8.31\text{J/mol} \cdot \text{K} * 313.15\text{K}} \quad (6)$$

$$= \frac{1.01 \times 10^2\text{N} \cdot \text{m}}{2602.2765\text{J/mol}} = \boxed{0.038812\text{mol}} \quad (7)$$

### 2.2 Solution (b)

The ideal gas law (or an equivalent) will be used here. The number of moles does not change here, neither does the gas constant  $R$ . We can use this to solve for the final value of the temperature.

$$\frac{p_1 V_1}{n_1 T_1} = \frac{p_2 V_2}{n_2 T_2} \quad (8)$$

$$\frac{p_1 V_1}{T_1} = \frac{p_2 V_2}{T_2} \quad (9)$$

$$T_2 = T_1 * \frac{p_2 V_2}{p_1 V_1} \quad (10)$$

We can substitute in values now.

$$T_2 = T_1 * \frac{p_2 V_2}{p_1 V_1} = 313.15 * \frac{1.06 \times 10^5 * 1500}{1.01 \times 10^5 * 1000} \quad (11)$$

$$= 313.15 * \frac{1.06 * 1.5}{1.01} = \boxed{492.97\text{K} \approx 220^\circ\text{C}} \quad (12)$$

### 3 Problem 5

The best laboratory vacuum has a pressure of about  $1.00 \times 10^{-18}$  atm, or  $1.01 \times 10^{-13}$  Pa. How many gas molecules are there per cubic centimeter in such a vacuum at 293 K?

#### 3.1 Solution

Use the ideal gas law, the version with Boltzmann's constant. We can solve for  $\frac{N}{V}$ .

$$pV = NkT \quad (13)$$

$$\frac{N}{V} = \frac{p}{kT} \quad (14)$$

From here, we can just plug and chug, so to speak.

$$\frac{N}{V} = \frac{1.01 \times 10^{-13} \text{ N/m}^2}{1.38 \times 10^{-23} \text{ N} \cdot \text{m/K} * 293 \text{ K}} = \frac{1.01 \times 10^{10}}{404.34} \text{ m}^{-3} \quad (15)$$

$$= 24978978.09 \times \text{m}^{-3} = \boxed{24.979 \text{ cm}^{-3}} \quad (16)$$

## 4 Problem 7

Suppose 1.80 mol of an ideal gas is taken from a volume of 3.00 m<sup>3</sup> to a volume of 1.50 m<sup>3</sup> via an isothermal compression at 30°C. (a) How much energy is transferred as heat during the compression, and (b) is the transfer to or from the gas?

### 4.1 Solution (a)

Energy transfered can be thought of as work. We have a formula for work done by an ideal gas.

$$W = nRT \ln \left( \frac{V_f}{V_i} \right) \quad (17)$$

We can plug and chug into this.

$$T_K = T_C + 273.15\text{K} = 30^\circ\text{C} + 273.15\text{K} = 303.15\text{K} \quad (18)$$

$$W = (1.80\text{mol})(8.31\text{J/mol})(303.15\text{K}) \ln \left( \frac{1.5\text{m}^3}{3.0\text{m}^3} \right) \quad (19)$$

$$= 4534.5177\text{J} * (-0.693147) = -3143.088\text{J} \quad (20)$$

The energy transfered is the absolute value of this, which would be 3143.088J.

### 4.2 Solution (b)

This is a volume compression process. The total energy in the system would remain constant, so  $Q = W$  by the first law of thermodynamics. This means  $Q < 0$ , so the energy is transfered from the gas as heat.

## 5 Problem 9

An automobile tire has a volume of  $1.64 \times 10^{-2}\text{m}^3$  and contains air at a gauge pressure (pressure above atmospheric pressure) of 165 kPa when the temperature is  $0.00^\circ\text{C}$ . What is the gauge pressure of the air in the tires when its temperature rises to  $27.0^\circ\text{C}$  and its volume increases to  $1.67 \times 10^{-2}\text{m}^3$ ? Assume atmospheric pressure is  $1.01 \times 10^5\text{Pa}$ .

### 5.1 Solution

We can here use the ideal gas law.

$$pV = nRT \quad (21)$$

$$\frac{p_1 V_1}{T_1} = \frac{p_2 V_2}{T_2} \quad (22)$$

With our equation, we solve for  $p_f$ .

$$p_f = p_i \frac{V_i T_f}{V_f T_i} \quad (23)$$

This can be “plugged-and-chugged” to find our final answer. We should also bear in mind that the gauge pressure should be added to the atmospheric pressure to find the overall pressure. Do not forget to convert to Kelvin.

$$T_{iK} = 0^\circ\text{C} + 273\text{K} = 273\text{K} \quad (24)$$

$$T_{fK} = 27^\circ\text{C} + 273\text{K} = 300\text{K} \quad (25)$$

$$p_i = p_{\text{gauge}} + p_{\text{atm}} = 165 \times 10^3\text{Pa} + 1.01 \times 10^5\text{Pa} = 266 \times 10^3\text{Pa} \quad (26)$$

$$p_f = p_i \frac{V_i T_f}{V_f T_i} = 266 \times 10^3\text{Pa} * \frac{(1.64 \times 10^{-2}\text{m}^3)(300\text{K})}{(1.67 \times 10^{-2}\text{m}^3)(273\text{K})} \quad (27)$$

$$= 266 \times 10^3\text{Pa} * \frac{492}{455.91} = 287 \times 10^3\text{Pa} \quad (28)$$

This can in turn be turned into the gauge pressure (our final answer) by subtracting the atmospheric pressure.

$$p_{\text{gauge};f} = 287 \times 10^3\text{Pa} - 1.01 \times 10^5\text{Pa} = \boxed{1.86 \times 10^5\text{Pa}} \quad (29)$$

## 6 Problem 11

Air that initially occupies  $0.140\text{m}^3$  at a gauge pressure of  $103.0\text{kPa}$  is expanded isothermally to a pressure of  $101.3\text{kPa}$  and then cooled at constant pressure until it reaches its initial volume. Compute the work done by the air. Gauge pressure is the difference between the actual pressure and atmospheric pressure.

### 6.1 Solution

Isothermal expansion means that there is no net change in energy, so  $Q_{in} = W_{out}$  (a.k.a.  $Q_{in} = -W_{in}$ ) as volume increases. There is a formula for work we can use.

$$W = \int_{V_i}^{V_f} p(V) dV \quad (30)$$

The problem outright states that it is separated into two parts: the isothermal expansion at changing pressure and the cooling at constant pressure. We can separate our equation into these two parts and find useful equations for each. The integrals will be separated at a point of  $V = V_{max}$  since that is the point where the volume reaches its highest. We can call the final pressure value of  $101.3\text{kPa}$  the value  $p_r$ , with no meaning to its name.

$$W = \int_{V_i}^{V_f} p(V) dV = \int_{V_i}^{V_{max}} p_1(V) dV + \int_{V_{max}}^{V_f} p_2(V) dV \quad (31)$$

$$= \int_{V_i}^{V_{max}} \frac{nRT_i}{V} dV + \int_{V_{max}}^{V_f} p_r dV \quad (32)$$

$$= nRT_i \ln\left(\frac{V_{max}}{V_i}\right) + \int_{V_{max}}^{V_f} p_r dV \quad (33)$$

$$= p_i V_i \ln\left(\frac{V_{max}}{V_i}\right) + p_r (V_f - V_{max}) \quad (34)$$

This requires the initial and maximum magnitudes of the volume, of which we only have one. We can use the ideal gas law to find that which we need. First solve for the final volume, given  $n$ ,  $R$ , and  $T$  are constant. Do not forget to convert gauge pressure to overall pressure by adding the atmospheric pressure ( $1.01 \times 10^5\text{Pa}$ ). We can assume that the final pressure

is not gauge pressure.

$$p_i = p_1 + p_{atm} = 103.0 \times 10^3 + 1.01 \times 10^5 \text{Pa} = 2.04 \times 10^5 \text{Pa} \quad (35)$$

$$p_f = p_2 = 101.3 \times 10^3 \text{Pa} \quad (36)$$

$$pV = nRT \quad (37)$$

$$p_i V_i = p_f V_f = p_f V_{max} \quad (38)$$

$$V_{max} = V_i * \frac{p_i}{p_f} = 0.140 \text{m}^3 * \frac{2.04 \times 10^5 \text{Pa}}{1.013 \times 10^5 \text{Pa}} = 0.281935 \text{m}^3 \quad (39)$$

This can in turn be applied to our above equation for the work done by the gas.

$$W = p_i V_i \ln \left( \frac{V_{max}}{V_i} \right) + \int_{V_{max}}^{V_f} p_r dV \quad (40)$$

$$= 0.140 \text{m}^3 * 2.04 \times 10^5 \text{Pa} * \ln \left( \frac{0.281935 \text{m}^3}{0.140 \text{m}^3} \right) + \int_{0.281935 \text{m}^3}^{0.140 \text{m}^3} 1.013 \times 10^5 \text{Pa} dV \quad (41)$$

$$= 28560 * 0.7000 + 101300 * (-0.1419) = 19992.96 \text{J} - 14378 \text{J} \quad (42)$$

$$= \boxed{5614.95 \text{J}} \quad (43)$$

## 7 Problem 13

A sample of an ideal gas is taken through the cyclic process  $abca$  shown in Fig. 19-20. The scale of the vertical axis is set by  $p_b = 7.5 \text{ kPa}$  and  $p_{ac} = 2.5 \text{ kPa}$ . At point  $a$ ,  $T = 200 \text{ K}$ . (a) How many moles of gas are in the sample? What are (b) the temperature of the gas at point  $b$ , (c) the temperature of the gas at point  $c$ , and (d) the net energy added to the gas as heat during the cycle?




Figure 19-20 Problem 13.

### 7.1 Solution (a)

We can use the ideal gas law. We need to solve for  $n$  at point  $a$ .

$$pV = nRT \rightarrow n = \frac{pV}{RT} \quad (44)$$

From here, we plug in values to find  $n$ .

$$n = \frac{(2.5 \text{ kPa}) * (1.0 \text{ m}^3)}{R(200 \text{ K})} = \frac{2.5 \times 10^3 \text{ J}}{1662 \text{ J/mol}} = \boxed{1.504 \text{ mol}} \quad (45)$$

### 7.2 Solution (b)

We can use the ideal gas law again, this time comparing two situations and finding the final temperature.

$$\frac{p_i V_i}{T_i} = \frac{p_f V_f}{T_f} \rightarrow T_f = T_i * \frac{p_f V_f}{p_i V_i} \quad (46)$$

From here, we plug in values to find the final temperature.

$$T_f = T_i * \frac{p_f V_f}{p_i V_i} = 200 \text{ K} * \frac{7.5 \text{ kPa} * 3.0 \text{ m}^3}{2.5 \text{ kPa} * 1.0 \text{ m}^3} \quad (47)$$

$$= 200 \text{ K} * 9 = \boxed{1800 \text{ K}} \quad (48)$$



### 7.3 Solution (c)

Use the ideal gas law again. This time, the pressure does not change.

$$\frac{V_i}{T_i} = \frac{V_f}{T_f} \rightarrow T_f = T_i * \frac{V_f}{V_i} \quad (49)$$

Now, we plug in to find  $T_f$ .

$$T_f = T_i * \frac{V_f}{V_i} = 1800 \text{ K} * \frac{1.0 \text{ m}^3}{3.0 \text{ m}^3} = \boxed{600 \text{ K}} \quad (50)$$

### 7.4 Solution (d)

Let's start by calculating the total work done by the gas. We can separate it based on the phases. The phase between points b and c has no change in volume, so no work is done.

$$W = \int p(V) dV \quad (51)$$

$$= \int_{1.0\text{m}^3}^{3.0\text{m}^3} V * 2500 \text{ Pa/m}^3 dV + \int_{3.0\text{m}^3}^{3.0\text{m}^3} p_{bc} dV + \int_{3.0\text{m}^3}^{1.0\text{m}^3} 2500 \text{ Pa} dV \quad (52)$$

$$= (1250 \text{ Pa/m}^3) \left( \frac{(3.0\text{m}^3)^2 - (1.0\text{m}^3)^2}{2} \right) + 2500 \text{ Pa} * (-2.0\text{m}^3) \quad (53)$$

$$= 4 * 2500 \text{ J} - 2 * 2500 \text{ J} = 10000 - 5000 \text{ J} = 5000 \text{ J} \quad (54)$$

Since this is a cyclical process, we can apply the first law of thermodynamics.

$$\Delta E_{\text{int}} = 0 = Q_{\text{in}} - W_{\text{out}} \quad (55)$$

$$Q_{\text{in}} = W_{\text{out}} = \boxed{5000 \text{ J}} \quad (56)$$

## 8 Problem 17

Container A in Fig. 19-22 holds an ideal gas at a pressure of  $5.0 \times 10^5 \text{ Pa}$  and a temperature of  $300 \text{ K}$ . It is connected by a thin tube (and a closed valve) to container B, with four times the volume of A. Container B holds the same ideal gas at a pressure of  $1.0 \times 10^5 \text{ Pa}$  and a temperature of  $400 \text{ K}$ . The valve is opened to allow the pressures to equalize, but the temperature of each container is maintained. What then is the pressure?




Figure 19-22 Problem 17.

### 8.1 Solution

Let's put together a formulas for the number of moles of the ideal gas in each container (A and B). We can use the Ideal Gas Law for this.

$$n_A = \frac{p_A V_A}{RT_A} = \frac{5.0 \times 10^5 \text{ Pa} * V_A}{R * 300 \text{ K}} = \frac{5 \times 10^3}{3} * \frac{V_A}{R} \quad (57)$$

$$n_B = \frac{p_B V_B}{RT_B} = \frac{1.0 \times 10^5 \text{ Pa} * 4V_A}{R * 400 \text{ K}} = 1 \times 10^3 * \frac{V_A}{R} \quad (58)$$

We can combine these to find the total number of moles.

$$n_{\Sigma} = n_A + n_B = \left( \frac{5}{3} + 1 \right) \times 10^3 * \frac{V_A}{R} = \frac{8}{3} \times 10^3 * \frac{V_A}{R} \quad (59)$$

We also know the total volume that there is.

$$V_{\Sigma} = \sum V_i = V_A + V_B = V_A + 4V_A = 5V_A \quad (60)$$

We can also calculate the average temperature of the cobined system, weighted by the volumes.

$$T_{\text{avg}} = \frac{\sum V_i * T_i}{\sum V_i} = \frac{V_A * T_A + V_B * T_B}{5V_A} \quad (61)$$

$$= \frac{V_A * 300 \text{ K} + 4V_A * 400 \text{ K}}{5V_A} \quad (62)$$

$$= \frac{300 \text{ K} + 1600 \text{ K}}{5} = \frac{1900 \text{ K}}{5} = 380 \text{ K} \quad (63)$$

We have our final values for the number of moles, volume, and temperature, not to mention that  $R$  is a constant. We can put these into the ideal gas law to find the pressure at this point, assuming the pressure to be even.

$$pV = nRT \quad (64)$$

$$p = \frac{nRT}{V} = \frac{\frac{8}{3} \times 10^3 * \frac{V_A}{R} * R * 380 \text{ K}}{5V_A} \quad (65)$$

$$= \left( \frac{8}{15} \times 10^3 * 380 \right) \text{ Pa} = \boxed{2.03 \times 10^5 \text{ Pa}} \quad (66)$$

## 9 Problem 19

(a) Compute the rms speed of a nitrogen molecule at 20.0°C. The molar mass of nitrogen molecules ( $N_2$ ) is given in Table 19-1. At what temperatures will the rms speed be (b) half that value and (c) twice that value?

### 9.1 Solution (a)

As stated in table 19-1, the molar mass of Nitrogen gas ( $N_2$ ) molecules is  $28.0 \times 10^{-3} \text{ kg/mol}$ . We can use that to find the molecule's rms (root-mean-square) speed.

$$T_K = T_{\text{°C}} + 273\text{K} = 293\text{K} \quad (67)$$

$$v_{\text{rms}} = \sqrt{\frac{3RT}{M}} = \sqrt{\frac{3 * 8.31 \text{ J/mol} \cdot \text{K} * 293 \text{ K}}{28.0 \times 10^{-3} \text{ kg/mol}}} \quad (68)$$

$$= \sqrt{\frac{7304.49}{28.0}} \times 10^3 \text{ m}^2/\text{s}^2 = \boxed{510.8 \text{ m/s}} \quad (69)$$

### 9.2 Solution (b)

We can use the same equation, this time solving for the temperature.

$$\frac{v_{\text{rms}}}{2} = \sqrt{\frac{3 * 8.31 \text{ J/mol} \cdot \text{K} * T}{28.0 \times 10^{-3} \text{ kg/mol}}} \quad (70)$$

$$\frac{3 * 8.31 \text{ J/mol} \cdot \text{K} * 293 \text{ K}}{28.0 \times 10^{-3} \text{ kg/mol}} * \frac{1}{4} = \frac{3 * 8.31 \text{ J/mol} \cdot \text{K} * T}{28.0 \times 10^{-3} \text{ kg/mol}} \quad (71)$$

$$T = \frac{293 \text{ K}}{4} = 73.25 \text{ K} = \boxed{-199.75 \text{ °C}} \quad (72)$$

### 9.3 Solution (c)

The same equation can be used.

$$2v_{\text{rms}} = \sqrt{\frac{3 * 8.31 \text{ J/mol} \cdot \text{K} * T}{28.0 \times 10^{-3} \text{ kg/mol}}} \quad (73)$$

$$\frac{3 * 8.31 \text{ J/mol} \cdot \text{K} * 293 \text{ K}}{28.0 \times 10^{-3} \text{ kg/mol}} * 4 = \frac{3 * 8.31 \text{ J/mol} \cdot \text{K} * T}{28.0 \times 10^{-3} \text{ kg/mol}} \quad (74)$$

$$T = 4 * 293 \text{ K} = 1172 \text{ K} = \boxed{899^\circ\text{C}} \quad (75)$$

## 10 Problem 23

A beam of hydrogen molecules ( $H_2$ ) is directed toward a wall, at an angle of  $55^\circ$  with the normal to the wall. Each molecule in the beam has a speed of 1.0 km/s and a mass of  $3.3 \times 10^{-24}$  g. The beam strikes the wall over an area of  $2.0 \text{ cm}^2$ , at the rate of  $10^{23}$  molecules per second. What is the beam's pressure on the wall?

### 10.1 Solution

This solution may be difficult to understand because  $p$  represents both pressure and momentum. We need more letters. Maybe we could look at some language with a lot of different letters/characters. My Chinese teacher says their language has about 1500. We have a formula for pressure exerted by an ideal gas.

$$p = \frac{nMv_{\text{rms}}^2}{3V} = \frac{F}{A} \quad (76)$$

We assume that any collision of a molecule with a wall is fully elastic, so we can find a formula for the change in the vertical momentum. This would only be for a single particle and we will only be considering the horizontal velocity.

$$v_i = v_{\text{rms}} * \sin(55^\circ) \quad (77)$$

$$v_f = -v_i \quad (78)$$

$$\Delta p = p_f - p_i = mv_f - mv_i \quad (79)$$

$$= 2mv_f = -2m * v_{\text{rms}} * \cos(55^\circ) \quad (80)$$

This can be multiplied by the rate at which the molecules hit the wall to find the total force, which is itself constant.

$$F = p * \frac{dN}{dt} = -2m * v_{\text{rms}} * \cos(55^\circ) * 10^{23} \text{ s}^{-1} \quad (81)$$

We can apply this to our equation for the pressure. That would be the final equation that we can plug values into.

$$p = \frac{F}{A} = -\frac{2m * v_{\text{rms}} * \cos(55^\circ) * 10^{23} \text{ s}^{-1}}{A} \quad (82)$$

$$= -\frac{2 * 3.3 \times 10^{-27} \text{ kg} * 1000 \text{ m/s} * \cos(55^\circ) * 10^{23} \text{ s}^{-1}}{2.0 \times 10^{-4} \text{ m}^2} \quad (83)$$

$$= -1893 \text{ Pa} \quad (84)$$

Since pressure is always positive, we can finalize this by taking the absolute value to get our final answer. Said answer would be  $\boxed{1893 \text{ Pa}}$ .

## 11 Problem 25

Determine the average value of the translational kinetic energy of the gas's molecules of an ideal gas at temperatures (a) 0.00°C and (b) 100°C. What is the translational kinetic energy per mole of an ideal gas at (c) 0.00°C and (d) 100°C?

### 11.1 Solution (a)

The average translational kinetic energy is determined by a simple formula, which we can use here.

$$K = \frac{3}{2}kT = \frac{3}{2} * 1.38 \times 10^{-23} \text{ J/K} * 273 \text{ K} = \boxed{565.11 \times 10^{-23} \text{ J}} \quad (85)$$

### 11.2 Solution (b)

Same strategy.

$$K = \frac{3}{2}kT = \frac{3}{2} * 1.38 \times 10^{-23} \text{ J/K} * 373 \text{ K} = \boxed{772.11 \times 10^{-23} \text{ J}} \quad (86)$$

### 11.3 Solution (c)

The kinetic energy of a system (in this case a mole of atoms) would be the sum of the kinetic energy of its parts. Each mole contains Avogadro's number ( $N_A$ ) atoms, each with a kinetic energy of its own, and the average translational kinetic energy of each atom is defined by the formula  $K = \frac{3}{2}kT$ . To get the total kinetic energy in one mole of an ideal gas, we can multiply the kinetic energy in each molecule by the number of molecules in a mole.

$$K_{\text{per mole}} = \frac{3}{2}kT \times N_A \quad (87)$$

Since  $k = \frac{R}{N_A}$ , we also know that  $R = k * N_A$ , which we can use.

$$K_{\text{per mole}} = \frac{3}{2}RT = \frac{3}{2} * 8.31 \text{ J/mol} \cdot \text{K} * 273 \text{ K} \quad (88)$$

$$= 3402.945 \text{ J/mol} = \boxed{3403 \text{ J/mol}} \quad (89)$$



## 11.4 Solution (d)

Same strategy.

$$K_{\text{per mole}} = \frac{3}{2}RT = \frac{3}{2} * 8.31 \text{ J/mol} \cdot \text{K} * 373 \text{ K} \quad (90)$$

$$= 4649.445 \text{ J/mol} = \boxed{4649 \text{ J/mol}} \quad (91)$$

## 12 Problem 27

Water standing in the open at 32.0°C evaporates because of the escape of some of the surface molecules. The heat of vaporization (539 cal/g) is approximately equal to  $\varepsilon n$ , where  $\varepsilon$  is the average energy of the escaping molecules and  $n$  is the number of molecules per gram. (a) Find  $\varepsilon$ . (b) What is the ratio of  $\varepsilon$  to the average kinetic energy of H<sub>2</sub>O molecules, assuming the latter is related to temperature in the same way as it is for gases?

### 12.1 Solution (a)

We can assume that  $\varepsilon n = 539 \text{ cal/g}$ . The number of molecules per gram can be calculated by using the molar mass of the water vapor (MM = 18.0 × 10<sup>-3</sup> kg/mol).

$$n = \frac{1}{\text{MM}} \times N_A = \frac{1 \text{ mol}}{18.0 \text{ g}} * 6.022 \times 10^{23} = 3.3456 \times 10^{22} \text{ g}^{-1} \quad (92)$$

Using this, we can find the value of  $\varepsilon$ .

$$\varepsilon n = 539 \text{ cal/g} \quad (93)$$

$$\varepsilon = \frac{539 \text{ cal/g}}{3.3456 \times 10^{22} \text{ g}^{-1}} = \boxed{1.611 \times 10^{-20} \text{ cal/molecule}} \quad (94)$$

### 12.2 Solution (b)

The average kinetic energy is calculatable by a formula.

$$K = \frac{3}{2} kT \quad (95)$$

This can be used in a ratio, bearing in mind that 32.0°C = 32.0°C + 273 K = 305 K.

$$\frac{\varepsilon}{K} = \frac{\varepsilon}{\frac{3}{2} kT} = \frac{1.611 \times 10^{-20} \text{ cal/molecule}}{\frac{3}{2} * 1.38 \times 10^{-23} \text{ J/K} * 305 \text{ K}} \quad (96)$$

$$= \frac{6.740 \times 10^{-20} \text{ J/molecule}}{\frac{3}{2} * 1.38 \times 10^{-23} \text{ J/K} * 305 \text{ K}} \quad (97)$$

$$= \boxed{10.67} \quad (98)$$

## 13 Problem 31

In a certain particle accelerator, protons travel around a circular path of diameter 23.0 m in an evacuated chamber, whose residual gas is at 295 K and  $1.00 \times 10^{-6}$  torr pressure. (a) Calculate the number of gas molecules per cubic centimeter at this pressure. (b) What is the mean free path of the gas molecules if the molecular diameter is  $2.00 \times 10^{-8}$  cm?

### 13.1 Solution (a)

The ideal gas law would be our friend, and we will assume this to be an ideal gas, where we seek to find the number of molecules per volume, better expressed as  $\frac{N}{V}$ . Also don't forget to convert pressure from torr to pascals.

$$p = 1.00 \times 10^{-6} \text{ torr} = 1.00 \times 10^{-6} \text{ mm Hg} \quad (99)$$

$$= 0.100 \times 10^{-6} \text{ cm Hg} = 133.30 \times 10^{-6} \text{ Pa} \quad (100)$$

$$pV = NkT \quad (101)$$

$$\frac{N}{V} = \frac{p}{kT} = \frac{133.30 \times 10^{-6} \text{ Pa}}{1.38 \times 10^{-23} \text{ J/K} * 295 \text{ K}} \quad (102)$$

$$= 3.274 \times 10^{16} \text{ molecule/m}^3 \times \frac{10 \times 10^{-6} \text{ m}^3}{1 \text{ cm}^3} \quad (103)$$

$$= \boxed{3.274 \times 10^{10} \text{ molecule/cm}^3} \quad (104)$$

### 13.2 Solution (b)

There's an equation for this. It's kind of ugly to be completely honest, but it's usable.

$$\lambda = \frac{1}{\sqrt{2}\pi d^2 N/V} = \frac{1}{\sqrt{2}\pi (2.00 \times 10^{-8} \text{ cm})^2 * 3.274 \times 10^{10} \text{ cm}^{-3}} \quad (105)$$

$$= 17187 \text{ cm} = \boxed{171.87 \text{ m}} \quad (106)$$

## 14 Problem 35

Ten particles are moving with the following speeds: four at 200m/s, two at 500m/s, and four at 600m/s. Calculate their (a) average and (b) rms speeds. (c) Is  $v_{rms} > v_{avg}$ ?

### 14.1 Solution (a)

We'll just do a simple average.

$$v_{avg} = \frac{\sum_{i=1}^n v_i}{n} = \frac{4 * 200 + 2 * 500 + 4 * 600}{10} \quad (107)$$

$$= 4 * 20 + 2 * 50 + 4 * 60 \quad (108)$$

$$= 80 + 100 + 240 = \boxed{420\text{m/s}} \quad (109)$$

### 14.2 Solution (b)

#### 14.2.1 Attempt 1 (Incorrect)

There's a formula for  $v_{rms}$ .

$$v_{rms} = \sqrt{\frac{3RT}{M}} \quad (110)$$

$$v_{rms}^2 = \frac{3RT}{M} \quad (111)$$

I don't know about you, but it reminds me of a formula for the average velocity.

$$v_{avg} = \sqrt{\frac{8RT}{\pi M}} \quad (112)$$

$$v_{avg}^2 = \frac{8RT}{\pi M} = \frac{8}{3\pi} \cdot \frac{3RT}{M} \quad (113)$$

$$\frac{3RT}{M} = \frac{3\pi}{8} v_{avg}^2 \quad (114)$$

We could use the transitive property on this.

$$v_{rms}^2 = \frac{3\pi}{8} v_{avg}^2 = \frac{3\pi}{8} * (420 \text{ m/s})^2 = 22050 * 3\pi = 66150\pi \text{ m}^2/\text{s}^2 \quad (115)$$

$$v_{rms} = \sqrt{66150\pi \text{ m}^2/\text{s}^2} = 455.9 \text{ m/s} \quad (116)$$

The error made here was assuming the gas to be an ideal gas, which is required for some of our equations.

### 14.2.2 Attempt 2

Just brute force it.

$$v_{\text{rms}}^2 = \frac{4 * 200^2 + 2 * 500^2 + 4 * 600^2}{10} \quad (117)$$

$$= \frac{4 * 40000 + 2 * 250000 + 4 * 360000}{10} \quad (118)$$

$$= \frac{160000 + 500000 + 1440000}{10} \quad (119)$$

$$= \frac{2100000 \text{ m}^2/\text{s}^2}{10} = 210000 \text{ m}^2/\text{s}^2 \quad (120)$$

$$v_{\text{rms}} = \boxed{458 \text{ m/s}} \quad (121)$$

### 14.3 Solution (c)

It's just by a difference of 38 m/s, but  $\boxed{yes}$ .

## 15 Problem 37

### 15.1 Solution

## 16 Problem 39

### 16.1 Solution

## 17 Problem 43

The temperature of 3.00 mol of an ideal diatomic gas is increased by  $40.0\text{ }^{\circ}\text{C}$  without the pressure of the gas changing. The molecules in the gas rotate but do not oscillate. (a) How much energy is transferred to the gas as heat? (b) What is the change in the internal energy of the gas? (c) How much work is done by the gas? (d) By how much does the rotational kinetic energy of the gas increase?

### 17.1 Solution



## 18 Problem 45

### 18.1 Solution

## 19 Problem 47

### 19.1 Solution

## 20 Problem 51

When 1.0 mol of oxygen ( $O_2$ ) gas is heated at constant pressure starting at  $0^\circ\text{C}$ , how much energy must be added to the gas as heat to double its volume? The molecules rotate but do not oscillate.

### 20.1 Solution

## 21 Problem 55

### 21.1 Solution

## 22 Problem 57

### 22.1 Solution

## **23 Problem 59**

### **23.1 Solution**

## **24 Problem 63**

### **24.1 Solution**

## 25 Problem 69

### 25.1 Solution



## 26 Problem 75

### 26.1 Solution

## 27 Problem 77

### 27.1 Solution

# Contents

<b>1</b>	<b>Problem 1</b>	<b>1</b>
1.1	Solution . . . . .	1
<b>2</b>	<b>Problem 3</b>	<b>2</b>
2.1	Solution (a) . . . . .	2
2.2	Solution (b) . . . . .	2
<b>3</b>	<b>Problem 5</b>	<b>3</b>
3.1	Solution . . . . .	3
<b>4</b>	<b>Problem 7</b>	<b>4</b>
4.1	Solution (a) . . . . .	4
4.2	Solution (b) . . . . .	4
<b>5</b>	<b>Problem 9</b>	<b>5</b>
5.1	Solution . . . . .	5
<b>6</b>	<b>Problem 11</b>	<b>6</b>
6.1	Solution . . . . .	6
<b>7</b>	<b>Problem 13</b>	<b>8</b>
7.1	Solution (a) . . . . .	8
7.2	Solution (b) . . . . .	8
7.3	Solution (c) . . . . .	9
7.4	Solution (d) . . . . .	9
<b>8</b>	<b>Problem 17</b>	<b>10</b>
8.1	Solution . . . . .	10
<b>9</b>	<b>Problem 19</b>	<b>12</b>
9.1	Solution (a) . . . . .	12
9.2	Solution (b) . . . . .	12
9.3	Solution (c) . . . . .	13
<b>10</b>	<b>Problem 23</b>	<b>14</b>
10.1	Solution . . . . .	14

<b>11 Problem 25</b>	<b>16</b>
11.1 Solution (a) . . . . .	16
11.2 Solution (b) . . . . .	16
11.3 Solution (c) . . . . .	16
11.4 Solution (d) . . . . .	17
<b>12 Problem 27</b>	<b>18</b>
12.1 Solution (a) . . . . .	18
12.2 Solution (b) . . . . .	18
<b>13 Problem 31</b>	<b>19</b>
13.1 Solution (a) . . . . .	19
13.2 Solution (b) . . . . .	19
<b>14 Problem 35</b>	<b>20</b>
14.1 Solution (a) . . . . .	20
14.2 Solution (b) . . . . .	20
14.2.1 Attempt 1 (Incorrect) . . . . .	20
14.2.2 Attempt 2 . . . . .	21
14.3 Solution (c) . . . . .	21
<b>15 Problem 37</b>	<b>22</b>
15.1 Solution . . . . .	22
<b>16 Problem 39</b>	<b>23</b>
16.1 Solution . . . . .	23
<b>17 Problem 43</b>	<b>24</b>
17.1 Solution . . . . .	24
<b>18 Problem 45</b>	<b>25</b>
18.1 Solution . . . . .	25
<b>19 Problem 47</b>	<b>26</b>
19.1 Solution . . . . .	26
<b>20 Problem 51</b>	<b>27</b>
20.1 Solution . . . . .	27

<b>21 Problem 55</b>	<b>28</b>
21.1 Solution . . . . .	28
<b>22 Problem 57</b>	<b>29</b>
22.1 Solution . . . . .	29
<b>23 Problem 59</b>	<b>30</b>
23.1 Solution . . . . .	30
<b>24 Problem 63</b>	<b>31</b>
24.1 Solution . . . . .	31
<b>25 Problem 69</b>	<b>32</b>
25.1 Solution . . . . .	32
<b>26 Problem 75</b>	<b>33</b>
26.1 Solution . . . . .	33
<b>27 Problem 77</b>	<b>34</b>
27.1 Solution . . . . .	34