

# Proof: Accelerating Approximate Aggregation Queries with Expensive Predicates

Daniel Kang<sup>\*,†</sup>, John Guibas<sup>\*,†</sup>, Peter Bailis<sup>†</sup>, Tatsunori Hashimoto<sup>†</sup>, Yi Sun<sup>‡</sup>, Matei Zaharia<sup>†</sup>  
Stanford University<sup>†</sup>, University of Chicago<sup>‡</sup>

February 28, 2021

## Abstract

Given a dataset  $\mathcal{D}$ , we are interested in computing the mean of a subset of  $\mathcal{D}$  which matches a predicate. ABAE leverages stratified sampling and proxy models to efficiently compute this statistic given a sampling budget  $N$ . In this document, we theoretically analyze ABAE and show that the MSE of the estimate decays at rate  $O(N_1^{-1} + N_2^{-1} + N_1^{1/2}N_2^{-3/2})$ , where  $N = K \cdot N_1 + N_2$  for some integer constant  $K$  and  $K \cdot N_1$  and  $N_2$  represent the number of samples used in Stage 1 and Stage 2 of ABAE respectively. Hence, if a constant fraction of the total sample budget  $N$  is allocated to each stage, we will achieve a mean squared error of  $O(N^{-1})$  which matches the rate of mean squared error of the optimal stratified sampling algorithm given a priori knowledge of the predicate positive rate and standard deviation per stratum.

Symbol	Description
$\mathcal{D}$	dataset of records
$O(x)$	predicate function
$f(x)$	statistic function
$K$	number of strata
$S$	set of $K$ equal sized stratum formed from $\mathcal{D}$
$X_{k,i}$	subgaussian r.v. in $S_k$ and the $i$ 'th i.i.d predicate-matching sample
$p_k$	predicate positive rate in stratum $k$
$\mu_k$	$E[X_k]$
$\sigma_k^2$	$Var[X_k]$
$C_k^{(\mu)}$	upper bound on $\mu_k$
$C_k^{(\sigma)}$	upper bound on $\sigma_k$
$p_{\text{all}}$	$\sum_i^K p_k$
$w_k$	$p_k/p_{\text{all}}$
$\mu_{\text{all}}$	$E[X]$ , also given by $\sum_{k=1}^K w_k \mu_k$
$N_1$	sampling budget we allocate per stratum during 1
$N_2$	sampling we allocate during 2 in total
$N$	total sampling budget ( $K \cdot N_1 + N_2 = N$ )
$T_k$	fraction of our $N_2$ samples we allocate to a stratum during 2
$B_k^{(1)}$	number of predicate-matching samples we get in Stage 1
$B_k^{(2)}$	number of predicate-matching samples we get in Stage 2
$p_*$	cutoff value used when we split cases where $p_k$ is small and large
$F_k$	we say that $B_k^{(2)} \geq F_k$ with high probability

Table 1: Summary of notation.

# 1 Setup

Given a dataset  $\mathcal{D} = \{x_1, \dots, x_{|\mathcal{D}|}\}$ , a statistic  $f : \mathcal{D} \rightarrow \mathbb{R}$ , and a predicate  $O : \mathcal{D} \rightarrow \{0, 1\}$ , we want to estimate  $\mathbb{E}_{x \sim \mathcal{D}}[f(x) | O(x) = 1] = \mu_{\text{all}}$  given a sampling budget  $N$ , which limits the number of times the predicate  $O(x)$  can be computed. We measure the quality of an estimator for  $\mu_{\text{all}}$  by measuring the mean squared error (MSE) or  $\mathbb{E}[(\hat{\mu}_{\text{all}} - \mu_{\text{all}})^2]$ .

To estimate this mean efficiently, we use stratified sampling. In stratified sampling, the data is split into  $K$  strata or disjoint subsets of the dataset. In other words, given a dataset  $\mathcal{D}$ , we stratify it into equal-sized strata  $S_1, \dots, S_K$  such that they are disjoint and span the entirety of  $\mathcal{D}$ . Given these strata, denote  $\mu_k, p_k, \sigma_k$  as the per-stratum mean of records which match the predicate, the per-stratum predicate positive rate, and the per-stratum standard deviation. Then,  $\mu_{\text{all}} = \sum_{k=1}^K \mu_k p_k / \sum_{k=1}^K p_k$ .

We make the following assumptions on our dataset. First, we assume that the distribution of the data we sample from is subgaussian. This is so that we can obtain concentration inequalities on estimates of variables such as  $\mu_k$  and  $\sigma_k$ . We represent the samples from a stratum by a subgaussian random variable  $X_{k,i}$  where  $k$  denotes the stratum and  $i$  denotes the  $i$ 'th independent and identically distributed (i.i.d.) sample. Second, we assume that  $\mathbb{E}[|X_{k,i}|] \leq C_k^{(\mu)}$  and  $\text{Var}[X_{k,i}] \leq C_k^{(\sigma^2)}$ . Lastly, we assume that  $p_{\text{all}} = \sum_{k=1}^K p_k \geq C^{(p_{\text{all}})}$ .

---

## Algorithm 1 Pseudocode for ABAE.

---

```

1: function ABAE( $\mathcal{D}, K, N_1, N_2$ )
2:    $S_1, \dots, S_K \leftarrow \text{Stratify}(\mathcal{D}, K)$ 
3:   for each  $k$  in  $[1, \dots, K]$  do
4:      $R_k^{(1)} \leftarrow \text{Sample}(S_k, N_1)$ 
5:      $X_k^{(1)} \leftarrow \{f(x) \mid x \in R_k^{(1)}, O(x) = 1\}$ 
6:      $B_k^{(1)} \leftarrow |X_k^{(1)}|$ 
7:      $\hat{\mu}_k \leftarrow \sum_{i=1}^{B_k^{(1)}} X_{k,i}^{(1)} / B_k^{(1)}$  if  $B_k^{(1)} > 0$  else 0
8:      $\hat{p}_k \leftarrow B_k^{(1)} / |R_k^{(1)}|$ 
9:      $\hat{\sigma}_k^2 \leftarrow \sum_{i=1}^{B_k^{(1)}} (X_{k,i}^{(1)} - \hat{\mu}_k)^2 / (B_k^{(1)} - 1)$  if  $B_k^{(1)} > 1$  else 0
10:  end for
11:  for each  $k$  in  $[1, \dots, K]$  do
12:     $\hat{T}_k \leftarrow \sqrt{\hat{p}_k \hat{\sigma}_k} / \sum_{i=1}^K \sqrt{\hat{p}_i \hat{\sigma}_i}$ 
13:     $R_k^{(2)} \leftarrow \text{Sample}(S_k, \lceil N_2 \hat{T}_k \rceil)$ 
14:     $R_k^{(2)} \leftarrow R_k^{(1)} + R_k^{(2)}$  if SampleReuse else  $R_k^{(2)}$ 
15:     $X_k^{(2)} \leftarrow \{f(x) \mid x \in R_k^{(2)}, O(x) = 1\}$ 
16:     $B_k^{(2)} \leftarrow |X_k^{(2)}|$ 
17:     $\hat{p}_k \leftarrow B_k^{(2)} / |R_k^{(2)}|$  if SampleReuse else  $\hat{p}_k$ 
18:     $\hat{\mu}_k \leftarrow \sum_{i=1}^{B_k^{(2)}} X_{k,i}^{(2)} / B_k^{(2)}$  if  $B_k^{(2)} > 0$  else 0
19:  end for
20:  return  $\sum_{k=1}^K \hat{p}_k \hat{\mu}_k / \sum_{k=1}^K \hat{p}_k$ 
21: end function

```

---

For the purpose of this proof, *SampleReuse* is *False*. We show that *SampleReuse* set to *True* does not change the error of ABAE asymptotically via Corollary X. Furthermore, note that on line 13 we round up the number of allocated samples  $N_2 \hat{T}_k$ . This rounding ensures that our bounds hold; however, this means that we may use up to  $N_2 + K$  samples. Hence, proof statements involving  $N$  are only accurate up to  $K$  samples.

## 2 Algorithm

Refer to Algorithm 1 for pseudocode of ABAE and Table 1 for a description of notation used throughout the document. ABAE creates the strata by splitting the dataset evenly between  $K$  strata by ordering the records by proxy score. ABAE will then perform a two-stage sampling procedure. ABAE first estimates  $p_k$  and  $\sigma_k$ , which are the key quantities for determining optimal allocation by sampling  $N_1$  samples from each of the  $K$  strata. Then, ABAE will allocate the remaining samples proportional to our estimates of the optimal allocation. ABAE will return the estimate  $\sum_k \hat{p}_k \hat{\mu} / \sum_k \hat{p}_k$  as the approximate answer.

## 3 Analysis (Optimal Stratified Sampling Allocation)

In this section, we assume that the true values of  $p_k$  and  $\sigma_k$  are known, and derive the optimal allocation according to knowledge of these quantities. We also assume that the sample selection process is deterministic and returns a number of draws equal to the expected number of draws which satisfy the predicate. We also derive an upper bound on the MSE based on this optimal allocation. To avoid fractional samples, we round up samples as appropriate.

Although the true  $p_k$  and  $\sigma_k$  are unknown in this setting, our algorithm is based on estimating them in Stage 1 and then allocating samples in Stage 2 based on those estimates. We will then later show in a future section that the estimates made in Stage 1 are tight enough with high probability to achieve matching asymptotic rates of error with respect to the optimal allocation.

**Proposition 3.1.** *Let  $T_k$  represent the fraction of our sampling budget we want to allocate to stratum  $S_k$ . Suppose that  $p_k$  and  $\sigma_k$  are known,  $\mathbb{E}[\hat{\mu}_k] = \mu_k$ , and that we get  $B_k = \lceil p_k T_k N \rceil$  samples for all  $k$ . Then, the choice of  $T_k$  such that  $T_k > 0$  and  $\sum_{k=1}^K T_k = 1$  which minimizes the loss function  $\mathcal{L}(T_1, \dots, T_K) = \sum_{k=1}^K \frac{w_k^2 \sigma_k^2}{p_k T_k N}$  of the unbiased estimator  $\hat{\mu}_{all} = \sum_{k=1}^K w_k \hat{\mu}_k$  is:*

$$T_k^* = \frac{\sqrt{p_k} \sigma_k}{\sum_{i=1}^K \sqrt{p_i} \sigma_i}. \quad (1)$$

In addition, the choice  $T_k = T_k^*$  will result in the following upper bound on the MSE:

$$E^* = \mathbb{E}[(\hat{\mu}_{all} - \mu_{all})^2 | B_k = \lceil p_k T_k^* N \rceil] \leq \frac{1}{N p_{all}^2} \cdot \left( \sum_{k=1}^K \sqrt{p_k} \sigma_k \right)^2. \quad (2)$$

*Proof.* We can decompose the MSE into bias and variance. Note that the bias is zero due to the strata-specific estimates of the mean being unbiased.

$$\mathbb{E}[(\hat{\mu}_{all} - \mu_{all})^2 | B_k = \lceil p_k T_k N \rceil] = \mathbb{E}[\hat{\mu}_{all} - \mu_{all}]^2 + \text{Var}[\hat{\mu}_{all}] = \text{Var} \left[ \sum_{k=1}^K w_k \hat{\mu}_k \right] \leq \mathcal{L}(T_1, \dots, T_K) = \sum_{k=1}^K \frac{w_k^2 \sigma_k^2}{p_k T_k N}. \quad (3)$$

We will use Lagrange multipliers to minimize the the loss function  $\mathcal{L}$  with respect to  $T_1, \dots, T_K$  under the constraint  $g$  which enforces that  $\sum_{k=1}^K T_k = 1$ .

$$\mathcal{L}(T_1, \dots, T_K) = \sum_{k=1}^K \frac{w_k^2 \sigma_k^2}{p_k T_k N}, \quad g(T_1, \dots, T_K) = \sum_{k=1}^K T_k - 1. \quad (4)$$

$$\frac{\partial \mathcal{L}}{\partial T_k} + \lambda \frac{\partial g}{\partial T_k} = 0 \longrightarrow \frac{w_k^2 \sigma_k^2}{p_k T_k^2 N} = \lambda \quad (5)$$

$$T_k = \frac{w_k \sigma_k}{\sqrt{p_k \lambda N}} \quad (6)$$

$$\sum_{k=1}^K T_k = \sum_{k=1}^K \frac{w_k \sigma_k}{\sqrt{p_k \lambda N}} = 1 \quad (7)$$

$$\sum_{k=1}^K \frac{w_k \sigma_k}{\sqrt{p_k N}} = \sqrt{\lambda} \quad (8)$$

$$\left( \sum_{k=1}^K \frac{w_k \sigma_k}{\sqrt{p_k N}} \right)^2 = \lambda. \quad (9)$$

Thus, the optimal choices of  $T_1, \dots, T_K$  to minimize  $\mathcal{L}$  under the constraint  $g$  is:

$$T_k^* = \frac{w_k \sigma_k}{\sqrt{p_k \lambda N}} \quad (10)$$

$$= \frac{w_k \sigma_k}{\sqrt{p_k N} \left( \sum_{i=1}^K \frac{w_i \sigma_i}{\sqrt{p_i N}} \right)} \quad (11)$$

$$= \frac{\sqrt{p_k} \sigma_k}{\sum_{i=1}^K \sqrt{p_i} \sigma_i}. \quad (12)$$

We will now derive an upper bound on the MSE given  $T_k = T_k^*$ . To deal with fractional samples, we round up  $p_k T_k^* N$  or the number of draws each strata gets, which means that the bound derived below is still valid.

$$\mathbb{E}[(\hat{\mu}_{\text{all}} - \mu_{\text{all}})^2 | B_k = \lceil p_k T_k^* N \rceil] \leq \sum_{k=1}^K \frac{w_k^2 \sigma_k^2}{p_k T_k^* N} \quad (13)$$

$$= \sum_{k=1}^K \frac{w_k^2 \sigma_k^2}{p_k \frac{\sqrt{p_k} \sigma_k}{\sum_{i=1}^K \sqrt{p_i} \sigma_i} N} \quad (14)$$

$$= \frac{1}{p_{\text{all}}^2 N} \cdot \sum_{k=1}^K \frac{p_k^2 \sigma_k^2}{p_k \frac{\sqrt{p_k} \sigma_k}{\sum_{i=1}^K \sqrt{p_i} \sigma_i}} \quad (15)$$

$$= \frac{1}{p_{\text{all}}^2 N} \cdot \sum_{k=1}^K \frac{\sqrt{p_k} \sigma_k}{\frac{1}{\sum_{i=1}^K \sqrt{p_i} \sigma_i}} \quad (16)$$

$$= \frac{1}{p_{\text{all}}^2 N} \cdot \left( \sum_{k=1}^K \sqrt{p_k} \sigma_k \right)^2 \quad \square$$

## 4 Analysis

Our goal is to prove that with arbitrarily small failure probability  $\delta$  over Stage 1, the expectation of the mean squared error of ABAE over Stage 2 is

$$\mathbb{E}[(\hat{\mu}_{\text{all}} - \mu_{\text{all}})^2] \leq O(N_1^{-1}) + O(N_2^{-1}) + O(N_1^{1/2}N_2^{-3/2}). \quad (17)$$

Our proof strategy is outlined below.

1. Obtain high probability bounds over all the random variables in Stage 1. Namely,  $p_k$ ,  $w_k$ ,  $\sigma_k$ ,  $B_k^{(1)}$ , and  $T_k$ .
2. Assuming the worst-case values described by these high probability bounds, we will upper bound the mean squared error of our algorithm in expectation over Stage 2.

### 4.1 Big-O Definition

To clarify our usage of Big-O, we will formally define it as:

$$T(\dots) = O(f(\dots)) \iff \exists C, N_0 \text{ s.t. } \forall N \geq N_0, 0 \leq T(\dots) \leq C \cdot f(\dots). \quad (18)$$

We use Big-O notation with respect to  $N_1$ ,  $N_2$ ,  $p_k$ , and  $\delta$ . Hence,  $C$  and  $N_0$  must be independent of those quantities.

### 4.2 Stage 1

We will obtain high probability bounds on all the random variables in Stage 1. Namely,  $p_k$ ,  $w_k$ ,  $\sigma_k$ ,  $B_k^{(1)}$ , and  $T_k$ .

**Lemma 4.1** ( $\hat{p}_k$  Upper Bound). *For small  $\delta > 0$ , the following holds for all  $k$  simultaneously,*

$$\sqrt{\hat{p}_k} \leq \sqrt{p_k + \sqrt{\frac{2 \ln(1/\delta) p_k}{N_1}}} = \sqrt{p_k} \sqrt{1 + O\left(\sqrt{\frac{\ln(1/\delta)}{p_k N_1}}\right)}. \quad (19)$$

*Proof.* We apply Chung and Lu [2002] reproduced as Lemma 5.1 to upper bound  $\hat{p}_k$ . We set  $Y_i \sim \text{Bernoulli}(p_k)$  and  $a_i = \frac{1}{N_1}$  for all  $i \in [N_1]$ . Thus,  $\nu = \sum_{i=1}^{N_1} a_i^2 p_i = \frac{p_k}{N_1}$  and  $Y = \sum_{i=1}^{N_1} a_i Y_i = \hat{p}_k$ . Then, according to the Lemma,

$$P(\hat{p}_k \geq p_k + \lambda) \leq \exp\left(\frac{-\lambda^2}{2\left(\frac{p_k}{N_1} + \frac{\lambda}{3N_1}\right)}\right). \quad (20)$$

We will set  $\delta = \exp\left(-\lambda^2 / \left(2\left(\frac{p_k}{N_1} + \frac{\lambda}{3N_1}\right)\right)\right)$  and  $\lambda = \left(\sqrt{18N_1 p_k \ln(1/\delta)} + \ln(1/\delta)\right) / (3N_1)$ . Thus, with probability at least  $1 - \delta$ ,

$$\hat{p}_k \leq p_k + \frac{\sqrt{18N_1 p_k \ln(1/\delta)} + \ln(1/\delta)}{3N_1} \quad (21)$$

$$\leq p_k + \frac{\sqrt{18N_1 p_k \ln(1/\delta)}}{3N_1} = p_k + \sqrt{\frac{2 \ln(1/\delta) p_k}{N_1}}. \quad (22)$$

With probability at least  $1 - \delta$ ,

$$\sqrt{\hat{p}_k} \leq \sqrt{p_k + \sqrt{\frac{2 \ln(1/\delta) p_k}{N_1}}} \leq \sqrt{p_k} \cdot \sqrt{1 + \sqrt{\frac{2 \ln(1/\delta)}{p_k N_1}}} = \sqrt{p_k} \sqrt{1 + O\left(\sqrt{\frac{\ln(1/\delta)}{p_k N_1}}\right)}. \quad (23)$$

We union bound across all strata  $S_1, \dots, S_k$ . As a result, the bound above holds for all  $k$  simultaneously with probability at least  $1 - K\delta$ .  $\square$

**Lemma 4.2** ( $\hat{p}_k$  Lower Bound). *For small  $\delta > 0$ , the following holds for all  $k$  simultaneously,*

$$\sqrt{\hat{p}_k} \geq \sqrt{p_k - \sqrt{\frac{2 \ln(1/\delta) p_k}{N_1}}} = \sqrt{p_k} \sqrt{1 - O\left(\sqrt{\frac{\ln(1/\delta)}{p_k N_1}}\right)}. \quad (24)$$

*Proof.* We apply Chung and Lu [2002] reproduced as Lemma 5.1 to lower bound  $\hat{p}_k$ . We set  $X_i \sim \text{Bernoulli}(p_k)$  and  $a_i = \frac{1}{N_1}$  for all  $i \in [N_1]$ . Thus,  $\nu = \sum_{i=1}^{N_1} a_i^2 p_i = \frac{p_k}{N_1}$  and  $X = \sum_{i=1}^{N_1} a_i X_i = \hat{p}_k$ . Then, according to the Lemma:

$$P(\hat{p}_k < p_k - \lambda) \leq \exp\left(\frac{-\lambda^2}{2\left(\frac{p}{N_1}\right)}\right). \quad (25)$$

We will set  $\delta = \exp\left(\frac{-\lambda^2}{2\left(\frac{p}{N_1}\right)}\right)$  and  $\lambda = \frac{\sqrt{p}}{\sqrt{N_1}} \cdot \sqrt{2 \ln(1/\delta)}$ . Hence, with probability at least  $1 - \delta$  (where  $\delta$  is probability of failure):

$$\hat{p}_k \geq p_k - \frac{\sqrt{p_k}}{\sqrt{N_1}} \cdot \sqrt{2 \ln(1/\delta)} = p_k - \sqrt{\frac{2 \ln(1/\delta) p_k}{N_1}}. \quad (26)$$

In other words, with probability at least  $1 - \delta$ ,

$$\sqrt{\hat{p}_k} \geq \sqrt{p_k - \sqrt{\frac{2 \ln(1/\delta) p_k}{N_1}}} \geq \sqrt{p_k} \sqrt{1 - \sqrt{\frac{2 \ln(1/\delta)}{p_k N_1}}} \quad (27)$$

$$= \sqrt{p_k} \sqrt{1 - O\left(\sqrt{\frac{\ln(1/\delta)}{p_k N_1}}\right)}. \quad (28)$$

We then apply the union bound, so the above holds for all  $k$  simultaneously with probability at least  $1 - K\delta$ .  $\square$

**Lemma 4.3** ( $\hat{w}_k$  Upper and Lower Bounds). *For small  $\delta > 0$ , the bounds of Lemma 4.1 and Lemma 4.2 hold and*

$$w_k - O\left(\frac{\ln(1/\delta) \sqrt{p_k}}{\sqrt{N_1}}\right) \leq \frac{p_k - \sqrt{\frac{2 \ln(1/\delta) p_k}{N_1}}}{p_{all} + \sum_{k=1}^K \sqrt{\frac{2 \ln(1/\delta) p_k}{N_1}}} \leq \hat{w}_k \leq \frac{p_k + \sqrt{\frac{2 \ln(1/\delta) p_k}{N_1}}}{p_{all} - \sum_{k=1}^K \sqrt{\frac{2 \ln(1/\delta) p_k}{N_1}}} \leq w_k + O\left(\frac{\ln(1/\delta) \sqrt{p_k}}{\sqrt{N_1}}\right). \quad (29)$$

*Proof.* Recall that  $\hat{w}_k = \hat{p}_k / \hat{p}_{\text{all}}$ , where  $\hat{p}_{\text{all}} = \sum_{k=1}^K \hat{p}_k$ . (29) holds on the event that the bounds of Lemma 4.1 and Lemma 4.2 hold for all  $k$  simultaneously, which occurs with probability at least  $1 - 2K\delta$  according to the union bound. For the remainder of the proof of this Lemma, we condition on this event and prove (29). Specifically, the condition is that for each  $k$ ,

$$p_k - \sqrt{\frac{2 \ln(1/\delta) p_k}{N_1}} \leq \hat{p}_k \leq p_k + \sqrt{\frac{2 \ln(1/\delta) p_k}{N_1}}. \quad (30)$$

Hence, by summing these inequalities across  $k$ , we have that with probability at least  $1 - 2K\delta$ ,

$$p_{\text{all}} - \sum_{k=1}^K \sqrt{\frac{2 \ln(1/\delta) p_k}{N_1}} \leq \hat{p}_{\text{all}} \leq p_{\text{all}} + \sum_{k=1}^K \sqrt{\frac{2 \ln(1/\delta) p_k}{N_1}}. \quad (31)$$

We will now maximize or minimize the numerator and denominator appropriately in  $\hat{w}_k = \hat{p}_k / \hat{p}_{\text{all}}$ .

$$\frac{p_k - \sqrt{\frac{2 \ln(1/\delta) p_k}{N_1}}}{p_{\text{all}} + \sum_{k=1}^K \sqrt{\frac{2 \ln(1/\delta) p_k}{N_1}}} \leq \hat{w}_k \leq \frac{p_k + \sqrt{\frac{2 \ln(1/\delta) p_k}{N_1}}}{p_{\text{all}} - \sum_{k=1}^K \sqrt{\frac{2 \ln(1/\delta) p_k}{N_1}}}. \quad (32)$$

We will now simplify this bound with Big O. Note that  $p_{\text{all}} \geq C^{(p_{\text{all}})}$ .

$$\hat{w}_k \leq \frac{p_k + \sqrt{\frac{2 \ln(1/\delta) p_k}{N_1}}}{p_{\text{all}} - \sum_{k=1}^K \sqrt{\frac{2 \ln(1/\delta) p_k}{N_1}}} \quad (33)$$

$$= \frac{\frac{1}{p_{\text{all}}}}{\frac{1}{p_{\text{all}}}} \cdot \frac{p_k}{p_{\text{all}} - \sum_{k=1}^K \sqrt{\frac{2 \ln(1/\delta) p_k}{N_1}}} + \frac{\frac{1}{p_{\text{all}}}}{\frac{1}{p_{\text{all}}}} \cdot \frac{\sqrt{\frac{2 \ln(1/\delta) p_k}{N_1}}}{p_{\text{all}} - \sum_{k=1}^K \sqrt{\frac{2 \ln(1/\delta) p_k}{N_1}}} \quad (34)$$

$$= \frac{w_k}{1 - \sum_{k=1}^K \sqrt{\frac{2 \ln(1/\delta) p_k}{p_{\text{all}}^2 N_1}}} + \frac{\sqrt{\frac{2 \ln(1/\delta) p_k}{p_{\text{all}}^2 N_1}}}{1 - \sum_{k=1}^K \sqrt{\frac{2 \ln(1/\delta) p_k}{p_{\text{all}}^2 N_1}}} \quad (35)$$

$$\leq \left( w_k + \sqrt{\frac{2 \ln(1/\delta) p_k}{p_{\text{all}}^2 N_1}} \right) \cdot \left( 1 + \sum_{k=1}^K \sqrt{\frac{2 \ln(1/\delta) p_k}{p_{\text{all}}^2 N_1}} \right). \quad (36)$$

$$\leq \left( w_k + \sqrt{\frac{2 \ln(1/\delta) p_k}{p_{\text{all}}^2 N_1}} \right) \cdot \left( 1 + O \left( \sqrt{\frac{\ln(1/\delta)}{N_1}} \right) \right) \quad (37)$$

$$\leq w_k + \sqrt{\frac{2 \ln(1/\delta) p_k}{p_{\text{all}}^2 N_1}} + O \left( \frac{p_k \sqrt{\ln(1/\delta)}}{\sqrt{N_1}} \right) + O \left( \frac{\ln(1/\delta) \sqrt{p_k}}{N_1} \right) \quad (38)$$

$$\leq w_k + O \left( \frac{\sqrt{\ln(1/\delta) p_k}}{\sqrt{N_1}} \right) + O \left( \frac{p_k \sqrt{\ln(1/\delta)}}{\sqrt{N_1}} \right) + O \left( \frac{\ln(1/\delta) \sqrt{p_k}}{N_1} \right) \quad (39)$$

$$\leq w_k + O \left( \frac{\ln(1/\delta) \sqrt{p_k}}{\sqrt{N_1}} \right). \quad (40)$$

We repeat these steps for the lower bound. Thus,

$$w_k - O\left(\frac{\ln(1/\delta)\sqrt{p_k}}{\sqrt{N_1}}\right) \leq \hat{w}_k \leq w_k + O\left(\frac{\ln(1/\delta)\sqrt{p_k}}{\sqrt{N_1}}\right). \quad (41)$$

□

**Lemma 4.4** ( $B_k^{(1)}$  Lower Bound). *For small  $\delta > 0$ , the following is true for all  $k$ ,*

$$B_k^{(1)} \geq p_k N_1 - \sqrt{2 \ln(1/\delta) p_k N_1}. \quad (42)$$

*Proof.* We apply Tarjan [2009] reproduced as Lemma 5.2 to lower bound  $B_k^{(1)}$ . Note that  $B_k^{(1)} \sim \text{Binomial}(N_1, p_k)$  and that equivalently  $B_k^{(1)} = \sum_{i=1}^{N_1} Y_i$  where  $Y_i \sim \text{Bernoulli}(p_k)$ . Hence,

$$P\left(B_k^{(1)} \leq (1 - \epsilon)(p_k N_1)\right) \leq \exp\left(\frac{-\epsilon^2 p_k N_1}{2}\right). \quad (43)$$

We set  $\delta = \exp\left(\frac{-\epsilon^2 p_k N_1}{2}\right)$  and  $\epsilon = \sqrt{\frac{2 \ln(1/\delta)}{p_k N_1}}$ . Thus, with probability at least  $1 - \delta$ :

$$B_k^{(1)} \geq p_k N_1 - \sqrt{2 \ln(1/\delta) p_k N_1}. \quad (44)$$

We then union bound across all  $k$ . As a result, with probability at least  $1 - K\delta$ , (44) holds for all  $k$  simultaneously. □

**Lemma 4.5** ( $\sigma_k$  Upper and Lower Bound for  $B_k^{(1)} \geq 2$ ). *For small  $\delta > 0$ , the following holds for all  $k$  such that  $B_k^{(1)} \geq 2$ .*

$$\sigma_k \cdot \sqrt{1 - O\left(\sqrt{\frac{\ln(1/\delta)}{p_k N_1}}\right)} \leq \hat{\sigma}_k \leq \sigma_k \cdot \sqrt{1 + O\left(\sqrt{\frac{\ln(1/\delta)}{p_k N_1}}\right)}. \quad (45)$$

*Proof.* To obtain a concentration inequality on  $\hat{\sigma}_k$ , we will apply bounded differences on the unbiased sample variance estimator for  $B_k^{(1)} > 2$ , which is:

$$\hat{\sigma}_k^2 = \frac{1}{B_k^{(1)} - 1} \sum_{i=1}^{B_k^{(1)}} (X_{k,i} - \hat{\mu}_k)^2. \quad (46)$$

Note that the unbiased sampled variance is a U-Statistic  $U_n$  that arises from taking  $g(X_{k,i}, X_{k,j}) = \frac{1}{2}(X_{k,i} - X_{k,j})^2$  (Ferguson [2005] reproduced as Lemma 5.3). For bounded U-Statistics, we can apply bounded differences to obtain a concentration bound (Rinaldo [2018] reproduced as Lemma 5.4). Namely, where  $g(X_{k,i}, X_{k,j}) \leq b$ ,

$$P(|U_n - \mathbb{E}[U_n]| \geq t) \leq 2 \exp\left(-\frac{nt^2}{8b^2}\right). \quad (47)$$

We use this concentration inequality with  $n = B_k^{(1)}$  and  $g(X_{k,i}, X_{k,j}) \leq b = C_k^{(\mu^2)}$ . Hence,

$$P(|\hat{\sigma}_k^2 - \sigma_k^2| \geq t) \leq 2 \exp\left(-\frac{B_k^{(1)} t^2}{8 C_k^{(\mu^4)}}\right). \quad (48)$$



We then set  $2\delta = 2\exp(-\frac{B_k^{(1)}t^2}{8C_k^{(\mu^4)}})$  and  $t = \sqrt{8\ln(1/\delta)C_k^{(\mu^4)}/B_k^{(1)}}$ . Thus, with probability at least  $1 - 2\delta$ ,

$$\sigma_k^2 - \sqrt{\frac{8\ln(1/\delta)C_k^{(\mu^4)}}{B_k^{(1)}}} \leq \hat{\sigma}_k^2 \leq \sigma_k^2 + \sqrt{\frac{8\ln(1/\delta)C_k^{(\mu^4)}}{B_k^{(1)}}}. \quad (49)$$

Using Lemma 4.4, we have  $B_k^{(1)} \geq p_k N_1 - \sqrt{2\ln(1/\delta)p_k N_1}$  with probability at least  $1 - K\delta$ .

$$\sigma_k^2 - \sqrt{\frac{8\ln(1/\delta)C_k^{(\mu^4)}}{p_k N_1 - \sqrt{2\ln(1/\delta)p_k N_1}}} \leq \hat{\sigma}_k^2 \leq \sigma_k^2 + \sqrt{\frac{8\ln(1/\delta)C_k^{(\mu^4)}}{p_k N_1 - \sqrt{2\ln(1/\delta)p_k N_1}}} \quad (50)$$

$$\sigma_k^2 - \sqrt{\frac{8\ln(1/\delta)C_k^{(\mu^4)}}{p_k N_1 \left(1 - \sqrt{\frac{2\ln(1/\delta)}{p_k N_1}}\right)}} \leq \hat{\sigma}_k^2 \leq \sigma_k^2 + \sqrt{\frac{8\ln(1/\delta)C_k^{(\mu^4)}}{p_k N_1 \left(1 - \sqrt{\frac{2\ln(1/\delta)}{p_k N_1}}\right)}} \quad (51)$$

$$\sigma_k^2 - \sqrt{\frac{8\ln(1/\delta)C_k^{(\mu^4)}}{p_k N_1} \cdot \left(1 + \sqrt{\frac{2\ln(1/\delta)}{p_k N_1}}\right)} \leq \hat{\sigma}_k^2 \leq \sigma_k^2 + \sqrt{\frac{8\ln(1/\delta)C_k^{(\mu^4)}}{p_k N_1} \cdot \left(1 + \sqrt{\frac{2\ln(1/\delta)}{p_k N_1}}\right)} \quad (52)$$

$$\sigma_k^2 - \sqrt{\frac{8\ln(1/\delta)C_k^{(\mu^4)}}{p_k N_1} + \frac{8\sqrt{2}\ln(1/\delta)^{3/2}}{p_k^{3/2} N_1^{3/2}}} \leq \hat{\sigma}_k^2 \leq \sigma_k^2 + \sqrt{\frac{8\ln(1/\delta)C_k^{(\mu^4)}}{p_k N_1} + \frac{8\sqrt{2}\ln(1/\delta)^{3/2}}{p_k^{3/2} N_1^{3/2}}} \quad (53)$$

$$\sigma_k^2 - \sqrt{O\left(\frac{1}{p_k N_1}\right)} \leq \hat{\sigma}_k^2 \leq \sigma_k^2 + \sqrt{O\left(\frac{1}{p_k N_1}\right)} \quad (54)$$

$$\sqrt{\sigma_k^2 - \sqrt{O\left(\frac{1}{p_k N_1}\right)}} \leq \hat{\sigma}_k \leq \sqrt{\sigma_k^2 + \sqrt{O\left(\frac{1}{p_k N_1}\right)}} \quad (55)$$

$$\sigma_k \cdot \sqrt{1 - O\left(\frac{1}{\sqrt{p_k N_1}}\right)} \leq \hat{\sigma}_k \leq \sigma_k \cdot \sqrt{1 + O\left(\frac{1}{\sqrt{p_k N_1}}\right)}. \quad (56)$$

We union bound across the strata and intersection bound our application of Lemma 4, and get that the above holds with probability at least  $(1 - 2K\delta)(1 - K\delta) = 1 - 3K\delta + 2K^2\delta^2 \geq 1 - 3K\delta$ . We impose the condition that  $\sigma_k \neq 0$  on this derivation because we divide by  $\sigma_k$  in (55). However, if  $\sigma_k = 0$ , then  $\hat{\sigma}_k = \sigma_k$ , so the bound still holds.  $\square$

**Lemma 4.6** ( $\sqrt{p_k}\sigma_k$  Upper Bound for  $B_k^{(1)} < 2$ ). *For  $B_k^{(1)} < 2$ , the following holds for all  $k$ :*

$$\sqrt{\hat{p}_k}\hat{\sigma}_k \leq \sqrt{p_k}\sigma_k + O\left(\frac{1}{\sqrt{N_1}}\right). \quad (57)$$

*Proof.* In the cases where  $B_k^{(1)} < 2$ , we will upper bound  $\sqrt{\hat{p}_k}\hat{\sigma}_k$  directly instead of combining our concentration inequalities for  $\sqrt{\hat{p}_k}$  and  $\hat{\sigma}_k$  separately.

$$\sqrt{\hat{p}_k}\hat{\sigma}_k \leq \sqrt{\frac{2}{N_1}}\hat{\sigma}_k \leq \sqrt{p_k}\sigma_k + \sqrt{\frac{2}{N_1}}C_k^{(\sigma)} \leq \sqrt{p_k}\sigma_k + O\left(\frac{1}{\sqrt{N_1}}\right). \quad (58)$$

$\square$

**Lemma 4.7** ( $T_k$  Lower Bound). *For small  $\delta > 0$ , the following is true simultaneously for all  $k$  where  $p_k > p_* = \frac{2(\sqrt{\ln(1/\delta)+1})}{N_1}$ ,*

$$\hat{T}_k \geq T_k \left( 1 - O \left( \sqrt{\frac{\ln(1/\delta)}{p_k N_1}} \right) \right). \quad (59)$$

*Proof.* Our goal is to lower bound  $\hat{T}_k = \frac{\sqrt{\hat{p}_k} \hat{\sigma}_k}{\sum_{i=1}^K \sqrt{\hat{p}_i} \hat{\sigma}_i}$ . To accomplish this, we will upper bound the denominator and lower bound the numerator. To lower bound the numerator, we will apply our concentration inequalities for  $\sqrt{\hat{p}_k}$  and  $\hat{\sigma}_k$  from Lemma 4.2 and Lemma 4.5. We only lower bound for  $k$  where  $B_k^{(1)} \geq 2$ . Through our high probability lower bound on  $B_k^{(1)}$ , we know that for  $B_k^{(1)} \geq 2$  to be true we can assume  $p_k > \frac{2(\sqrt{\ln(1/\delta)+1})}{N_1}$ . We denote  $p_* = \frac{2(\sqrt{\ln(1/\delta)+1})}{N_1}$ . This allows us to satisfy the conditions of Lemma 4.5.

$$\sqrt{\hat{p}_k} \sigma_k \geq \sqrt{p_k} \sigma_k \sqrt{1 - O \left( \sqrt{\frac{\ln(1/\delta)}{p_k N_1}} \right)} \sqrt{1 - O \left( \sqrt{\frac{\ln(1/\delta)}{p_k N_1}} \right)} \quad (60)$$

$$\geq \sqrt{p_k} \sigma_k \left( 1 - O \left( \sqrt{\frac{\ln(1/\delta)}{p_k N_1}} \right) \right) \geq \sqrt{p_k} \sigma_k - O \left( \sqrt{\frac{\ln(1/\delta)}{N_1}} \right). \quad (61)$$

To maximize the denominator, we also apply Lemma 4.1 and split the cases where  $p_k > p_*$  and where  $p_k \leq p_*$ .

$$\sum_{i=1}^K \sqrt{\hat{p}_i} \hat{\sigma}_i \leq \sum_{p_k > p_*}^K \sqrt{p_i} \sigma_i \sqrt{1 + O \left( \sqrt{\frac{\ln(1/\delta)}{p_i N_1}} \right)} \sqrt{1 + O \left( \sqrt{\frac{\ln(1/\delta)}{p_i N_1}} \right)} + \sum_{p_k \leq p_*}^K \sqrt{p_i} \sigma_i + O \left( \frac{1}{\sqrt{N_1}} \right) \quad (62)$$

$$\leq \sum_{p_k < p_*}^K \sqrt{p_i} \sigma_i \left( 1 + O \left( \sqrt{\frac{\ln(1/\delta)}{p_i N_1}} \right) \right) + \sum_{p_k \leq p_*}^K \sqrt{p_i} \sigma_i + O \left( \frac{1}{\sqrt{N_1}} \right) \quad (63)$$

$$\leq \sum_{i=1}^K \sqrt{p_i} \sigma_i + O \left( \sqrt{\frac{\ln(1/\delta)}{N_1}} \right). \quad (64)$$

Thus, for  $p_k > p_* = \frac{2(\sqrt{\ln(1/\delta)+1})}{N_1}$ ,

$$\hat{T}_k = \frac{\sqrt{\hat{p}_k} \hat{\sigma}_k}{\sum_{i=1}^K \sqrt{\hat{p}_i} \hat{\sigma}_i} \geq \frac{\sqrt{p_k} \sigma_k - O \left( \sqrt{\frac{\ln(1/\delta)}{N_1}} \right)}{\sum_{i=1}^K \sqrt{p_i} \sigma_i + O \left( \sqrt{\frac{\ln(1/\delta)}{N_1}} \right)} = \frac{\frac{1}{\sum_{i=1}^K \sqrt{p_i} \sigma_i}}{\frac{1}{\sum_{i=1}^K \sqrt{p_i} \sigma_i}} \cdot \frac{\sqrt{p_k} \sigma_k - O \left( \sqrt{\frac{\ln(1/\delta)}{N_1}} \right)}{\sum_{i=1}^K \sqrt{p_i} \sigma_i + O \left( \sqrt{\frac{\ln(1/\delta)}{N_1}} \right)} \quad (65)$$

$$\geq \frac{T_k - O \left( \sqrt{\frac{\ln(1/\delta)}{N_1}} \right)}{1 + O \left( \sqrt{\frac{\ln(1/\delta)}{N_1}} \right)} \geq \left( T_k - O \left( \sqrt{\frac{\ln(1/\delta)}{N_1}} \right) \right) \left( 1 - O \left( \sqrt{\frac{\ln(1/\delta)}{N_1}} \right) \right) \quad (66)$$

$$\geq T_k - O \left( \sqrt{\frac{\ln(1/\delta)}{N_1}} \right) - O \left( \sqrt{\frac{p_k \ln(1/\delta)}{N_1}} \right) + O \left( \frac{\ln(1/\delta)}{N_1} \right) \quad (67)$$

$$\geq T_k - O \left( \sqrt{\frac{\ln(1/\delta)}{N_1}} \right) \geq T_k \left( 1 - O \left( \sqrt{\frac{\ln(1/\delta)}{p_k N_1}} \right) \right) \quad \square$$

Through the union bound on Lemma 4.1, Lemma 4.2, Lemma 4.5, (73) occurs with probability at least  $1 - 5K\delta$ .

### 4.3 Stage 2

Recall that we use high probability bounds over Stage 1 and take the expectation over Stage 2. In this section, we derive bounds on quantities in Stage 2 that will help us upper bound the mean squared error of our algorithm.

**Lemma 4.8** ( $B_k^{(2)}$  Lower Bound). *If Lemma 4.7 holds, the following is true for all  $k$  where  $p_k > p_* = \frac{2(\sqrt{\ln(1/\delta)}+1)}{N_1}$  with probability at least  $1 - \gamma$ ,*

$$B_k^{(2)} \geq p_k T_k N_2 \left( 1 - O\left(\sqrt{\frac{1}{p_k N_1}}\right) - O\left(\sqrt{\frac{\ln(1/\gamma)}{N_2 p_k^{3/2}}}\right) \right). \quad (68)$$

*Proof.* Recall that  $B_k^{(2)} \sim \text{Binomial}(N_2 \hat{T}_k, p_k)$ . Thus, using Tarjan [2009] reproduced as Lemma 5.2, we have that:

$$P\left(B_k^{(2)} \leq (1 - \epsilon) \left(p_k \hat{T}_k N_2\right)\right) \leq \exp\left(\frac{-\epsilon^2 p_k \hat{T}_k N_2}{2}\right). \quad (69)$$

We set  $\delta = \exp\left(\frac{-\epsilon^2 p_k \hat{T}_k N_2}{2}\right)$  and  $\epsilon = \sqrt{\frac{2 \ln(1/\delta)}{p_k \hat{T}_k N_2}}$ . Thus, we have that with probability  $1 - 6K\delta$ :

$$B_k^{(2)} \geq p_k \hat{T}_k N_2 \left( 1 - \sqrt{\frac{2 \ln(1/\delta)}{p_k \hat{T}_k N_2}} \right) \geq p_k \hat{T}_k N_2 - \sqrt{2 \ln(1/\delta) p_k \hat{T}_k N_2} \geq p_k \hat{T}_k N_2 \left( 1 - \sqrt{\frac{2 \ln(1/\delta)}{p_k \hat{T}_k N_2}} \right). \quad (70)$$

We lower bound the number of draws  $\hat{T}_k N_2$  by conditioning on Lemma 4.7 which says that  $\hat{T}_k \geq T_k \left( 1 - O\left(\sqrt{\frac{\ln(1/\delta)}{p_k N_1}}\right) \right)$ . Thus, with probability at least  $1 - \gamma$ ,

$$\geq p_k T_k N_2 \left( 1 - O\left(\sqrt{\frac{\ln(1/\delta)}{p_k N_1}}\right) \right) \left( 1 - \sqrt{\frac{2 \ln(1/\delta)}{p_k T_k N_2 \left( 1 - O\left(\sqrt{\frac{\ln(1/\delta)}{p_k N_1}}\right) \right)}} \right) \quad (71)$$

$$\geq p_k T_k N_2 \left( 1 - O\left(\sqrt{\frac{\ln(1/\delta)}{p_k N_1}}\right) - \sqrt{\frac{2 \ln(1/\delta)}{p_k T_k N_2 \left( 1 - O\left(\frac{\ln(1/\delta)}{\sqrt{p_k N_1}}\right) \right)}} \right) \quad (72)$$

$$\geq p_k T_k N_2 \left( 1 - O\left(\sqrt{\frac{\ln(1/\delta)}{p_k N_1}}\right) - \sqrt{\frac{2 \ln(1/\delta)}{p_k T_k N_2} \cdot \left( 1 + O\left(\frac{\ln(1/\delta)}{\sqrt{p_k N_1}}\right) \right)} \right) \quad (73)$$

$$\geq p_k T_k N_2 \left( 1 - O\left(\sqrt{\frac{\ln(1/\delta)}{p_k N_1}}\right) - \sqrt{\frac{2 \ln(1/\delta)}{p_k T_k N_2} + O\left(\frac{\ln(1/\delta) \ln(1/\gamma)}{N_2 p_k^2 \sqrt{N_1}}\right)} \right) \quad (74)$$

$$\geq p_k T_k N_2 \left( 1 - O\left(\sqrt{\frac{1}{p_k N_1}}\right) - O\left(\sqrt{\frac{\ln(1/\gamma)}{N_2 p_k^{3/2}}}\right) \right) \quad \square$$

We use the fact that  $p_k > p_* = 1/N_1$ .

**Lemma 4.9** ( $w_k^2 P(B_k^{(2)} = 0)$  Upper Bound). *If Lemma 4.7 holds for  $p_k > p_*$  and if  $N_2 = \omega(N_1^{3/4})$ ,*

$$p_k^x P(B_k^{(2)} = 0) \leq O\left(\frac{1}{N_1^{x/2}}\right) + O\left(\frac{\sqrt{N_1}}{N_2^2}\right). \quad (75)$$

*Proof.*

$$p_k^x P(B_k^{(2)} = 0) = p_k(1 - p_k)^{\lceil \hat{T}_k N_2 \rceil} \leq p_k(1 - p_k)^{\hat{T}_k N_2}. \quad (76)$$

If  $p_k \leq p_* = \frac{2(\sqrt{\ln(1/\delta)+1})}{N_1}$ , ,

$$p_k^x(1 - p_k)^{\hat{T}_k N_2} \leq O\left(\frac{1}{N_1^{x/2}}\right). \quad (77)$$

Else if  $p_k > p_*$ , which satisfies the conditions of Lemma 4.7,

$$p_k^x(1 - p_k)^{\hat{T}_k N_2} \leq (1 - p_k)^{T_k N_2 \left(1 - O\left(\frac{1}{\sqrt{p_k N_1}}\right)\right)} \quad (78)$$

$$\leq (1 - p_k)^{N_2 \left(O(\sqrt{p_k}) - O\left(\frac{1}{\sqrt{N_1}}\right)\right)} \quad (79)$$

$$\leq (1 - p_k)^{O\left(\frac{N_2}{N_1^{1/4}}\right)}. \quad (80)$$

We assume that  $N_2 = \omega(N_1^{3/4})$  which means that (80) is decreasing exponentially. Hence, we can say that

$$(1 - p_k)^{O\left(\frac{N_2}{N_1^{1/4}}\right)} = O\left(\frac{\sqrt{N_1}}{N_2^2}\right).$$

$$p_k^x P(B_k^{(2)} = 0) \leq O\left(\frac{1}{N_1^{x/2}}\right) + O\left(\frac{\sqrt{N_1}}{N_2^2}\right) \quad \square$$

## 4.4 Final Result

**Theorem 1.** *For small  $\delta > 0$  over Stage 1 (Lemma 4.1, Lemma 4.2, Lemma 4.3, Lemma 4.4, Lemma 4.5, Lemma 4.6, and Lemma 4.7) and if  $N_2 = \omega(N_1^{3/4})$ , the expectation of the mean squared error of ABAE over Stage 2 is*

$$\mathbb{E}[(\hat{\mu}_{all} - \mu_{all})^2] \leq O(N_1^{-1}) + O(N_2^{-1}) + O(N_1^{1/2} N_2^{-3/2}). \quad (81)$$

*Proof.* By Lemma 4.1, Lemma 4.2, Lemma 4.3, Lemma 4.4, Lemma 4.5, Lemma 4.6, and Lemma 4.7, we have that for small  $\delta > 0$ :

$$\sqrt{p_k} \sqrt{1 - O\left(\sqrt{\frac{\ln(1/\delta)}{p_k N_1}}\right)} \leq \sqrt{\hat{p}_k} \leq \sqrt{p_k} \sqrt{1 + O\left(\sqrt{\frac{\ln(1/\delta)}{p_k N_1}}\right)}. \quad (82)$$

$$w_k - O\left(\frac{\ln(1/\delta)\sqrt{p_k}}{\sqrt{N_1}}\right) \leq \hat{w}_k \leq w_k + O\left(\frac{\ln(1/\delta)\sqrt{p_k}}{\sqrt{N_1}}\right) \quad (83)$$

$$B_k^{(1)} \geq p_k N_1 - \sqrt{2 \ln(1/\delta) p_k N_1}. \quad (84)$$

$$\sigma_k \cdot \sqrt{1 - O\left(\sqrt{\frac{\ln(1/\delta)}{p_k N_1}}\right)} \leq \hat{\sigma}_k \leq \sigma_k \cdot \sqrt{1 + O\left(\sqrt{\frac{\ln(1/\delta)}{p_k N_1}}\right)}. \quad (85)$$

$$\hat{T}_k \geq T_k \left(1 - O\left(\sqrt{\frac{\ln(1/\delta)}{p_k N_1}}\right)\right). \quad (86)$$

Let  $\mathcal{E}$  be the event that all these inequalities are satisfied. For the remainder of the proof, we condition on  $\mathcal{E}$  and take the expectation over the randomness in Stage 2. We now begin to decompose the mean squared error.

$$\mathbb{E}[(\hat{\mu}_{\text{all}} - \mu_{\text{all}})^2] = \mathbb{E}\left[\left(\sum_{k=1}^K \hat{w}_k \hat{\mu}_k - w_k \mu_k\right)^2\right]. \quad (87)$$

We know that  $\mathbb{E}[\hat{\mu}_k] = P(B_k^{(2)} > 0) \cdot \mu_k$ . As a result,  $\mathbb{E}[\hat{\mu}_k] + \mu_k P(B_k^{(2)} = 0) = \mu_k$ . Hence,

$$= \mathbb{E}\left[\left(\sum_{k=1}^K \hat{w}_k (\hat{\mu}_k - \mathbb{E}[\hat{\mu}_k]) + (\hat{w}_k - w_k) \mathbb{E}[\hat{\mu}_k] - w_k \mu_k P(B_k = 0)\right)^2\right] \quad (88)$$

We now use that  $\mathbb{E}[(A + B)^2] = \mathbb{E}[A^2 + 2AB + B^2]$ , so if  $\mathbb{E}[A] = 0$  then  $\mathbb{E}[(A + B)^2] = \mathbb{E}[A^2] + \mathbb{E}[B^2]$ . We set  $A = \hat{w}_k (\hat{\mu}_k - \mathbb{E}[\hat{\mu}_k])$  and  $B = (\hat{w}_k - w_k) \mathbb{E}[\hat{\mu}_k] - w_k \mu_k P(B_k = 0)$ . Note that  $\mathbb{E}[A] = 0$ . Thus,

$$= \mathbb{E}\left[\left(\sum_{k=1}^K \hat{w}_k (\hat{\mu}_k - \mathbb{E}[\hat{\mu}_k])\right)^2\right] + \mathbb{E}\left[\left(\sum_{k=1}^K (\hat{w}_k - w_k) \mathbb{E}[\hat{\mu}_k] - w_k \mu_k P(B_k = 0)\right)^2\right] \quad (89)$$

$$\leq \sum_{k=1}^K \hat{w}_k^2 \text{Var}(\hat{\mu}_k) + \left[\max_k C_k^{\mu_k^2}\right] \mathbb{E}\left[\left(\sum_{k=1}^K w_k P(B_k^{(2)} = 0) + (\hat{w}_k - w_k)\right)^2\right]. \quad (90)$$

We will now separately upper bound the two terms in this expression. For the first term, notice that

$$\text{Var}[\hat{\mu}_k] = \mathbb{E}[\hat{\mu}_k^2] - \mathbb{E}[\hat{\mu}_k]^2 \quad (91)$$

$$= P(B_k^{(2)} > 0) \left(\mathbb{E}\left[\frac{\sigma_k^2}{B_k^{(2)}} | B_k^{(2)} > 0\right] + \mu_k^2\right) - P(B_k^{(2)} > 0)^2 \mu_k^2 \quad (92)$$

$$= P(B_k^{(2)} > 0) \mathbb{E}\left[\frac{\sigma_k^2}{B_k^{(2)}} | B_k^{(2)} > 0\right] + P(B_k^{(2)} > 0)(1 - P(B_k^{(2)} > 0)) \mu_k^2 \quad (93)$$

$$= P(B_k^{(2)} > 0) \mathbb{E}\left[\frac{\sigma_k^2}{B_k^{(2)}} | B_k^{(2)} > 0\right] + P(B_k^{(2)} > 0) P(B_k^{(2)} = 0) \mu_k^2 \quad (94)$$

$$\leq \mathbb{E}\left[\frac{\sigma_k^2}{B_k^{(2)}} | B_k^{(2)} > 0\right] + P(B_k^{(2)} = 0) \mu_k^2 \quad (95)$$

$$\sum_{k=1}^K \hat{w}_k^2 \text{Var}[\hat{\mu}_k] \leq \sum_{k=1}^K \hat{w}_k^2 \left(\mathbb{E}\left[\frac{\sigma_k^2}{B_k^{(2)}} | B_k^{(2)} > 0\right] + P(B_k^{(2)} = 0) \mu_k^2\right) \quad (96)$$

$$\leq \sum_{k=1}^K \left(O(p_k^2) + O\left(\frac{p_k^{3/2}}{\sqrt{N_1}}\right)\right) \left(\mathbb{E}\left[\frac{\sigma_k^2}{B_k^{(2)}} | B_k^{(2)} > 0\right] + P(B_k^{(2)} = 0) \mu_k^2\right) \quad (97)$$

$$\leq \sum_{k=1}^K \left(O(p_k^2) + O\left(\frac{p_k^{3/2}}{\sqrt{N_1}}\right)\right) \left(\mathbb{E}\left[\frac{\sigma_k^2}{B_k^{(2)}} | B_k^{(2)} > 0\right] + P(B_k^{(2)} = 0) \mu_k^2\right) \quad (98)$$

We split the cases where  $p_k$  is small and  $p_k$  is large via  $p_* = \frac{2(\sqrt{\ln(1/\delta)+1})}{N_1}$ . This value of  $p_*$  is chosen as according to the conditions of Lemma 4.7.

$$\leq \sum_{p_k > p_*}^K \left( O(p_k^2) + O\left(\frac{p_k^{3/2}}{\sqrt{N_1}}\right) \right) \left( \mathbb{E}\left[\frac{\sigma_k^2}{B_k^{(2)}} | B_k^{(2)} > 0\right] + P(B_k^{(2)} = 0)\mu_k^2 \right) + \sum_{p_k \leq p_*}^K O\left(\frac{1}{N_1}\right) \quad (99)$$

Note that  $p_k^2 P(B_k^{(2)} = 0) \leq O\left(\frac{1}{N_1}\right) + O\left(\frac{\sqrt{N_1}}{N_2^2}\right)$  according to Lemma 4.9.

$$\leq \sum_{p_k > p_*}^K \left( O(p_k^2) + O\left(\frac{p_k^{3/2}}{\sqrt{N_1}}\right) \right) \mathbb{E}\left[\frac{\sigma_k^2}{B_k^{(2)}} | B_k^{(2)} > 0\right] + O\left(\frac{1}{N_1}\right) + O\left(\frac{\sqrt{N_1}}{N_2^2}\right) \quad (100)$$

We will now apply Lemma 4.8 to upper bound  $\mathbb{E}\left[\frac{\sigma_k^2}{B_k^{(2)}} | B_k^{(2)} > 0\right]$ . Namely, we will split the expectation into two cases, one where  $B_k^{(2)} \geq F_k$  and another where  $B_k < F_k$ . Specifically, we apply it such that  $B_k^{(2)} \geq F_k = p_k T_k N_2 \left(1 - O\left(\sqrt{\frac{1}{p_k N_1}}\right) - O\left(\sqrt{\frac{\ln(1/\gamma)}{N_2 p_k^{3/2}}}\right)\right)$  with failure probability  $\gamma = \frac{1}{e\sqrt{N_1}} \leq O\left(\frac{1}{N_1}\right)$ .

$$\leq \sum_{p_k > p_*}^K \left( O(p_k^2) + O\left(\frac{p_k^{3/2}}{\sqrt{N_1}}\right) \right) \left( \mathbb{E}\left[\frac{\sigma_k^2}{B_k^{(2)}} | B_k^{(2)} > 0\right] \right) + O\left(\frac{1}{N_1}\right) + O\left(\frac{\sqrt{N_1}}{N_2^2}\right) \quad (101)$$

$$\leq \sum_{p_k > p_*}^K \left( O(p_k^2) + O\left(\frac{p_k^{3/2}}{\sqrt{N_1}}\right) \right) \left( P(B_k^{(2)} > F_k) \mathbb{E}\left[\frac{\sigma_k^2}{B_k^{(2)}} | B_k^{(2)} > F_k\right] + O\left(\frac{1}{N_1}\right) C_k^{\sigma^2} \right) + O\left(\frac{1}{N_1}\right) + O\left(\frac{\sqrt{N_1}}{N_2^2}\right) \quad (102)$$

$$\leq \sum_{p_k > p_*}^K \left( O(p_k^2) + O\left(\frac{p_k^{3/2}}{\sqrt{N_1}}\right) \right) \left( \mathbb{E}\left[\frac{\sigma_k^2}{B_k^{(2)}} | B_k^{(2)} > F_k\right] + O\left(\frac{1}{N_1}\right) C_k^{\sigma^2} \right) + O\left(\frac{1}{N_1}\right) + O\left(\frac{\sqrt{N_1}}{N_2^2}\right) \quad (103)$$

$$\leq \sum_{p_k > p_*}^K \left( O(p_k^2) + O\left(\frac{p_k^{3/2}}{\sqrt{N_1}}\right) \right) \mathbb{E}\left[\frac{\sigma_k^2}{B_k^{(2)}} | B_k^{(2)} > F_k\right] + O\left(\frac{1}{N_1}\right) + O\left(\frac{\sqrt{N_1}}{N_2^2}\right) \quad (104)$$

$$\leq \sum_{p_k > p_*}^K \left( O(p_k^2) + O\left(\frac{p_k^{3/2}}{\sqrt{N_1}}\right) \right) \frac{\sigma_k^2}{p_k T_k N_2 \left(1 - O\left(\sqrt{\frac{1}{p_k N_1}}\right) - O\left(\sqrt{\frac{\ln(1/\gamma)}{N_2 p_k^{3/2}}}\right)\right)} + O\left(\frac{1}{N_1}\right) + O\left(\frac{\sqrt{N_1}}{N_2^2}\right) \quad (105)$$

$$\leq \sum_{p_k > p_*}^K \left( O\left(\frac{\sqrt{p_k}}{N_2}\right) + O\left(\frac{1}{N_2 \sqrt{N_1}}\right) \right) \left( 1 + O\left(\sqrt{\frac{1}{p_k N_1}}\right) + O\left(\sqrt{\frac{\ln(1/\gamma)}{N_2 p_k^{3/2}}}\right) \right) + O\left(\frac{1}{N_1}\right) + O\left(\frac{\sqrt{N_1}}{N_2^2}\right) \quad (106)$$

$$\leq \sum_{p_k > p_*}^K \left( O\left(\frac{\sqrt{p_k}}{N_2}\right) + O\left(\frac{1}{N_2 \sqrt{N_1}}\right) \right) \left( 1 + O\left(\sqrt{\frac{1}{p_k N_1}}\right) + O\left(\sqrt{\frac{\ln(1/\gamma)}{N_2 p_k^{3/2}}}\right) \right) + O\left(\frac{1}{N_1}\right) + O\left(\frac{\sqrt{N_1}}{N_2^2}\right) \quad (107)$$

$$\leq \sum_{p_k > p_*}^K O\left(\frac{1}{N_2} + \frac{1}{N_2 \sqrt{N_1}} + \frac{\sqrt{N_1}}{N_2 \sqrt{N_2}}\right) + O\left(\frac{1}{N_2 \sqrt{N_1}}\right) \left( 1 + O\left(\sqrt{\frac{1}{p_k N_1}}\right) + O\left(\sqrt{\frac{\ln(1/\gamma)}{N_2 p_k^{3/2}}}\right) \right) + O\left(\frac{1}{N_1}\right) + O\left(\frac{\sqrt{N_1}}{N_2^2}\right) \quad (108)$$

$$\leq \sum_{p_k > p_*}^K O\left(\frac{1}{N_2} + \frac{\sqrt{N_1}}{N_2 \sqrt{N_2}}\right) + O\left(\frac{1}{N_2 \sqrt{N_1}} + \frac{1}{N_2 \sqrt{N_1}} + \frac{\sqrt{N_1}}{N_2 \sqrt{N_2}}\right) + O\left(\frac{1}{N_1}\right) + O\left(\frac{\sqrt{N_1}}{N_2^2}\right) \quad (109)$$

$$\leq O\left(\frac{1}{N_1}\right) + O\left(\frac{1}{N_2}\right) + O\left(\frac{\sqrt{N_1}}{N_2 \sqrt{N_2}}\right) \quad (110)$$

We will now upper bound the second term in the expression on (90).

$$\mathbb{E} \left[ \left( \sum_{k=1}^K (\hat{w}_k - w_k) \mathbb{E}[\hat{\mu}_k] - w_k \mu_k P(B_k = 0) \right)^2 \right] \leq \left[ \max_k C_k^{\mu_k^2} \right] \mathbb{E} \left[ \left( \sum_{k=1}^K w_k P(B_k^{(2)} = 0) + (\hat{w}_k - w_k) \right)^2 \right] \quad (111)$$

$$\leq \left[ \max_k C_k^{\mu_k^2} \right] \mathbb{E} \left[ \left( O\left(\frac{\sqrt{N_1}}{N_2}\right) + O\left(\frac{1}{\sqrt{N_1}}\right) \right)^2 \right] \quad (112)$$

$$\leq O\left(\frac{1}{N_1}\right) + O\left(\frac{1}{N_2}\right) \quad (113)$$

Thus, by combining (110) and (113), we finally have that

$$\mathbb{E}[(\hat{\mu}_{\text{all}} - \mu_{\text{all}})^2 | \mathcal{E}] = \mathbb{E} \left[ \left( \sum_{k=1}^K \hat{w}_k \hat{\mu}_k - w_k \mu_k \right)^2 \right] \quad (114)$$

$$= \sum_{k=1}^K \hat{w}_k^2 \text{Var}(\hat{\mu}_k) + \left[ \max_k C_k^{\mu_k^2} \right] \mathbb{E} \left[ \left( \sum_{k=1}^K w_k P(B_k^{(2)} = 0) + (\hat{w}_k - w_k) \right)^2 \right] \quad (115)$$

$$\leq \left[ O\left(\frac{1}{N_1}\right) + O\left(\frac{1}{N_2}\right) + O\left(\frac{\sqrt{N_1}}{N_2 \sqrt{N_2}}\right) \right] + \left[ O\left(\frac{1}{N_1}\right) + O\left(\frac{1}{N_2}\right) \right] \quad (116)$$

$$\leq O(N_1^{-1}) + O(N_2^{-1}) + O(N_1^{1/2} N_2^{-3/2}) \quad \square$$

## 4.5 Sample Reuse

Note that our algorithm shown in Algorithm 1 involves a parameter *SampleReuse*, which decides whether we reuse samples from Stage 1 in Stage 2 and vice-versa. In our analysis, we assumed that *SampleReuse* was false. However, we show that *SampleReuse* strictly reduces the error of ABAE.

**Corollary 4.1.** *If SampleReuse is true, ABAE still achieves the the following upper bound on the the mean squared error for small  $\delta > 0$  over Stage 1 and expectation over Stage 2,*

$$\mathbb{E}[(\hat{\mu}_{\text{all}} - \mu_{\text{all}})^2] \leq O(N_1^{-1}) + O(N_2^{-1}) + O(N_1^{1/2} N_2^{-3/2}). \quad (117)$$

*Proof.* The high probability bounds we have on  $\hat{p}_k$ ,  $\hat{w}_k$ ,  $B_k^{(1)}$ ,  $\hat{\sigma}_k$ ,  $\hat{T}_k$  still hold with sample reuse, so the rates presented in Theorem 1 still hold. Furthermore, note that the estimate  $\hat{\mu}_k$  is actually a stochastic sum where the number of summands are dependent on Stage 1. As a result, the samples from Stage 1 and Stage 2 are actually identical and independently distributed, so we can reuse samples when estimating  $\hat{\mu}_k$ . Although not reflected in the bounds presented for  $\hat{p}_k$ ,  $\hat{w}_k$ , and  $\hat{\mu}_k$ , the error is further reduced by better estimates produced by sample reuse in practice.  $\square$

## 4.6 Edge Cases

We address the following edge cases in our algorithm and analysis.

1. If the variance in a stratum is estimated to be zero, this will result in  $\hat{T}_k = 0$ . In this scenario, we can simply use a simple sample from Stage 1 to represent the constant  $\mu_k$ .
2. If the predicate positive rate or the variance in every stratum is estimated to be zero, this will result in  $\hat{T}_k = 0$  for all  $k$ . In this scenario, we set  $T_k = 1/K$  for each  $k$ .
3. ...

## 5 Appendix

**Lemma 5.1** (Chung and Lu [2002]). *Let  $X_1, \dots, X_n$  be independent random variables with:*

$$P(X_i = 1) = p_i, \quad P(X_i = 0) = 1 - p_i \quad (118)$$

*For  $X = \sum_{i=1}^n a_i X_i$ , we have  $\mathbb{E}[X] = \sum_{i=1}^n a_i p_i$  and we have define  $\nu = \sum_{i=1}^n a_i^2 p_i$ . Then we have:*

$$P(x < \mathbb{E}[X] - \lambda) \leq e^{-\lambda^2/2\nu} \quad (119)$$

$$P(X > \mathbb{E}[X] + \lambda) \leq e^{-\frac{\lambda^2}{2(\nu + a\lambda/3)}} \quad (120)$$

where  $a = \max\{a_1, \dots, a_n\}$

**Lemma 5.2** (Tarjan [2009]). *Let  $X_1, \dots, X_n$  be independent random variables. They do not need to be from the same distribution. Assume that  $0 \leq X_i \leq 1$  for each  $i$ . Let  $X = X_1 + \dots + X_n$ . Write  $\mu = \mathbb{E}[X] = \mathbb{E}[X_1] + \dots + \mathbb{E}[X_n]$ . Then, for any  $\epsilon \geq 0$ ,*

$$P(X \geq (1 + \epsilon)\mu) \leq \exp\left(-\frac{\epsilon^2}{2 + \epsilon}\mu\right) \quad (121)$$

$$P(X \leq (1 - \epsilon)\mu) \leq \exp\left(-\frac{\epsilon^2}{2}\mu\right) \quad (122)$$

**Lemma 5.3** (Ferguson [2005]). *Let  $X_1, \dots, X_n$  be i.i.d. For a U-statistic of order 2 in the form  $U_n = \frac{1}{\binom{n}{2}} \sum_{i < j} g(X_i, X_j)$ , if  $g(X_i, X_j) = \frac{1}{2}(X_i - X_j)^2$ ,*

$$\mathbb{E}[U_n] = \text{Var}[X] \quad (123)$$

**Lemma 5.4** (Rinaldo [2018]). *Let  $X_1, \dots, X_n$  be i.i.d. For a U-statistic of order 2 in the form  $U_n = \frac{1}{\binom{n}{2}} \sum_{i < j} g(X_i, X_j)$  with an upper bound  $g(X_i, X_j) \leq b$ ,*

$$P(|U_n - \mathbb{E}[U_n]| \geq t) \leq 2 \exp\left(-\frac{nt^2}{8b^2}\right) \quad (124)$$

## References

- Fan Chung and Linyuan Lu. Connected components in random graphs with given expected degree sequences. *Annals of combinatorics*, 6(2):125–145, 2002.
- Thomas Ferguson. U-statistics notes for statistics 200c, spring 2005. [http://www.stat.cmu.edu/~arinaldo/Teaching/36710/F18/Scribed\\_Lectures/Sep24.pdf](http://www.stat.cmu.edu/~arinaldo/Teaching/36710/F18/Scribed_Lectures/Sep24.pdf), 2005. [Online; accessed 11-Feb-2021].
- Alessandro Rinaldo. 36-710: Advanced statistical theory lecture 7: September 24. [http://www.stat.cmu.edu/~arinaldo/Teaching/36710/F18/Scribed\\_Lectures/Sep24.pdf](http://www.stat.cmu.edu/~arinaldo/Teaching/36710/F18/Scribed_Lectures/Sep24.pdf), 2018. [Online; accessed 11-Feb-2021].
- Robert Tarjan. 15-359: Probability and computing lecture 10: More chernoff bounds, sampling, and the chernoff + union bound method. <http://aiweb.techfak.uni-bielefeld.de/content/bworld-robot-control-software/>, 2009. [Online; accessed 11-Feb-2021].