Lecture 12 - Inverse and Implicit Differentiation

Garrett Credi

10/20/2026

Updates

Curve information forthcoming - in contact with course coordinator about adjusting final grade distribution.
Regrade requests for Exam 1 closed.

Quiz Time!

Let's get into neat rows, spread out, and slay this quiz!

Inverse and Implicit Differentiation

The Road So Far

Recall we're trying to build a *synthetic* approach to studying derivatives:

The Road So Far

Recall we're trying to build a *synthetic* approach to studying derivatives:
Atomic Rules:

The Road So Far

Recall we're trying to build a *synthetic* approach to studying derivatives:
Atomic Rules:

- Power Rule: $\frac{d}{dx}x^n$
- Trigonometric Rules: $\frac{d}{dx}\sin(x)$ etc.
- Exponential/Logarithm Rules: $\frac{d}{dx}e^x$, $\frac{d}{dx}\ln(x)$

Combination Rules:

The Road So Far

Recall we're trying to build a *synthetic* approach to studying derivatives:
Atomic Rules:

- Power Rule: $\frac{d}{dx}x^n$
- Trigonometric Rules: $\frac{d}{dx}\sin(x)$ etc.
- Exponential/Logarithm Rules: $\frac{d}{dx}e^x$, $\frac{d}{dx}\ln(x)$

Combination Rules:

- Sum/Difference Rule: $\frac{d}{dx}(f(x) \pm g(x))$
- Product Rule: $\frac{d}{dx}f(x)g(x)$
- Chain Rule: $\frac{d}{dx}g(f(x))$
- Constant Multiple Rule: $\frac{d}{dx}c \cdot f(x)$
- Quotient Rule: $\frac{d}{dx}\frac{f(x)}{g(x)}$
- Inverse Function Rule: $\frac{d}{dx}f^{-1}(x)$

Explicit vs. Implicit Definitions I

Up until now we've been dealing with *explicit* definitions of various functions and have been using those *explicit* definitions to compute their derivatives.

Explicit vs. Implicit Definitions I

Up until now we've been dealing with *explicit* definitions of various functions and have been using those *explicit* definitions to compute their derivatives.
An *explicit* definition of a function is one where we defined $f(x)$ by *explicitly* saying where a given input x maps to. E.g.:

Explicit vs. Implicit Definitions I

Up until now we've been dealing with *explicit* definitions of various functions and have been using those *explicit* definitions to compute their derivatives.
An *explicit* definition of a function is one where we defined $f(x)$ by *explicitly* saying where a given input x maps to. E.g.:

- $f(x) = x^2$
- $g(x) = \sin(x)$
- $h(x) = \sqrt{x^2 + 2x}$
- $j(x) = e^{\sin(x)}$
- $k(x) = (x + \sin(x))^{\frac{1}{2}}$
- etc

Explicit vs. Implicit Definitions I

Up until now we've been dealing with *explicit* definitions of various functions and have been using those *explicit* definitions to compute their derivatives.
An *explicit* definition of a function is one where we defined $f(x)$ by *explicitly* saying where a given input x maps to. E.g.:

- $f(x) = x^2$
- $g(x) = \sin(x)$
- $h(x) = \sqrt{x^2 + 2x}$
- $j(x) = e^{\sin(x)}$
- $k(x) = (x + \sin(x))^{\frac{1}{2}}$
- etc

This is not the only way in which we can define functions!

Explicit vs Implicit Definitions II

We could instead define relations *implicitly* - i.e. by writing out some equation that we want the function to satisfy, e.g.:

Explicit vs Implicit Definitions II

We could instead define relations *implicitly* - i.e. by writing out some equation that we want the function to satisfy, e.g.:

- $x^2 + y^2 = 1$
- $y^2 = x^3 + 2x + 1$
- $\arctan(\frac{y}{x}) = \frac{\pi}{4}$
- $\sin(x) + \cos(y) = 1$
- $x^2 - y^2 = 2$
- $x^3 + y^3 = 1$

Explicit vs Implicit Definitions II

We could instead define relations *implicitly* - i.e. by writing out some equation that we want the function to satisfy, e.g.:

- $x^2 + y^2 = 1$
- $y^2 = x^3 + 2x + 1$
- $\arctan(\frac{y}{x}) = \frac{\pi}{4}$
- $\sin(x) + \cos(y) = 1$
- $x^2 - y^2 = 2$
- $x^3 + y^3 = 1$

Just as we can *graph* a function: i.e. we plot all the points $(x, f(x))$,

Explicit vs Implicit Definitions II

We could instead define relations *implicitly* - i.e. by writing out some equation that we want the function to satisfy, e.g.:

- $x^2 + y^2 = 1$
- $y^2 = x^3 + 2x + 1$
- $\arctan(\frac{y}{x}) = \frac{\pi}{4}$
- $\sin(x) + \cos(y) = 1$
- $x^2 - y^2 = 2$
- $x^3 + y^3 = 1$

Just as we can *graph* a function: i.e. we plot all the points $(x, f(x))$, we can plot some implicit equation $F(x, y) = 0$ by plotting all the points (x, y) so that $F(x, y) = 0$.

Explicit vs Implicit Definitions II

We could instead define relations *implicitly* - i.e. by writing out some equation that we want the function to satisfy, e.g.:

- $x^2 + y^2 = 1$
- $y^2 = x^3 + 2x + 1$
- $\arctan(\frac{y}{x}) = \frac{\pi}{4}$
- $\sin(x) + \cos(y) = 1$
- $x^2 - y^2 = 2$
- $x^3 + y^3 = 1$

Just as we can *graph* a function: i.e. we plot all the points $(x, f(x))$, we can plot some implicit equation $F(x, y) = 0$ by plotting all the points (x, y) so that $F(x, y) = 0$.
When is this useful?

Applications of Implicit Relations

Oftentimes, real world process don't have simple ways of relating one quantity to another quantity.

Applications of Implicit Relations

Oftentimes, real world process don't have simple ways of relating one quantity to another quantity.
Consider: conservation of momentum/energy in physics:

$$\sum_{i=1}^{n} m_i v_i^{init} = \sum_{i=1}^{n} m_i v_i^{final} \quad (1)$$

$$\sum_{i=1}^{n} \frac{1}{2} m_i (v_i^{init})^2 = \sum_{i=1}^{n} \frac{1}{2} m_i (v_i^{final})^2 \quad (2)$$

Applications of Implicit Relations

Oftentimes, real world process don't have simple ways of relating one quantity to another quantity.
Consider: conservation of momentum/energy in physics:

$$\sum_{i=1}^{n} m_i v_i^{init} = \sum_{i=1}^{n} m_i v_i^{final} \quad (1)$$

$$\sum_{i=1}^{n} \frac{1}{2} m_i (v_i^{init})^2 = \sum_{i=1}^{n} \frac{1}{2} m_i (v_i^{final})^2 \quad (2)$$

The relation between initial and final velocities here are not a simple input-to-output relationship -

Applications of Implicit Relations

Oftentimes, real world process don't have simple ways of relating one quantity to another quantity.
Consider: conservation of momentum/energy in physics:

$$\sum_{i=1}^{n} m_i v_i^{init} = \sum_{i=1}^{n} m_i v_i^{final} \quad (1)$$

$$\sum_{i=1}^{n} \frac{1}{2} m_i (v_i^{init})^2 = \sum_{i=1}^{n} \frac{1}{2} m_i (v_i^{final})^2 \quad (2)$$

The relation between initial and final velocities here are not a simple input-to-output relationship - they are equations that need to be enforced.

Applications of Implicit Relations

Oftentimes, real world process don't have simple ways of relating one quantity to another quantity.
Consider: conservation of momentum/energy in physics:

$$\sum_{i=1}^{n} m_i v_i^{init} = \sum_{i=1}^{n} m_i v_i^{final} \quad (1)$$

$$\sum_{i=1}^{n} \frac{1}{2} m_i (v_i^{init})^2 = \sum_{i=1}^{n} \frac{1}{2} m_i (v_i^{final})^2 \quad (2)$$

The relation between initial and final velocities here are not a simple input-to-output relationship - they are equations that need to be enforced.
Also: predator-prey models, resource allocation, budgeting, etc.

Inverse Functions as a Relation

To address our final desired synthetic rule (inverse function rule)

Inverse Function

Inverse Functions as a Relation

To address our final desired synthetic rule (inverse function rule) we can also see that inverse functions are defined via an implicit rule:

Inverse Function

Inverse Functions as a Relation

To address our final desired synthetic rule (inverse function rule) we can also see that inverse functions are defined via an implicit rule:

Inverse Function

For a function $f(x)$ with domain $D \subseteq \mathbb{R}$ and range $R \subseteq \mathbb{R}$, the inverse function $f^{-1}(x)$ is the function that:

Inverse Functions as a Relation

To address our final desired synthetic rule (inverse function rule) we can also see that inverse functions are defined via an implicit rule:

Inverse Function

For a function $f(x)$ with domain $D \subseteq \mathbb{R}$ and range $R \subseteq \mathbb{R}$, the inverse function $f^{-1}(x)$ is the function that:

- Has domain R (domain of inverse is range of original)

Inverse Functions as a Relation

To address our final desired synthetic rule (inverse function rule) we can also see that inverse functions are defined via an implicit rule:

Inverse Function

For a function $f(x)$ with domain $D \subseteq \mathbb{R}$ and range $R \subseteq \mathbb{R}$, the inverse function $f^{-1}(x)$ is the function that:

- Has domain R (domain of inverse is range of original)
- Has range D (range of inverse is domain of original)

Inverse Functions as a Relation

To address our final desired synthetic rule (inverse function rule) we can also see that inverse functions are defined via an implicit rule:

Inverse Function

For a function $f(x)$ with domain $D \subseteq \mathbb{R}$ and range $R \subseteq \mathbb{R}$, the inverse function $f^{-1}(x)$ is the function that:

- Has domain R (domain of inverse is range of original)
- Has range D (range of inverse is domain of original)
- Satisfies the two relations:
  1. $f(f^{-1}(x)) = x$
  2. $f^{-1}(f(x)) = x$

Inverse Functions as a Relation

To address our final desired synthetic rule (inverse function rule) we can also see that inverse functions are defined via an implicit rule:

Inverse Function

For a function $f(x)$ with domain $D \subseteq \mathbb{R}$ and range $R \subseteq \mathbb{R}$, the inverse function $f^{-1}(x)$ is the function that:

- Has domain R (domain of inverse is range of original)
- Has range D (range of inverse is domain of original)
- Satisfies the two relations:
  1. $f(f^{-1}(x)) = x$
  2. $f^{-1}(f(x)) = x$

  i.e. $f(x)$ and $f^{-1}(x)$ "undo" eachother.

Examples of Inverse Functions

For some examples:

Examples of Inverse Functions

For some examples:

1. If $f(x) = x^2$ with domain *the positive real numbers* and range *the positive real numbers*

Examples of Inverse Functions

For some examples:

1. If $f(x) = x^2$ with domain *the positive real numbers* and range *the positive real numbers* $f^{-1}(x) = \sqrt{x}$ is the positive square root.

Examples of Inverse Functions

For some examples:

1. If $f(x) = x^2$ with domain *the positive real numbers* and range *the positive real numbers* $f^{-1}(x) = \sqrt{x}$ is the positive square root.
2. If $f(x) = x^2$ with domain *the negative real numbers* and range *the positive real numbers*

Examples of Inverse Functions

For some examples:

1. If $f(x) = x^2$ with domain *the positive real numbers* and range *the positive real numbers* $f^{-1}(x) = \sqrt{x}$ is the positive square root.
2. If $f(x) = x^2$ with domain *the negative real numbers* and range *the positive real numbers* $f^{-1}(x) = -\sqrt{x}$ is the negative square root.

Examples of Inverse Functions

For some examples:

1. If $f(x) = x^2$ with domain *the positive real numbers* and range *the positive real numbers* $f^{-1}(x) = \sqrt{x}$ is the positive square root.
2. If $f(x) = x^2$ with domain *the negative real numbers* and range *the positive real numbers* $f^{-1}(x) = -\sqrt{x}$ is the negative square root.
3. If $f(x) = \sin(x)$ with domain $[0, 2\pi)$ and range $[-1, 1]$

Examples of Inverse Functions

For some examples:

1. If $f(x) = x^2$ with domain *the positive real numbers* and range *the positive real numbers* $f^{-1}(x) = \sqrt{x}$ is the positive square root.
2. If $f(x) = x^2$ with domain *the negative real numbers* and range *the positive real numbers* $f^{-1}(x) = -\sqrt{x}$ is the negative square root.
3. If $f(x) = \sin(x)$ with domain $[0, 2\pi)$ and range $[-1, 1]$ $f^{-1}(x) = \arcsin(x)$

Examples of Inverse Functions

For some examples:

1. If $f(x) = x^2$ with domain *the positive real numbers* and range *the positive real numbers* $f^{-1}(x) = \sqrt{x}$ is the positive square root.
2. If $f(x) = x^2$ with domain *the negative real numbers* and range *the positive real numbers* $f^{-1}(x) = -\sqrt{x}$ is the negative square root.
3. If $f(x) = \sin(x)$ with domain $[0, 2\pi)$ and range $[-1, 1]$ $f^{-1}(x) = \arcsin(x)$
4. If $f(x) = e^x$ with domain $\mathbb{R}$ and range $(0, \infty)$

Examples of Inverse Functions

For some examples:

1. If $f(x) = x^2$ with domain *the positive real numbers* and range *the positive real numbers* $f^{-1}(x) = \sqrt{x}$ is the positive square root.
2. If $f(x) = x^2$ with domain *the negative real numbers* and range *the positive real numbers* $f^{-1}(x) = -\sqrt{x}$ is the negative square root.
3. If $f(x) = \sin(x)$ with domain $[0, 2\pi)$ and range $[-1, 1]$ $f^{-1}(x) = \arcsin(x)$
4. If $f(x) = e^x$ with domain $\mathbb{R}$ and range $(0, \infty)$ $f^{-1}(x) = \ln(x)$.

Inverse Differentiation I

Question: Using the relation

$$f(f^{-1}(x)) = x \tag{3}$$

how might we relate the derivative of $f(x)$ with the derivative of it's inverse function $f^{-1}(x)$?

Inverse Differentiation I

Question: Using the relation

$$f(f^{-1}(x)) = x \tag{3}$$

how might we relate the derivative of $f(x)$ with the derivative of it's inverse function $f^{-1}(x)$?
Answer: Chain Rule!

Inverse Differentiation I

Question: Using the relation

$$f(f^{-1}(x)) = x \tag{3}$$

how might we relate the derivative of $f(x)$ with the derivative of it's inverse function $f^{-1}(x)$?
Answer: Chain Rule!

$$f(f^{-1}(x)) = x \tag{4}$$

Differentiate both sides (5)

Inverse Differentiation I

Question: Using the relation

$$f(f^{-1}(x)) = x \tag{3}$$

how might we relate the derivative of $f(x)$ with the derivative of it's inverse function $f^{-1}(x)$?
Answer: Chain Rule!

$$f(f^{-1}(x)) = x \tag{4}$$

Differentiate both sides (5)

$$\frac{d}{dx} f(f^{-1}(x)) = \frac{d}{dx} x \tag{6}$$

Inverse Differentiation I

Question: Using the relation

$$f(f^{-1}(x)) = x \tag{3}$$

how might we relate the derivative of $f(x)$ with the derivative of it's inverse function $f^{-1}(x)$?
Answer: Chain Rule!

$$f(f^{-1}(x)) = x \tag{4}$$

Differentiate both sides (5)

$$\frac{d}{dx} f(f^{-1}(x)) = \frac{d}{dx} x \tag{6}$$

$$\left(\frac{d}{dx} f^{-1}(x)\right) \cdot f'(f^{-1}(x)) = 1 \tag{7}$$

Inverse Differentiation I

Question: Using the relation

$$f(f^{-1}(x)) = x \tag{3}$$

how might we relate the derivative of $f(x)$ with the derivative of it's inverse function $f^{-1}(x)$?
Answer: Chain Rule!

$$f(f^{-1}(x)) = x \tag{4}$$

Differentiate both sides (5)

$$\frac{d}{dx} f(f^{-1}(x)) = \frac{d}{dx} x \tag{6}$$

$$\left(\frac{d}{dx} f^{-1}(x)\right) \cdot f'(f^{-1}(x)) = 1 \tag{7}$$

$$\frac{d}{dx} f^{-1}(x) = \frac{1}{f'(f^{-1}(x))} \tag{8}$$

Inverse Differentiation II

Inverse Differentiation

If $f(x)$ and $f^{-1}(x)$ are an inverse pair of functions then:

Inverse Differentiation II

Inverse Differentiation

If $f(x)$ and $f^{-1}(x)$ are an inverse pair of functions then:

$$\frac{d}{dx}f^{-1}(x) = \frac{1}{f'(f^{-1}(x))} \tag{9}$$

Inverse Differentiation II

Inverse Differentiation

If $f(x)$ and $f^{-1}(x)$ are an inverse pair of functions then:

$$\frac{d}{dx}f^{-1}(x) = \frac{1}{f'(f^{-1}(x))} \tag{9}$$

Inverse Differentiation III

Ex. $\frac{d}{dx}\ln(x)$.

Derivative of Natural Logarithm

Inverse Differentiation III

Ex. $\dfrac{d}{dx}\ln(x)$.

$\ln(x)$ is inverse to $e^x \implies f(x) = e^x, f^{-1}(x) = \ln(x)$.

Derivative of Natural Logarithm

Inverse Differentiation III

Ex. $\frac{d}{dx}\ln(x)$.

$\ln(x)$ is inverse to $e^x \implies f(x) = e^x, f^{-1}(x) = \ln(x)$.

$$\frac{d}{dx}\ln(x) = \frac{1}{\left(\frac{d}{du}e^u\right)|_{u=\ln(x)}}$$

Derivative of Natural Logarithm

Inverse Differentiation III

Ex. $\frac{d}{dx}\ln(x)$.

$\ln(x)$ is inverse to $e^x \implies f(x) = e^x, f^{-1}(x) = \ln(x)$.

$$\frac{d}{dx}\ln(x) = \frac{1}{\left(\frac{d}{du}e^u\right)|_{u=\ln(x)}} = \frac{1}{(e^u)|_{u=\ln(x)}} \tag{10}$$

Derivative of Natural Logarithm

Inverse Differentiation III

Ex. $\frac{d}{dx}\ln(x)$.

$\ln(x)$ is inverse to $e^x \implies f(x) = e^x, f^{-1}(x) = \ln(x)$.

$$\frac{d}{dx}\ln(x) = \frac{1}{\left(\frac{d}{du}e^u\right)|_{u=\ln(x)}} = \frac{1}{(e^u)|_{u=\ln(x)}} \tag{10}$$
$$= \frac{1}{e^{\ln(x)}}$$

Derivative of Natural Logarithm

Inverse Differentiation III

Ex. $\frac{d}{dx}\ln(x)$.

$\ln(x)$ is inverse to $e^x \implies f(x) = e^x, f^{-1}(x) = \ln(x)$.

$$\frac{d}{dx}\ln(x) = \frac{1}{\left(\frac{d}{du}e^u\right)|_{u=\ln(x)}} = \frac{1}{(e^u)|_{u=\ln(x)}} \tag{10}$$

$$= \frac{1}{e^{\ln(x)}} = \frac{1}{x} \tag{11}$$

Derivative of Natural Logarithm

Inverse Differentiation III

Ex. $\frac{d}{dx}\ln(x)$.
$\ln(x)$ is inverse to $e^x \implies f(x) = e^x, f^{-1}(x) = \ln(x)$.

$$\frac{d}{dx}\ln(x) = \frac{1}{\left(\frac{d}{du}e^u\right)|_{u=\ln(x)}} = \frac{1}{(e^u)|_{u=\ln(x)}} \tag{10}$$

$$= \frac{1}{e^{\ln(x)}} = \frac{1}{x} \tag{11}$$

Derivative of Natural Logarithm

$$\frac{d}{dx}\ln(x) = \frac{1}{x} \tag{12}$$

Inverse Differentiation IV

Ex. $\dfrac{d}{dx}\arcsin(x)$.

Inverse Differentiation IV

Ex. $\frac{d}{dx}\arcsin(x)$.

$$\frac{d}{dx}\arcsin(x) = \frac{1}{\left(\frac{d}{du}\sin(u)\right)|_{u=\arcsin(x)}}$$

Inverse Differentiation IV

Ex. $\frac{d}{dx}\arcsin(x)$.

$$\frac{d}{dx}\arcsin(x) = \frac{1}{\left(\frac{d}{du}\sin(u)\right)|_{u=\arcsin(x)}} = \frac{1}{(\cos(u))|_{u=\arcsin(x)}} \tag{13}$$

$$= \cos(\arcsin(x))$$

Inverse Differentiation IV

Ex. $\frac{d}{dx}\arcsin(x)$.

$$\frac{d}{dx}\arcsin(x) = \frac{1}{\left(\frac{d}{du}\sin(u)\right)|_{u=\arcsin(x)}} = \frac{1}{(\cos(u))|_{u=\arcsin(x)}} \tag{13}$$

$$= \cos(\arcsin(x)) = \frac{1}{\sqrt{1-x^2}} \tag{14}$$

Inverse Differentiation IV

Ex. $\frac{d}{dx}\arcsin(x)$.

$$\frac{d}{dx}\arcsin(x) = \frac{1}{\left(\frac{d}{du}\sin(u)\right)|_{u=\arcsin(x)}} = \frac{1}{(\cos(u))|_{u=\arcsin(x)}} \tag{13}$$

$$= \cos(\arcsin(x)) = \frac{1}{\sqrt{1-x^2}} \tag{14}$$

$$\frac{d}{dx}\arccos(x) = \frac{1}{-\sin(\arccos(x))}$$

Inverse Differentiation IV

Ex. $\frac{d}{dx}\arcsin(x)$.

$$\frac{d}{dx}\arcsin(x) = \frac{1}{\left(\frac{d}{du}\sin(u)\right)|_{u=\arcsin(x)}} = \frac{1}{(\cos(u))|_{u=\arcsin(x)}} \tag{13}$$

$$= \cos(\arcsin(x)) = \frac{1}{\sqrt{1-x^2}} \tag{14}$$

$$\frac{d}{dx}\arccos(x) = \frac{1}{-\sin(\arccos(x))} = -\frac{1}{\sqrt{1-x^2}} \tag{15}$$

Inverse Differentiation IV

Ex. $\frac{d}{dx}\arcsin(x)$.

$$\frac{d}{dx}\arcsin(x) = \frac{1}{\left(\frac{d}{du}\sin(u)\right)|_{u=\arcsin(x)}} = \frac{1}{(\cos(u))|_{u=\arcsin(x)}} \tag{13}$$

$$= \cos(\arcsin(x)) = \frac{1}{\sqrt{1-x^2}} \tag{14}$$

$$\frac{d}{dx}\arccos(x) = \frac{1}{-\sin(\arccos(x))} = -\frac{1}{\sqrt{1-x^2}} \tag{15}$$

$$\frac{d}{dx}\arctan(x) = \frac{1}{\sec^2(\arctan(x))}$$

Inverse Differentiation IV

Ex. $\frac{d}{dx}\arcsin(x)$.

$$\frac{d}{dx}\arcsin(x) = \frac{1}{\left(\frac{d}{du}\sin(u)\right)|_{u=\arcsin(x)}} = \frac{1}{(\cos(u))|_{u=\arcsin(x)}} \tag{13}$$

$$= \cos(\arcsin(x)) = \frac{1}{\sqrt{1-x^2}} \tag{14}$$

$$\frac{d}{dx}\arccos(x) = \frac{1}{-\sin(\arccos(x))} = -\frac{1}{\sqrt{1-x^2}} \tag{15}$$

$$\frac{d}{dx}\arctan(x) = \frac{1}{\sec^2(\arctan(x))} = \frac{1}{1+x^2} \tag{16}$$

Inverse Differentiation V

Derivatives of Inverse Trigonometric Functions

$$\frac{d}{dx}\arcsin(x) = \frac{1}{\sqrt{1-x^2}} \tag{17}$$

$$\frac{d}{dx}\arccos(x) = -\frac{1}{\sqrt{1-x^2}} \tag{18}$$

$$\frac{d}{dx}\arctan(x) = \frac{1}{1+x^2} \tag{19}$$

Implicit Differentiation I

What about for the other kinds of implicit relations?

Implicit Differentiation I

What about for the other kinds of implicit relations?

Q: At a point (x, y) on the circle of radius 1, what is the slope of the line tangent to the circle at that point?

Implicit Differentiation I

What about for the other kinds of implicit relations?

Q: At a point (x, y) on the circle of radius 1, what is the slope of the line tangent to the circle at that point?

Circle of radius 1 $\implies x^2 + y^2 = 1$.

$$x^2 + y^2 = 1 \tag{20}$$

Differentiate both sides w.r.t. x (21)

Implicit Differentiation I

What about for the other kinds of implicit relations?

Q: At a point (x, y) on the circle of radius 1, what is the slope of the line tangent to the circle at that point?

Circle of radius 1 $\implies x^2 + y^2 = 1$.

$$x^2 + y^2 = 1 \tag{20}$$

$$\text{Differentiate both sides w.r.t. } x \tag{21}$$

$$\frac{d}{dx}(x^2 + y^2) = \frac{d}{dx}1 \tag{22}$$

Implicit Differentiation I

What about for the other kinds of implicit relations?

Q: At a point (x, y) on the circle of radius 1, what is the slope of the line tangent to the circle at that point?

Circle of radius 1 $\implies x^2 + y^2 = 1$.

$$x^2 + y^2 = 1 \tag{20}$$

Differentiate both sides w.r.t. x (21)

$$\frac{d}{dx}(x^2 + y^2) = \frac{d}{dx}1 \tag{22}$$

$$\frac{d}{dx}(x^2) + \frac{d}{dx}(y^2) = 0 \tag{23}$$

Implicit Differentiation I

What about for the other kinds of implicit relations?

Q: At a point (x, y) on the circle of radius 1, what is the slope of the line tangent to the circle at that point?

Circle of radius 1 $\implies x^2 + y^2 = 1$.

$$x^2 + y^2 = 1 \tag{20}$$

Differentiate both sides w.r.t. x (21)

$$\frac{d}{dx}(x^2 + y^2) = \frac{d}{dx}1 \tag{22}$$

$$\frac{d}{dx}(x^2) + \frac{d}{dx}(y^2) = 0 \tag{23}$$

$$2x + \frac{d}{dx}(y(x)^2) = 0 \tag{24}$$

Implicit Differentiation I

What about for the other kinds of implicit relations?

Q: At a point (x, y) on the circle of radius 1, what is the slope of the line tangent to the circle at that point?

Circle of radius 1 $\implies x^2 + y^2 = 1$.

$$x^2 + y^2 = 1 \tag{20}$$

Differentiate both sides w.r.t. x (21)

$$\frac{d}{dx}(x^2 + y^2) = \frac{d}{dx}1 \tag{22}$$

$$\frac{d}{dx}(x^2) + \frac{d}{dx}(y^2) = 0 \tag{23}$$

$$2x + \frac{d}{dx}(y(x)^2) = 0 \tag{24}$$

$$2x + 2y \cdot \frac{dy}{dx} = 0$$

Implicit Differentiation I

What about for the other kinds of implicit relations?

Q: At a point (x, y) on the circle of radius 1, what is the slope of the line tangent to the circle at that point?

Circle of radius 1 $\implies x^2 + y^2 = 1$.

$$x^2 + y^2 = 1 \tag{20}$$

Differentiate both sides w.r.t. x (21)

$$\frac{d}{dx}(x^2 + y^2) = \frac{d}{dx}1 \tag{22}$$

$$\frac{d}{dx}(x^2) + \frac{d}{dx}(y^2) = 0 \tag{23}$$

$$2x + \frac{d}{dx}(y(x)^2) = 0 \tag{24}$$

$$2x + 2y \cdot \frac{dy}{dx} = 0 \quad 2y\frac{dy}{dx} = -2x$$

Implicit Differentiation I

What about for the other kinds of implicit relations?

Q: At a point (x, y) on the circle of radius 1, what is the slope of the line tangent to the circle at that point?

Circle of radius 1 $\implies x^2 + y^2 = 1$.

$$x^2 + y^2 = 1 \tag{20}$$

Differentiate both sides w.r.t. x (21)

$$\frac{d}{dx}(x^2 + y^2) = \frac{d}{dx}1 \tag{22}$$

$$\frac{d}{dx}(x^2) + \frac{d}{dx}(y^2) = 0 \tag{23}$$

$$2x + \frac{d}{dx}(y(x)^2) = 0 \tag{24}$$

$$2x + 2y \cdot \frac{dy}{dx} = 0 \quad 2y\frac{dy}{dx} = -2x \quad \frac{dy}{dx} = -\frac{x}{y} \tag{25}$$

Implicit Differentiation II

The core idea is that in an implicit relation $F(x, y) = 0$,

Implicit Differentiation II

The core idea is that in an implicit relation $F(x, y) = 0$, we can *always* treat one of the variables as a function of the other (fancy reason: *implicit function theorem*)

Implicit Differentiation II

The core idea is that in an implicit relation $F(x, y) = 0$, we can *always* treat one of the variables as a function of the other (fancy reason: *implicit function theorem*) and thus can differentiate implicit relations just by using the chain rule!

Implicit Differentiation II

The core idea is that in an implicit relation $F(x, y) = 0$, we can *always* treat one of the variables as a function of the other (fancy reason: *implicit function theorem*) and thus can differentiate implicit relations just by using the chain rule!
We'll see more application oriented examples when we do related rates soon :)

Worksheet Time

Let’s get into groups and work on the worksheet!

End of Class

That’s all folks!