



XXIV.—*Four new South American curculionidæ (Col.)*

Guy A.K.MarshallC.M.G.D.Sc.F.R.S.

To cite this article: Guy A.K.MarshallC.M.G.D.Sc.F.R.S. (1929) XXIV.—*Four new South American curculionidæ (Col.)*, Journal of Natural History, 4:21, 264-270, DOI: [10.1080/00222932908673050](https://doi.org/10.1080/00222932908673050)

To link to this article: <https://doi.org/10.1080/00222932908673050>



Published online: 26 Aug 2009.



Submit your article to this journal [↗](#)



Article views: 9



View related articles [↗](#)

- Fig. 20. *Hoplichthys langsdorffia*, $\times 3$.
 Fig. 21. *Congiopodus leucopæcilus*, $\times 3$.
 Fig. 22. *Cottus gobio*, $\times 4\frac{1}{2}$.
 Fig. 23. — *scorpius*, $\times 2$.
 Fig. 24. — *bubalis*, $\times 2\frac{1}{2}$.
 Fig. 25. *Neophrynichthys latus*, $\times 3$.
 Fig. 26. *Agonus cataphractus*, $\times 4$.
 Fig. 27. *Cyclopterus lumpus*, $\times 6$.
 Fig. 28. *Cephalacanthus orientalis*, $\times 4$.
 Fig. 29. *Gastrosteus aculeatus*, $\times 20$.

XXIV.—Four new South American Curculionidæ (Col.).

By GUY A. K. MARSHALL, C.M.G., D.Sc., F.R.S.

Subfamily *HYPERINÆ*.

Phelypera pachiræ, sp. n.

♂ ♀. Derm black; pronotum with black scaling and the following markings formed of fulvous scales: on each side a broad outwardly-angulated stripe from apex to base, two small isolated spots in the middle of the disk, some variable spots along the apical margin, and outside the longitudinal stripes some variable patches, which are often united to the stripe so as to form an 8-shaped marking; scutellum black; elytra irregularly variegated with black and fulvous scaling; venter with yellowish-grey scaling, with two black patches on the apical margin of each ventrite on either side, and an indefinite blackish median stripe on ventrites 3 and 4 (5 and 6).

Head closely punctate, with black recumbent hairs and few pale ones on each side, the frons about as wide as the funicle. *Rostrum* stout, gradually widening from base to apex, finely and closely punctate (subconfluently in the basal half), and with an obsolescent median carina, which is more distinct in ♂. *Antennæ* with the two basal joints of the funicle subequal, 3 and 4 equal and somewhat longer than broad, 5–7 moniliform. *Prothorax* transverse, rounded at the sides, subparallel-sided from the base to the middle, thence narrowed to the apex; the base arcuate with the external angles slightly projecting outwardly, the apex truncate; the dorsum gently convex longitudinally and sloping forwards, closely and finely punctate throughout, without any median furrow

or carina. *Elytra* oblong-ovate, with the shoulders prominent; the shallow striae with distinct separated punctures, the intervals broad and almost flat. *Legs* with black scales sparsely intermingled with pale ones, except on the base half of the posterior pairs of femora, where the scales are all pale.

Length 6.6–8.4 mm., breadth 3.6–4.5 mm.

BRAZIL: Bahia, 3 ♂♂, 7 ♀♀ (*Dr. G. Bondar*).

Recorded as damaging the leaves of a tree, *Pachira insignis*.

Very closely allied to *P. schuppeli*, Boh., 1834, which is, however, a rather narrower insect with a red-brown integument; the rostrum is clothed with pale, instead of black, hairs; in the antennæ joints 3–7 of the funicle are all more or less transverse; the colouring of the pronotum is fulvous with three transverse rows each containing four blackish spots; the scutellum is pale, etc.

Subfamily CRYPTORRHYNCHINÆ.

Conotrachelus myrciariæ, sp. n.

♂ ♀. Derm red-brown, with rather thin narrow scaling; the head fulvous; the prothorax with mingled greyish-white and fulvous scales, and with a large indefinite subtriangular darker area in the middle of the base, on which the scales are very sparse and fulvous only; the pleuræ and front coxæ with much denser and broader whitish scales; the elytra sparsely but fairly evenly covered with narrow greyish-white scales, which become fulvous near the base and apex, and with a denser whitish transverse band on the shoulders between intervals 5 and 7; the sternum with sparse narrow pale scales, but the mesepisternum with an oblique stripe of dense, broader, creamy-white scales; the venter with very sparse, pale, short, recumbent setæ.

Head with coarse subconfluent punctures throughout; the frons narrower than the base of the rostrum and with a shallow median fovea; the eyes not completely covered in repose. *Rostrum* of ♂ moderately long and slender, longer than the pronotum, gently curved, and distinctly 5-carinate nearly to the antennæ, which are inserted quite close to the apex in both sexes; rostrum of ♀ similar, except that the carinæ are restricted to the basal half. *Antennæ* elongate, the two basal joints of the funicle equal, 3 longer than 4 and both longer than broad, 5 and 6 as long as broad, and 7 transverse. *Prothorax* slightly broader than long, rounded at the sides, widest about the middle, but little narrowed

behind and with the basal angles somewhat projecting, strongly narrowed in front and with a very shallow subapical constriction; the dorsum convex, coarsely and subconfluently punctate throughout, with a low smooth carina from the apex almost to the base. *Scutellum* ovate, punctate, almost bare. *Elytra* subtriangular, 1.7 times as wide at the rectangular shoulders as the base of the prothorax, gradually narrowed from the shoulders to the apex, and shallowly bisinuate at the base; the punctures deep and distinct, gradually diminishing behind, and each containing a small, narrow, pale scale; intervals 3, 5, 7, 9 carinate throughout, 3 being the highest, 8 carinate from about one-fourth to three-fourths, and 10 for a short distance near the base. *Legs* with shallow indefinite punctation, which becomes coarse and longitudinally confluent at the base of the femora; the pale scaling fairly dense on the femora and sparse on the tibiae; the femora with a large triangular tooth and a small distal one, the hind tibiae of ♂ only with a double apical mucro; all the tibiae, and especially the front pair, with the dorsal margin sinuate just before the apex, so that the apical angle projects. *Sternum*: front coxae almost contiguous; the mesosternum between the coxae sloping and broadly sinuate.

Length 6.0–6.3 mm., breadth 3.0–3.3 mm.

BRAZIL: Bahia, 1 ♂, 1 ♀, 1924 (*Dr. G. Bondar*).

The larvæ of this species feed in the fruits of a wild Myrtaceous tree, *Myrciaria cauliflora*.

In general facies very similar to *C. psidii*, Mshl. (Bull. Ent. Res. xiii. 1922, p. 64, pl. i. fig. 1), but in that species the carina on interval 3 is twice interrupted, the eyes are covered by the postocular lobes, the femora have only a single tooth, etc.

Genus HUARUCUS, nov.

Head not concealed by the pronotum; frons narrower than the base of the rostrum; eyes with a marginal furrow posteriorly and widely separated beneath. *Rostrum* curved; scrobes gradually sloping to beneath the rostrum at the base. *Antennæ* with the scape slender, abruptly clavate, and not quite reaching the eye; funicle 7-jointed; club elliptical, distinctly and transversely segmented. *Prothorax* with distinct postocular lobes. *Scutellum* small but distinct in ♀, but liable to disappear in ♂. *Elytra* ovate, a little wider basally than the base of the prothorax, without any trace of a humeral callus; stria 10 abbreviated. *Wings* rudimentary.

Legs approximately equal in length; femora slender, sub-linear and not toothed, the hind pair not exceeding the apex of the elytra; tibiæ compressed, curved at the base, sub-carinate dorsally, but without lateral carinæ, uncinatæ at the apex and with a double tuft of setæ at the inner apical angle, which bears a stout tooth also in the front pair only; tarsi with joint 1 a little longer than 2+3, 2 subtriangular and as long as broad, the claws free and with a basal lateral seta. *Sternum*: the pectoral canal bounded behind by the mesosternum, its apical margin not overhanging, but sloping backwards and reaching the posterior edge of the median coxæ; the metasternum shorter than the median coxæ, the metepisterna broad and parallel-sided. *Venter* with the intercoxal process ogival and as broad as a coxa; ventrite 2 (4) longer than 3 (5).

Genotype, *Huarucus cacti*, sp. n.


A somewhat isolated genus, which in Faust's key to the South American genera of Cryptorrhynchinae (Stett. Ent. Ztg. 1896, p. 49) would come into the section containing *Parisacalles*, Fst., and *Echinotus*, Fst. The former differs, *inter alia*, in having toothed femora, a median projection in the base of the pronotum, the elytra with only 9 striæ, and the intercoxal process of the venter truncate. *Echinonotus* differs in having the rostrum straight and dorso-ventrally compressed, with the scrobes entirely lateral, and the pronotum concealing the head from above and with an angular emargination in the middle of the base.

Huarucus cacti, sp. n.

♂ ♀. Colour dull black, with rather thin greyish-white or pale fawn scaling above and greyish-white scales beneath.

Head with large close punctures throughout, each containing a scale. *Rostrum* gently curved, parallel-sided for the most part, but widening slightly at both base and apex, the antennæ inserted at about the middle in both sexes; dorsum in ♀ with close, but separated, punctures throughout, except laterally at the base where the punctures are coarser and confluent; dorsum in ♂ much more coarsely and confluent punctate, especially on the apical half, which is more or less subgranulate, and with a shallow longitudinal sulcus just above the scrobe. *Antennæ* with the scape almost bare and the funicle with recumbent setiform scales; the joints of the funicle in order of length: 1, 2, 3, 4, (5, 6, 7), joints 1-4 being longer than broad, 5 and 6 as long as broad, and 7 slightly transverse. *Prothorax* as long as broad in ♀

and usually slightly transverse in ♂, strongly rounded at the sides, widest well behind the middle, narrowing to the base and more so anteriorly, with a shallow subapical constriction; the basal margin truncate and twice as wide as the apex, which is gently arcuate dorsally, the postocular lobes with a dense fringe of short setæ; the dorsum gently convex longitudinally and highest in front of the middle, densely set



Huarucus cacti, sp. n.

throughout with large punctures, with only a much abbreviated median carina (sometimes obsolete); each of the dorsal punctures containing a small ovate or lanceolate scale, those on the pleuræ with much larger and mostly subcircular scales. *Elytra* narrowly ovate or subelliptical, truncate at the base, with the external angles projecting very slightly forwards, widest about the middle, jointly and obtusely rounded at the apex, and with a distinct subapical

lateral constriction; the striæ shallow, containing large quadrate or transverse punctures, which are separated by narrow septa and rapidly diminish behind; the intervals narrow and each with a sparse row of indistinct punctures, intervals 3, 5, 7 carinate and much higher than the others, the lateral intervals also carinate but lower; the scales confined mostly to the sulci, leaving the higher intervals bare; the punctures on the intervals containing inconspicuous recumbent setæ. *Legs* with large shallow separated punctures containing ovate grey scales; the front tibiæ feebly denticulate internally, especially in ♂, in which they are more curved towards the apex; the tarsi densely squamose above, the clothing beneath consisting of dense blackish hairs on joints 1 and 2, but yellow and spongy on 3.

Length 4.2–6.0 mm., *breadth* 9.0–12.6 mm.

PERU: Huaruco, 7 ♂ ♂, 4 ♀ ♀, on cactus, 12. xii. 1928 (R. C. Mundell).

Type, ♂, in the British Museum; cotypes in the United States National Museum.

I am indebted to Mr. H. Morrison, of the U.S. National Museum, for the opportunity of describing this interesting species.

Subfamily BARIDINÆ.

Microstrates ypsilon, sp. n.

♂ ♀. Dorsum black, the lateral and apical margins of the pronotum gradually turning to red-brown; elytra, which are sometimes piceous or red-brown (? immature), with a common basal Y-shaped patch of pale yellowish or brassy scales, which at the base extends on each elytron to stria 3, the stem of the Y reaching only to stria 1 and ending a little before the middle; the pygidium red-brown; the rostrum of ♂ black and reddish at apex, of ♀ testaceous-brown and darker at apex; the lower surface red-brown, with pale scaling that is not very dense; the legs and antennæ paler.

Head opaque, coriaceous, the frons with very shallow reticulate punctation. *Rostrum* of ♂ slightly longer than a front femur, gently curved, very gradually narrowing from base to apex, with longitudinally confluent punctation dorsally and becoming finely carinate laterally; rostrum of ♀ much narrower, almost straight, cylindrical (except for the very short basal dilatation), shiny and entirely impunctate, except for some shallow confluent punctation at the extreme base. *Antennæ* with joint 1 of the funicle nearly as long as 2 + 3 + 4, 2 being shorter than usual, 3–6 subequal and about

as long as broad, 7 a little longer and broader. *Prothorax* slightly longer than broad, gently rounded at the sides, widest at about one-fourth from the base, thence much narrowed in front, with a shallow subapical constriction; the base subtruncate, with the carinate margin interrupted for a short distance in the middle; the apex half the width of the base and produced in the middle into an obtuse horn-like prominence that projects horizontally well beyond the head; the dorsum (except for an indefinite abbreviated impunctate median line) with shallow punctures separated in the middle of the disk by about the length of their own diameters, but becoming longitudinally confluent laterally and in front; the sides of the dorsum sparsely covered with small separated pale scales lying transversely; immediately below this a bare stripe, the rest of the pleuræ being sparsely squamose. *Scutellum* trapezoidal, widest behind, with a few rugose punctures. *Elytra* narrowly ovate, only slightly wider than the prothorax, flattened; the deep striæ with shallow distant punctures, the broad intervals each with a row of small rasp-like punctures, each containing a microscopic recumbent seta. *Legs* with sparse pale scaling; front tibiæ of ♂ with a long fringe of pale hairs on the lower edge, but no fringe on the femora.

Length 3·3–4·8 mm., *breadth* 1·2–1·6 mm.

BRAZIL: Bahia, 7 ♂ ♂, 9 ♀ ♀, in flowers of *Cocos coronata*, 1928 (*Dr. G. Bondar*).

The genotype, *M. cucullus*, Boh., 1844, which is the only other species known to me, is a much larger and proportionately broader insect (6 mm. by 2·5 mm.), has a longer and more elevated prothoracic process, and a dense patch of scales in the basal angles of the thorax; the elytra have only a small patch of scales at the base of interval 2, and the intervals bear numerous simple punctures; and joint 2 of the funicle is proportionately much longer, joint 1 being equal to 2 + 3.

XXV.—*Two Fossil Frogs from the Lower Miocene of Europe.* By H. W. PARKER, B.A.

(Published by permission of the Trustees of the British Museum.)

DIRECT palæontological evidence with a bearing on the phylogeny or former distribution of the Salientia is, unfortunately, very scanty, and consequently every fossil frog, no matter how fragmentary or badly preserved, is worthy of the