

37181: WEEK 3: EUCLIDEAN ALGORITHM, SET THEORY

A/Prof Murray Elder, UTS
Wednesday 7 August 2019

PLAN

- introduction to set theory notation
- Division and remainder lemma
- Euclidean algorithm
- power set

SET THEORY

A *set* is a well-defined collection of objects.¹ The objects are called *elements* of the set, or *members* of the set.

¹Carefully defining what *well-defined* means will take us beyond the scope of this course, into axiomatic set theory and foundations of mathematics.

SET THEORY

A *set* is a well-defined collection of objects.¹ The objects are called *elements* of the set, or *members* of the set.

We can represent a set using brackets, for example
 $A = \{1, 2, a, 5, c, 3\}.$

¹Carefully defining what *well-defined* means will take us beyond the scope of this course, into axiomatic set theory and foundations of mathematics.

SET THEORY

A *set* is a well-defined collection of objects.¹ The objects are called *elements* of the set, or *members* of the set.

We can represent a set using brackets, for example
 $A = \{1, 2, a, 5, c, 3\}$.

The elements are the five symbols you see listed inside the brackets. We could also describe a set using variables satisfying some conditions, for example:

$$A = \{x \mid ((x \in \mathbb{N}) \wedge (1 \leq x \leq 5) \wedge (x \neq 4)) \vee (x = a) \vee (x = c)\}.$$

¹Carefully defining what *well-defined* means will take us beyond the scope of this course, into axiomatic set theory and foundations of mathematics.

SET THEORY

A *set* is a well-defined collection of objects.¹ The objects are called *elements* of the set, or *members* of the set.

We can represent a set using brackets, for example
 $A = \{1, 2, a, 5, c, 3\}$.

The elements are the five symbols you see listed inside the brackets. We could also describe a set using variables satisfying some conditions, for example:

$$A = \{x \mid ((x \in \mathbb{N}) \wedge (1 \leq x \leq 5) \wedge (x \neq 4)) \vee (x = a) \vee (x = c)\}.$$

The set $\{1, 5, 3, c, a, 1, 2\}$ is the same as the set A , since a set is defined only by the elements it contains, no matter how they are listed or displayed.

¹Carefully defining what *well-defined* means will take us beyond the scope of this course, into axiomatic set theory and foundations of mathematics.

The notation $x \in A$ means x is an element of A and $x \notin A$ means $\neg(x \in A)$.

The notation $x \in A$ means x is an element of A and $x \notin A$ means $\neg(x \in A)$.

Formally, if A, B are sets we define $A = B$ if

$$\forall x[x \in A \leftrightarrow x \in B]$$

Eg:

- $A = \{x \mid x \in \mathbb{Q}, x < 0\}$

- $B = \{y \mid y \in \mathbb{R}, y^2 = 2\}$

Test: where does $-\sqrt{2}$ live?

Eg:

- $A = \{x \mid x \in \mathbb{Q}, x < 0\}$
- $B = \{y \mid y \in \mathbb{R}, y^2 = 2\}$

Test: where does $-\sqrt{2}$ live?

Definition

- $A \cap B = \{x \mid x \in A \wedge x \in B\}$ (intersection)
- $A \cup B = \{x \mid x \in A \vee x \in B\}$ (union)

Eg:

- $A = \{x \mid x \in \mathbb{Q}, x < 0\}$
- $B = \{y \mid y \in \mathbb{R}, y^2 = 2\}$

Test: where does $-\sqrt{2}$ live?

Definition

- $A \cap B = \{x \mid x \in A \wedge x \in B\}$ (intersection)
- $A \cup B = \{x \mid x \in A \vee x \in B\}$ (union)

Note the similarity of notation for \cap and \wedge , and \cup and \vee .

SET THEORY

Eg:

- $A = \{x \mid x \in \mathbb{Q}, x < 0\}$
- $B = \{y \mid y \in \mathbb{R}, y^2 = 2\}$

Test: where does $-\sqrt{2}$ live?

Definition

- $A \cap B = \{x \mid x \in A \wedge x \in B\}$ (intersection)
- $A \cup B = \{x \mid x \in A \vee x \in B\}$ (union)

Note the similarity of notation for \cap and \wedge , and \cup and \vee .

In our Eg: $A \cap B =$

YOUR TURN

Let $A = \{a, b, c, d, e\}$, $B = \{b, d, e\}$, $C = \{f, g, a\}$. Find

1. $(A \cup B) \cap (A \cup C)$
2. $A \cap (B \cup C)$

Let $A = \{a, b, c, d, e\}$, $B = \{b, d, e\}$, $C = \{f, g, a\}$. Find

1. $(A \cup B) \cap (A \cup C)$
2. $A \cap (B \cup C)$

A pictorial way to do this exercise is to draw a *Venn diagram*.

If A, B are sets then $A \setminus B = \{x \mid x \in A \wedge x \notin B\}$.

Eg: $A = \{a, b, c, d, e\}, B = \{b, d, e\}, C = \{f, g, a\}$. Find

1. $A \setminus B$
2. $A \setminus C$

MORE NOTATION

If A, B are sets we say A is a *subset* of B if $\forall x \in A, x \in B$, or $(x \in A) \rightarrow (x \in B)$. Notation $A \subseteq B$.

MORE NOTATION

If A, B are sets we say A is a *subset* of B if $\forall x \in A, x \in B$, or $(x \in A) \rightarrow (x \in B)$. Notation $A \subseteq B$.

The notation $A \subsetneq B$ means *strictly contains*:

$$((x \in A) \rightarrow (x \in B)) \wedge (\exists y[y \in B \wedge y \notin A]).$$

MORE NOTATION

If A, B are sets we say A is a *subset* of B if $\forall x \in A, x \in B$, or $(x \in A) \rightarrow (x \in B)$. Notation $A \subseteq B$.

The notation $A \subsetneq B$ means *strictly contains*:

$$((x \in A) \rightarrow (x \in B)) \wedge (\exists y[y \in B \wedge y \notin A]).$$

So $\mathbb{N} \subsetneq \mathbb{Z} \subsetneq \mathbb{Q} \subsetneq \mathbb{R}$.

MORE NOTATION

If A, B are sets we say A is a *subset* of B if $\forall x \in A, x \in B$, or $(x \in A) \rightarrow (x \in B)$. Notation $A \subseteq B$.

The notation $A \subsetneq B$ means *strictly contains*:

$$((x \in A) \rightarrow (x \in B)) \wedge (\exists y[y \in B \wedge y \notin A]).$$

So $\mathbb{N} \subsetneq \mathbb{Z} \subsetneq \mathbb{Q} \subsetneq \mathbb{R}$.

Let \mathcal{U} be some large “universal” set, so we assume all sets we speak about are subsets of \mathcal{U} . Then $\bar{A} = \{x \mid x \notin A\} = \mathcal{U} \setminus A$ means the set of elements in \mathcal{U} that are not in A .

LOGIC VS. SET THEORY

There is a strong connection to the logic we covered before. We have three operations on sets: $\cap, \cup, -$ which we can use to build new sets from old ones, and in logic we have three connectives \wedge, \vee, \neg .

LOGIC VS. SET THEORY

There is a strong connection to the logic we covered before. We have three operations on sets: $\cap, \cup, ^c$ which we can use to build new sets from old ones, and in logic we have three connectives \wedge, \vee, \neg .

Recall the tautologies in logic such as

$$\neg(p \wedge q) \leftrightarrow \neg p \vee \neg q$$

In set theory we could consider sets

$$\overline{A \cap B} \text{ and } \overline{A} \cup \overline{B}.$$

LOGIC VS. SET THEORY

There is a strong connection to the logic we covered before. We have three operations on sets: $\cap, \cup, -$ which we can use to build new sets from old ones, and in logic we have three connectives \wedge, \vee, \neg .

Recall the tautologies in logic such as

$$\neg(p \wedge q) \leftrightarrow \neg p \vee \neg q$$

In set theory we could consider sets

$$\overline{A \cap B} \text{ and } \overline{A} \cup \overline{B}.$$

How do we show two sets are the same? We show they contain exactly the same elements.

LOGIC VS. SET THEORY

There is a strong connection to the logic we covered before. We have three operations on sets: $\cap, \cup, -$ which we can use to build new sets from old ones, and in logic we have three connectives \wedge, \vee, \neg .

Recall the tautologies in logic such as

$$\neg(p \wedge q) \leftrightarrow \neg p \vee \neg q$$

In set theory we could consider sets

$$\overline{A \cap B} \text{ and } \overline{A} \cup \overline{B}.$$

How do we show two sets are the same? We show they contain exactly the same elements.

Formally, if A, B are sets we define $A = B$ if

$$\forall x [x \in A \leftrightarrow x \in B]$$

DE MORGAN (SET VERSION)

Lemma

$$\overline{A \cap B} = \bar{A} \cup \bar{B}.$$

DE MORGAN (SET VERSION)

Lemma

$$\overline{A \cap B} = \bar{A} \cup \bar{B}.$$

The proof goes: pick some arbitrary element of the LHS.

DE MORGAN (SET VERSION)

Lemma

$$\overline{A \cap B} = \bar{A} \cup \bar{B}.$$

The proof goes: pick some arbitrary element of the LHS.

Show it belongs to the RHS.

Lemma

$$\overline{A \cap B} = \bar{A} \cup \bar{B}.$$

The proof goes: pick some arbitrary element of the LHS.

Show it belongs to the RHS.

Since we picked an arbitrary thing, this shows everything in the LHS is also in the RHS, so $\text{LHS} \subseteq \text{RHS}$.

Lemma

$$\overline{A \cap B} = \bar{A} \cup \bar{B}.$$

The proof goes: pick some arbitrary element of the LHS.

Show it belongs to the RHS.

Since we picked an arbitrary thing, this shows everything in the LHS is also in the RHS, so $\text{LHS} \subseteq \text{RHS}$.

Repeat to get $\text{RHS} \subseteq \text{LHS}$, then $\text{LHS} = \text{RHS}$.

DE MORGAN (SET VERSION)

Lemma

$$\overline{A \cap B} = \bar{A} \cup \bar{B}.$$

Proof. Suppose $x \in \overline{A \cap B}$.

DE MORGAN (SET VERSION)

Lemma

$$\overline{A \cap B} = \bar{A} \cup \bar{B}.$$

Proof. Suppose $x \in \overline{A \cap B}$.

Then x is not in $A \cap B$.

DE MORGAN (SET VERSION)

Lemma

$$\overline{A \cap B} = \bar{A} \cup \bar{B}.$$

Proof. Suppose $x \in \overline{A \cap B}$.

Then x is not in $A \cap B$.

Now either $x \in A$ or not. If $x \in A$ then since $x \notin A \cap B$ we must have x is not in B .

DE MORGAN (SET VERSION)

Lemma

$$\overline{A \cap B} = \bar{A} \cup \bar{B}.$$

Proof. Suppose $x \in \overline{A \cap B}$.

Then x is not in $A \cap B$.

Now either $x \in A$ or not. If $x \in A$ then since $x \notin A \cap B$ we must have x is not in B .

So either $x \in \bar{A}$ or $x \in \bar{B}$, so $x \in \bar{A} \cup \bar{B}$.

DE MORGAN (SET VERSION)

Lemma

$$\overline{A \cap B} = \bar{A} \cup \bar{B}.$$

Proof. Suppose $x \in \overline{A \cap B}$.

Then x is not in $A \cap B$.

Now either $x \in A$ or not. If $x \in A$ then since $x \notin A \cap B$ we must have x is not in B .

So either $x \in \bar{A}$ or $x \in \bar{B}$, so $x \in \bar{A} \cup \bar{B}$.

Thus

$$\overline{A \cap B} \subseteq \bar{A} \cup \bar{B}.$$

YOUR TURN

Next, start over and suppose $x \in \overline{A} \cup \overline{B}$.

Thus

$$\overline{A} \cup \overline{B} \subseteq \overline{A \cap B}.$$

Since each set is contained in the other, they are equal.



Show that for any sets $A, B, C \subseteq \mathcal{U}$

$$A \cap (B \cup C) = (A \cap B) \cup (A \cap C).$$

VENN DIAGRAMS ARE NOT PROOFS

Note: a *Venn diagram* can be useful to check if a statement about sets looks correct, or to find a counterexample.

But drawing a picture of a Venn diagram does not constitute a proof
– you must do the LHS, RHS proof.

VENN DIAGRAMS ARE NOT PROOFS

Note: a *Venn diagram* can be useful to check if a statement about sets looks correct, or to find a counterexample.

But **drawing a picture of a Venn diagram does not constitute a proof**
– you must do the LHS, RHS proof.

Eg: check if you think $A \cup (B \cap C) = (A \cup B) \cap C$ is true or not.

PAUSE

RECALL

An element s in a subset $S \subseteq \mathbb{N}$ is called a *first element* in S if $s \leq x$ for every $x \in S$.

Eg: $\{5, 4, 6, 7\}$ has a first element, 4.

RECALL

An element s in a subset $S \subseteq \mathbb{N}$ is called a *first element* in S if $s \leq x$ for every $x \in S$.

Eg: $\{5, 4, 6, 7\}$ has a first element, 4.

Lemma

First elements are unique.

RECALL

An element s in a subset $S \subseteq \mathbb{N}$ is called a *first element* in S if $s \leq x$ for every $x \in S$.

Eg: $\{5, 4, 6, 7\}$ has a first element, 4.

Lemma

First elements are unique. (So we can say “the” first element).

RECALL

An element s in a subset $S \subseteq \mathbb{N}$ is called a *first element* in S if $s \leq x$ for every $x \in S$.

Eg: $\{5, 4, 6, 7\}$ has a first element, 4.

Lemma

First elements are unique. (So we can say “the” first element).

Axiom (Well ordering principle)

Every non-empty subset of \mathbb{N} has a first element.

axiom = fact which does not follow from other facts.

APPLICATION: DIVISION AND REMAINDER

Lemma

Let $n, d \in \mathbb{Z}$ with $d > 0$. Then there exist $q, r \in \mathbb{Z}$ with $0 \leq r < d$ such that $n = qd + r$.

Proof:

APPLICATION: DIVISION AND REMAINDER

Lemma

Let $n, d \in \mathbb{Z}$ with $d > 0$. Then there exist $q, r \in \mathbb{Z}$ with $0 \leq r < d$ such that $n = qd + r$.

Proof: Define $M = \{n - qd \mid q \in \mathbb{Z}\}$.

APPLICATION: DIVISION AND REMAINDER

Lemma

Let $n, d \in \mathbb{Z}$ with $d > 0$. Then there exist $q, r \in \mathbb{Z}$ with $0 \leq r < d$ such that $n = qd + r$.

Proof: Define $M = \{n - qd \mid q \in \mathbb{Z}\}$. Then $M \cap \mathbb{N}$ is a subset of \mathbb{N} .

APPLICATION: DIVISION AND REMAINDER

Lemma

Let $n, d \in \mathbb{Z}$ with $d > 0$. Then there exist $q, r \in \mathbb{Z}$ with $0 \leq r < d$ such that $n = qd + r$.

Proof: Define $M = \{n - qd \mid q \in \mathbb{Z}\}$. Then $M \cap \mathbb{N}$ is a subset of \mathbb{N} .

It is non-empty because if $n \geq 0$ you can take $q = 0$ and if $n < 0$ take $q = 100n$ (which is a negative number, so $-qd$ is a big positive number).

APPLICATION: DIVISION AND REMAINDER

Lemma

Let $n, d \in \mathbb{Z}$ with $d > 0$. Then there exist $q, r \in \mathbb{Z}$ with $0 \leq r < d$ such that $n = qd + r$.

Proof: Define $M = \{n - qd \mid q \in \mathbb{Z}\}$. Then $M \cap \mathbb{N}$ is a subset of \mathbb{N} .

It is non-empty because if $n \geq 0$ you can take $q = 0$ and if $n < 0$ take $q = 100n$ (which is a negative number, so $-qd$ is a big positive number).

Therefore by the well ordering principle $M \cap \mathbb{N}$ has a first element, call it r .

Since $r \in M \cap \mathbb{N}$ we have $r \geq 0$ and $r = n - qd$ for some $q \in \mathbb{Z}$.

Since $r \in M \cap \mathbb{N}$ we have $r \geq 0$ and $r = n - qd$ for some $q \in \mathbb{Z}$.

If $r \geq d$ (for contradiction) then $r - d \geq 0$ and $r - d = n - (q + 1)d$ so belongs to $M \cap \mathbb{N}$, and is smaller than r , contradicting our choice of r as first element. \square

APPLICATION OF DIVISION LEMMA

Definition

Let $a, b \in \mathbb{Z}$. Then $d \in \mathbb{N}$ is called the *greatest common divisor* of a and b if $d \mid a$, $d \mid b$, and if $c \mid a$, $c \mid b$ then $c \mid d$.

Eg: compute

- $\gcd(3, 9)$
- $\gcd(6, 8)$

APPLICATION OF DIVISION LEMMA

Definition

Let $a, b \in \mathbb{Z}$. Then $d \in \mathbb{N}$ is called the *greatest common divisor* of a and b if $d \mid a$, $d \mid b$, and if $c \mid a$, $c \mid b$ then $c \mid d$.

Eg: compute

- $\gcd(3, 9)$
- $\gcd(6, 8)$

The following algorithm claims to compute \gcd . It is called the *Euclidean algorithm*. We should not believe this claim, until we know how to prove algorithms are correct (lecture 6):

1. stops 2. gives the correct output

Input 54, 187.

Use the lemma to write $187 = q_1 \cdot 54 + r_1$.

Input 54, 187.

Use the lemma to write $187 = q_1 \cdot 54 + r_1$.

Use the lemma to write $54 = q_2 \cdot r_1 + r_2$.

Repeat until you get $r_i = 0$.

YOUR TURN

Input 154, 287.

Use the lemma to write $287 = q \cdot 154 + r$.

Repeat until you get $r = 0$.

ONE MORE PROOF

Lemma

Let $n, d \in \mathbb{Z}$ with $d > 0$. Then there exist unique integers q, r with $0 \leq r < d$ such that $n = qd + r$.

Proof.

ONE MORE PROOF

Lemma

Let $n, d \in \mathbb{Z}$ with $d > 0$. Then there exist unique integers q, r with $0 \leq r < d$ such that $n = qd + r$.

Proof.

We already proved some q, r values exist. Suppose they are not unique.

ONE MORE PROOF

Lemma

Let $n, d \in \mathbb{Z}$ with $d > 0$. Then there exist unique integers q, r with $0 \leq r < d$ such that $n = qd + r$.

Proof.

We already proved some q, r values exist. Suppose they are not unique.

Then we have q_1, q_2, r_1, r_2 and $n = q_1d + r_1 = q_2d + r_2$ so

ONE MORE PROOF

Lemma

Let $n, d \in \mathbb{Z}$ with $d > 0$. Then there exist unique integers q, r with $0 \leq r < d$ such that $n = qd + r$.

Proof.

We already proved some q, r values exist. Suppose they are not unique.

Then we have q_1, q_2, r_1, r_2 and $n = q_1d + r_1 = q_2d + r_2$ so $r_1 - r_2 = d(q_2 - q_1)$.

This means d divides $r_1 - r_2$, but since they are both between 0 and $d - 1$ we must have $r_1 - r_2 = 0$, so $r_1 = r_2$ and then $q_1 - q_2 = 0$ so $q_1 = q_2$.



BACK TO THE DEFINITION OF “SET”

The next exercise explains why *well-defined collection of objects* is not quite good enough.

BACK TO THE DEFINITION OF “SET”

The next exercise explains why *well-defined collection of objects* is not quite good enough.

Let $P(S)$ be the property (of sets) that S does not contain itself. For example, $P(\mathbb{N})$ is true because \mathbb{N} contains numbers, it does not contain sets so it cannot contain itself.

BACK TO THE DEFINITION OF “SET”

The next exercise explains why *well-defined collection of objects* is not quite good enough.

Let $P(S)$ be the property (of sets) that S does not contain itself. For example, $P(\mathbb{N})$ is true because \mathbb{N} contains numbers, it does not contain sets so it cannot contain itself.

Another example: the *empty set* \emptyset is the set that has no elements, $\emptyset = \{\}$. So it contains nothing so cannot contain itself.

BACK TO THE DEFINITION OF “SET”

The next exercise explains why *well-defined collection of objects* is not quite good enough.

Let $P(S)$ be the property (of sets) that S does not contain itself. For example, $P(\mathbb{N})$ is true because \mathbb{N} contains numbers, it does not contain sets so it cannot contain itself.

Another example: the *empty set* \emptyset is the set that has no elements, $\emptyset = \{\}$. So it contains nothing so cannot contain itself.

(a) Give some more examples.

BACK TO THE DEFINITION OF “SET”

Consider the set of all abstract concepts. Call it \mathcal{A} . Then \mathcal{A} contains things like art, postmodernism, democracy, imaginary numbers.

BACK TO THE DEFINITION OF “SET”

Consider the set of all abstract concepts. Call it \mathcal{A} . Then \mathcal{A} contains things like art, postmodernism, democracy, imaginary numbers.

(b) Which is true: $\mathcal{A} \in \mathcal{A}$ or $\mathcal{A} \notin \mathcal{A}$?

BACK TO THE DEFINITION OF “SET”

Consider the set of all abstract concepts. Call it \mathcal{A} . Then \mathcal{A} contains things like art, postmodernism, democracy, imaginary numbers.

(b) Which is true: $\mathcal{A} \in \mathcal{A}$ or $\mathcal{A} \notin \mathcal{A}$?

Let $\mathcal{S} = \{S \mid P(S)\}$ be the set of all sets that do not contain themselves.

BACK TO THE DEFINITION OF “SET”

Consider the set of all abstract concepts. Call it \mathcal{A} . Then \mathcal{A} contains things like art, postmodernism, democracy, imaginary numbers.

(b) Which is true: $\mathcal{A} \in \mathcal{A}$ or $\mathcal{A} \notin \mathcal{A}$?

Let $\mathcal{S} = \{S \mid P(S)\}$ be the set of all sets that do not contain themselves.

So $\mathbb{N} \in \mathcal{S}$ and $\mathcal{A} \notin \mathcal{S}$.

BACK TO THE DEFINITION OF “SET”

Consider the set of all abstract concepts. Call it \mathcal{A} . Then \mathcal{A} contains things like art, postmodernism, democracy, imaginary numbers.

(b) Which is true: $\mathcal{A} \in \mathcal{A}$ or $\mathcal{A} \notin \mathcal{A}$?

Let $\mathcal{S} = \{S \mid P(S)\}$ be the set of all sets that do not contain themselves.

So $\mathbb{N} \in \mathcal{S}$ and $\mathcal{A} \notin \mathcal{S}$.

(c) Which is true: $\mathcal{S} \in \mathcal{S}$ or $\mathcal{S} \notin \mathcal{S}$?

BACK TO THE DEFINITION OF “SET”

Consider the set of all abstract concepts. Call it \mathcal{A} . Then \mathcal{A} contains things like art, postmodernism, democracy, imaginary numbers.

(b) Which is true: $\mathcal{A} \in \mathcal{A}$ or $\mathcal{A} \notin \mathcal{A}$?

Let $\mathcal{S} = \{S \mid P(S)\}$ be the set of all sets that do not contain themselves.

So $\mathbb{N} \in \mathcal{S}$ and $\mathcal{A} \notin \mathcal{S}$.

(c) Which is true: $\mathcal{S} \in \mathcal{S}$ or $\mathcal{S} \notin \mathcal{S}$?

The moral of this story: you cannot define a set using a condition, in general. *i.e.* $\{x \mid P(x)\}$ may not actually be a well-defined collection of objects.

POWER SET

Let A be a set. Then (axiom)

$$\mathcal{P}(A) = \{B \mid B \subseteq A\}$$

is a set. Its called the *power set* of A .

POWER SET

Let A be a set. Then (axiom)

$$\mathcal{P}(A) = \{B \mid B \subseteq A\}$$

is a set. Its called the *power set* of A .

Questions:

- is $\emptyset \in \mathcal{P}(A)$?
- is $A \in \mathcal{P}(A)$?
- is $\mathcal{P}(A) \in \mathcal{P}(A)$?

POWER SET

Let A be a set. Then (axiom)

$$\mathcal{P}(A) = \{B \mid B \subseteq A\}$$

is a set. Its called the *power set* of A .

Questions:

- is $\emptyset \in \mathcal{P}(A)$?
- is $A \in \mathcal{P}(A)$?
- is $\mathcal{P}(A) \in \mathcal{P}(A)$?

Another axiom: \emptyset is a set.

POWER SET

Let A be a set. Then (axiom)

$$\mathcal{P}(A) = \{B \mid B \subseteq A\}$$

is a set. Its called the *power set* of A .

Questions:

- is $\emptyset \in \mathcal{P}(A)$?
- is $A \in \mathcal{P}(A)$?
- is $\mathcal{P}(A) \in \mathcal{P}(A)$?

Another axiom: \emptyset is a set.

What can you build with just these two axioms?

YOUR TURN

- Given $A = \{1, 2, 3\}$ is a set, what is $\mathcal{P}(A)$?

YOUR TURN

- Given $A = \{1, 2, 3\}$ is a set, what is $\mathcal{P}(A)$?
- Prove that if A is a set then $A \subseteq \mathcal{P}(A)$

Next lecture:

- induction
- correctness of computer code
- relations and functions