

Q1. 22 January Shift 2

Identify the correct statements :

- A. Hydrated salts can be used as primary standard.
- B. Primary standard should not undergo any reaction with air.
- C. Reactions of primary standard with another substance should be instantaneous and stoichiometric.
- D. Primary standard should not be soluble in water.
- E. Primary standard should have low relative molar mass.

Choose the correct answer from the options given below :

- (1) A, B, C and E Only
- (2) A, B and E Only
- (3) A, B and C Only
- (4) D and E Only

Q2. 23 January Shift 2

200 cc of $x \times 10^{-3}$ M potassium dichromate is required to oxidise 750 cc of 0.6 M Mohr's salt solution in acidic medium. Here $x =$ ____.

Q3. 24 January Shift 1

X and Y are the number of electrons involved, respectively during the oxidation of I^- to I_2 and S^{2-} to S by acidified $K_2Cr_2O_7$. The value of $X + Y$ is ____.

Q4. 24 January Shift 2

One mole of Cl_2 (g) was passed into 2 L of cold 2 M KOH solution. After the reaction, the concentrations of Cl^- , ClO^- and OH^- are respectively (assume volume remains constant)

- (1) 0.5M, 0.5M, 1M
- (2) 0.75M, 0.75M, 1M
- (3) 0.5M, 0.5M, 0.5M
- (4) 1M, 1M, 1M

Q5. 28 January Shift 1

500 mL of 1.2 M KI solution is mixed with 500 mL of 0.2M $KMnO_4$ solution in basic medium. The liberated iodine was titrated with standard 0.1M $Na_2S_2O_3$ solution in the presence of starch indicator till the blue color disappeared. The volume (in L) of $Na_2S_2O_3$ consumed is _____. (Nearest integer)

ANSWER KEYS

1. (3)

2. 375

3. 12

4. (1)

5. 3