

All My Work

Your saved learning record for Cellular Reproduction.

7 completed activity runs. Required checks and formative practice are recorded separately.

Unfinished work: 0 required check(s), 0 practice run(s). Resume them from their original sections; they are not counted as completed.

Writing and flash-card self-assessments are not automatically graded. Completion is not a mastery certificate.

Alternation of generations

9/17/2026, 8:41:29 PM · Elapsed 0m 0s · Formative practice

Review submitted responses and attempts

Label A

Attempt 1: Sporophyte (2n) — Correct

Reference: Sporophyte (2n)

Label B

Attempt 1: Meiosis — Correct

Reference: Meiosis

Label C

Attempt 1: Spore (n) — Correct

Reference: Spore (n)

Label D

Attempt 1: Gametophyte (n) — Correct

Reference: Gametophyte (n)

Label E

Attempt 1: Gametes (n) — Correct

Reference: Gametes (n)

Label F

Attempt 1: Fertilization — Correct

Reference: Fertilization

Nondisjunction: compare the gamete outcomes

9/17/2026, 8:41:29 PM · Elapsed 0m 0s · Formative practice

Review submitted responses and attempts

Label A

Attempt 1: Meiosis I nondisjunction — Correct

Reference: Meiosis I nondisjunction

Label B

Attempt 1: Meiosis II nondisjunction — Correct

Reference: Meiosis II nondisjunction

Crossing over involves non-sister chromatids

9/17/2026, 8:41:29 PM · Elapsed 0m 0s · Formative practice

Review submitted responses and attempts

Label A

Attempt 1: Non-sister chromatids — Correct

Reference: Non-sister chromatids

Label B

Attempt 1: Parental chromatid — Correct

Reference: Parental chromatid

Label C

Attempt 1: Recombinant chromatid — Correct

Reference: Recombinant chromatid

Label D

Attempt 1: Parental chromatid — Correct

Reference: Parental chromatid

Meiosis: separate homologues, then sisters

9/17/2026, 8:41:28 PM · Elapsed 0m 0s · Formative practice

Review submitted responses and attempts

Label A

Attempt 1: Replicated diploid cell — Correct

Reference: Replicated diploid cell

Label B

Attempt 1: Haploid cells after meiosis I — Correct

Reference: Haploid cells after meiosis I

Label C

Attempt 1: Haploid cells after meiosis II — Correct

Reference: Haploid cells after meiosis II

Mitosis in a model cell: $2n = 4$

9/17/2026, 8:41:28 PM · Elapsed 0m 0s · Formative practice

Review submitted responses and attempts

Label A

Attempt 1: Prophase — Correct

Reference: Prophase

Label B

Attempt 1: Metaphase — Correct

Reference: Metaphase

Label C

Attempt 1: Anaphase — Correct

Reference: Anaphase

Label D

Attempt 1: Telophase — Correct

Reference: Telophase

One chromosome before and after replication

9/17/2026, 8:41:28 PM · Elapsed 0m 0s · Formative practice

Review submitted responses and attempts

Label A

Attempt 1: One unreplicated chromosome — Correct

Reference: One unreplicated chromosome

Label B

Attempt 1: Sister chromatid — Correct

Reference: Sister chromatid

Label C

Attempt 1: Sister chromatid — Correct

Reference: Sister chromatid

Label D

Attempt 1: Centromere region — Correct

Reference: Centromere region

Chromosomes, DNA and chromosome sets

9/17/2026, 8:41:25 PM · Elapsed 0m 0s · Required check

Review submitted responses and attempts

First-try multiple choice: 1/2. Writing is not automatically graded.

Which statement best describes homologous chromosomes?

Options: They have corresponding genes but may carry different alleles · They must have identical DNA sequences throughout · They exist only in bacterial cells · They are always two chromatids joined at one centromere

Attempt 1: They must have identical DNA sequences throughout — Incorrect

Attempt 2: They have corresponding genes but may carry different alleles — Correct

A diploid species has $2n = 10$. How many chromosome types are present in one normal gamete?

Options: 2 · 5 · 10 · 20

Attempt 1: 5 — Correct

Compare chromatin and condensed chromosomes without suggesting that a cell gains DNA only when it divides.

Real-origin saved Chapter 16 work. Chromatin is DNA associated with proteins. It is often less condensed between divisions and becomes tightly packaged into visible chromosomes during division. The change is packaging of existing genetic material, not DNA appearing for the first time.

Submitted writing; human review required.

Explain why a homologous chromosome pair is not the same as a pair of sister chromatids.

Homologous chromosomes typically originate from the two parents and carry corresponding genes, potentially with different alleles. Sister chromatids are the replicated copies of one chromosome and remain associated until separation.

Submitted writing; human review required.

Frayer work

No Frayer words selected.

Collected optional responses

No optional written responses collected.

Textbook question work

Draft and saved question responses. Optional and ungraded.

In Text · p. 551 · question 1 — Saved

Real-origin saved Chapter 16 work. Chromatin is DNA associated with proteins. It is often less condensed between divisions and becomes tightly packaged into visible

chromosomes during division. The change is packaging of existing genetic material, not DNA appearing for the first time. Edited.

Supported and independent practice