



Reviews



Multivariate Linear Regression



Size in feet ² (x)	Price (\$) in 1000's (y)
2104	460
1416	232
1534	315
852	178
...	...

Size (feet ²)	Number of bedrooms	Number of floors	Age of home (years)	Price (\$1000)
2104	5	1	45	460
1416	3	2	40	232
1534	3	2	30	315
852	2	1	36	178
...



Hypothesis:

$$h_{\theta}(x) = \theta^T x = \theta_0 x_0 + \theta_1 x_1 + \theta_2 x_2 + \cdots + \theta_n x_n$$

Parameters:

$$\theta_0, \theta_1, \dots, \theta_n$$

Cost Function:

$$J(\theta_0, \theta_1, \dots, \theta_n) = \frac{1}{2m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2$$

Gradient Descent:

$$\theta_j := \theta_j - \alpha \frac{\partial}{\partial \theta_j} J(\theta_0, \dots, \theta_n)$$

Tips for Gradient Descent

(1) Feature Scaling

Idea: Make sure features are on a similar scale.


E.g. X_1 = size (0-2000 feet²)

X_2 = number of bedrooms (1-5)

↓ Normalize

E.g. X_1 = size (0-2000 feet²) / 2000

X_2 = number of bedrooms (1-5) / 5



(2) Making sure gradient descent is working correctly

Idea: Plot it



$$\frac{1}{2m} \sum_{i=1}^m (h_\theta(x^{(i)}) - y^{(i)})^2$$

Gradient Descent 梯度下降

$$\theta_j := \theta_j - \alpha \frac{\partial}{\partial \theta_j} J(\theta_0, \dots, \theta_n)$$

Normal Equation 正规方程

$$\theta = (X^T X)^{-1} X^T y$$



Normal Equation 正规方程

	Size (feet ²)	Number of bedrooms	Number of floors	Age of home (years)	Price (\$1000)
x_0	x_1	x_2	x_3	x_4	
1	2104	5	1	45	460
1	1416	3	2	40	232
1	1534	3	2	30	315
1	852	2	1	36	178

$$X = \begin{bmatrix} 1 & 2104 & 5 & 1 & 45 \\ 1 & 1416 & 3 & 2 & 40 \\ 1 & 1534 & 3 & 2 & 30 \\ 1 & 852 & 2 & 1 & 36 \end{bmatrix}$$

$$y = \begin{bmatrix} 460 \\ 232 \\ 315 \\ 178 \end{bmatrix}$$

$$\theta = (X^T X)^{-1} X^T y$$



Classification



Bi-Classification

- Email: Spam / Not Spam?
- Online Transactions: Fraudulent (Yes / No)?
- Tumor: Malignant / Benign ?

$$y \in \{0, 1\}$$

Multi-Classification

Linear Regression

$h_\theta(x)$ can be > 1 or < 0

Logistic Regression

$0 \leq h_\theta(x) \leq 1$

Classification!

Logistic Regression


$$0 \leq h_{\theta}(x) \leq 1$$

$$h_{\theta}(x) = g(\theta^T x)$$

$$g(z) = \frac{1}{1 + e^{-z}}$$


Sigmoid function
Logistic function

estimated probability
that $y = 1$ on input x




Decision Boundary

estimated probability
that $y = 1$ on input x



If $h \geq 0.5$, predict $y = 1$

If $h < 0.5$, predict $y = 0$




$$h_\theta(x) = g(\theta_0 + \theta_1 x_1 + \theta_2 x_2)$$

$-3, 1, 1$

Predict " $y = 1$ " if $-3 + x_1 + x_2 \geq 0$

Non-linear decision boundaries



$$h_{\theta}(x) = g(\theta_0 + \theta_1 x_1 + \theta_2 x_2 + \theta_3 x_1^2 + \theta_4 x_2^2)$$

-1, 0, 0, 1, 1

More complex decision boundary

$$h_{\theta}(x) = g(\theta_0 + \theta_1 x_1 + \theta_2 x_2 + \theta_3 x_1^2 + \theta_4 x_1^2 x_2 + \theta_5 x_1^2 x_2^2 + \theta_6 x_1^3 x_2 + \dots)$$



Cost Function

Training set

$$\{(x^{(1)}, y^{(1)}), (x^{(2)}, y^{(2)}), \dots, (x^{(m)}, y^{(m)})\}$$

Cost Function
in Linear Regression



$$J(\theta) = \frac{1}{m} \sum_{i=1}^m \frac{1}{2} (h_\theta(x^{(i)}) - y^{(i)})^2$$

Cost Function
in Logistic Regression

$$\text{Cost}(h_\theta(x), y) = \begin{cases} -\log(h_\theta(x)) & \text{if } y = 1 \\ -\log(1 - h_\theta(x)) & \text{if } y = 0 \end{cases}$$

$$\text{Cost}(h_\theta(x), y) = \begin{cases} -\log(h_\theta(x)) & \text{if } y = 1 \\ -\log(1 - h_\theta(x)) & \text{if } y = 0 \end{cases}$$

$$\text{Cost}(h_\theta(x), y) = -\log(h_\theta(x)) * y -\log(1 - h_\theta(x)) * (1-y)$$





Gradient Descent

$$J(\theta) = -\frac{1}{m} \left[\sum_{i=1}^m y^{(i)} \log h_\theta(x^{(i)}) + (1 - y^{(i)}) \log (1 - h_\theta(x^{(i)})) \right]$$

Want $\min_{\theta} J(\theta)$:



Repeat {

$$\theta_j := \theta_j - \alpha \sum_{i=1}^m (h_\theta(x^{(i)}) - y^{(i)}) x_j^{(i)}$$

}

(simultaneously update all θ_j)

One-Vs-ALL






2021 Spring: Machine Learning


Regularized Linear/Logistic Regression

Lecturer: Min Lu
Email: lumin.vis@gmail.com



Size of house

$$\theta_0 + \theta_1 x + \theta_2 x^2$$



Size of house

$$\theta_0 + \theta_1 x + \theta_2 x^2 + \theta_3 x^3 + \theta_4 x^4$$



Suppose we penalize and make these two small

$$\min_{\theta} \frac{1}{2m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2 + 1000 * \theta_3^2 + 1000 * \theta_4^2$$

Regularization

Small values for parameters

- “Simpler” hypothesis
- Less prone to overfitting

$$J(\theta) = \frac{1}{2m} \left[\sum_{i=1}^m (h_\theta(x^{(i)}) - y^{(i)})^2 + \lambda \sum_{j=1}^n \theta_j^2 \right]$$

$$\min_{\theta} J(\theta)$$

Regularization
parameter

Regularized Gradient descent

Repeat {

$$\theta_0 := \theta_0 - \alpha \frac{1}{m} \sum_{i=1}^m (h_\theta(x^{(i)}) - y^{(i)}) x_0^{(i)}$$

$$\theta_j := \theta_j - \alpha \frac{1}{m} \sum_{i=1}^m (h_\theta(x^{(i)}) - y^{(i)}) x_j^{(i)}$$




$$\theta_j := \theta_j \boxed{\left(1 - \alpha \frac{\lambda}{m}\right)} - \alpha \frac{1}{m} \sum_{i=1}^m (h_\theta(x^{(i)}) - y^{(i)}) x_j^{(i)}$$

}

Regularized Normal Equation

$$\theta = \left(X^T X + \lambda \begin{bmatrix} 0 & & & \\ & 1 & & \\ & & 1 & \\ & & & \ddots \\ & & & 1 \end{bmatrix} \right)^{-1} X^T y$$

$$X = \begin{bmatrix} (x^{(1)})^T \\ \vdots \\ (x^{(m)})^T \end{bmatrix} \quad y = \begin{bmatrix} y^{(1)} \\ \vdots \\ y^{(m)} \end{bmatrix}$$



hypothesis function

$$h_\theta(x) = g(\theta_0 + \theta_1 x_1 + \theta_2 x_1^2 + \theta_3 x_1^2 x_2 + \theta_4 x_1^2 x_2^2 + \theta_5 x_1^2 x_2^3 + \dots)$$

cost function

$$J(\theta) = - \left[\frac{1}{m} \sum_{i=1}^m y^{(i)} \log h_\theta(x^{(i)}) + (1 - y^{(i)}) \log (1 - h_\theta(x^{(i)})) \right]$$

Regularized Gradient descent

Repeat {

$$\theta_0 := \theta_0 - \alpha \frac{1}{m} \sum_{i=1}^m (h_\theta(x^{(i)}) - y^{(i)}) x_0^{(i)}$$

$$\theta_j := \theta_j - \alpha \frac{1}{m} \sum_{i=1}^m (h_\theta(x^{(i)}) - y^{(i)}) x_j^{(i)}$$



$$\theta_j := \theta_j \boxed{\left(1 - \alpha \frac{\lambda}{m}\right)} - \alpha \frac{1}{m} \sum_{i=1}^m \boxed{(h_\theta(x^{(i)}) - y^{(i)})} x_j^{(i)}$$

}