


Monte Carlo Diagnostics


Method: Monte Carlo
Fit method: leastsq
Parameters: 11
Requested samples: 500
Completed samples: 500
chisqr mean: 763.35
chisqr median: 765.19
chisqr min: 651.43
chisqr max: 830.64


Correlation threshold: |r| >= 0.5
[KEX_AB] vs [PB]: r=-0.520
[R2_A, NUC->55N, B0->600.3MHZ] vs [R2_A, NUC->65N, B0->600.3MHZ]: r=0.513



Sample distribution: [DW_AB, NUC->14N]
95% percentile
median
Density
0
5
10
15
20
6.56
6.58
6.60
6.62
6.64
6.66
6.68
6.70
6.72
[DW_AB, NUC->14N]



Sample distribution: [DW_AB, NUC->55N]
95% percentile
median
Density
20.0
17.5
15.0
12.5
10.0
7.5
5.0
2.5
0.0
4.64
4.66
4.68
4.70
4.72
4.74
4.76
[DW_AB, NUC->55N]



Sample distribution: [DW_AB, NUC->65N]
95% percentile
median
Density
0.0
2.5
5.0
7.5
10.0
12.5
15.0
17.5
4.04
4.06
4.08
4.10
4.12
4.14
[DW_AB, NUC->65N]



Sample distribution: [KEX_AB]
95% percentile
median
Density
0.00
0.01
0.02
0.03
0.04
400
410
420
430
440
450
460
[KEX_AB]



Sample distribution: [PB]
95% percentile
median
Density
10000
8000
6000
4000
2000
0
0.003400
0.003425
0.003450
0.003475
0.003500
0.003525
0.003550
[PB]



Sample distribution: [R1_A, NUC->14N, B0->600.3MHZ]
95% percentile
median
Density
0
20
40
60
80
1.950
1.955
1.960
1.965
1.970
1.975
1.980
[R1_A, NUC->14N, B0->600.3MHZ]



Sample distribution: [R1_A, NUC->55N, B0->600.3MHZ]
95% percentile
median
Density
0
10
20
30
40
50
60
70
80
1.585
1.590
1.595
1.600
1.605
1.610
[R1_A, NUC->55N, B0->600.3MHZ]



Sample distribution: [R1_A, NUC->65N, B0->600.3MHZ]
95% percentile
median
Density
0
20
40
60
80
100
120
1.595
1.600
1.605
1.610
1.615
[R1_A, NUC->65N, B0->600.3MHZ]



Sample distribution: [R2_A, NUC->14N, B0->600.3MHZ]
95% percentile
median
Density
0
1
2
3
4
5
10.45
10.50
10.55
10.60
10.65
10.70
10.75
10.80
10.85
[R2_A, NUC->14N, B0->600.3MHZ]


Sample distribution: [R2_A, NUC->55N, B0->600.3MHZ]
95% percentile
median
Density
0
1
2
3
4
5
6
7
9.75
9.80
9.85
9.90
9.95
10.00
[R2_A, NUC->55N, B0->600.3MHZ]


Sample distribution: [R2_A, NUC->65N, B0->600.3MHZ]
95% percentile
median
Density
0
2
4
6
8
10
12
9.200
9.225
9.250
9.275
9.300
9.325
9.350
9.375
9.400
[R2_A, NUC->65N, B0->600.3MHZ]


chisqr distribution
median
Samples
chisqr
0
10
20
30
40
50
60
70
80
650
675
700
725
750
775
800
825


Sample pair: [KEX_AB] vs [PB] (r = -0.520)
0.00356
0.00354
0.00352
0.00350
0.00348
0.00346
0.00344
0.00342
0.00340
[PB]
400
410
420
430
440
450
[KEX_AB]
12
10
8
6
4
2
0
Samples


pair: [R2_A, NUC->55N, B0->600.3MHZ] vs [R2_A, NUC->65N, B0->600.3MHZ] (r = 0.513
[R2_A, NUC->65N, B0->600.3MHZ]
9.375
9.350
9.325
9.300
9.275
9.250
9.225
9.75
9.80
9.85
9.90
9.95
10.00
[R2_A, NUC->55N, B0->600.3MHZ]
Samples
20
15
10
5
0