

NN_DebarajBarua_NareshKumarGurulingan_061117

November 11, 2017

1 Hochschule Bonn-Rhein-Sieg

2 Neural Networks, WS17/18

3 Assignment 05 (06-november-2017)

3.1 Debaraj Barua, Naresh Kumar Gurulingan

```
In [2]: import numpy as np
import matplotlib.pyplot as plt
import sympy as sp
import IPython
sp.init_printing()
```

3.2 Question 1:

Read Haykin NN edition 2 : chapter 3.1-3.5. Make a summary

3.3 Question 2 (Haykin Edition:2, Ex: 3.1):

Explore the method of steepest descent involving a single weight w by considering the following cost function:

$$\varepsilon(w) = \frac{1}{2}\sigma^2 - r_{xd}w + \frac{1}{2}r_xw^2$$

where, σ^2, r_{xd} and r_x are constants.

Answer 2:

```
In [3]: s,r_xd,r_x,w=sp.symbols('s,r_xd,r_x,w')

e=1./2.*s**2-r_xd*w+1./2.*r_x*w**2

nabla=sp.diff(e,w)
nabla
```

Out [3] :

$$1.0r_xw - r_{xd}$$

So, Gradient,

$$\begin{aligned}\nabla \varepsilon(w) &= \frac{\partial \varepsilon(w)}{\partial w} \\ &= -r_{xd} + r_x w\end{aligned}$$

According to steepest descent method, weight vector is updated in the direction of the steepest descent, i.e., in the direction opposite to the gradient vector.

So, $\Delta w = -\eta \nabla \varepsilon(w)$, where η is the learning rate

Now, updated weight is:

$$\begin{aligned}w(n+1) &= w(n) + \Delta w \\ &= w(n) - \eta(-r_{xd} + r_x w)\end{aligned}$$

3.4 Question 3 (Haykin Edition:2, Ex: 3.2):

Consider the cost function

$$\varepsilon(w) = \frac{1}{2}\sigma^2 - r_{xd}^T w + \frac{1}{2}w^T R_x w$$

where, σ^2, r_{xd} and r_x are constants.

$$r_{xd} = \begin{bmatrix} 0.8182 \\ 0.354 \end{bmatrix}$$
$$R_x = \begin{bmatrix} 1 & 0.8182 \\ 0.8182 & 1 \end{bmatrix}$$

3.4.1 3(a)

Find the optimum value of w^* for which $\varepsilon(w)$ reaches it's minimum value.

Answer 3(a): $\varepsilon(w)$ will reach maximum value when it's Jacobian will be zero.

```
In [11]: s=sp.symbols('s')
w=sp.MatrixSymbol('w',2,1)
r_xd=sp.Matrix([[0.8182],[0.354]])
R_x=sp.Matrix([[1,0.8182],[0.8182,1]])

e=1./2.*s**2-r_xd.T.dot(sp.Matrix(w))+1./2.*sp.Matrix(w).T.dot(R_x.dot(sp.Matrix(w)))
e=sp.Matrix([e])
symbolic_jacobian=e.jacobian(w)
print "Jacobian Matrix is:"
symbolic_jacobian
```

Jacobian Matrix is:

Out[11]:

$$\begin{bmatrix} -0.8182 + 0.5(2w_{0,0} + 1.6364w_{1,0}) & -0.354 + 0.5(1.6364w_{0,0} + 2w_{1,0}) \end{bmatrix}$$

```
In [12]: print "optimum values of weight, w*:"  
         sp.Matrix([sp.solve(symbolic_jacobian, sp.Matrix(w))])
```

optimum values of weight, w*:

Out[12]:

$$\left[\{w_{0,0} : 1.59902944424901, \quad w_{1,0} : -0.954325891284541\} \right]$$

3.4.2 3(b)

Use the method of steepest descent to compute w^* for the following two values of learning rate parameter:

<p>(i) $\eta=0.3$ </p>


<p>(ii) $\eta=1.0$ </p>

For each case, plot the trajectory traced the evolution of the weight vector $w(n)$ in the W-Plane.


Note: The trajectories obtained for cases (i) and (ii) of part (b) should correspond to the picture in Fig. 3.2

```
In [187]: IPython.display.Image("images/HaykinFig3.2.png")
```

Out[187]:



(a)



(b)

FIGURE 3.2 Trajectory of the method of steepest descent in a two-dimensional space for two different values of learning-rate parameter: (a) $\eta = 0.3$, (b) $\eta = 1.0$. The coordinates w_1 and w_2 are elements of the weight vector \mathbf{w} .

Answer 3(b):

In [188]: *#Lambda function for substitution of weight into symbolic jacobian matrix.*
`delta_wt_func=sp.lambdify((w),symbolic_jacobian)`

```
def steepest_desc(weight, eta, max_iteration):
    found_optimum=False
    iteration=0
    wt_matrix=np.array(weight).T
```

```

print "Learning Rate: ", eta
while (found_optimum==False and iteration<max_iteration):
    iteration+=1
    delta_wt=delta_wt_func(weight)

    if(abs(delta_wt.all())<1e-5):
        """
        If absolute change in weight is less than 10^(-5),
        we consider we have reached the optimal solution
        """
        print "Reached optimum value"
        print "Absolute change in weight is less than 10^(-5), exiting loop."
        found_optimum=True
        break
    else:
        weight+=-eta*delta_wt.T
        wt_matrix=np.vstack((wt_matrix,weight.T))

if iteration==max_iteration:
    print "Max iteration reached!"
print "optimum values of weight, w*:",np.array(weight).T
print "Total iterations required:",iteration
print wt_matrix.shape
plt.plot(wt_matrix[:,0],wt_matrix[:,1])
plt.xlabel("w1(n)")
plt.ylabel("w2(n)")
plt.title(("Learning Rate",eta))
plt.show()


initial_weight = sp.Matrix([[0],[0]])
print initial_weight.shape
eta1 = 0.3
eta2 = 1.0
max_iteration =1000
steepest_desc(initial_weight,eta1,max_iteration)
steepest_desc(initial_weight,eta2,max_iteration)

```

```

(2, 1)
Learning Rate:  0.3
Reached optimum value
Absolute change in weight is less than 10^(-5), exiting loop.
optimum values of weight, w*: [[1.59897673946915 -0.954273186504682]]
Total iterations required: 181
(181, 2)

```



Learning Rate: 1.0


Reached optimum value

Absolute change in weight is less than 10^{-5} , exiting loop.

optimum values of weight, w*: $[[1.59899794732667 \ -0.954307093426509]]$

Total iterations required: 55

(55, 2)



3.5 Question 4 (Haykin Edition:2, Ex: 3.4):

The correlation matrix R_x of the input vector $x(n)$ in the LMS algorithm is defined by

$$R_x = \begin{bmatrix} 1 & 0.5 \\ 0.5 & 1 \end{bmatrix}$$

Define the range of values for the learning-rate parameter η of the LMS algorithm for it to be convergent in the mean square.

Answer 4: LMS algorithm is convergent in mean square in the below mentioned region:

$$0 < \eta < \frac{2}{\lambda_{max}}$$

Where, λ_{max} is the largest eigen value of the correlation matrix of the input vector.

```
In [191]: R_x=np.array([[1,0.5],[0.5,1]])

          eigval,eigvec=np.linalg.eig(R_x)
          max_eigvalue=eigval[np.argmax(eigval)]

          print "Range of values for learning rate: [0,",2./max_eigvalue,")"
```

Range of values for learning rate: [0, 1.33333333333]

3.6 Question 5 (Haykin Edition:2, Ex: 3.8):

The ensemble-averaged counterpart of the sum of error squares viewed as a cost function is the mean-square value of the error signal:

$$\begin{aligned} J(w) &= \frac{1}{2} E[e^2(n)] \\ &= \frac{1}{2} E[(d(n) - x^T(n)w)^2] \end{aligned}$$

3.6.1 5(a)

Assuming that the input vector $x(n)$ and the desired response $d(n)$ are drawn from a stationary environment, show that

$$J(w) = \frac{1}{2} \sigma_d^2 - r_{xd}^T w + \frac{1}{2} w^T R_x w$$

where,

$$\begin{aligned} \sigma_d^2 &= E[d^2(n)] \\ r_{xd} &= E[x(n)d(n)] \\ R_x &= E[x(n)x^T(n)] \end{aligned}$$

3.6.2 Answer:

Given:

$$\begin{aligned} J(w) &= \frac{1}{2} E[(d(n) - x^T(n)w)^2] \\ &= \frac{1}{2} E[d^2(n) - 2 \cdot d(n)x^T(n)w + (x^T(n)w)^2] \end{aligned}$$

Since expectation is a linear operator:

$$\begin{aligned} &= \frac{1}{2} E[d^2(n)] - \frac{1}{2} \cdot 2 \cdot E[d(n)x^T(n)w] + \frac{1}{2} E[(x^T(n)w)^2] \\ &= \frac{1}{2} E[d^2(n)] - E[d(n)x^T(n)w] + \frac{1}{2} E[(x^T(n)w)^T(x^T(n)w)] \\ &= \frac{1}{2} E[d^2(n)] - E[d(n)x^T(n)w] + \frac{1}{2} E[w^T x(n)x^T(n)w] \\ &= \frac{1}{2} E[d^2(n)] - E[d(n)x^T(n)]w + \frac{1}{2} w^T E[x(n)x^T(n)]w \end{aligned}$$

Substituting $\sigma_d^2, r_{xd}^T = E[d(n)x^T(n)]$ and R_x , we have:

$$J(w) = \frac{1}{2}\sigma_d^2 - r_{xd}^T w + \frac{1}{2}w^T R_x w$$

which is the required proof.

3.6.3 5(b)

For this cost function, show that the gradient vector and the Hessian matrix of $J(w)$ are as follows, respectively:

$$\begin{aligned} g &= -r_{xd} + R_x w \\ H &= R_x \end{aligned}$$

3.6.4 Answer:

The gradient vector is obtained by partial differentiation of $J(w)$ with respect to w :

$$\nabla J(w) = g = -r_{xd} + R_x w \quad (1)$$

The Hessian matrix is the partial derivative of gradient vector with respect to w :

$$H = \nabla g = R_x \quad (2)$$

3.6.5 5(c)

In the *LMS/Newton algorithm*, the gradient vector g is replaced by its instantaneous value (*Widrow and Stearns, 1985*). Show that this algorithm, incorporating a learning rate parameter η , is described by

$$\hat{w}(n+1) = \hat{w}(n) + \eta R_x^{-1} x(n)(d(n) - x^T(n)w(n))$$

The inverse of the correlation matrix R_x , assumed to be positive definite, is calculated ahead of time.

3.6.6 Answer:

From Newton's Method:

$$\Delta w = -H^{-1}g \quad (3)$$

For the given cost function, from 5(b), we know that Hessian matrix is:

$$H = R_x \quad (4)$$

We also know the instantaneous value of the gradient vector is given by:

$$\hat{g} = -x(n)(e(n)) \quad (5)$$

$$\hat{g} = -x(n)(d(n) - x^T(n)w(n)) \quad (6)$$

Therefore,

$$\Delta w = -R_x^{-1}(-x(n)(d(n) - x^T(n)w(n))) \quad (7)$$

$$\Delta w = R_x^{-1}x(n)(d(n) - x^T(n)w(n)) \quad (8)$$

Hence, after incorporating learning rate η ,

$$\hat{w}(n+1) = \hat{w}(n) + \eta R_x^{-1}x(n)(d(n) - x^T(n)w(n))$$

which is the required proof.
