


Recap:

Fish sorting problem / classification

Binary



④ dist $(\underline{z}_{qj}, \pi_d)$ \rightarrow confidence
 \doteq
d-dim



$$\frac{\underline{w^T z_{qj} + w_0}}{\|w\|}$$

x_1 lies on Td

$\checkmark \quad w^T x_1 + w_0 = 0$

$\checkmark \quad -\|w\| \|x_1\| \cos \theta + w_0 = 0$

$(8^\circ) > \theta > 90^\circ$

$\cos \theta < 0$

$\cos(180 - \theta) = -\cos(\theta)$

$\checkmark \quad Anvesha$

$(geom + algebra)$

$\checkmark \quad$

$w^T x_1 + w_0 = 0$

$w^T x_2 + w_0 = 0$

$\cos \theta (\|w\| \|x_1\| + \|w\| \|x_2\|) < 0$

$\cos \theta (\|w\| \|x_1\| + \|w\| \|x_2\|) < 0$

$Td: w^T x + w_0 = 0$

$w^T x_2 + w_0 \leq 0$

≤ 0

1


2

3/3

(Q) * SVM

d-dim

$$\left\{ \begin{array}{l} \text{normal} \\ \pi_1: \underline{\underline{\omega^T x + w_1}} = 0 \\ \pi_2: \underline{\underline{\omega^T x + w_2}} = 0 \end{array} \right.$$



$$d = \frac{\|\omega_2 - \omega_1\|}{\|\omega\|}$$

$\xrightarrow{\text{abs}}$


$\rightarrow \|\omega_2 - \omega_1\|$

\rightarrow

$\|\omega_2 - \omega_1\|$

 $\|\omega\|$

- ① $\pi_1 \parallel \pi_2$
- ② $\text{dist}(\pi_1, \pi_2)$



3 Types → $T_1, T_2 \text{ & } T_3$

Simplest way

$\pi_1 M_1 \rightarrow T_1 \text{ or not}$

$\pi_2 M_2 \rightarrow T_2 \text{ or not}$


$\pi_3 M_3 \rightarrow T_3 \text{ or not}$

one vs rest

(Q)

Plane \rightarrow ✓ normal or \perp to π_d (w)

dist (o, π_d) : $\frac{w_o}{\|w\|}$




can we define $a \pi_d =$

$d(o, \pi_d)$ &

one direction that
is \parallel to the π_d


$$\begin{aligned}\mathbf{x}_1 \cdot \mathbf{x}_2 &= \|\mathbf{x}_1\| \|\mathbf{x}_2\| \cos \theta \\ &= \|\mathbf{x}_1\| \|\mathbf{x}_2\|\end{aligned}$$



Vector: lines

✓ Unique \perp direction

3D




$$\begin{aligned} w \perp w_1 \\ w \perp w_2 \end{aligned}$$

:
→ on the plane
direction
x parallel to π_d

w, w_d
↓
Tl d






$\omega \perp \pi\theta$


$\omega_1 \parallel \pi\theta$

$\omega_2 \perp \omega_1$

$\omega_2 \not\parallel \pi\theta$



Unit -vector



$$\|\omega\| = 10 \text{ (let)}$$

$\hat{\omega}$ has same dir as ω

$$\|\hat{\omega}\| = 1$$

Diagram illustrating the components of a vector w :

Top part: A vector w is shown as a green circle with a pink arrow pointing from the origin. A red box encloses the vector, and a green arrow points down to its center. The text "2-dim" is written next to it.

Middle part: The vector w is divided by its norm $\|w\|$ to obtain a unit vector \hat{w} , which is shown as a green circle with a pink arrow pointing upwards.

Bottom part: The norm $\|w\|$ is calculated as the square root of the sum of the squares of the components: $\|w\| = \sqrt{w_1^2 + w_2^2 + \dots + w_d^2}$.

Equations:

$$\frac{w}{\|w\|} = [w_1 \ w_2 \ \dots \ w_d]^T$$
$$\|w\| = \sqrt{\frac{w_1^2}{l^2} + \frac{w_2^2}{l^2} + \dots + \frac{w_d^2}{l^2}}$$

$$\|\vec{\omega}\| = \sqrt{\left(\frac{\omega_1}{\omega}\right)^2 + \left(\frac{\omega_2}{\omega}\right)^2 + \dots + \left(\frac{\omega_d}{\omega}\right)^2}$$

$$= \sqrt{\frac{\omega_1^2}{\omega_1^2 + \omega_2^2 + \dots} + \frac{\omega_2^2}{\omega_1^2 + \omega_2^2 + \dots} + \dots}$$

$$= \sqrt{1 + 1}$$


Q

$$\omega = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$$

$$\hat{\omega} = \begin{bmatrix} \frac{1}{\sqrt{14}} \\ \frac{2}{\sqrt{14}} \\ \frac{3}{\sqrt{14}} \end{bmatrix}$$


Diagram showing the vector ω originating from the origin O and pointing towards the vector $\hat{\omega}$.

$$\|\omega\| = \sqrt{1+4+9} = \sqrt{14}.$$





(Q)



$$\begin{aligned}x_2' &= \text{Proj}_{x_1}(x_2) \\&= \text{len} \times \text{direclm} \\&\Rightarrow \|x_2\| \cos \theta \cdot \frac{x_1}{\|x_1\|}\end{aligned}$$






$$\mathbf{x}_1 + \mathbf{x}_2 = \begin{bmatrix} x_{11} \\ x_{12} \\ \vdots \\ x_{1d} \end{bmatrix} + \begin{bmatrix} x_{21} \\ x_{22} \\ \vdots \\ x_{2d} \end{bmatrix} = \begin{bmatrix} x_{11} + x_{21} \\ x_{12} + x_{22} \\ \vdots \\ x_{1d} + x_{2d} \end{bmatrix}$$

addition / sub → component-wise

$x_1 \in \mathbb{R}^d$ $K \neq d$ $x_2 \in \mathbb{R}^K$ $x_1 + x_2 = ?$

~~dot-product~~

2 unit vec



{ Geom: angle

between 2 vec

ML \rightarrow planes; dir; dist

len of proj of x_1 onto x_2

$$l = \|x_1\| \cos \theta = \frac{x_1 \cdot x_2}{\|x_2\|}$$

$$\cos \theta = \frac{x_1 \cdot x_2}{\|x_1\| \|x_2\|}$$


$$\frac{\hat{x}_1 \cdot \hat{x}_2}{\|\hat{x}_1\| \|\hat{x}_2\|} = \cos \theta$$

$$\checkmark \theta = \cos^{-1} (\hat{x}_1 \cdot \hat{x}_2)$$



planes (\overline{Td}) → Simplest sep in d -dim space

\rightsquigarrow Sep → Circles/spheres / hyper spheres




circle:

(2D)

not often
used in ML

→ planes with
transformations

(log-transform)



$$\sqrt{x^2 + y^2} = r$$

$$x^2 + y^2 = r^2$$


circle through origin

$d\text{-dim:}$

{Center $\in \mathbb{R}^d$
radius $= \gamma$

$$\|\vec{cx}\| = \sqrt{\begin{bmatrix} x_1 - c_1 \\ x_2 - c_2 \\ \vdots \\ x_d - c_d \end{bmatrix}^T \begin{bmatrix} x_1 - c_1 \\ x_2 - c_2 \\ \vdots \\ x_d - c_d \end{bmatrix}} = \gamma$$

$S_d:$



any x on the S_d

$$x(x_1, x_2, \dots, x_d)$$

$x \in \mathbb{R}^d$
 $c \in \mathbb{R}^d$

$$S_d: \|\vec{cx}\| = \gamma$$

$$\vec{oc} + \vec{cx} = \vec{ox}$$

$$\vec{cx} = \vec{ox} - \vec{oc}$$

$$\|\vec{cx}\| = \|\vec{ox} - \vec{oc}\| = \gamma$$

$$o = [0, p, \dots, 0]^T$$

$$\left\| \begin{bmatrix} x_1 - c_1 \\ x_2 - c_2 \\ \vdots \\ x_d - c_d \end{bmatrix} \right\| = \gamma$$

$[x_1 \dots x_d]^T$ on S^d

$$\Rightarrow \boxed{\sqrt{(x_1 - c_1)^2 + (x_2 - c_2)^2 + \dots + (x_d - c_d)^2} = \gamma}$$

⇒




S_d : Separation




inside / outside

$$\sum_{i=1}^d (x_i - c_i)^2 - r^2 = 0$$

\geq



$$\sum_{i=1}^d (x_i - c_i)^2 - r^2 \leq 0$$



✓ Bi^z-problem (Fish-Sorting (classfn))



intuition (geometry)



1st time

✓ { Learn
Math
(CG, LA, Calc)



solve the problem & write

ML


\rightarrow numerical

fish:

len, width;

width

d-dim space



s_1

s_2

s_3

s_h

Shape
 (S_1, S_2, S_3)

let

Categorical features (discrete r.v.)




\rightarrow Type! or $Type^2$

$$\left\{ \begin{array}{l} S_1=1 \\ S_2=2 \\ S_3=3 \end{array} \right.$$

problem: ordering
 artificially

$$\left\{ \begin{array}{l} S_1=2 \\ S_3=100 \\ S_2=1000 \end{array} \right.$$

3-models:




dim-~~of~~ My Space

len ; wid ;


numerical
len, width; shape
 S_1, S_2, S_3

5-dim

d -dim space



5-dim Space







Q

$x_2 = \langle l, w, s \rangle$

\downarrow


$[l, 2, 0, 1, 0] \quad J^T$

One-hot encoding



Given:

$$\mathcal{D} = \left\{ (x_i, y_i) \right\}_{i=1}^{n=1000}; \quad \begin{array}{l} \text{d-dim} \\ \text{feature} \\ \text{vector} \end{array}$$




One strategy:
Brute force Extremely
(Terrible)

$(d+1) \rightarrow$ Variables

$$\text{TD} : \quad \mathbf{w}^T \mathbf{x} + w_0 \quad \downarrow \quad \downarrow \\ d\text{-dim}$$

∞ -many combinations




\max dist of points from TD

3D-space

$l,$ \downarrow $1\text{D}\text{im}$ color
 c_1, c_2 \downarrow $2\text{D}\text{im}$


$$\begin{aligned}x_1 &= 10, c_1 \\x_2 &= 5, c_2 \rightarrow (5, 0, 1)\end{aligned}$$



fish \rightarrow color
 c_1, c_2, c_3


$x_1 \rightarrow c_1 \rightarrow (1, 0, 0)$


$x_2 \rightarrow c_3 \rightarrow (0, 0, 1)$











For all -ve labelled points ($y_i < 0$)

$$\textcircled{1} \rightarrow w^T x_j + w_0 < 0$$

$$\textcircled{2} d_j = \frac{w^T x_j + w_0}{\|w\|}$$

d_j 's are -ve

d_j 's \rightarrow large -ve values

+Ve labelled: $y_i = +1$

$$\frac{w^T x_i + w_0}{\|w\|} \cdot y_i \xrightarrow{+1} \text{maximal}$$

-Ve labelled: $y_i = -1$

$$\frac{w^T x_i + w_0}{\|w\|} \cdot y_i \xrightarrow{-1} \text{maximal}$$

Find
Td s.t

for each point x_i


$$z_i = \frac{\underline{w^T x_i + w_0}}{\|w\|} \cdot y_i + t_1 / -1$$

$x_i \rightarrow y_i$

Maximal
(as large as possible)

$$\mathcal{D} = \left\{ (x_i, y_i) \right\}_{i=1}^{n=1000} \quad \dots \quad \}$$

params
(args)



Simultaneously

$$\text{Max } (z_1 + z_2 + \dots + z_n) = \text{Max}_{\underline{w}, \underline{w_0}} \sum_{i=1}^n z_i$$

∇w

∇w_0

$\nabla \theta$

$\text{Max}_{\underline{w}, \underline{w_0}} \sum_{i=1}^n y_i (\underline{w}^T \underline{x}_i + \underline{w_0})$

$\nabla \|\underline{w}\|$

optimization

Q

Max z_1, z_2, \dots, z_n Simultaneously

$$\checkmark \text{ Max } \sum_{i=1}^n z_i$$

Q Why not

$$\max z_1 \cdot z_2 \cdot z_3 \cdots z_n$$

\downarrow

$$\max \prod_{i=1}^n z_i$$

x₃ lies on
Tl

$\mathcal{D} = \{(x_i, y_i)\}_{i=1}^n$

$$\max_{w, w_0} f(w, w_0)$$

$$f(w, w_0) = \sum_{i=1}^n y_i \frac{w^T x_i + w_0}{\|w\|}$$


dotted

\downarrow

$$\max_{w, w_0} f(w, w_0)$$

\mathcal{D} $\xrightarrow{\text{d+var}}$ d+var

$$= \min_{w, w_0} -f(w, w_0)$$



one-var

$$f(x) = x^2 - 2x + 5$$

find x that minimizes $f(x)$

Calculus in one-var
multivariable calc

11/15 8/2 h⁻¹

$$\min_x f(x) = x^2 - 2x + 5$$

①

$$\frac{df(x)}{dx} = 2x - 2 = 0$$

\Rightarrow

$$x = \underline{\underline{1}}$$


②

$$\frac{d^2f(x)}{dx^2} = \underline{\underline{2}} > 0$$

minimal-val of
 $f(x)$


next class


why



Maxima & minima

g3
class





You have left the meeting

We get frequent requests for your notes!

Notes written by you helps in understanding the topic better. You can upload the notes in two simple steps mentioned below

- 1 Scan the QR code with your iPad**
Scanner should be present in the top menu on your iPad
 - 2 Upload Notes on the generated link**
All notes uploaded will be visible in the saved version of this session



- OR

Drag and drop files or [click here to upload](#)

Files Uploaded from your computer appear here