

This is a repository copy of *Individual differences and the multidimensional nature of face perception*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/185951/>

Version: Accepted Version

Article:

White, David and Burton, A. Mike orcid.org/0000-0002-2035-2084 (2022) Individual differences and the multidimensional nature of face perception. *Nature Reviews Psychology*. pp. 287-300. ISSN 2731-0574

<https://doi.org/10.1038/s44159-022-00041-3>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

1 **Individual differences and the multidimensional nature of face perception**

2 **David White¹ and A. Mike Burton^{1, 2, *}**

3 ¹School of Psychology, UNSW Sydney, Australia

4 ²Department of Psychology, University of York, York, UK

5 ³Faculty of Society & Design, Bond University, Gold Coast, Australia

6

7 * e-mail: mike.burton@york.ac.uk

8

9 **Acknowledgements**

10 DW's contribution was supported by funding from the Australian Research Council (Future Fellowship FT200100353, Discovery Project DP190100957).

11 **Author contributions**

12 The authors contributed equally to all aspects of the article.

13 **Competing interests**

14 The authors declare no competing interests

15 **Peer review information**

16 *Nature Reviews Psychology* thanks [Referee#1 name], [Referee#2 name] and the other, anonymous, reviewer(s) for their contribution to the peer review of this
17 work.

18 **Publisher's note**

19 Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

ABSTRACT

Face perception is critical to social interactions, yet people vary in how easily they can recognise their friends, verify an identification document, or notice someone's smile. There are widespread differences in people's abilities to recognise faces and research has particularly focused on exceptionally good or poor recognition performance. In this Review, we synthesise literature on individual differences in face processing across different tasks including identification and estimates of emotional state and social attributes. The individual differences approach has considerable untapped potential for theoretical progress in understanding the perceptual and cognitive organisation of face processing. This approach also has practical consequences — for example, in determining who is best suited to check passports. We also discuss the underlying structural and anatomical predictors of face perception ability. Furthermore, we highlight problems of measurement that pose challenges for the effective study of individual differences. Finally, we note that research in individual differences rarely addresses perception of familiar faces. Despite people's everyday experience of being 'good' or 'bad' with faces, a theory of how people recognise their friends remains elusive.

[H1] Introduction

Faces provide many types of information. Using faces, people can recognise others they know and also quite accurately estimate the age, gender or health of strangers and friends. Faces can also be used to judge transient states, for example someone's mood, focus of attention, or speech patterns. The multiple sources of information available in a face are critical for social behaviour, enabling people to identify someone as family, friend or colleague, and to decide whether they should speak to, hug, or stay away from them. These decisions are made quickly and easily, often without reflection. However, these remarkable abilities in face processing are not distributed equally across people.

Widespread differences in face perception between individuals have become an important research focus for three reasons. First, everyday experience suggests that some people are better at face perception than others, and some people have strong beliefs about whether they are 'good with faces'. Second, given the wide variety of information available in a face, much theoretical work focuses on whether the processes underlying different perceptual decisions are independent¹. Individual differences techniques are well-suited to addressing these questions in both face processing^{2,3}, and other areas of cognition research^{4,5}. Third, face processing research has a good track record of using converging evidence to build theory⁶⁻⁸. Individual differences methodology can be recruited alongside evidence from experimental psychology, neuroscience, neuropsychology and computational modelling to make significant progress in the field.

Research on individual differences in face processing has tended to focus on identity processing. Tests have been designed that measure performance on a relatively small, narrowly defined set of tasks, which often do not capture the richness of daily life. Nevertheless, differences in performance reveal a multidimensional face processing system and its links to broader perceptual and cognitive processes.

In this Review, we explore why some people are better at certain face tasks than others, and whether this variation helps to uncover the fundamental processes involved. We summarise the approaches to measuring individual differences and the instruments developed across the broad range of face perception tasks. We then consider the notion of 'holistic' processing — the tendency for faces to be perceived as unitary objects rather than as a collection of parts⁹ — and describe how studies of individual differences inform the representational codes supporting face perception. We highlight the differences between perceiving familiar and unfamiliar faces, and point out the relative dearth of individual differences studies of familiar face processing. Finally, we review the practical issues that emerge for face processing in professional settings. Overall, we provide a snapshot of an approach that contributes to our understanding of face perception and offers opportunities for insights that complement converging evidence in the field.

Throughout the Review, we use the term ‘face recognition’ to denote the process by which someone is identified from their face. Recognition of identity is just one component of ‘face perception’, a term that covers face processing for multiple purposes, including decisions about somebody’s expression, age, or attractiveness, as well as their identity. We use the term ‘face processing’ to refer to any of the perceptual, cognitive or neural processes underpinning face perception for any purpose. Finally, person perception refers to the processing of perceptual information across a whole person including, for example, someone’s voice or gait.

[H1] Converging research on face perception

Whereas the majority of individual differences research focuses on face recognition¹⁰, there is also individual variation in other aspects of face perception^{11,12}. Knowing how these different abilities co-vary in the population can help reveal the structural and processing constraints that shape the face processing system.

The traditional research emphasis in face perception has been on performance in clinical populations or group-level analysis of average measures, in particular to detect transient changes induced by experimental manipulation. By contrast, a focus on the natural variation between people’s face processing abilities represents a fundamental shift in research focus. Although individual differences research presents challenges for measuring natural variation, it also adds novel tools. As in other areas of psychology, a single face perception study rarely fully resolves an issue through a conclusive experiment or simulation. Instead, converging evidence is highly valued, and this is the spirit in which individual differences techniques are becoming popular. Here we describe three major approaches taken by individual differences researchers.

[H3] Extreme abilities

Much of the individual differences research literature derives from the observation that there are some individuals who are unusually poor or unusually good at face recognition tasks. In addition to acquired prosopagnosia (the inability to recognise faces as a result of brain damage), there are also people who show poor recognition ability throughout life in the absence of known organic cause, a condition known as developmental prosopagnosia^{13–15} (or sometimes ‘congenital prosopagnosia’: see Box 1). These individuals typically report an inability to recognise familiar people on the basis of their face alone, sometimes termed ‘face blindness’. Research on developmental prosopagnosia is complemented by studies on ‘super-recognisers’^{16–18}, individuals with extremely high face recognition ability relative to

the average person. These people often report that they can recognise previously seen faces in challenging conditions, for example in poor lighting, and despite not having encountered them for many years.

Recruiting participant groups with extreme abilities enables group-level comparisons between these extreme profiles and average performers. This approach differs substantially from traditional neuropsychological case studies in which an individual is compared to a single control individual or a control group. Group studies on participants selected from extremes of the ability spectrum can offer high statistical power for detecting differences and have the potential to provide insight into the fundamental nature of that ability. This approach can be used to establish dissociations between different subtypes of face processing ability¹⁹, or between face processing and other types of ability. However, the increased power gained by group comparisons relies on a degree of homogeneity within groups and people with developmental prosopagnosia can sometimes display rather diverse symptoms.

[H3] Variation within the normal range

There is growing awareness of the value of individual differences research across the entire range of face processing abilities. Whereas face perception variability is sometimes studied to identify target groups (for example, to establish a range of abilities from which one might select people to perform face-related tasks such as checking passports) it is more commonly used to establish associations or independence between variables of interest. The main tool for establishing the relationship between two tasks is correlation – deriving a quantitative measure of the association between performance on two or more different tasks. This method raises issues of the reliability and validity of particular scales of test processing. The composition of the tests themselves is critical to scientific progress.

Associations between tests of face perception and other performance measures provide an opportunity to understand the relationships between fundamental processes that are not easily uncovered using traditional group-level analysis. For example, if people who are good at face recognition also turn out to be good at voice recognition, that would provide evidence for some commonality in the processing required of these tasks. Association studies between face processing and broader cognitive tasks – for example IQ tests – can address the key question of whether individual differences in face processing reflect more domain-general differences in perceptual and cognitive abilities^{20,21}.

[H3] Structural Variation

The third major approach to individual differences in face perception is to link quantifiable structural

variation – for example in genetic factors or brain physiology – to differences in face perception abilities. A series of twin studies has demonstrated that certain face perception abilities – particularly those related to recognition – are highly heritable^{22–25}. Along with the evidence from extreme performers, the twin data has been taken to support a stable, trait-like ability underpinning performance on some face perception tasks. Interestingly, this heritability is not observed for some other face perception abilities, including those related to social attributions, for example ratings of perceived trustworthiness¹². Patterns of findings in which certain abilities are strongly heritable but others are not are a good example of how individual differences research can be used to constrain models of the face processing network in general. For example, heritable face identity recognition abilities signal a structural basis that is consistent with this being a discrete processing module.

In addition to genetic factors, individual differences have also been instructive for understanding the relation between face processing abilities and variations in neural physiology. For example, better face processing abilities have been linked to increased grey matter volume in certain regions^{26–28}, as well as to neural activity^{29–31} and connectivity between regions of the face processing network³⁰ (Box 2). Relatively small participant group sizes in neurophysiological studies means that some of this evidence is preliminary, but cumulative evidence from studies of individual differences provides a promising approach linking face processing abilities to their neural substrates.

Finally, researchers have examined the relationship between face processing and certain pathologies. For example, those with low levels of ability in social communication characteristic of Autism Spectrum Disorder tend to perform poorly on a variety of face perception tasks³² including identity³³ and emotion recognition³⁴. People with Autism Spectrum Disorder also show abnormal patterns of attention to faces³², as well as divergence from typical cognitive³⁵ and neurophysiological³⁶ markers of face processing. Whether these are directly linked to reduced social communication abilities in Autism Spectrum Disorder, or to other symptomatic sensory atypicalities, is not clear³⁷. Establishing the complex causal links between face processing and more general communicative abilities therefore remains a challenging task²⁵.

Many of the questions addressed in studies of individual differences remain unresolved. But accumulating evidence from the various strands we have outlined in this section demonstrates the power of this approach to bring together research from cognitive psychology, genetics neuroscience and psychopathology. By observing associations between these diverse measures of individual differences, a multidimensional system of related face processing abilities begins to emerge, and its mapping to structural properties can be revealed.

[H1] Face processing abilities

A longstanding issue in philosophy, psychology, and neuroscience is the extent to which individual differences in performance on cognitive tasks have a common underlying driver, for example IQ. Alternative ‘modular’ accounts hold that certain encapsulated processes such as early visual processing operate relatively independently³⁸. Face perception provides an important example of modularity in higher-level processing, with converging evidence pointing to a degree of encapsulation: Individual differences in face recognition tasks are independent of general intelligence³⁹ and to some extent general visual processing^{22,40–43}. These findings from the individual differences literature concur with long-established evidence from other research traditions including neuropsychology⁴⁴, neuroscience⁴⁵ and behavioural group studies⁴⁶ showing dissociations between face perception and other types of visual processing.

In the field of face perception the modularity issue arises in an often polarised debate about the functional specialisation of face processing relative to other high-level visual processing tasks^{46–48}. In fact, individual difference studies show a moderate, rather than sharp, dissociation between face and visual object perception tasks^{20,40}. Some studies report significant associations between face and general object perception ability (r ranging from .00 to .37)^{20,40,42,49,50} whereas face recognition⁴² and perception²¹ do not correlate with non-visual aspects of intelligence. However, associations between face and object recognition are consistently weaker than those between different face (r ranging from .20 to .65)^{20,22,50–54}, or object recognition tasks (r ranging from .27 to .68)^{42,49}. This pattern is consistent with the observations that two thirds of people with developmental prosopagnosia have impaired object recognition abilities⁴³ abilities³⁹ and super-recognisers outperform control groups on non-face object processing tasks¹⁶. Overall, these patterns of associations are not consistent with a strictly modular face processing system, suggesting a graded rather than absolute distinction between face and object perception abilities.

[H3] Measuring face processing

There is a range of face processing tasks used to study individual differences, including recognition of identity, facial expression, and inferences about personal characteristics (Table 1). Tasks measuring recognition of identity are highly over-represented compared to other aspects of face perception. Diverse task formats have been used to study aspects of face processing (Fig. 1). Probing a particular ability with a range of tests is valuable to gain converging evidence. High levels of convergence suggest an underlying common ability, in contrast to highly task-specific abilities. But diversity in task format can also pose problems for inferences about the relatedness between abilities – lack of

association could be attributed to differences in superficial aspects of the tasks rather than the underlying abilities.

Three key psychometric properties constrain the individual differences approach. First, test-retest reliability, which is the correlation between test scores on the same test across two separate test sessions. This property is critical for interpretation because it places an upper bound on the associations between tests – associations between tests cannot exceed associations between a test and itself. However, estimates of test-retest reliability are not available for most tests. The second property, convergent validity, is the correlation between different tasks that ostensibly measure the same thing. This property is similarly critical⁴² because it establishes that the common variance in test scores is attributable to an underlying ability recruited by the tests, rather than due to artifacts of any particular test – for example the specific images that were used to create it. Third, external validity relates to whether the tests measure what they are intended to measure, by correlating test scores with performance outside the laboratory. In the context of face processing, externally valid tests capture abilities as they are used in daily life, rather than reflecting the highly-specific context of psychological assessment. Thus, whereas test-retest reliability can be measured using a single test (Table 1), convergent and external validity are contingent on comparison with other tests.

[H3] Facial identity

The most common tests of face recognition require participants to remember previously seen faces (for example, ‘which of these faces did you see earlier?’) or to match faces (for example, ‘do these two photos show the same person?’). The results of facial identity processing tests show a wide diversity in abilities across people^{55–57} but an individual’s score is highly stable over time, with test-retest correlations typically above .7 (Table 1)^{22,51,58}. Face recognition is heritable^{22–24}, with correlations of 0.7 between scores of monozygotic twins, compared to 0.29 for dizygotic twins²², and estimates of heritability ranging from 68 to 97%^{22,24}. Thus, identity processing can be measured reliably and apparently taps a stable underlying dimension. Yet this evidence is based predominantly on a single test of unfamiliar face memory, the Cambridge Face Memory Test, CFMT⁵⁵. It is therefore important to confirm that this trait generalises beyond a specific test.

Regarding convergent validity of face identification tests, associations persist despite substantial changes in task format. Similar results are observed despite differences in memory demands^{20,56}, retention interval⁵⁹, and types of response (for example, naming or multiple choice⁴²). The term f – measured by associations between face tasks – is defined as a general factor underlying face identity processing ability²¹. The reported correlation between tests of face memory and matching is typically in

the range of .5 to .7^{21,51}. It has been estimated that f can account for up to 25% of the variance across face recognition tests – including the popular CFMT, Before They Were Famous Test (BFTWF), and Glasgow Face Matching Test (GFMT)²⁰ – and it has been linked to particular polymorphisms from a genome-wide association study^{55,60}. Whereas these face tasks tend to correlate, reports typically show no reliable associations between performance on these tasks and visual processing tasks for other objects. This pattern is consistent with research in object recognition reporting a general factor accounting for shared variance in novel object processing tasks (denoted o ⁴⁹). The general object factor shows relatively weak association with the CFMT ($r = 0.28$), providing converging evidence that face identity processing is somewhat isolated from more general object processing ability.

Another key question is whether face recognition is a unitary ability across familiar and unfamiliar faces. The study of identification commonly focuses on unfamiliar faces, despite the fact that recognition of familiar people is an important component of daily life (only 3 out of 17 face recognition tests use familiar faces, Table 1). This under-representation may be due to recognition of familiar faces being generally easier than recognizing unfamiliar faces, so it is challenging to design discriminating tasks. Furthermore, there is no common set of faces that are familiar to all individuals. People are highly bound to their cultures, age and social groups, each having distinct sets of familiar faces such as celebrities, politicians, and famous athletes⁶¹. There are large behavioural differences between the perception of familiar and unfamiliar faces, which have been argued to reflect qualitatively different processing^{62–64}. Some authors have reported an absence of correlation between familiar and unfamiliar face recognition using matching tasks⁶⁵ but these tests tend to give near-perfect results for familiar faces, limiting the measurement of their association. Studies using tests of familiar face naming tend to find significant associations between familiar and unfamiliar face recognition^{20,42,66}, consistent with the idea that face identity processing is a relatively coherent ability.

Self-report measures of face recognition ability correlate very highly with each other ($r = .82$)⁶⁷, but tend to predict test performance less well (r ranging from .14 to .52)^{68–72}. Ad-hoc (non-psychometric) tests of people's insights into their face recognition abilities have reported even smaller associations to actual performance ($r = .13$ to .26)⁷¹. This modest relationship between self-report and face identity tests could be due to a general lack of metacognitive insight⁷³. Another possibility is that tests of face recognition do not capture the processes involved in everyday recognition. This might also explain relatively low correlations between standardised tests and more naturalistic learning and recognition tasks, for example between CFMT scores and viewers' recognition of faces from the TV show Game of Thrones ($r = 0.45$)^{74,75}. Thus, although face identity processing tests show good internal, convergent, and divergent validity, important questions remain regarding their external validity. Establishing

external validity is critical if face identity tests are to be used for selecting good face recognisers in professional settings¹⁸, and so we return to this issue later (see Practical Implications).

[H3] Expressions and impressions

The study of emotion perception in faces has been dominated by a debate about whether a few ‘basic’ emotions are expressed and perceived similarly by all humans^{76–78}. However, using the individual differences approach, some studies of subtle variations in expression recognition show reliable between-person differences in the ability to judge emotion^{11,79–81}. In contrast to identity processing, the pattern of correlation on tests of emotion recognition is highly sensitive to changes in the particular emotions being expressed² and the task format¹¹. This pattern points to a lack of convergence onto a unitary process for the visual analysis of facial expression. Instead, it seems that the recognition of different emotions (for example, happiness or fear) call on somewhat different abilities^{2,3,82}.

One focus of individual differences research has been social and aesthetic judgments made to unfamiliar faces (that is, the faces of people unknown to the viewers). When asked to judge the trustworthiness or dominance of a face, viewers tend to agree with each other, even though these judgements are typically not accurate indicators of a person’s true character^{83–86}. People asked to judge the attractiveness of a face, show moderate levels of agreement, albeit lower than their ratings of trustworthiness/dominance^{87–89}. Despite some agreements between people in making social judgments, there remains some variation, pointing to idiosyncrasies in individual perceptions, which appear to be relatively stable over time^{12,89}.

Unlike face identity processing, variation in social judgements is not associated with genotype. A large-scale twin study showed that trustworthiness judgements of unfamiliar faces vary more strongly with the viewers’ personal experiences than their genetics¹². Individual variation in aesthetic judgements of attractiveness is also associated more strongly with environment than genes⁸⁹. The implication of current research is that social judgements are a product of social learning^{78,84,90}, and researchers are beginning to propose mechanistic accounts of this learning at an individual level. For example, trustworthiness and attractiveness judgments appear to be linked to transitory changes such as smiling or warm expressions^{91–93}. This observation reflects an emerging view that social signals from faces are intertwined in real world tasks, limiting the external validity of many lab-based tasks that use artificial faces or highly standardised images^{94–96}.

[H3] Multimodal person perception

In daily life, people perceive information about each other using multiple sources, including cues from voices, bodies, clothing, and context. Lab-based experiments using isolated faces can obscure the fact that recognising or making a perceptual judgement about someone usually involves many cues presented together and often in redundant combinations. For example, a viewer might recognise a friend from their face, their walking style, a particular jacket, and the fact that they arrive at an arranged meeting on time (Fig. 2). Variation in real life person perception might incorporate differences in all these dimensions too – each requires cognitive and perceptual decisions, and so they may be subject to individual variation between people.

Within the visual domain, the ability to recognise facial identity is only weakly correlated with the ability to process cues from bodies and movement⁹⁷, suggesting somewhat separable processes. Beyond vision, there are widespread differences in people's ability to recognise voices⁹⁸⁻¹⁰⁰ over and above general differences in auditory perception¹⁰¹. On tests of identity using matching and similarity tasks, there is some evidence for an association between recognising faces and recognising voices¹⁰² and some individuals have high ability levels in both¹⁰³. Associations are typically quite small (r ranging from .24 to .41)¹⁰² and so cross-modal mechanisms do not seem to underpin all voice recognition. The relatively weak association between face and voice recognition is further supported by the report of individuals with developmental prosopagnosia but intact familiar voice recognition¹⁰⁴.

There are also associations between decisions about attractiveness from voices and faces (r from .15 to .34)^{105,106}. The perception of attractiveness appears to be multimodal and can be influenced by olfactory cues¹⁰⁷. Well-established individual differences in olfactory sensory apparatus¹⁰⁸ and effects of scent on other impression judgments¹⁰⁷ highlight the potential for individual differences in the associations between olfactory and visual cues to social judgments.

Individual differences in judgements of emotion from faces and voices also appear to show common processing across modalities^{3,11} and this association extends to tactile perception, elicited by the touch of another person¹⁰⁹. Connolly and colleagues^{2,3} identify shared variability for tests of expression recognition accuracy from both face and voice stimuli, which they describe as a 'supra-modal' factor underlying emotion perception. This factor is related to the ability to introspect on one's own emotional state, which varies dimensionally in the typical population³. The association between face processing and social abilities in the general population has implications for the diagnosis of pathology (Box 1), for example emotion processing impairments in psychopathy^{110,111} and autism¹¹².

Studying individual differences in face processing abilities has contributed to a greater understanding of the subtle ways in which the different aspects of face processing are related. Associations between performance on multiple face tasks, including identity recognition and

expression perception, points to some shared underlying processes. For emotion perception particularly, these processes appear to be multi-modal, incorporating vision, audition and touch. However, it has also been possible to establish some key differences between different face processing abilities, for example the strong genetic component underlying identity recognition, but not social judgements.

[H1] Underlying representations

Research on the associations between different abilities, as described in the previous section, is complemented by a parallel focus on the cognitive mechanisms. Models of face perception posit multiple (serial or parallel) processing stages between visual input and perceptual decisions⁶. Neural models instantiate these networks in connected brain regions⁷ (Box 2). In the study of individual differences, these networks are revealed as systems of related abilities that provide converging sources of information to support perception. Individual differences research tends to report some overlap in people's abilities in recognition and emotion perception^{2,3,11,113–115}. This pattern suggests some shared representational resource between abilities to recognise identities and emotions.

Patterns of association and dissociation help clarify the modular structure and representations of the face processing system. For example, research on developmental prosopagnosia has found that some individuals show impaired gender discrimination¹¹⁶ but spared facial age estimation^{19,117}. This pattern provides good evidence that perception of age is not dependent on identity or gender perception.

These studies begin from an observed data pattern to determine the underlying cognitive processes. Working in the reverse direction, an understanding of the underlying representations used by the face processing system can help to explain observed differences in performance. We examine this latter approach next.

[H3] Holistic processing

Holistic processing refers to the idea that perception of a whole object (or Gestalt) has precedence over perception of its parts. Faces are widely believed to be perceived more holistically than other types of objects⁹. Individual facial features (for example, eyes or noses) are easier to remember when embedded in a face than in isolation¹¹⁸. Similarly, when the top and bottom halves of two different faces are aligned to form a new face, the composed face appears as a new identity¹¹⁹.

The importance of holistic face processing has led to the hypothesis that the extent to which different people process faces holistically might underpin differences in their face processing

ability^{10,59,120,121}. However, this hypothesis is not well supported among individuals with face processing in the typical range. One of the most popular measures of holistic processing is the Composite Face Effect¹¹⁹, a phenomenon in which the top and bottom halves of two different faces are aligned and tend to fuse perceptually into a single new face. This fusion impedes the separate processing of the face halves compared to when they are not aligned. Some studies show weak-to-moderate correlation between the composite face effect and performance on the CFMT^{59,121,122}, but others have found no association with CFMT^{21,123} or other face recognition tasks^{59,124}. The performance of individuals with developmental prosopagnosia also provides mixed evidence for an association between holistic processing and face recognition ability. Some studies have found slightly poorer holistic processing in individuals with developmental prosopagnosia by comparison to controls^{125–127} while others have found no difference^{124,128–130}.

Another challenge to the use of holistic representations as an explanation for differences in face processing is that a person's ability to perform face perception tasks from whole images is highly correlated with their ability to recognise isolated face features¹³¹. Some individuals with developmental prosopagnosia have equivalent impairment on face recognition from isolated features and from whole faces¹⁰⁴, and a hallmark of super-recognisers is their ability to identify faces from relatively limited local face information^{132,133}. Furthermore, recognition is less impacted by distortions that change the spatial layout of facial features in high compared to low performers within the typical range¹³⁴, and those at the top of the typical range are less sensitive to changes in global shape of a face^{135,136}. These findings suggest the need for a more sophisticated understanding of the representational differences underlying face processing ability¹³⁷.

Progress in this area also relies on a greater understanding of the tasks themselves. Problems of measurement have dominated research on holistic processing for over a decade^{59,121,138,139} and the challenge of developing valid and reliable measures appears intractable¹³¹. Even the best-established measures of holistic processing suffer from very low reliability and do not correlate with differences in face recognition performance despite best practice in psychometric approach^{131,140,141}. These observations might signal a problem with the construct of holistic processing itself. Other measures of holistic processing, including face inversion effects (faces are harder to process upside down) and part-whole effects (recognition of isolated features is easier when they are embedded in a face) also correlate very poorly with each other¹²⁴. This pattern is perhaps symptomatic of a broader lack of clarity in operationalising processing mechanisms in the field^{137,142}. Substantial methodological and conceptual challenges need to be overcome to understand how differences in underlying representations give rise to differences in ability.

429 [H3] Unfamiliar and familiar faces

430 Research on individual differences has overwhelmingly examined recognition of unfamiliar faces, yet
 431 the faces of familiar people comprise much of personal daily experience. Group-level evidence shows
 432 that familiarity is directly related to recognition success, viewers are much better at recognizing
 433 familiar than unfamiliar faces^{8,143–145} and higher levels of familiarity exert more powerful modulating
 434 effects on neural responses^{64,146–150}. If differences between familiar face representations are important
 435 for performance within an individual, these differences could also be important between individuals.

436 Performance in recognising famous faces is moderately related to performance on an unfamiliar face
 437 test (CFMT), with correlations ranging from .55⁴² to .33²⁰. There is also some evidence that the
 438 representations underlying familiar and unfamiliar face recognition tasks are distinct. Whereas high
 439 performers on a famous face test were less reliant on global face shape than low performers, high
 440 performers on the CFMT (unfamiliar faces) showed the opposite pattern and were more reliant on
 441 global face shape¹³⁶.

442 For familiar faces, idiosyncratic cues contribute to the representation of identity^{151,152}. One face
 443 might be recognised from a characteristic smirk, another from distinctive facial speech movements.
 444 Analyses of multiple images of the same person reveal not only consistent differences between people,
 445 but also idiosyncratic within-person variability^{153,154}. To become familiar with a new face, one needs to
 446 experience the range over which that face can vary^{155,156}. This multidimensional view of familiar face
 447 representations has implications for individual differences in responses to unfamiliar faces too. People
 448 who are particularly skilled at unfamiliar face recognition recruit elaborate semantic and emotional
 449 representations more commonly used for familiar face processing^{29,30}. In this way, the difficulty of
 450 unfamiliar face recognition may be alleviated to some extent in skilled viewers.

451 Representations of the same faces, both familiar and unfamiliar, also diverge considerably across
 452 individuals. For example, participants disagree entirely which images of unfamiliar faces look most
 453 similar to one another¹⁵⁷, and there are large differences in the photos that people report as showing the
 454 best likeness of a familiar face¹⁵⁸. Research on the representations underlying these differences is rare
 455 and it remains puzzling why different viewers show different patterns of similarity between the same
 456 familiar faces.

457 In summary, researchers have sought to explain differences in performance on face tasks through
 458 differences in viewers' underlying representations. The degree to which people tend to use holistic
 459 processing was once thought to be a good candidate to explain variation in face recognition
 460 performance, but the evidence is weak. Differences between recognition of familiar and unfamiliar face

recognition offer some promise for understanding the relationship between people's representations and their performance. Further exploration of this relationship will require detailed investigation of people's idiosyncratic representations of the faces they know.

[H1] Practical implications

In addition to providing theoretical understanding, studying individual differences brings practical implications. For example, face perception in disorders influencing social cognition has clinical relevance (Box 1) and social consequences for individuals with these disorders^{159–162}. There are also clear societal implications of individual differences in face recognition. The outcomes of face identification decisions in security and forensic settings can often be profound – impacting civil liberties and even leading to wrongful convictions – and the science of individual differences can help address these problems.

Identity-checking

Tasks that involve checking the identity of unfamiliar people are known to be difficult and error-prone. Error rates of 20-30 percent are common in studies asking viewers to match two different photos of the same person, taken on different occasions, even when using high quality images taken in good lighting¹⁶³. This difficulty extends to professionals who conduct daily face matching, such as in passport control or forensic face identification. In a meta-analysis of 29 comparisons between professional groups and participants from the general population on tests of unfamiliar face identity matching, 40% of tests showed equivalent face matching accuracy in these groups¹⁶⁴. High error rates were found in staff performing a variety of important identity verification roles in border control¹⁶⁵, government offices¹⁶⁶, passport issuance^{167,168}, police departments¹⁶⁹, security firms¹⁷⁰ and banks¹⁷¹. Simply performing identification tasks in daily work is not sufficient for expertise. Furthermore, current approaches to training in many professional settings are ineffective (Box 3).

The discovery of reliable individual differences in face recognition provides one means of addressing this problem. It is becoming increasingly popular to select people for specialist face identification roles on the basis of their natural ability as measured by standard tests. This strategy has been used by the London Metropolitan Police^{172,173} and the Australian Passport Office¹⁶⁸, with groups selected for high face performance showing 10-20% gains in accuracy over control groups.

Forensic face identification

Facial forensic examiners – who analyse similarities and differences between face images to provide evidentiary reports for police investigations and criminal trials – outperform standard participant groups by roughly the same margin^{57,164,174,175} as selectively-recruited, but untrained, super-recognisers. In contrast to super-recognisers' quick and intuitive recognition ability, forensic abilities are founded on years of deliberate training in comparing images of unfamiliar faces¹⁷⁶ and involve slow, analytic comparison^{174,175}.

Forensic identifications are also made by eye-witnesses. These are highly vulnerable to error, with meta-analysis suggesting that 50% of eyewitness lineup selections are wrong¹⁷⁷. Given the range of face recognition abilities in the general population, it is likely that a large proportion of errors are made by people with relatively poor face recognition abilities. Researchers have examined the use of tests of face identification to screen eyewitnesses, an approach that pre-dates broader interest in individual differences¹⁷⁸. Face recognition tests can be used to predict eyewitness errors¹⁷⁹, by allowing law-enforcement officers to weigh witnesses' identifications against their objective abilities. Furthermore, eyewitnesses are often overconfident, and tests can establish whether particular individuals tend to over-estimate their recognition performance, providing a level of credibility to testimony^{73,180,181}.

The potential for individual difference research to improve accuracy in real-world tasks relies on reliable and valid tests. From an applied perspective, valid tests must correspond with real-world tasks. As a basic example, a face memory test might not be an optimal measure for professionals who are required to match but not remember faces. However, there is sufficient task diversity among relevant practitioners to present a nontrivial challenge in choosing tests for specific professional contexts¹⁸. Forensic identifications made from CCTV involve a complex set of cognitive demands¹⁸² and might incorporate cues beyond the face including behaviour, gait, or clothing. The use of these cues these might each represent separate skills⁹⁷. Preliminary evidence suggests that face recognition tests are not especially reliable predictors of performance on CCTV monitoring tasks^{75,183} indicating that basic understanding of skills underpinning accuracy on different real-world identification tasks is lacking.

The challenges in forensic face identification echo the lack of diversity in measures for effectively capturing everyday abilities. Batteries of face tests that target distinct subskills provide an alternative to reducing ability to a single test score^{18,52}, and might provide the necessary flexibility to capture the multidimensional nature of person identification for both applied and theoretical use.

Human-AI collaboration

Face recognition in applied settings increasingly relies on combined processing by humans and technology. Deep neural network approaches to facial recognition have been highly successful and the

best-performing systems are now as accurate as both super-recognisers and facial forensic examiners⁵⁷. Such automated processes are used for passport control in some countries as well as police searches for suspects in image surveillance¹⁸⁴. Critically, in many of these applications, the technology does not replace human processing but rather presents operators with arrays of potential matches for follow-up. This procedure automatically makes easy match decisions, leaving more difficult matches to human reviewers and error rates in human review can be as high as 50%¹⁶⁸. Thus, this type of forensic identification can be problematic in the same way as traditional identification processes, such as eyewitness lineups¹⁷⁷.

Selecting people with the necessary skills to review matches generated by facial recognition technology is a potential way to reduce error rates. Moreover, it appears that personnel selection can be tailored to the specific face recognition algorithms that are being used. Statistical aggregation of the decisions made by algorithm and high performing humans produces accuracy that exceeds either algorithms or humans alone⁵⁷. This statistical combination benefit is driven by independent processes recruited by algorithms and human perceivers. Given the present revival of interest in deep learning networks as models of face processing^{185,186}, evaluating similarities and differences between human and machine processing can also lead to theoretical advances.

[H1] Summary and future directions

Individual differences research is a complementary approach to traditional group studies for understanding face perception. In a field that has traditionally drawn on converging evidence, individual differences research enables new questions to be asked and can address some long-standing issues. Although there has been considerable research focusing on people with extreme levels of ability (individuals with developmental prosopagnosia and super-recognisers), there is considerable potential for broader scientific progress across the full scale of abilities.

The study of individual differences has also highlighted some major problems in the field of face perception. Perhaps the most significant of these is the problem of measurement. The construction of reliable and valid tests lies at the heart of an individual differences research programme, but tests of face perception remain comparatively weak in these properties. Without reliable tests, it is impossible to draw valid conclusions. Although the construction of new tests remains a challenge, it would be relatively straightforward for researchers to only use tests with published test-reliability measures. The problem of reliability in psychological measures is not specific to the study of face processing, but the problem seems particularly acute in this field because there are multiple tests for measuring each aspect of face processing (Table 1).

559 The issue of measurement has also highlighted another problem with the theory-led approach to
560 some face recognition questions. A good example is the widespread view that holistic perceptual
561 processing underlies face perception. However, the set of tests used to measure holistic processing
562 correlate very poorly with each other¹²⁴ - a key finding that has perhaps not yet had the influence it
563 deserves. Theoretical statements based on holistic processing are common and the field has perhaps
564 been too willing to adopt these generalisations without clear operationalisation¹⁴². At the very least,
565 holistic processing accounts of face perception should specify the relevant measure of holistic
566 processing¹³¹.

567 Another major challenge remains in eliciting general principles of face perception while
568 acknowledging that every person has different experience with faces. Developing methods for studying
569 variation in familiar face recognition will be a major challenge, given the highly idiosyncratic set of
570 personally familiar faces and the laborious processes required to tailor experimental materials to
571 individual participants^{64,148,149,187}. For example, one person might not recognise Barack Obama and
572 another might not recognise Kim Kardashian – discrepancies that often lead to mutual disbelief.
573 Approaches targeting specific cohorts of TV viewers who have comparable perceptual exposure could
574 offer a promising solution to this methodological problem⁷⁴. Most studies of familiar face perception
575 treat familiarity as a binary categorisation (familiar/unfamiliar), a methodological constraint which has,
576 to some extent, obscured our understanding using traditional experimental approaches¹⁵⁴. It remains to
577 be seen whether individual differences approaches, which conceptually differentiate between people,
578 can be harnessed to capture natural idiosyncrasy.

579 In this Review, we have emphasised the implications of multiple sources of information for face
580 perception. We have shown how individual differences approaches shed light on the perceptual
581 architecture necessary to use faces in the flexible ways that humans do. Yet, the biggest unsolved
582 problem in face perception remains how someone recognises the people they know.

References

1. Calder, A. J. & Young, A. W. Understanding the recognition of facial identity and facial expression. *Nat. Rev. Neurosci.* **6**, 641–651, DOI: [10.1038/nrn1724](https://doi.org/10.1038/nrn1724) (2005).
2. Connolly, H. L., Young, A. W. & Lewis, G. J. Recognition of facial expression and identity in part reflects a common ability, independent of general intelligence and visual short-term memory. *Cogn. Emot.* **33**, 1–10, DOI: [10.1080/02699931.2018.1535425](https://doi.org/10.1080/02699931.2018.1535425) (2018).
3. Connolly, H. L., Lefevre, C. E., Young, A. W. & Lewis, G. J. Emotion recognition ability: Evidence for a supramodal factor and its links to social cognition. *Cognition* **197**, 104166, DOI: [10.1016/j.cognition.2019.104166](https://doi.org/10.1016/j.cognition.2019.104166) (2020).
4. Vogel, E. K., McCollough, A. W. & Machizawa, M. G. Neural measures reveal individual differences in controlling access to working memory. *Nature* **438**, 500–503, DOI: [10.1038/nature04171](https://doi.org/10.1038/nature04171) (2005).
5. Kanai, R. & Rees, G. The structural basis of inter-individual differences in human behaviour and cognition. *Nat. Rev. Neurosci.* **12**, 231–242, DOI: [10.1038/nrn3000](https://doi.org/10.1038/nrn3000) (2011).
6. Bruce, V. & Young, A. Understanding face recognition. *Br. J. Psychol.* **77**, 305–327, DOI: [10.1111/j.2044-8295.1986.tb02199.x](https://doi.org/10.1111/j.2044-8295.1986.tb02199.x) (1986).
7. Haxby, J. V. *et al.* The distributed human neural system for face perception. *Trends Cogn. Sci.* **4**, 223–233, DOI: [10.1016/s1364-6613\(00\)01482-0](https://doi.org/10.1016/s1364-6613(00)01482-0) (2000).
8. Jenkins, R., White, D., Montfort, X. V. & Burton, A. M. Variability in photos of the same face. *Cognition* **121**, 313–323, DOI: [10.1016/j.cognition.2011.08.001](https://doi.org/10.1016/j.cognition.2011.08.001) (2011).
9. Farah, M. J., Wilson, K. D., Drain, M. & Tanaka, J. N. What Is “Special” About Face Perception? *Psychol. Rev.* **105**, 482–498, DOI: [10.1037/0033-295x.105.3.482](https://doi.org/10.1037/0033-295x.105.3.482) (1998).
10. Wilmer, J. B. Individual Differences in Face Recognition: A Decade of Discovery. *Curr. Dir. Psychol. Sci.* **26**, 225–230, DOI: [10.1177/0963721417710693](https://doi.org/10.1177/0963721417710693) (2017).
11. Palermo, R., O’Connor, K. B., Davis, J. M., Irons, J. & McKone, E. New Tests to Measure Individual Differences in Matching and Labelling Facial Expressions of Emotion, and Their Association with Ability to Recognise Vocal Emotions and Facial Identity. *PLoS ONE* **8**, e68126, DOI: [10.1371/journal.pone.0068126](https://doi.org/10.1371/journal.pone.0068126) (2013).
12. Sutherland, C. A. M. *et al.* Individual differences in trust evaluations are shaped mostly by

environments, not genes. *Proc. Natl. Acad. Sci.* **117**, 10218–10224, DOI:

[10.1073/pnas.1920131117](https://doi.org/10.1073/pnas.1920131117) (2020).

13. McConachie, H. R. Developmental Prosopagnosia. A Single Case Report. *Cortex* **12**, 76–82, DOI: [10.1016/s0010-9452\(76\)80033-0](https://doi.org/10.1016/s0010-9452(76)80033-0) (1976).

14. Behrmann, M. & Avidan, G. Congenital prosopagnosia: face-blind from birth. *Trends Cogn. Sci.* **9**, 180–187, DOI: [10.1016/j.tics.2005.02.011](https://doi.org/10.1016/j.tics.2005.02.011) (2005).

15. Dobel, C., Bölte, J., Aicher, M. & Schweinberger, S. R. Prosopagnosia Without Apparent Cause: Overview and Diagnosis of Six Cases. *Cortex* **43**, 718–733, DOI: [10.1016/s0010-9452\(08\)70501-x](https://doi.org/10.1016/s0010-9452(08)70501-x) (2007).

16. Bobak, A. K., Bennetts, R. J., Parris, B. A., Jansari, A. & Bate, S. An in-depth cognitive examination of individuals with superior face recognition skills. *Cortex* **82**, 48–62, DOI: [10.1016/j.cortex.2016.05.003](https://doi.org/10.1016/j.cortex.2016.05.003) (2016).

17. Russell, R., Duchaine, B. & Nakayama, K. Super-recognizers: People with extraordinary face recognition ability. *Psychon. Bull. & Rev.* **16**, 252–257, DOI: [10.3758/pbr.16.2.252](https://doi.org/10.3758/pbr.16.2.252) (2009).

18. Ramon, M., Bobak, A. K. & White, D. Super-recognizers: From the lab to the world and back again. *Br. J. Psychol.* **110**, 461–479, DOI: [10.1111/bjop.12368](https://doi.org/10.1111/bjop.12368) (2019).

19. Chatterjee, G. & Nakayama, K. Normal facial age and gender perception in developmental prosopagnosia. *Cogn. Neuropsychol.* **29**, 482–502, DOI: [10.1080/02643294.2012.756809](https://doi.org/10.1080/02643294.2012.756809) (2012).

20. McCaffery, J. M., Robertson, D. J., Young, A. W. & Burton, A. M. Individual differences in face identity processing. *Cogn. Res. Princ. Implic.* **3**, 21, DOI: [10.1186/s41235-018-0112-9](https://doi.org/10.1186/s41235-018-0112-9) (2018).

21. Verhallen, R. J. *et al.* General and specific factors in the processing of faces. *Vis. Res.* **141**, 217–227, DOI: [10.1016/j.visres.2016.12.014](https://doi.org/10.1016/j.visres.2016.12.014) (2017).

22. Wilmer, J. B. *et al.* Human face recognition ability is specific and highly heritable. *Proc. Natl. Acad. Sci.* **107**, 5238–5241, DOI: [10.1073/pnas.0913053107](https://doi.org/10.1073/pnas.0913053107) (2010).

23. Zhu, Q. *et al.* Heritability of the Specific Cognitive Ability of Face Perception. *Curr. Biol.* **20**, 137–142, DOI: [10.1016/j.cub.2009.11.067](https://doi.org/10.1016/j.cub.2009.11.067) (2010).

24. Shakeshaft, N. G. & Plomin, R. Genetic specificity of face recognition. *Proc. Natl. Acad. Sci.* **112**, 12887–12892, DOI: [10.1073/pnas.1421881112](https://doi.org/10.1073/pnas.1421881112) (2015).

25. Lewis, G. J., Shakeshaft, N. G. & Plomin, R. Face Identity Recognition and the Social Difficulties Component of the Autism-Like Phenotype: Evidence for Phenotypic and Genetic Links. *J. Autism*

Dev. Disord. **48**, 2758–2765, DOI: [10.1007/s10803-018-3539-4](https://doi.org/10.1007/s10803-018-3539-4) (2018).

26. Garrido, L. *et al.* Voxel-based morphometry reveals reduced grey matter volume in the temporal cortex of developmental prosopagnosics. *Brain* **132**, 3443–3455, DOI: [10.1093/brain/awp271](https://doi.org/10.1093/brain/awp271) (2009).
27. McGugin, R. W., Newton, A. T., Tamber-Rosenau, B., Tomarken, A. & Gauthier, I. Thickness of Deep Layers in the Fusiform Face Area Predicts Face Recognition. *J. Cogn. Neurosci.* **32**, 1316–1329, DOI: [10.1162/jocn_a_01551](https://doi.org/10.1162/jocn_a_01551) (2020).
28. Jin, Z. *et al.* Impaired face recognition is associated with abnormal gray matter volume in the posterior cingulate cortex in congenital amusia. *Neuropsychologia* **156**, 107833, DOI: [10.1016/j.neuropsychologia.2021.107833](https://doi.org/10.1016/j.neuropsychologia.2021.107833) (2021).
29. Ramot, M., Walsh, C. & Martin, A. Multifaceted Integration: Memory for Faces Is Subserved by Widespread Connections between Visual, Memory, Auditory, and Social Networks. *J. Neurosci.* **39**, 4976–4985, DOI: [10.1523/jneurosci.0217-19.2019](https://doi.org/10.1523/jneurosci.0217-19.2019) (2019).
30. Elbich, D. B. & Scherf, S. Beyond the FFA: Brain-behavior correspondences in face recognition abilities. *NeuroImage* **147**, 409–422, DOI: [10.1016/j.neuroimage.2016.12.042](https://doi.org/10.1016/j.neuroimage.2016.12.042) (2017).
31. Weibert, K. & Andrews, T. J. Activity in the right fusiform face area predicts the behavioural advantage for the perception of familiar faces. *Neuropsychologia* **75**, 588–596, DOI: [10.1016/j.neuropsychologia.2015.07.015](https://doi.org/10.1016/j.neuropsychologia.2015.07.015) (2015).
32. Webb, S. J., Neuhaus, E. & Faja, S. Face perception and learning in autism spectrum disorders. *The Q. J. Exp. Psychol.* **70**, 1–17, DOI: [10.1080/17470218.2016.1151059](https://doi.org/10.1080/17470218.2016.1151059) (2016).
33. Hedley, D., Brewer, N. & Young, R. Face recognition performance of individuals with Asperger syndrome on the Cambridge face memory test. *Autism Res.* **4**, 449–455, DOI: [10.1002/aur.214](https://doi.org/10.1002/aur.214) (2011).
34. Uljarevic, M. & Hamilton, A. Recognition of Emotions in Autism: A Formal Meta-Analysis. *J. Autism Dev. Disord.* **43**, 1517–1526, DOI: [10.1007/s10803-012-1695-5](https://doi.org/10.1007/s10803-012-1695-5) (2013).
35. Behrmann, M. *et al.* Configural processing in autism and its relationship to face processing. *Neuropsychologia* **44**, 110–129, DOI: [10.1016/j.neuropsychologia.2005.04.002](https://doi.org/10.1016/j.neuropsychologia.2005.04.002) (2006).
36. Harms, M. B., Martin, A. & Wallace, G. L. Facial Emotion Recognition in Autism Spectrum Disorders: A Review of Behavioral and Neuroimaging Studies. *Neuropsychol. Rev.* **20**, 290–322,

DOI: [10.1007/s11065-010-9138-6](https://doi.org/10.1007/s11065-010-9138-6) (2010).

37. Pellicano, E. & Burr, D. When the world becomes ‘too real’: a Bayesian explanation of autistic perception. *Trends Cogn. Sci.* **16**, 504–510, DOI: [10.1016/j.tics.2012.08.009](https://doi.org/10.1016/j.tics.2012.08.009) (2012).
38. Pylyshyn, Z. Is vision continuous with cognition?: The case for cognitive impenetrability of visual perception. *Behav. Brain Sci.* **22**, 341–365, DOI: [10.1017/s0140525x99002022](https://doi.org/10.1017/s0140525x99002022) (1999).
39. Gignac, G. E., Shankaralingam, M., Walker, K. & Kilpatrick, P. Short-term memory for faces relates to general intelligence moderately. *Intelligence* **57**, 96–104, DOI: [10.1016/j.intell.2016.05.001](https://doi.org/10.1016/j.intell.2016.05.001) (2016).
40. Dennett, H. W. *et al.* The Cambridge Car Memory Test: A task matched in format to the Cambridge Face Memory Test, with norms, reliability, sex differences, dissociations from face memory, and expertise effects. *Behav. Res. Methods* **44**, 587–605, DOI: [10.3758/s13428-011-0160-2](https://doi.org/10.3758/s13428-011-0160-2) (2012).
41. Richler, J. J., Wilmer, J. B. & Gauthier, I. General object recognition is specific: Evidence from novel and familiar objects. *Cognition* **166**, 42–55, DOI: [10.1016/j.cognition.2017.05.019](https://doi.org/10.1016/j.cognition.2017.05.019) (2017).
42. Wilmer, J. B. *et al.* Capturing specific abilities as a window into human individuality: The example of face recognition. *Cogn. Neuropsychol.* **29**, 360–392, DOI: [10.1080/02643294.2012.753433](https://doi.org/10.1080/02643294.2012.753433) (2012).
43. Geskin, J. & Behrmann, M. Congenital prosopagnosia without object agnosia? A literature review. *Cogn. Neuropsychol.* **35**, 1–51, DOI: [10.1080/02643294.2017.1392295](https://doi.org/10.1080/02643294.2017.1392295) (2017).
44. McNeil, J. E. & Warrington, E. K. Prosopagnosia: A face-specific disorder. *The Q. J. Exp. Psychol. Sect. A* **46**, 1–10, DOI: [10.1080/14640749308401064](https://doi.org/10.1080/14640749308401064) (1993).
45. Kanwisher, N. & Yovel, G. The fusiform face area: a cortical region specialized for the perception of faces. *Philos. Transactions Royal Soc. B: Biol. Sci.* **361**, 2109–2128, DOI: [10.1098/rstb.2006.1934](https://doi.org/10.1098/rstb.2006.1934) (2006).
46. McKone, E., Kanwisher, N. & Duchaine, B. C. Can generic expertise explain special processing for faces? *Trends Cogn. Sci.* **11**, 8–15, DOI: [10.1016/j.tics.2006.11.002](https://doi.org/10.1016/j.tics.2006.11.002) (2007).
47. Kanwisher, N. Domain specificity in face perception. *Nat. Neurosci.* **3**, 759–763, DOI: [10.1038/77664](https://doi.org/10.1038/77664) (2000).
48. Gauthier, I., Curran, T., Curby, K. M. & Collins, D. Perceptual interference supports a non-

modular account of face processing. *Nat. Neurosci.* **6**, 428–432, DOI: [10.1038/nn1029](https://doi.org/10.1038/nn1029) (2003).

49. Richler, J. J. *et al.* Individual Differences in Object Recognition. *Psychol. Rev.* **126**, 226–251, DOI: [10.1037/rev0000129](https://doi.org/10.1037/rev0000129) (2019).

50. Dunn, J. D., Summersby, S., Towler, A., Davis, J. P. & White, D. UNSW Face Test: A screening tool for super-recognizers. *PLOS ONE* **15**, e0241747, DOI: [10.1371/journal.pone.0241747](https://doi.org/10.1371/journal.pone.0241747) (2020).

51. Balsdon, T., Summersby, S., Kemp, R. I. & White, D. Improving face identification with specialist teams. *Cogn. Res. Princ. Implic.* **3**, 25, DOI: [10.1186/s41235-018-0114-7](https://doi.org/10.1186/s41235-018-0114-7) (2018).

52. Fysh, M. C., Stacchi, L. & Ramon, M. Differences between and within individuals, and subprocesses of face cognition: implications for theory, research and personnel selection. *Royal Soc. Open Sci.* **7**, 200233, DOI: [10.1098/rsos.200233](https://doi.org/10.1098/rsos.200233) (2020).

53. Bobak, A. K., Pampoulov, P. & Bate, S. Detecting Superior Face Recognition Skills in a Large Sample of Young British Adults. *Front. Psychol.* **7**, 1378, DOI: [10.3389/fpsyg.2016.01378](https://doi.org/10.3389/fpsyg.2016.01378) (2016).

54. Stantic, M. *et al.* The Oxford Face Matching Test: A non-biased test of the full range of individual differences in face perception. *Behav. Res. Methods* 1–16, DOI: [10.3758/s13428-021-01609-2](https://doi.org/10.3758/s13428-021-01609-2) (2021).

55. Duchaine, B. & Nakayama, K. The Cambridge Face Memory Test: Results for neurologically intact individuals and an investigation of its validity using inverted face stimuli and prosopagnosic participants. *Neuropsychologia* **44**, 576–585, DOI: [10.1016/j.neuropsychologia.2005.07.001](https://doi.org/10.1016/j.neuropsychologia.2005.07.001) (2006).

56. Burton, A. M., White, D. & McNeill, A. The Glasgow Face Matching Test. *Behav. Res. Methods* **42**, 286–291, DOI: [10.3758/brm.42.1.286](https://doi.org/10.3758/brm.42.1.286) (2010).

57. Phillips, P. J. *et al.* Face recognition accuracy of forensic examiners, superrecognizers, and face recognition algorithms. *Proc. Natl. Acad. Sci.* **115**, 201721355, DOI: [10.1073/pnas.1721355115](https://doi.org/10.1073/pnas.1721355115) (2018).

58. White, D., Guilbert, D., Varela, V. P. L., Jenkins, R. & Burton, A. M. GFMT2: A psychometric measure of face matching ability. *Behav. Res. Methods* 1–9, DOI: [10.3758/s13428-021-01638-x](https://doi.org/10.3758/s13428-021-01638-x) (2021).

59. Richler, J. J., Cheung, O. S. & Gauthier, I. Holistic Processing Predicts Face Recognition. *Psychol. Sci.* **22**, 464–471, DOI: [10.1177/0956797611401753](https://doi.org/10.1177/0956797611401753) (2011).

60. Verhallen, R. J. *et al.* An online version of the Mooney Face Test: phenotypic and genetic associations.

63, 19–25, DOI: [10.1016/j.neuropsychologia.2014.08.011](https://doi.org/10.1016/j.neuropsychologia.2014.08.011) (2014).

- 61.** Jenkins, R., Dowsett, A. J. & Burton, A. M. How many faces do people know? *Proc. Royal Soc. B* **285**, 20181319, DOI: [10.1098/rspb.2018.1319](https://doi.org/10.1098/rspb.2018.1319) (2018).
- 62.** Johnston, R. A. & Edmonds, A. J. Familiar and unfamiliar face recognition: A review. *Memory* **17**, 577–596, DOI: [10.1080/09658210902976969](https://doi.org/10.1080/09658210902976969) (2009).
- 63.** Collins, E., Robinson, A. K. & Behrmann, M. Distinct neural processes for the perception of familiar versus unfamiliar faces along the visual hierarchy revealed by EEG. *NeuroImage* **181**, 120–131, DOI: [10.1016/j.neuroimage.2018.06.080](https://doi.org/10.1016/j.neuroimage.2018.06.080) (2018).
- 64.** Wiese, H. *et al.* A Robust Neural Index of High Face Familiarity. *Psychol. Sci.* **30**, 261–272, DOI: [10.1177/0956797618813572](https://doi.org/10.1177/0956797618813572) (2019).
- 65.** Megreya, A. M. & Burton, A. M. Unfamiliar faces are not faces: Evidence from a matching task. *Mem. & Cogn.* **34**, 865–876, DOI: [10.3758/bf03193433](https://doi.org/10.3758/bf03193433) (2006).
- 66.** Bate, S. *et al.* Objective Patterns of Face Recognition Deficits in 165 Adults with Self-Reported Developmental Prosopagnosia. *Brain Sci.* **9**, 133, DOI: [10.3390/brainsci9060133](https://doi.org/10.3390/brainsci9060133) (2019).
- 67.** Matsuyoshi, D. & Watanabe, K. People have modest, not good, insight into their face recognition ability: a comparison between self-report questionnaires. *Psychol. Res.* **85**, 1713–1723, DOI: [10.1007/s00426-020-01355-8](https://doi.org/10.1007/s00426-020-01355-8) (2021).
- 68.** Bobak, A. K., Mileva, V. R. & Hancock, P. J. Facing the facts: Naive participants have only moderate insight into their face recognition and face perception abilities. *Q. J. Exp. Psychol.* **72**, 872–881, DOI: [10.1177/1747021818776145](https://doi.org/10.1177/1747021818776145) (2018).
- 69.** Gray, K. L. H., Bird, G. & Cook, R. Robust associations between the 20-item prosopagnosia index and the Cambridge Face Memory Test in the general population. *Royal Soc. Open Sci.* **4**, 160923, DOI: [10.1098/rsos.160923](https://doi.org/10.1098/rsos.160923) (2017).
- 70.** Ventura, P., Livingston, L. A. & Shah, P. Adults have moderate-to-good insight into their face recognition ability: Further validation of the 20-item Prosopagnosia Index in a Portuguese sample. *Q. J. Exp. Psychol.* **71**, 2677–2679, DOI: [10.1177/1747021818765652](https://doi.org/10.1177/1747021818765652) (2018).
- 71.** Palermo, R. *et al.* Do people have insight into their face recognition abilities? *The Q. J. Exp. Psychol.* **70**, 1–16, DOI: [10.1080/17470218.2016.1161058](https://doi.org/10.1080/17470218.2016.1161058) (2017).
- 72.** Arizpe, J. M. *et al.* Self-reported face recognition is highly valid, but alone is not highly discriminative of prosopagnosia- level performance on objective assessments. *Behav. Res. Methods*

51, 1102–1116, DOI: [10.3758/s13428-018-01195-w](https://doi.org/10.3758/s13428-018-01195-w) (2019).

73. Zhou, X. & Jenkins, R. Dunning–Kruger effects in face perception. *Cognition* **203**, 104345, DOI: [10.1016/j.cognition.2020.104345](https://doi.org/10.1016/j.cognition.2020.104345) (2020).
74. Devue, C., Wride, A. & Grimshaw, G. M. New Insights on Real-World Human Face Recognition. *J. Exp. Psychol. Gen.* **148**, 994–1007, DOI: [10.1037/xge0000493](https://doi.org/10.1037/xge0000493) (2018).
75. Thielgen, M. M., Schade, S. & Bosé, C. Face processing in police service: the relationship between laboratory-based assessment of face processing abilities and performance in a real-world identity matching task. *Cogn. Res. Princ. Implic.* **6**, 54, DOI: [10.1186/s41235-021-00317-x](https://doi.org/10.1186/s41235-021-00317-x) (2021).
76. Ekman, P. & Friesen, W. V. Constants across cultures in the face and emotion. *J. Pers. Soc. Psychol.* **17**, 124–129, DOI: [10.1037/h0030377](https://doi.org/10.1037/h0030377) (1971).
77. Jack, R. E., Garrod, O. G. B., Yu, H., Caldara, R. & Schyns, P. G. Facial expressions of emotion are not culturally universal. *Proc. Natl. Acad. Sci.* **109**, 7241–7244, DOI: [10.1073/pnas.1200155109](https://doi.org/10.1073/pnas.1200155109) (2012).
78. Mesquita, B., Boiger, M. & Leersnyder, J. D. The cultural construction of emotions. *Curr. Opin. Psychol.* **8**, 31–36, DOI: [10.1016/j.copsyc.2015.09.015](https://doi.org/10.1016/j.copsyc.2015.09.015) (2016).
79. Olderbak, S. & Wilhelm, O. Emotion Perception and Empathy: An Individual Differences Test of Relations. *Emotion* **17**, 1092–1106, DOI: [10.1037/emo0000308](https://doi.org/10.1037/emo0000308) (2017).
80. Lázaro, E. *et al.* Instrument for Assessing the Ability to Identify Emotional Facial Expressions in Healthy Children and in Children With ADHD: The FEEL Test. *J. Atten. Disord.* **23**, 563–569, DOI: [10.1177/1087054716682335](https://doi.org/10.1177/1087054716682335) (2019).
81. Baron-Cohen, S., Wheelwright, S. & Jolliffe, and Therese. Is There a "Language of the Eyes"? Evidence from Normal Adults, and Adults with Autism or Asperger Syndrome. *Vis. Cogn.* **4**, 311–331, DOI: [10.1080/713756761](https://doi.org/10.1080/713756761) (1997).
82. Calder, A. J., Ewbank, M. & Passamonti, L. Personality influences the neural responses to viewing facial expressions of emotion. *Philos. Transactions Royal Soc. B: Biol. Sci.* **366**, 1684–1701, DOI: [10.1098/rstb.2010.0362](https://doi.org/10.1098/rstb.2010.0362) (2011).
83. Willis, J. & Todorov, A. First Impressions. *Psychol. Sci.* **17**, 592–598, DOI: [10.1111/j.1467-9280.2006.01750.x](https://doi.org/10.1111/j.1467-9280.2006.01750.x) (2005).
84. Zebrowitz, L. A. & Montepare, J. M. First impressions from facial appearance cues. In *First impressions*, 171—204 (Guilford Publications, 2008).

85. Oosterhof, N. N. & Todorov, A. The functional basis of face evaluation. *Proc. Natl. Acad. Sci.* **105**, 11087–11092, DOI: [10.1073/pnas.0805664105](https://doi.org/10.1073/pnas.0805664105) (2008).
86. Todorov, A., Olivola, C. Y., Dotsch, R. & Mende-Siedlecki, P. Social Attributions from Faces: Determinants, Consequences, Accuracy, and Functional Significance. *Annu. Rev. Psychol.* **66**, 519–545, DOI: [10.1146/annurev-psych-113011-143831](https://doi.org/10.1146/annurev-psych-113011-143831) (2015).
87. Hehman, E., Sutherland, C. A. M., Flake, J. K. & Slepian, M. L. The Unique Contributions of Perceiver and Target Characteristics in Person Perception. *J. Pers. Soc. Psychol.* **113**, 513–529, DOI: [10.1037/pspa0000090](https://doi.org/10.1037/pspa0000090) (2017).
88. Hehman, E., Stoller, R. M., Freeman, J. B., Flake, J. K. & Xie, S. Y. Toward a comprehensive model of face impressions: What we know, what we do not, and paths forward. *Soc. Pers. Psychol. Compass* **13**, e12431, DOI: [10.1111/spc3.12431](https://doi.org/10.1111/spc3.12431) (2019).
89. Germine, L. *et al.* Individual Aesthetic Preferences for Faces Are Shaped Mostly by Environments, Not Genes. *Curr. Biol.* **25**, 2684–2689, DOI: [10.1016/j.cub.2015.08.048](https://doi.org/10.1016/j.cub.2015.08.048) (2015).
90. Zebrowitz, L. A. First Impressions From Faces. *Curr. Dir. Psychol. Sci.* **26**, 237–242, DOI: [10.1177/0963721416683996](https://doi.org/10.1177/0963721416683996) (2017).
91. Sutherland, C. A. M., Rhodes, G., Burton, N. S. & Young, A. W. Do facial first impressions reflect a shared social reality? *Br. J. Psychol.* **111**, 215–232, DOI: [10.1111/bjop.12390](https://doi.org/10.1111/bjop.12390) (2020).
92. Sofer, C., Dotsch, R., Wigboldus, D. H. J. & Todorov, A. What Is Typical Is Good. *Psychol. Sci.* **26**, 39–47, DOI: [10.1177/0956797614554955](https://doi.org/10.1177/0956797614554955) (2014).
93. Sutherland, C. A. M., Young, A. W. & Rhodes, G. Facial first impressions from another angle: How social judgements are influenced by changeable and invariant facial properties. *Br. J. Psychol.* **108**, 397–415, DOI: [10.1111/bjop.12206](https://doi.org/10.1111/bjop.12206) (2017).
94. Sutherland, C. A. *et al.* Social inferences from faces: Ambient images generate a three-dimensional model. *Cognition* **127**, 105–118, DOI: [10.1016/j.cognition.2012.12.001](https://doi.org/10.1016/j.cognition.2012.12.001) (2013).
95. Todorov, A. & Porter, J. M. Misleading First Impressions. *Psychol. Sci.* **25**, 1404–1417, DOI: [10.1177/0956797614532474](https://doi.org/10.1177/0956797614532474) (2014).
96. White, D., Sutherland, C. A. M. & Burton, A. L. Choosing face: The curse of self in profile image selection. *Cogn. Res. Princ. Implic.* **2**, 23, DOI: [10.1186/s41235-017-0058-3](https://doi.org/10.1186/s41235-017-0058-3) (2017).
97. Noyes, E., Hill, M. Q. & O’Toole, A. J. Face recognition ability does not predict person

identification performance: using individual data in the interpretation of group results. *Cogn. Res. Princ. Implic.* **3**, 23, DOI: [10.1186/s41235-018-0117-4](https://doi.org/10.1186/s41235-018-0117-4) (2018).

98. Mühl, C., Sheil, O., Jarutyte, L. & Bestelmeyer, P. E. G. The Bangor Voice Matching Test: A standardized test for the assessment of voice perception ability. *Behav. Res. Methods* **50**, 2184–2192, DOI: [10.3758/s13428-017-0985-4](https://doi.org/10.3758/s13428-017-0985-4) (2018).

99. Aglieri, V. *et al.* The Glasgow Voice Memory Test: Assessing the ability to memorize and recognize unfamiliar voices. *Behav. Res. Methods* **49**, 97–110, DOI: [10.3758/s13428-015-0689-6](https://doi.org/10.3758/s13428-015-0689-6) (2017).

100. Lavan, N., Burston, L. F. K. & Garrido, L. How many voices did you hear? Natural variability disrupts identity perception from unfamiliar voices. *Br. J. Psychol.* **110**, 576–593, DOI: [10.1111/bjop.12348](https://doi.org/10.1111/bjop.12348) (2019).

101. Bestelmeyer, P. E. & Mühl, C. Individual differences in voice adaptability are specifically linked to voice perception skill. *Cognition* **210**, 104582, DOI: [10.1016/j.cognition.2021.104582](https://doi.org/10.1016/j.cognition.2021.104582) (2021).

102. Johnson, J., McGettigan, C. & Lavan, N. Comparing unfamiliar voice and face identity perception using identity sorting tasks. *Q. J. Exp. Psychol.* **73**, 1537–1545, DOI: [10.1177/1747021820938659](https://doi.org/10.1177/1747021820938659) (2020).

103. Jenkins, R. E. *et al.* Are super-face-recognisers also super-voice-recognisers? Evidence from cross-modal identification tasks. *Appl. Cogn. Psychol.* **35**, 590–605, DOI: [10.1002/acp.3813](https://doi.org/10.1002/acp.3813) (2021).

104. Tsantani, M. & Cook, R. Normal recognition of famous voices in developmental prosopagnosia. *Sci. Reports* **10**, 19757, DOI: [10.1038/s41598-020-76819-3](https://doi.org/10.1038/s41598-020-76819-3) (2020).

105. Fraccaro, P. J. *et al.* Correlated Male Preferences for Femininity in Female Faces and Voices. *Evol. Psychol.* **8**, 147470491000800311, DOI: [10.1177/147470491000800311](https://doi.org/10.1177/147470491000800311) (2010).

106. Yovel, G. & Belin, P. A unified coding strategy for processing faces and voices. *Trends Cogn. Sci.* **17**, 263–271, DOI: [10.1016/j.tics.2013.04.004](https://doi.org/10.1016/j.tics.2013.04.004) (2013).

107. Spence, C. The scent of attraction and the smell of success: crossmodal influences on person perception. *Cogn. Res. Princ. Implic.* **6**, 46, DOI: [10.1186/s41235-021-00311-3](https://doi.org/10.1186/s41235-021-00311-3) (2021).

108. Secundo, L. *et al.* Individual olfactory perception reveals meaningful nonolfactory genetic information. *Proc. Natl. Acad. Sci.* **112**, 8750–8755, DOI: [10.1073/pnas.1424826112](https://doi.org/10.1073/pnas.1424826112) (2015).

109. Schirmer, A. & Adolphs, R. Emotion Perception from Face, Voice, and Touch: Comparisons and Convergence. *Trends Cogn. Sci.* **21**, 216–228, DOI: [10.1016/j.tics.2017.01.001](https://doi.org/10.1016/j.tics.2017.01.001) (2017).

110. Dawel, A., O’Kearney, R., McKone, E. & Palermo, R. Not just fear and sadness: Meta-analytic

evidence of pervasive emotion recognition deficits for facial and vocal expressions in psychopathy. *Neurosci. & Biobehav. Rev.* **36**, 2288–2304, DOI: [10.1016/j.neubiorev.2012.08.006](https://doi.org/10.1016/j.neubiorev.2012.08.006) (2012).

111. Dadds, M. R., Kimonis, E. R., Schollar-Root, O., Moul, C. & Hawes, D. J. Are impairments in emotion recognition a core feature of callous–unemotional traits? testing the primary versus secondary variants model in children. *Dev. psychopathology* **30**, 67–77 (2018).

112. Bird, G. & Viding, E. The self to other model of empathy: Providing a new framework for understanding empathy impairments in psychopathy, autism, and alexithymia. *Neurosci. & Biobehav. Rev.* **47**, 520–532, DOI: [10.1016/j.neubiorev.2014.09.021](https://doi.org/10.1016/j.neubiorev.2014.09.021) (2014).

113. Rhodes, G. *et al.* How distinct is the coding of face identity and expression? Evidence for some common dimensions in face space. *Cognition* **142**, 123–137, DOI: [10.1016/j.cognition.2015.05.012](https://doi.org/10.1016/j.cognition.2015.05.012) (2015).

114. Hildebrandt, A., Schacht, A., Sommer, W. & Wilhelm, O. Measuring the speed of recognising facially expressed emotions. *Cogn. & Emot.* **26**, 650–666, DOI: [10.1080/02699931.2011.602046](https://doi.org/10.1080/02699931.2011.602046) (2012).

115. Herzmann, G., Kunina, O., Sommer, W. & Wilhelm, O. Individual Differences in Face Cognition: Brain–Behavior Relationships. *J. Cogn. Neurosci.* **22**, 571–589, DOI: [10.1162/jocn.2009.21249](https://doi.org/10.1162/jocn.2009.21249) (2010).

116. Esins, J., Schultz, J., Stemper, C., Kennerknecht, I. & Bülthoff, I. Face Perception and Test Reliabilities in Congenital Prosopagnosia in Seven Tests. *i-Perception* **7**, 2041669515625797, DOI: [10.1177/2041669515625797](https://doi.org/10.1177/2041669515625797) (2016).

117. Thomas, A. L., Lawler, K., Olson, I. R. & Aguirre, G. K. The Philadelphia Face Perception Battery. *Arch. Clin. Neuropsychol.* **23**, 175–187, DOI: [10.1016/j.acn.2007.10.003](https://doi.org/10.1016/j.acn.2007.10.003) (2008).

118. Tanaka, J. W. & Farah, M. J. Parts and wholes in face recognition. *The Q. J. Exp. Psychol. Sect. A* **46**, 225–245, DOI: [10.1080/14640749308401045](https://doi.org/10.1080/14640749308401045) (1993).

119. Young, A. W., Hellawell, D. & Hay, D. C. Configurational Information in Face Perception. *Perception* **16**, 747–759, DOI: [10.1068/p160747](https://doi.org/10.1068/p160747) (1987).

120. Yovel, G., Wilmer, J. B. & Duchaine, B. What can individual differences reveal about face processing? *Front. Hum. Neurosci.* **8**, 562, DOI: [10.3389/fnhum.2014.00562](https://doi.org/10.3389/fnhum.2014.00562) (2014).

121. DeGutis, J., Mercado, R. J., Wilmer, J. & Rosenblatt, A. Individual Differences in Holistic Processing Predict the Own-Race Advantage in Recognition Memory. *PLoS ONE* **8**, e58253, DOI: [10.1371/journal.pone.0058253](https://doi.org/10.1371/journal.pone.0058253) (2013).

122. Wang, R., Li, J., Fang, H., Tian, M. & Liu, J. Individual Differences in Holistic Processing Predict

Face Recognition Ability. *Psychol. Sci.* **23**, 169–177, DOI: [10.1177/0956797611420575](https://doi.org/10.1177/0956797611420575) (2011).

123. Konar, Y., Bennett, P. J. & Sekuler, A. B. Holistic Processing Is Not Correlated With Face-
Identification Accuracy. *Psychol. Sci.* **21**, 38–43, DOI: [10.1177/0956797609356508](https://doi.org/10.1177/0956797609356508) (2009).

124. Rezlescu, C., Susilo, T., Wilmer, J. B. & Caramazza, A. The Inversion, Part-Whole, and
Composite Effects Reflect Distinct Perceptual Mechanisms With Varied Relationships to Face
Recognition. *J. Exp. Psychol. Hum. Percept. Perform.* **43**, 1961–1973, DOI: [10.1037/xhp0000400](https://doi.org/10.1037/xhp0000400)
(2017).

125. Palermo, R. *et al.* Impaired holistic coding of facial expression and facial identity in congenital
prosopagnosia. *Neuropsychologia* **49**, 1226–1235, DOI: [10.1016/j.neuropsychologia.2011.02.021](https://doi.org/10.1016/j.neuropsychologia.2011.02.021)
(2011).

126. Avidan, G., Tanzer, M. & Behrmann, M. Impaired holistic processing in congenital prosopagnosia.
Neuropsychologia **49**, 2541–2552, DOI: [10.1016/j.neuropsychologia.2011.05.002](https://doi.org/10.1016/j.neuropsychologia.2011.05.002) (2011).

127. Liu, T. T. & Behrmann, M. Impaired holistic processing of left-right composite faces in congenital
prosopagnosia. *Front. Hum. Neurosci.* **8**, 750, DOI: [10.3389/fnhum.2014.00750](https://doi.org/10.3389/fnhum.2014.00750) (2014).

128. Biotti, F. & Cook, R. Impaired perception of facial emotion in developmental prosopagnosia.
Cortex **81**, 126–136, DOI: [10.1016/j.cortex.2016.04.008](https://doi.org/10.1016/j.cortex.2016.04.008) (2016).

129. Susilo, T. *et al.* Face recognition impairments despite normal holistic processing and face space
coding: Evidence from a case of developmental prosopagnosia. *Cogn. Neuropsychol.* **27**, 636–664,
DOI: [10.1080/02643294.2011.613372](https://doi.org/10.1080/02643294.2011.613372) (2010).

130. Ulrich, P. I. N. *et al.* Perceptual and Memorial Contributions to Developmental Prosopagnosia. *Q. J. Exp. Psychol.* **70**, 298–315, DOI: [10.1080/17470218.2016.1177101](https://doi.org/10.1080/17470218.2016.1177101) (2017).

131. Sunday, M. A., Richler, J. J. & Gauthier, I. Limited evidence of individual differences in holistic
processing in different versions of the part-whole paradigm. *Attention, Perception, & Psychophys.*
79, 1453–1465, DOI: [10.3758/s13414-017-1311-z](https://doi.org/10.3758/s13414-017-1311-z) (2017).

132. Royer, J., Blais, C., Gosselin, F., Duncan, J. & Fiset, D. When Less Is More: Impact of Face
Processing Ability on Recognition of Visually Degraded Faces. *J. Exp. Psychol. Hum. Percept. Perform.* **41**, 1179–1183, DOI: [10.1037/xhp0000095](https://doi.org/10.1037/xhp0000095) (2015).

133. Dunn, J. D. *et al.* Face information sampling in super-recognizers. *PsyArXiv* DOI:
[10.31234/osf.io/z2k4a](https://doi.org/10.31234/osf.io/z2k4a) (2021).

134. Itz, M. L., Schweinberger, S. R. & Kaufmann, J. M. Familiar Face Priming: The Role of Second-
Order Configuration and Individual Face Recognition Abilities. *Perception* **47**, 185–196, DOI:

924 [10.1177/0301006617742069](https://doi.org/10.1177/0301006617742069) (2018).

925 135. Kaufmann, J. M., Schulz, C. & Schweinberger, S. R. High and low performers differ in the use of
926 shape information for face recognition. *Neuropsychologia* **51**, 1310–1319, DOI:
927 [10.1016/j.neuropsychologia.2013.03.015](https://doi.org/10.1016/j.neuropsychologia.2013.03.015) (2013).

928 136. Itz, M. L., Golle, J., Luttman, S., Schweinberger, S. R. & Kaufmann, J. M. Dominance of texture
929 over shape in facial identity processing is modulated by individual abilities. *Br. J. Psychol.* **108**,
930 369–396, DOI: [10.1111/bjop.12199](https://doi.org/10.1111/bjop.12199) (2017).

931 137. Avidan, G. & Behrmann, M. Spatial Integration in Normal Face Processing and Its Breakdown in
932 Congenital Prosopagnosia. *Annu. Rev. Vis. Sci.* **7**, 1–21, DOI: [10.1146/annurev-vision-113020-
933 012740](https://doi.org/10.1146/annurev-vision-113020-012740) (2021).

934 138. Rossion, B. The composite face illusion: A whole window into our understanding of holistic face
935 perception. *Vis. Cogn.* **21**, 139–253, DOI: [10.1080/13506285.2013.772929](https://doi.org/10.1080/13506285.2013.772929) (2013).

936 139. Richler, J. J. & Gauthier, I. When intuition fails to align with data: A reply to Rossion (2013). *Vis.*
937 *Cogn.* **21**, 254–276, DOI: [10.1080/13506285.2013.796035](https://doi.org/10.1080/13506285.2013.796035) (2013).

938 140. Richler, J. J., Floyd, R. J. & Gauthier, I. About-face on face recognition ability and holistic
939 processing. *J. Vis.* **15**, 15–15, DOI: [10.1167/15.9.15](https://doi.org/10.1167/15.9.15) (2015).

940 141. Ross, D. A., Richler, J. J. & Gauthier, I. Reliability of composite-task measurements of holistic
941 face processing. *Behav. Res. Methods* **47**, 736–743, DOI: [10.3758/s13428-014-0497-4](https://doi.org/10.3758/s13428-014-0497-4) (2015).

942 142. Burton, A. M., Schweinberger, S. R., Jenkins, R. & Kaufmann, J. M. Arguments Against a
943 Configural Processing Account of Familiar Face Recognition. *Perspectives on Psychol. Sci.* **10**,
944 482–496, DOI: [10.1177/1745691615583129](https://doi.org/10.1177/1745691615583129) (2015).

945 143. Clutterbuck, R. & Johnston, R. A. Exploring Levels of Face Familiarity by Using an Indirect Face-
946 Matching Measure. *Perception* **31**, 985–994, DOI: [10.1068/p3335](https://doi.org/10.1068/p3335) (2002).

947 144. Burton, A. M., Wilson, S., Cowan, M. & Bruce, V. Face Recognition in Poor-Quality Video:
948 Evidence From Security Surveillance. *Psychol. Sci.* **10**, 243–248, DOI: [10.1111/1467-9280.00144](https://doi.org/10.1111/1467-9280.00144)
949 (1998).

950 145. Bruce, V., Henderson, Z., Newman, C. & Burton, A. M. Matching Identities of Familiar and
951 Unfamiliar Faces Caught on CCTV Images. *J. Exp. Psychol. Appl.* **7**, 207–218, DOI:
952 [10.1037/1076-898x.7.3.207](https://doi.org/10.1037/1076-898x.7.3.207) (2001).

953 146. Ambrus, G. G., Eick, C. M., Kaiser, D. & Kovács, G. Getting to Know You: Emerging Neural

Representations during Face Familiarization. *The J. Neurosci.* **41**, 5687–5698, DOI: [10.1523/jneurosci.2466-20.2021](https://doi.org/10.1523/jneurosci.2466-20.2021) (2021).

147. Dalski, A., Kovács, G. & Ambrus, G. G. Evidence for a general neural signature of face familiarity. *bioRxiv*, DOI: [10.1101/2021.04.18.440317](https://doi.org/10.1101/2021.04.18.440317) (2021).

148. Ramon, M. & Gobbini, M. I. Familiarity matters: A review on prioritized processing of personally familiar faces. *Vis. Cogn.* **26**, 1–17, DOI: [10.1080/13506285.2017.1405134](https://doi.org/10.1080/13506285.2017.1405134) (2018).

149. Wiese, H. *et al.* Later but not early stages of familiar face recognition depend strongly on attentional resources: Evidence from event-related brain potentials. *Cortex* **120**, 147–158, DOI: [10.1016/j.cortex.2019.06.004](https://doi.org/10.1016/j.cortex.2019.06.004) (2019).

150. Kovács, G. Getting to Know Someone: Familiarity, Person Recognition, and Identification in the Human Brain. *J. Cogn. Neurosci.* **32**, 2205–2225, DOI: [10.1162/jocn_a_01627](https://doi.org/10.1162/jocn_a_01627) (2020).

151. Hildebrandt, A., Olderbak, S. & Wilhelm, O. Facial Emotion Expression, Individual Differences in. *Int. Encycl. Soc. & Behav. Sci. 2nd edition* 667–675, DOI: [10.1016/b978-0-08-097086-8.25008-3](https://doi.org/10.1016/b978-0-08-097086-8.25008-3) (2015).

152. Redfern, A. S. & Benton, C. P. Representation of facial identity includes expression variability. *Vis. Res.* **157**, 123–131, DOI: [10.1016/j.visres.2018.05.004](https://doi.org/10.1016/j.visres.2018.05.004) (2019).

153. Burton, A. M., Kramer, R. S. S., Ritchie, K. L. & Jenkins, R. Identity From Variation: Representations of Faces Derived From Multiple Instances. *Cogn. Sci.* **40**, 202–223, DOI: [10.1111/cogs.12231](https://doi.org/10.1111/cogs.12231) (2016).

154. Kramer, R. S., Young, A. W. & Burton, A. M. Understanding face familiarity. *Cognition* **172**, 46–58, DOI: [10.1016/j.cognition.2017.12.005](https://doi.org/10.1016/j.cognition.2017.12.005) (2018).

155. Ritchie, K. L. & Burton, A. M. Learning faces from variability. *The Q. J. Exp. Psychol.* **70**, 1–9, DOI: [10.1080/17470218.2015.1136656](https://doi.org/10.1080/17470218.2015.1136656) (2017).

156. Murphy, J., Ipser, A., Gaigg, S. B. & Cook, R. Exemplar Variance Supports Robust Learning of Facial Identity. *J. Exp. Psychol. Hum. Percept. Perform.* **41**, 577–581, DOI: [10.1037/xhp0000049](https://doi.org/10.1037/xhp0000049) (2015).

157. White, D., Burton, A. L. & Kemp, R. I. Not looking yourself: The cost of self-selecting photographs for identity verification. *Br. J. Psychol.* **107**, 359–373, DOI: [10.1111/bjop.12141](https://doi.org/10.1111/bjop.12141) (2016).

158. Ritchie, K. L., Kramer, R. S. & Burton, A. M. What makes a face photo a ‘good likeness’? *Cognition* **170**, 1–8, DOI: [10.1016/j.cognition.2017.09.001](https://doi.org/10.1016/j.cognition.2017.09.001) (2018).

- 985 159. Yardley, L., McDermott, L., Pisarski, S., Duchaine, B. & Nakayama, K. Psychosocial
986 consequences of developmental prosopagnosia: A problem of recognition. *J. Psychosom. Res.* **65**,
987 445–451, DOI: [10.1016/j.jpsychores.2008.03.013](https://doi.org/10.1016/j.jpsychores.2008.03.013) (2008).
- 988 160. Dalrymple, K. A. *et al.* “A room full of strangers every day”: The psychosocial impact of
989 developmental prosopagnosia on children and their families. *J. Psychosom. Res.* **77**, 144–150,
990 DOI: [10.1016/j.jpsychores.2014.06.001](https://doi.org/10.1016/j.jpsychores.2014.06.001) (2014).
- 991 161. Murray, E., Hills, P. J., Bennetts, R. J. & Bate, S. Identifying Hallmark Symptoms of
992 Developmental Prosopagnosia for Non-Experts. *Sci. Reports* **8**, 1690, DOI: [10.1038/s41598-018-](https://doi.org/10.1038/s41598-018-20089-7)
993 [20089-7](https://doi.org/10.1038/s41598-018-20089-7) (2018).
- 994 162. Adams, A., Hills, P. J., Bennetts, R. J. & Bate, S. Coping strategies for developmental
995 prosopagnosia. *Neuropsychol. Rehabil.* **30**, 1996–2015, DOI: [10.1080/09602011.2019.1623824](https://doi.org/10.1080/09602011.2019.1623824)
996 (2019).
- 997 163. Bindemann, M. *Forensic Face Matching* (Oxford University Press, USA, 2021).
- 998 164. White, D., Towler, A. & Kemp, I., R. Understanding professional expertise in unfamiliar face
999 matching. In *Forensic Face Matching*, 62–88 (Oxford University Press, 2020).
- 1000 165. Wirth, B. E. & Carbon, C.-C. An Easy Game for Frauds? Effects of Professional Experience and
1001 Time Pressure on Passport-Matching Performance. *J. Exp. Psychol. Appl.* **23**, 138–157, DOI:
1002 [10.1037/xap0000114](https://doi.org/10.1037/xap0000114) (2017).
- 1003 166. Heyer, R., Semmler, C. & Hendrickson, A. T. Humans and Algorithms for Facial Recognition:
1004 The Effects of Candidate List Length and Experience on Performance. *J. Appl. Res. Mem. Cogn.* **7**,
1005 597–609, DOI: [10.1016/j.jarmac.2018.06.002](https://doi.org/10.1016/j.jarmac.2018.06.002) (2018).
- 1006 167. White, D., Kemp, R. I., Jenkins, R., Matheson, M. & Burton, A. M. Passport Officers’ Errors in
1007 Face Matching. *PLoS ONE* **9**, e103510, DOI: [10.1371/journal.pone.0103510](https://doi.org/10.1371/journal.pone.0103510) (2014).
- 1008 168. White, D., Dunn, J. D., Schmid, A. C. & Kemp, R. I. Error Rates in Users of Automatic Face
1009 Recognition Software. *PLoS ONE* **10**, e0139827, DOI: [10.1371/journal.pone.0139827](https://doi.org/10.1371/journal.pone.0139827) (2015).
- 1010 169. Towler, A. *et al.* Do professional facial image comparison training courses work? *PLOS ONE* **14**,
1011 e0211037, DOI: [10.1371/journal.pone.0211037](https://doi.org/10.1371/journal.pone.0211037) (2019).
- 1012 170. Weatherford, D. R., Roberson, D. & Erickson, W. B. When experience does not promote
1013 expertise: security professionals fail to detect low prevalence fake IDs. *Cogn. Res. Princ. Implic.*
1014 **6**, 25, DOI: [10.1186/s41235-021-00288-z](https://doi.org/10.1186/s41235-021-00288-z) (2021).

- 1015 171. Papesh, M. H. Photo ID verification remains challenging despite years of practice. *Cogn. Res.*
1016 *Princ. Implic.* **3**, 19, DOI: [10.1186/s41235-018-0110-y](https://doi.org/10.1186/s41235-018-0110-y) (2018).
- 1017 172. Robertson, D. J., Noyes, E., Dowsett, A. J., Jenkins, R. & Burton, A. M. Face Recognition by
1018 Metropolitan Police Super-Recognisers. *PLOS ONE* **11**, e0150036, DOI:
1019 [10.1371/journal.pone.0150036](https://doi.org/10.1371/journal.pone.0150036) (2016).
- 1020 173. Davis, J. P., Lander, K., Evans, R. & Jansari, A. Investigating Predictors of Superior Face
1021 Recognition Ability in Police Super-recognisers. *Appl. Cogn. Psychol.* **30**, 827–840, DOI:
1022 [10.1002/acp.3260](https://doi.org/10.1002/acp.3260) (2016).
- 1023 174. White, D., Phillips, P. J., Hahn, C. A., Hill, M. & O'Toole, A. J. Perceptual expertise in forensic facial
1024 image comparison. *Proc. Royal Soc. B: Biol. Sci.* **282**, 20151292, DOI: [10.1098/rspb.2015.1292](https://doi.org/10.1098/rspb.2015.1292) (2015).
- 1025 175. Towler, A., White, D. & Kemp, R. I. Evaluating the Feature Comparison Strategy for Forensic
1026 Face Identification. *J. Exp. Psychol. Appl.* **23**, 47–58, DOI: [10.1037/xap0000108](https://doi.org/10.1037/xap0000108) (2017).
- 1027 176. Moreton, R., Havard, C., Strathie, A. & Pike, G. An international survey of applied face-matching
1028 training courses. *Forensic Sci. Int.* 110947, DOI: <https://doi.org/10.1016/j.forsciint.2021.110947> (2021).
- 1029 177. Steblay, N., Dysart, J., Fulero, S. & Lindsay, R. C. L. Eyewitness Accuracy Rates in Sequential
1030 and Simultaneous Lineup Presentations: A Meta-Analytic Comparison. *Law Hum. Behav.* **25**, 459–
1031 473, DOI: [10.1023/a:1012888715007](https://doi.org/10.1023/a:1012888715007) (2001).
- 1032 178. Geiselman, R. E. *et al.* Benton facial recognition test scores: Index of eyewitness accuracy. *Am. J.*
1033 *Forensic Psychol.* **19**, 77–88 (2001).
- 1034 179. Bindemann, M., Brown, C., Koyas, T. & Russ, A. Individual differences in face identification
1035 postdict eyewitness accuracy. *J. Appl. Res. Mem. Cogn.* **1**, 96–103, DOI:
1036 [10.1016/j.jarmac.2012.02.001](https://doi.org/10.1016/j.jarmac.2012.02.001) (2012).
- 1037 180. Grabman, J. H., Dobolyi, D. G., Berelovich, N. L. & Dodson, C. S. Predicting High Confidence
1038 Errors in Eyewitness Memory: The Role of Face Recognition Ability, Decision-Time, and
1039 Justifications. *J. Appl. Res. Mem. Cogn.* **8**, 233–243, DOI: [10.1016/j.jarmac.2019.02.002](https://doi.org/10.1016/j.jarmac.2019.02.002) (2019).
- 1040 181. Grabman, J. H. & Dodson, C. S. Stark Individual Differences: Face Recognition Ability Influences
1041 the Relationship Between Confidence and Accuracy in a Recognition Test of Game of Thrones
1042 Actors. *J. Appl. Res. Mem. Cogn.* **9**, 254–269, DOI: [10.1016/j.jarmac.2020.02.007](https://doi.org/10.1016/j.jarmac.2020.02.007) (2020).
- 1043 182. Hodgetts, H. M., Vachon, F., Chamberland, C. & Tremblay, S. See No Evil: Cognitive Challenges
1044 of Security Surveillance and Monitoring. *J. Appl. Res. Mem. Cogn.* **6**, 230–243, DOI:
1045 [10.1016/j.jarmac.2017.05.001](https://doi.org/10.1016/j.jarmac.2017.05.001) (2017).

- 1046 **183.** Davis, J. P., Forrest, C., Treml, F. & Jansari, A. Identification from CCTV: Assessing police
1047 super-recogniser ability to spot faces in a crowd and susceptibility to change blindness. *Appl.*
1048 *Cogn. Psychol.* **32**, 337–353, DOI: [10.1002/acp.3405](https://doi.org/10.1002/acp.3405) (2018).
- 1049 **184.** Garvie, C., Bedoya, A. & Frankle, J. The perpetual line-up. unregulated police face recognition in
1050 america. *Georgetown Law Center on Privacy & Technology* (2019).
- 1051 **185.** Blauch, N. M., Behrmann, M. & Plaut, D. C. Computational insights into human perceptual
1052 expertise for familiar and unfamiliar face recognition. *Cognition* **208**, 104341, DOI:
1053 [10.1016/j.cognition.2020.104341](https://doi.org/10.1016/j.cognition.2020.104341) (2021).
- 1054 **186.** Grossman, S. *et al.* Convergent evolution of face spaces across human face-selective neuronal
1055 groups and deep convolutional networks. *Nat. Commun.* **10**, 4934, DOI: [10.1038/s41467-019-](https://doi.org/10.1038/s41467-019-12623-6)
1056 [12623-6](https://doi.org/10.1038/s41467-019-12623-6) (2019).
- 1057 **187.** Natu, V. & O’Toole, A. J. The neural processing of familiar and unfamiliar faces: A review and
1058 synopsis. *Br. J. Psychol.* **102**, 726–747, DOI: [10.1111/j.2044-8295.2011.02053.x](https://doi.org/10.1111/j.2044-8295.2011.02053.x) (2011).
- 1059 **188.** Benton, A. & Allen, M. V. Impairment in Facial Recognition in Patients with Cerebral Disease.
1060 *Cortex* **4**, 344–IN1, DOI: [10.1016/s0010-9452\(68\)80018-8](https://doi.org/10.1016/s0010-9452(68)80018-8) (1968).
- 1061 **189.** Rossion, B. & Michel, C. Normative accuracy and response time data for the computerized Benton
1062 Facial Recognition Test (BFRT-c). *Behav. Res. Methods* **50**, 2442–2460, DOI: [10.3758/s13428-](https://doi.org/10.3758/s13428-018-1023-x)
1063 [018-1023-x](https://doi.org/10.3758/s13428-018-1023-x) (2018).
- 1064 **190.** Fysh, M. C. & Bindemann, M. The Kent Face Matching Test. *Br. J. Psychol.* **109**, 219–231, DOI:
1065 [10.1111/bjop.12260](https://doi.org/10.1111/bjop.12260) (2018).
- 1066 **191.** Dowsett, A. J. & Burton, A. M. Unfamiliar face matching: Pairs out-perform individuals and provide a
1067 route to training. *Br. J. Psychol.* **106**, 433–445, DOI: [10.1111/bjop.12103](https://doi.org/10.1111/bjop.12103) (2015).
- 1068 **192.** Stacchi, L., Huguenin-Elie, E., Caldara, R. & Ramon, M. Normative data for two challenging tests
1069 of face matching under ecological conditions. *Cogn. Res. Princ. Implic.* **5**, 8, DOI:
1070 [10.1186/s41235-019-0205-0](https://doi.org/10.1186/s41235-019-0205-0) (2020).
- 1071 **193.** Duchaine, B., Germine, L. & Nakayama, K. Family resemblance: Ten family members with
1072 prosopagnosia and within-class object agnosia. *Cogn. Neuropsychol.* **24**, 419–430, DOI:
1073 [10.1080/02643290701380491](https://doi.org/10.1080/02643290701380491) (2007).
- 1074 **194.** Belanova, E., Davis, J. P. & Thompson, T. Cognitive and neural markers of super-recognisers’
1075 face processing superiority and enhanced cross-age effect. *Cortex* **108**, 92–111, DOI:
1076 [10.1016/j.cortex.2018.07.008](https://doi.org/10.1016/j.cortex.2018.07.008) (2018).

195. Herzmann, G., Danthiir, V., Schacht, A., Sommer, W. & Wilhelm, O. Toward a comprehensive test battery for face cognition: Assessment of the tasks. *Behav. Res. Methods* **40**, 840–857, DOI: [10.3758/brm.40.3.840](https://doi.org/10.3758/brm.40.3.840) (2008).
196. Kennerknecht, I. *et al.* First report of prevalence of non-syndromic hereditary prosopagnosia (HPA). *Am. J. Med. Genet. Part A* **140A**, 1617–1622, DOI: [10.1002/ajmg.a.31343](https://doi.org/10.1002/ajmg.a.31343) (2006).
197. Shah, P., Gaule, A., Sowden, S., Bird, G. & Cook, R. The 20-item prosopagnosia index (PI20): a self-report instrument for identifying developmental prosopagnosia. *Royal Soc. Open Sci.* **2**, 140343, DOI: [10.1098/rsos.140343](https://doi.org/10.1098/rsos.140343) (2015).
198. Young, A., Perrett, D., Calder, A., Sprengelmeyer, R. & Ekman, P. Facial expressions of emotion: Stimuli and tests (feest), version for pc. *Thames Val. Test Co. Thurston, UK* (2002).
199. Young, A. W. *et al.* Facial expression megamix: Tests of dimensional and category accounts of emotion recognition. *Cognition* **63**, 271–313, DOI: [10.1016/s0010-0277\(97\)00003-6](https://doi.org/10.1016/s0010-0277(97)00003-6) (1997).
200. Cecilione, J. L. *et al.* Test–Retest Reliability of the Facial Expression Labeling Task. *Psychol. Assess.* **29**, 1537–1542, DOI: [10.1037/pas0000439](https://doi.org/10.1037/pas0000439) (2017).
201. Calvo, M. G. & Lundqvist, D. Facial expressions of emotion (KDEF): Identification under different display-duration conditions. *Behav. Res. Methods* **40**, 109–115, DOI: [10.3758/brm.40.1.109](https://doi.org/10.3758/brm.40.1.109) (2008).
202. Baron-Cohen, S., Wheelwright, S., Hill, J., Raste, Y. & Plumb, I. The “Reading the Mind in the Eyes” Test Revised Version: A Study with Normal Adults, and Adults with Asperger Syndrome or High-functioning Autism. *J. Child Psychol. Psychiatry* **42**, 241–251, DOI: [10.1111/1469-7610.00715](https://doi.org/10.1111/1469-7610.00715) (2001).
203. Fernández-Abascal, E. G., Cabello, R., Fernández-Berrocal, P. & Baron-Cohen, S. Test-retest reliability of the ‘Reading the Mind in the Eyes’ test: a one-year follow-up study. *Mol. Autism* **4**, 33, DOI: [10.1186/2040-2392-4-33](https://doi.org/10.1186/2040-2392-4-33) (2013).
204. Barton, J. J. & Corrow, S. L. The problem of being bad at faces. *Neuropsychologia* **89**, 119–124, DOI: [10.1016/j.neuropsychologia.2016.06.008](https://doi.org/10.1016/j.neuropsychologia.2016.06.008) (2016).
205. Noyes, E., Phillips, P. & O’Toole, A. What is a super-recogniser? In *Face processing: Systems, disorders and cultural differences*, 173–201 (Nova Science Publishers Inc, 2017).
206. Grand, R. L. *et al.* What aspects of face processing are impaired in developmental prosopagnosia? *Brain Cogn.* **61**, 139–158, DOI: [10.1016/j.bandc.2005.11.005](https://doi.org/10.1016/j.bandc.2005.11.005) (2006).
207. Schmalzl, L., Palermo, R. & Coltheart, M. Cognitive heterogeneity in genetically based

prosopagnosia: A family study. *J. Neuropsychol.* **2**, 99–117, DOI: [10.1348/174866407x256554](https://doi.org/10.1348/174866407x256554) (2008).

208. Plomin, R., Haworth, C. M. A. & Davis, O. S. P. Common disorders are quantitative traits. *Nat. Rev. Genet.* **10**, 872–878, DOI: [10.1038/nrg2670](https://doi.org/10.1038/nrg2670) (2009).

209. Germine, L. T., Duchaine, B. & Nakayama, K. Where cognitive development and aging meet: Face learning ability peaks after age 30. *Cognition* **118**, 201–210, DOI: [10.1016/j.cognition.2010.11.002](https://doi.org/10.1016/j.cognition.2010.11.002) (2011).

210. Lane, J. *et al.* Impacts of impaired face perception on social interactions and quality of life in age-related macular degeneration: A qualitative study and new community resources. *PLoS ONE* **13**, e0209218, DOI: [10.1371/journal.pone.0209218](https://doi.org/10.1371/journal.pone.0209218) (2018).

211. Werheid, K. & Clare, L. Are Faces Special in Alzheimer’s Disease? Cognitive Conceptualisation, Neural Correlates, and Diagnostic Relevance of Impaired Memory for Faces and Names. *Cortex* **43**, 898–906, DOI: [10.1016/s0010-9452\(08\)70689-0](https://doi.org/10.1016/s0010-9452(08)70689-0) (2007).

212. Kumfor, F. *et al.* Do I know you? Examining face and object memory in frontotemporal dementia. *Neuropsychologia* **71**, 101–111, DOI: [10.1016/j.neuropsychologia.2015.03.020](https://doi.org/10.1016/j.neuropsychologia.2015.03.020) (2015).

213. Hutchings, R., Palermo, R., Piguet, O. & Kumfor, F. Disrupted Face Processing in Frontotemporal Dementia: A Review of the Clinical and Neuroanatomical Evidence. *Neuropsychol. Rev.* **27**, 18–30, DOI: [10.1007/s11065-016-9340-2](https://doi.org/10.1007/s11065-016-9340-2) (2017).

214. Stantic’, M., Ichijo, E., Catmur, C. & Bird, G. Face memory and face perception in autism. *Autism* 136236132110276, DOI: [10.1177/13623613211027685](https://doi.org/10.1177/13623613211027685) (2021).

215. Bortolon, C., Capdevielle, D. & Raffard, S. Face recognition in schizophrenia disorder: A comprehensive review of behavioral, neuroimaging and neurophysiological studies. *Neurosci. & Biobehav. Rev.* **53**, 79–107, DOI: [10.1016/j.neubiorev.2015.03.006](https://doi.org/10.1016/j.neubiorev.2015.03.006) (2015).

216. Gray, H. M. & Tickle-Degnen, L. A Meta-Analysis of Performance on Emotion Recognition Tasks in Parkinson’s Disease. *Neuropsychology* **24**, 176–191, DOI: [10.1037/a0018104](https://doi.org/10.1037/a0018104) (2010).

217. Edens, J. F., Marcus, D. K., Lilienfeld, S. O. & Poythress, N. G. Psychopathic, Not Psychopath: Taxometric Evidence for the Dimensional Structure of Psychopathy. *J. Abnorm. Psychol.* **115**, 131–144, DOI: [10.1037/0021-843x.115.1.131](https://doi.org/10.1037/0021-843x.115.1.131) (2006).

218. Skeem, J. L., Polaschek, D. L. L., Patrick, C. J. & Lilienfeld, S. O. Psychopathic Personality. *Psychol. Sci. Public Interest* **12**, 95–162, DOI: [10.1177/1529100611426706](https://doi.org/10.1177/1529100611426706) (2011).

219. Fox, E. & Zougkou, K. Influence of Personality Traits on Processing of Facial Expressions. DOI:

1139 [10.1093/oxfordhb/9780199559053.013.0026](https://doi.org/10.1093/oxfordhb/9780199559053.013.0026) (2011).

1140 220. Megías-Robles, A. *et al.* The ‘Reading the mind in the Eyes’ test and emotional intelligence. *Royal*
1141 *Soc. Open Sci.* **7**, 201305, DOI: [10.1098/rsos.201305](https://doi.org/10.1098/rsos.201305) (2020).

1142 221. Meinhardt-Injac, B., Daum, M. M., Meinhardt, G. & Persike, M. The Two-Systems Account of
1143 Theory of Mind: Testing the Links to Social- Perceptual and Cognitive Abilities. *Front. Hum.*
1144 *Neurosci.* **12**, 25, DOI: [10.3389/fnhum.2018.00025](https://doi.org/10.3389/fnhum.2018.00025) (2018).

1145 222. Saxe, R., Brett, M. & Kanwisher, N. Divide and conquer: A defense of functional localizers.
1146 *NeuroImage* **30**, 1088–1096, DOI: [10.1016/j.neuroimage.2005.12.062](https://doi.org/10.1016/j.neuroimage.2005.12.062) (2006).

1147 223. Kanwisher, N. The Quest for the FFA and Where It Led. *J. Neurosci.* **37**, 1056–1061, DOI:
1148 [10.1523/jneurosci.1706-16.2016](https://doi.org/10.1523/jneurosci.1706-16.2016) (2017).

1149 224. Zhen, Z. *et al.* Quantifying interindividual variability and asymmetry of face-selective regions: A
1150 probabilistic functional atlas. *NeuroImage* **113**, 13–25, DOI: [10.1016/j.neuroimage.2015.03.010](https://doi.org/10.1016/j.neuroimage.2015.03.010)
1151 (2015).

1152 225. Berman, M. G. *et al.* Evaluating functional localizers: The case of the FFA. *NeuroImage* **50**, 56–
1153 71, DOI: [10.1016/j.neuroimage.2009.12.024](https://doi.org/10.1016/j.neuroimage.2009.12.024) (2010).

1154 226. Engell, A. D. & McCarthy, G. Probabilistic atlases for face and biological motion perception: An
1155 analysis of their reliability and overlap. *NeuroImage* **74**, 140–151, DOI:
1156 [10.1016/j.neuroimage.2013.02.025](https://doi.org/10.1016/j.neuroimage.2013.02.025) (2013).

1157 227. Golarai, G. *et al.* Differential development of high-level visual cortex correlates with category-
1158 specific recognition memory. *Nat. Neurosci.* **10**, 512–522, DOI: [10.1038/nn1865](https://doi.org/10.1038/nn1865) (2007).

1159 228. Jiang, X. *et al.* A quantitative link between face discrimination deficits and neuronal selectivity for faces
1160 in autism. *NeuroImage: Clin.* **2**, 320–331, DOI: [10.1016/j.nicl.2013.02.002](https://doi.org/10.1016/j.nicl.2013.02.002) (2013).

1161 229. Furl, N., Garrido, L., Dolan, R. J., Driver, J. & Duchaine, B. Fusiform Gyrus Face Selectivity
1162 Relates to Individual Differences in Facial Recognition Ability. *J. Cogn. Neurosci.* **23**, 1723–1740,
1163 DOI: [10.1162/jocn.2010.21545](https://doi.org/10.1162/jocn.2010.21545) (2011).

1164 230. McGugin, R. W. & Gauthier, I. The reliability of individual differences in face-selective responses
1165 in the fusiform gyrus and their relation to face recognition ability. *Brain Imaging Behav.* **10**, 707–
1166 718, DOI: [10.1007/s11682-015-9467-4](https://doi.org/10.1007/s11682-015-9467-4) (2016).

1167 231. Duchaine, B. C. & Nakayama, K. Developmental prosopagnosia: a window to content-specific face
1168 processing. *Curr. Opin. Neurobiol.* **16**, 166–173, DOI: [10.1016/j.conb.2006.03.003](https://doi.org/10.1016/j.conb.2006.03.003) (2006).

1169 232. Jiahui, G., Yang, H. & Duchaine, B. Developmental prosopagnosics have widespread selectivity

reductions across category-selective visual cortex. *Proc. Natl. Acad. Sci.* **115**, 201802246, DOI: [10.1073/pnas.1802246115](https://doi.org/10.1073/pnas.1802246115) (2018).

233. Avidan, G., Hasson, U., Malach, R. & Behrmann, M. Detailed Exploration of Face-related Processing in Congenital Prosopagnosia: 2. Functional Neuroimaging Findings. *J. Cogn. Neurosci.* **17**, 1150–1167, DOI: [10.1162/0898929054475145](https://doi.org/10.1162/0898929054475145) (2005).
234. Dubois, J. & Adolphs, R. Building a Science of Individual Differences from fMRI. *Trends Cogn. Sci.* **20**, 425–443, DOI: [10.1016/j.tics.2016.03.014](https://doi.org/10.1016/j.tics.2016.03.014) (2016).
235. Kaltwasser, L., Hildebrandt, A., Recio, G., Wilhelm, O. & Sommer, W. Neurocognitive mechanisms of individual differences in face cognition: A replication and extension. *Cogn. Affect. & Behav. Neurosci.* **14**, 861–878, DOI: [10.3758/s13415-013-0234-y](https://doi.org/10.3758/s13415-013-0234-y) (2014).
236. Xu, B., Liu-Shuang, J., Rossion, B. & Tanaka, J. Individual Differences in Face Identity Processing with Fast Periodic Visual Stimulation. *J. Cogn. Neurosci.* **29**, 1368–1377, DOI: [10.1162/jocn_a_01126](https://doi.org/10.1162/jocn_a_01126) (2017).
237. Stacchi, L., Liu-Shuang, J., Ramon, M. & Caldara, R. Reliability of individual differences in neural face identity discrimination. *NeuroImage* **189**, 468–475, DOI: [10.1016/j.neuroimage.2019.01.023](https://doi.org/10.1016/j.neuroimage.2019.01.023) (2019).
238. Towler, J., Fisher, K. & Eimer, M. The Cognitive and Neural Basis of Developmental Prosopagnosia. *Q. J. Exp. Psychol.* **70**, 316–344, DOI: [10.1080/17470218.2016.1165263](https://doi.org/10.1080/17470218.2016.1165263) (2016).
239. Avidan, G. *et al.* Selective Dissociation Between Core and Extended Regions of the Face Processing Network in Congenital Prosopagnosia. *Cereb. Cortex* **24**, 1565–1578, DOI: [10.1093/cercor/bht007](https://doi.org/10.1093/cercor/bht007) (2014).
240. Lohse, M. *et al.* Effective Connectivity from Early Visual Cortex to Posterior Occipitotemporal Face Areas Supports Face Selectivity and Predicts Developmental Prosopagnosia. *The J. Neurosci.* **36**, 3821–3828, DOI: [10.1523/jneurosci.3621-15.2016](https://doi.org/10.1523/jneurosci.3621-15.2016) (2016).
241. Rosenthal, G. *et al.* Altered topology of neural circuits in congenital prosopagnosia. *eLife* **6**, e25069, DOI: [10.7554/elife.25069](https://doi.org/10.7554/elife.25069) (2017).
242. Thomas, C. *et al.* Reduced structural connectivity in ventral visual cortex in congenital prosopagnosia. *Nat. Neurosci.* **12**, 29–31, DOI: [10.1038/nn.2224](https://doi.org/10.1038/nn.2224) (2009).
243. Gomez, J. *et al.* Functionally Defined White Matter Reveals Segregated Pathways in Human Ventral Temporal Cortex Associated with Category-Specific Processing. *Neuron* **85**, 216–227, DOI: [10.1016/j.neuron.2014.12.027](https://doi.org/10.1016/j.neuron.2014.12.027) (2015).

- 1201 **244.** Song, S. *et al.* Local but not long-range microstructural differences of the ventral temporal cortex
1202 in developmental prosopagnosia. *Neuropsychologia* **78**, 195–206, DOI:
1203 [10.1016/j.neuropsychologia.2015.10.010](https://doi.org/10.1016/j.neuropsychologia.2015.10.010) (2015).
- 1204 **245.** Wan, L. *et al.* Face-Blind for Other-Race Faces: Individual Differences in Other-Race Recognition
1205 Impairments. *J. Exp. Psychol. Gen.* **146**, 102–122, DOI: [10.1037/xge0000249](https://doi.org/10.1037/xge0000249) (2017).
- 1206 **246.** McKone, E. *et al.* A critical period for faces: Other-race face recognition is improved by childhood
1207 but not adult social contact. *Sci. Reports* **9**, 12820, DOI: [10.1038/s41598-019-49202-0](https://doi.org/10.1038/s41598-019-49202-0) (2019).
- 1208 **247.** DeGutis, J., Cohan, S. & Nakayama, K. Holistic face training enhances face processing in
1209 developmental prosopagnosia. *Brain* **137**, 1781–1798, DOI: [10.1093/brain/awu062](https://doi.org/10.1093/brain/awu062) (2014).
- 1210 **248.** Ellis, H. D. & Young, A. W. Training in face-processing skills for a child with acquired prosopagnosia.
1211 *Dev. Neuropsychol.* **4**, 283–294, DOI: [10.1080/87565648809540412](https://doi.org/10.1080/87565648809540412) (1988).
- 1212 **249.** DeGutis, J., Cohan, S., Kahn, D. A., Aguirre, G. K. & Nakayama, K. Facial expression training
1213 improves emotion recognition and changes neural tuning in a patient with acquired emotion
1214 recognition deficits and prosopagnosia. *J. Vis.* **13**, 993–993, DOI: [10.1167/13.9.993](https://doi.org/10.1167/13.9.993) (2013).
- 1215 **250.** Brunsdon, R., Coltheart, M., Nickels, L. & Joy, P. Developmental prosopagnosia: A case analysis
1216 and treatment study. *Cogn. Neuropsychol.* **23**, 822–840, DOI: [10.1080/026432905000441841](https://doi.org/10.1080/026432905000441841)
1217 (2006).
- 1218 **251.** Bate, S., Adams, A. & Bennetts, R. J. Guess Who? Facial Identity Discrimination Training
1219 Improves Face Memory in Typically Developing Children. *J. Exp. Psychol. Gen.* **149**, 901–913,
1220 DOI: [10.1037/xge0000689](https://doi.org/10.1037/xge0000689) (2020).
- 1221 **252.** Dolzycka, D., Herzmann, G., Sommer, W. & Wilhelm, O. Can Training Enhance Face Cognition
1222 Abilities in Middle-Aged Adults? *PLoS ONE* **9**, e90249, DOI: [10.1371/journal.pone.0090249](https://doi.org/10.1371/journal.pone.0090249)
1223 (2014).
- 1224 **253.** Dowsett, A. J., Sandford, A. & Burton, A. M. Face learning with multiple images leads to fast
1225 acquisition of familiarity for specific individuals. *Q. J. Exp. Psychol.* **69**, 1–10, DOI:
1226 [10.1080/17470218.2015.1017513](https://doi.org/10.1080/17470218.2015.1017513) (2016).
- 1227 **254.** Jeckeln, G., Hahn, C. A., Noyes, E., Cavazos, J. G. & O’Toole, A. J. Wisdom of the social versus
1228 non-social crowd in face identification. *Br. J. Psychol.* **109**, 724–735, DOI: [10.1111/bjop.12291](https://doi.org/10.1111/bjop.12291)
1229 (2018).
- 1230 **255.** White, D., Kemp, R. I., Jenkins, R. & Burton, A. M. Feedback training for facial image
1231 comparison. *Psychon. Bull. & Rev.* **21**, 100–106, DOI: [10.3758/s13423-013-0475-3](https://doi.org/10.3758/s13423-013-0475-3) (2014).

- 1232 **256.** Towler, A., Keshwa, M., Ton, B., Kemp, R. I. & White, D. Diagnostic Feature Training Improves Face
1233 Matching Accuracy. *J. Exp. Psychol. Learn. Mem. Cogn.* DOI: [10.1037/xlm0000972](https://doi.org/10.1037/xlm0000972) (2021).
- 1234 **257.** Young, A. W. & Burton, A. M. Are We Face Experts? *Trends Cogn. Sci.* **22**, 100–110, DOI:
1235 [10.1016/j.tics.2017.11.007](https://doi.org/10.1016/j.tics.2017.11.007) (2018).
- 1236 **258.** Andrews, S., Jenkins, R., Cursiter, H. & Burton, A. M. Telling faces together: Learning new faces
1237 through exposure to multiple instances. *Q. J. Exp. Psychol.* **68**, 2041–2050, DOI:
1238 [10.1080/17470218.2014.1003949](https://doi.org/10.1080/17470218.2014.1003949) (2015).
- 1239 **259.** Matthews, C. M. & Mondloch, C. J. Improving Identity Matching of Newly Encountered Faces:
1240 Effects of Multi-image Training. *J. Appl. Res. Mem. Cogn.* **7**, 280–290, DOI:
1241 [10.1016/j.jarmac.2017.10.005](https://doi.org/10.1016/j.jarmac.2017.10.005) (2018).
- 1242 **260.** Dunn, J. D., Kemp, R. I. & White, D. Search templates that incorporate within-face variation
1243 improve visual search for faces. *Cogn. Res. Princ. Implic.* **3**, 37, DOI: [10.1186/s41235-018-0128-1](https://doi.org/10.1186/s41235-018-0128-1)
1244 (2018).
- 1245
- 1246

Table 1. Tasks used to measure individual differences in face processing abilities. Only tasks that were specifically developed to test individual differences in face processing are included, where normative accuracy data is available based on non-clinical adult samples of more than 80 participants. Test-retest reliability is presented, and ranges indicate variable reliability in sub-measures reported. This table signals a maturing field of individual differences in face processing with progress in testing a range of face processing abilities, albeit most heavily concentrated in identity processing.

| Target ability | Task type | Task | Test-retest reliability |
|----------------|---------------------------|--|--------------------------|
| Identity | Perceptual matching | BFRT, Benton Face Recognition Test ^{188, 189} | - |
| | | Glasgow Face Matching Test ⁵⁶ | .77 ⁵⁴ |
| | | Glasgow Face Matching Test 2 ⁵⁸ | .79 ⁵⁸ |
| | | Kent Face Matching Test ¹⁹⁰ | .67 ⁵² |
| | | Models Matching Test ^{191, 52} | - |
| | | Oxford Face Matching Test ⁵⁴ | .75 ⁵⁴ |
| | | Yearbook Test ¹⁹² | - |
| | | 1-in-10 Matching Test ⁵² | - |
| | Perceptual discrimination | Cambridge Face Perception Test ^{53, 193} | - |
| | | Facial Identity Card Sorting Test ^{52, 192} | - |
| | Recognition memory | Adult/ Infant Face recognition Test ¹⁹⁴ | - |
| | | Cambridge Face Memory Test ⁵⁵ | .70 ²² |
| | | Cambridge Face Memory Test Extended ^{17, 53} | - |
| | | UNSW Face Test ⁵⁰ | .59 ⁵⁰ |
| | Naming | Bielefelder famous faces test (BFFT) ¹⁹⁵ | - |
| | | Before They Were Famous Test ^{17, 20} | - |
| | | Familiar Faces Memory Test ^{42, 72} | - |
| | Self report | Cambridge Face Memory Questionnaire ⁷² | - |
| | | Hong Kong Prosopagnosia Questionnaire ^{196, 67} | - |
| | | Prosopagnosia Index ¹⁹⁷ | .89 ⁵⁴ |
| | | Stirling Face Recognition Scale ⁶⁸ | - |
| Expressions | Perceptual matching | Emotion Matching Task ¹¹ | - |
| | Naming | Ekman 60 Faces ¹⁹⁸ | - |
| | | Emotion Hexagon Test ¹⁹⁹ | - |
| | | Facial Expression Labelling Test ²⁰⁰ | .39 - .85 ²⁰⁰ |
| | | Karolinska Directed Emotional Faces ²⁰¹ | - |
| | | Reading the Mind in the Eyes Test ²⁰² | .63 ²⁰³ |
| Impressions | Rating | Facial Impression Tests (Trustworthiness) ¹² | .73 ¹² |
| | | Facial Impression Tests (Dominance) ¹² | .58 ¹² |
| | | Facial Impression Tests (Attractiveness) ¹² | .50 ¹² |
| | | Individual Preference Test (Attractiveness) ⁸⁹ | .75 ⁸⁹ |
| | | Philadelphia Face Perception Battery (Attractiveness) ¹¹⁷ | .50 ¹¹⁷ |
| Demographics | Perceptual matching | Philadelphia Face Perception Battery (Age) ¹¹⁷ | .49 ¹¹⁷ |
| | Naming | Philadelphia Face Perception Battery (Gender) ¹¹⁷ | .37 ¹¹⁷ |

Figure 1. Taxonomy of tasks used to measure face processing abilities. (A) Perceptual matching involves deciding whether two or more images match on a given dimension (here: identity). (B) Perceptual discrimination requires comparing two or more images on a given dimension and either choosing most/least, or ranking from high to low (here: expression). (C) Recognition memory requires studying faces and some time later memory for the faces is tested. (D) Naming tasks require naming a person or labelling an expression, sometimes from a set of predefined labels. (E) Rating tasks ask participants to rate a single image on a dimension using a Likert scale. (F) Self-report measures ask participants about their face processing experiences in everyday life. Correlation of tests measuring the same ability across different task formats establishes convergent validity, but differences in task format can also interfere with measurement of association between different abilities.

Figure 2. Everyday decisions depend on rapid decoding of multidimensional facial cues. Everyday decisions are made in rich and dynamic environments where multiple cues from multiple senses are integrated and linked with complex social contexts. Coloured boxes list some of the ambient visual cues that might influence perceptual judgments on given dimensions in real world tasks. For example, a decision about where to sit on a bus might be contingent on both identifying your colleague and on whether her mood would be conducive to casual conversation (is she upset?). Indeed those cues might not be independent, if for example you have only encountered your colleague in a happy mood then her expression might influence the identity judgment itself. Situational contexts such as the bus route, and the clothes worn by the men who might be arguing, are also likely to influence judgments.

1297 **Box 1: Defining developmental prosopagnosia**

1298 Acquired prosopagnosia is characterised by impairment of face processing resulting from brain
1299 damage, but developmental prosopagnosia is not linked to known structural or genetic pathology.
1300 Nevertheless, poor face processing abilities can have severe negative impacts on social interactions^{159–}
1301 ¹⁶². The problem of diagnosis is therefore critical. It is unclear whether developmental prosopagnosia is
1302 better conceptualised as the low-end of the range of typical ability or as a condition in its own right,
1303 independent of typical variability^{204,205}. Understanding the dimensional structure of individual
1304 differences in face processing can help better define developmental prosopagnosia and its association
1305 with other conditions.

1306 No genetic markers have yet been identified for developmental prosopagnosia, and reported neural
1307 abnormalities vary between studies (Box 2). In the absence of reliable markers, the definition of
1308 developmental prosopagnosia is purely based on behavioural performance on tests of face identity
1309 processing or questionnaires probing everyday face recognition. Accurate diagnosis is therefore
1310 conditional on the psychometric properties of these measures.

1311 That some people with developmental prosopagnosia show impaired holistic processing but others
1312 do not might reflect ‘cognitive heterogeneity’ of the condition^{116,206}, which could signal a family of
1313 related subtypes of prosopagnosia rather than a unitary condition²⁰⁷. This proposal would be consistent
1314 with a genetic basis for the condition, despite the current absence of markers: many inherited disorders
1315 are end-points of quantitative dimensional traits determined by multiple genes exerting small effects,
1316 resulting in heterogeneity across a group of individuals^{196,204,208}.

1317 Defining developmental prosopagnosia as a condition is further complicated by age-related declines
1318 in ability^{50,209}, and the need to exclude the contributions of associated conditions. Some of these
1319 conditions do have a clear organic basis (for example, macular degeneration²¹⁰, Alzheimer’s
1320 pathologies²¹¹, frontotemporal dementia^{212,213}) and produce associated progressive deficits in face
1321 perception and memory abilities. The basis of other conditions is less well understood, for example
1322 Autism Spectrum Disorder (ASD)^{32,33,37}, and Schizophrenia²¹⁵. The complexity of these disorders
1323 involve social and perceptual deficits that are not specific to faces²⁵ and manifests as heterogeneity in
1324 the patterns of face processing impairment.

1325 When symptomatic of broader conditions, patterns of impairment reflect the multidimensionality of
1326 face processing abilities. Some disorders are associated with both impaired emotion and identity
1327 processing (Autism³⁴, Schizophrenia²¹⁵, Anxiety⁸²). Other conditions selectively impair expression
1328 recognition (Parkinsons²¹⁶, Psychopathy¹¹⁰). Individual difference studies can improve understanding of
1329 the links between emotion processing deficits⁸² and face abilities in the typical population. Aside from

1330 Parkinson's, these conditions involve traits that vary dimensionally in the typical population^{81,217,218} and
1331 so associated face processing impairments have implications for non-pathological variation^{3,25,219–}
1332 ²²¹.

Box 2: Neural bases of face recognition

Examining anatomical brain differences and their relation to different ability levels can help improve understanding of functional aspects of face processing abilities. Most studies on this topic have focused on differences in blood flow within face-selective regions, measured using functional Magnetic Resonance Imaging (fMRI). Some regions have been functionally defined as ‘face-selective’, and differential activation to faces and non-face objects can then be measured to capture face-selective responses at the individual level²²². Individual differences are found in the precise locations of these regions^{223,224}, and they are mostly stable over time within individuals^{225,226}.

The Fusiform Face Area (FFA) is a functionally defined area that selectively responds to images of faces across repeated brain scans (figure panel a, dark blue). Some studies show correlations between scores on face identification performance and FFA activation strength^{29,30} and region size^{30,227}. However a number of studies report no association^{228–230}. This inconsistency might be due in part to small sample sizes, which are not well suited to individual difference analysis. Some comparisons of FFA activation in people with developmental prosopagnosia to controls show reduced activity^{229,231,232}, but others show no difference²³³. Inconsistency might also arise from poor reliability of brain responses²³⁴. Test-retest reliability of FFA activation has not been examined rigorously, although one study does show relatively high stability in this measure over different presentations of faces in the same experimental session²³⁰.

An association has also been found between FFA grey matter volume and performance in face recognition^{26–28}. A small number of studies using electrophysiological recordings from the scalp (ERPs), have also reported correlations between face-specific components and face recognition performance^{115,235,236}. Despite high reliability of some ERP measures over repeated testing²³⁷, in each of these studies correlations between multiple face-selective ERP components were low ($r = .3$), and the degree to which the components were face-selective did not reliably distinguish developmental prosopagnosia from typical recognition abilities²³⁸.

The FFA is just one part of the neuronal network that has been identified as responding selectively to faces (see figure). But outside the FFA, the association between individual differences in face recognition and brain response in specific regions are relatively inconsistent across studies (light blue and gray in figure: Occipital Face Area, OFA; Anterior Temporal Lobe, ATL; Amygdala, AMG; Superior Temporal Sulcus, STS)^{29,30,239}. The degree of network connectivity, both within this core set of regions and beyond, correlates with measures of face recognition³⁰ and reduced communication between areas has been implicated in developmental prosopagnosia^{239–241}. The importance of interconnection is also supported by structural investigations of white matter connections between

1391 cortical areas (figure panel b, dark blue), with structural deficits of these fibre tracts reported in
1392 developmental prosopagnosia^{242–244}.
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418

Box 3: Training face recognition

The extent to which face recognition abilities can be improved with training has implications for understanding individual differences and plasticity. Face recognition ability does not develop fully until after the age of 30^{50,209} and people's history of perceptual exposure to faces influences their abilities^{245,246}. This flexibility in the face processing system could support training, and hence benefit people with developmental prosopagnosia and those using face recognition professionally.

[H1] Training impaired face abilities

Attempts to train face recognition abilities of adults with developmental prosopagnosia have been largely unsuccessful. One approach has been to train a holistic processing strategy when learning previously unfamiliar faces, but accuracy benefits from these methods are rarely found. Where they are reported, the benefits generalise poorly to faces not included in the training, and do not transfer to superficially different faces, for example photos taken with different cameras or lighting²⁴⁷. This poor generalisation limits the clinical benefit of training and is consistent with earlier failed attempts to improve face recognition performance in patients with acquired prosopagnosia^{248,249}.

Another approach is to encourage use of individual face features for identification of familiar faces. Many people with developmental prosopagnosia report using distinguishing facial features to identify familiar faces¹⁶². In a case study, researchers were able to teach children with developmental prosopagnosia to recognise familiar faces by memorising three distinctive features of each person's face²⁵⁰ and anecdotal evidence suggests that these improvements carried into daily life. Other studies have also produced promising results training children with developmental prosopagnosia²⁵¹, suggesting that treatment in early development could confer some benefit. However, training does not transfer well to more naturalistic task conditions, a finding that is consistent with attempts to train face recognition in the broader population²⁵².

[H1] Training typical face abilities

Training in applied settings tends to be tailored to the specific task of matching unfamiliar faces¹⁷⁶. A large-scale evaluation of professional training courses showed no learning beyond the specific faces used in each course¹⁶⁹. In laboratory studies, collaborative face matching decisions with another person^{253,254}, and accuracy feedback on decisions²⁵⁵ produce small benefits to accuracy. Improvements were specific to individuals with poorer recognition skills and were small in comparison to individual differences. A common element might be participants' realization that the task is more difficult than they expect it to be, leading them to more careful analysis. Some paradigms have successfully

1452 improved accuracy by directing participants' attention to diagnostic features^{[175,256](#)}, which would be
1453 consistent with the benefit of additional analysis.

1454 Given the very large benefits of familiarity for face recognition^{[62,257](#)}, another approach has been
1455 to develop familiar face representations. Substantial improvements are found when participants view
1456 multiple different photos of the same face^{[155,156,258–260](#)}, encouraging the formation of a coherent
1457 representation across variability. However, these benefits do not generalise to new faces^{[155,258](#)}, limiting
1458 their value in applied settings.

1459

1460